Title	Excretion of polycyclic aromatic hydrocarbon metabolites (OH-PAHs) in cattle urine in Ghana
Author(s)	Bortey-Sam, Nesta; Ikenaka, Yoshinori; Akoto, Osei; Nakayama, Shouta M. M.; Marfo, Jemima; Saengtienchai, Aksorn; Mizukawa, Hazuki; Ishizuka, Mayumi
Citation	Environmental Pollution, 218, 331-337 https://doi.org/10.1016/j.envpol.2016.07.008
Issue Date	2016-11
Doc URL	http://hdl.handle.net/2115/72006
Rights	©2016. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	Environ Pollut v.218 p.331-337.pdf

[Instructions for use](#)

Excretion of polycyclic aromatic hydrocarbon metabolites (OH-PAHs) in cattle urine in
Ghana

Nesta Bortey-Sam¹, Yoshinori Ikenaka^{1, 2}, Osei Akoto³, Shouta M.M. Nakayama¹, Jemima
Marfo¹, Aksorn Saengtienchai^{1, 4}, Hazuki Mizukawa⁵, Mayumi Ishizuka^{1*}

¹Laboratory of Toxicology, Department of Environmental Veterinary Science, Graduate
School of Veterinary Medicine, Hokkaido University, Kita 18, Nishi 9, Kita ku, Sapporo,
060–0818, Japan

²Water Research Group, Unit for Environmental Sciences and Management, North-West
University, Potchefstroom, South Africa

³Department of Chemistry, Kwame Nkrumah University of Science and Technology,
Kumasi, Ghana

⁴Department of Pharmacology, Faculty of Veterinary Medicine, Kasetsart University,
Bangkok, Thailand

⁵Department of Environmental Veterinary Science, Graduate School of Veterinary
Medicine, Hokkaido University, Kita 18, Nishi 9, Kita ku, Sapporo 060–0818, Japan

*Corresponding author: Mayumi Ishizuka, e-mail: ishizum@vetmed.hokudai.ac.jp
Laboratory of Toxicology, Department of Environmental Veterinary Sciences, Graduate
School of Veterinary Medicine, Hokkaido University, N18, W9, Kita-ku, Sapporo, 060–
0818, Japan. Tel: +81–11–706–6949; Fax: +81–11–706–5105

Abstract

Previous studies of polycyclic aromatic hydrocarbons (PAHs) in particulate matter, soils and livers of wild rats indicated that the city centre of Kumasi, Ghana has been severely polluted with high cancer potency. Cattle urine were therefore collected from Kumasi (urban) and Offinso (rural), Ghana: to determine concentrations of urinary PAH metabolites (OH-PAHs); and find their association with sex; and to estimate exposure of cattle to PAHs from the different sites. From the results, geometric mean concentrations (adjusted by specific gravity), GM_{SG} , showed that 2-OHNaphthalene (2-OHNap) was the most abundant OH-PAH in cattle urine from all study sites, and naphthalene-containing-mothballs might have contributed significantly to the levels. There was no significant difference between urinary OH-PAHs concentrations in cattle from urban and rural sites except for 2-OHPhe and 4-OHPhe, and similar to urban areas, rural sites could also be polluted with PAHs. GM_{SG} of 2-OHNap in cattle urine in Kokote (21.9 ± 6.51 ng/mL; a rural area), was significantly higher compared to the other sites followed by Oforikrom (4.15 ± 4.37 ng/mL; urban). The GM_{SG} concentration (ng/mL) of the sum of OH-PAHs decreased in the order, Kokote (44.7) > Oforikrom (7.87) > Saboa (6.98) > Santasi (6.68) > and Twumasen Estate (5.23). The high concentrations of urinary 2-OHNap, 2-3-OHFlu, 2-OHPhe, 3-OHPhe and 4-OHPhe in Kokote indicated high PAHs exposure to cattle in this area or different/specific source of PAHs exposure. GM_{SG} of 2-OHNap was significantly higher in male cattle compared to females while 1-9-OHPhe was significantly higher in females.

Capsule:

PAH metabolites were measured in cattle urine in urban and rural areas in Ghana; 2-OHNaphthalene (2-OHNap) was the most abundant PAH metabolites.

Keywords: OH-PAHs; Kumasi; Metabolites; Cattle; Urine

1. Introduction

Polycyclic aromatic hydrocarbons (PAHs) are pollutants formed during incomplete combustion of organic materials. They are found in vehicle exhaust, wood and cigarette smoke, and also in grilled foods. Human and animal exposure to PAHs occur mainly through inhalation of contaminated air or ingestion of soil, food and/or drinking contaminated water (Barranco et al., 2004; Dissanayake and Galloway, 2004). According to the Agency for Toxic Substances and Disease Registry's (ATSDR, 2013) priority list of hazardous chemicals, PAHs were classified as the 9th most hazardous chemical. In humans and animals, PAHs are metabolized by cytochrome P450 enzymes and excreted in urine. One of the major metabolites is monohydroxylated PAH (OH-PAHs) (Burczynski et al., 1998) and urinary levels of 1-hydroxypyrene has been used as biomarker of PAHs exposure (Bouchard and Viau, 1999; Jongeneelen, 2001). However, since ratios of different PAHs may vary depending on the source and personal enzymatic capacity, concentration profiles of multiple OH-PAHs biomarkers are necessary and required to assess the environmental exposure risk. Urinary metabolites of naphthalene, fluorene and phenanthrene are also commonly used as biomarkers to assess the exposure level and environmental risk (Li et al., 2008; Fan et al., 2012; 2012b).

The assessment of health risk to humans exposed to PAHs is primarily based on results from animal studies, which indicated that PAHs can produce carcinogenic and mutagenic effects. Recent studies also indicated that some PAH metabolites have strong correlation with atherosclerosis and cardiovascular diseases (Xu et al., 2010), and exposure of rats and mice to naphthalene caused nasal and bronchiolar tumors, respectively (NTP, 1992; 2000).

As a developing country, the economic and population growth rates in Ghana have seen tremendous increases over the past few years. The growing rate of industrialization is gradually leading to contamination and deterioration of the environment and pollution is likely to reach disturbing levels (Bortey-Sam et al., 2014). Studies by Bortey-Sam et al. (2013; 2014; 2015) in particulate matter (PM10) and soils indicated that the city centre of Kumasi, Ghana has been polluted with PAHs when compared with recommended levels, and fuel and wood/grass combustion were the dominant sources. The total benzo(a)pyrene equivalent concentration and estimated carcinogenicity of PAHs in PM10 and soil from the city centre was approximately 18 and 150 times higher, respectively, as compared to a pristine site. Rats were therefore used as sentinels to measure the environmental pollution state, and higher levels of PAHs were detected in the livers of wild rats in the city centre of Kumasi. Naphthalene was detected in 80% of those samples, and levels of phenanthrene and pyrene (the first and second most abundant, respectively) were significantly higher than other PAHs measured (Bortey-Sam et al., 2015b).

Based on the high levels and cancer potency of PAHs in PM10, soils and the levels found in livers of wild rats (Bortey-Sam et al. 2013; 2014; 2015; 2015b), cattle urine was collected because cattle is known to excrete large amount of PAH metabolites due to high

intake of the parent compound through feed or inhalation (Saengtienchai et al., 2014). PAHs in the atmosphere are known to settle in soil (Rey-Salgueiro et al. 2008) and this could increase the levels of exposure in Kumasi, Ghana, because these free-range cattle also pick food and/or water from the ground. Urinary levels of PAHs could be widely used as biological indicator of exposure (Jongeneelen, 2001), and there is limited/no data from literature that addresses the excretory levels of OH-PAHs in cattle in Ghana. The objectives of the present study were therefore: to determine the concentrations of OH-PAHs in cattle urine in Kumasi (urban) and Offinso (rural), Ghana; find the association between urinary OH-PAHs concentrations and sex; and to estimate cattle's exposure to PAHs from the different sites.

2. Materials and methods

2.1. Sampling

In August 2014, urine samples of healthy cattle (West African Shorthorn) were randomly collected from 5 communities in Kumasi and Offinso, both in the Ashanti Region of Ghana. Offinso is about 33 km from the city centre of Kumasi (Fig. 1). Samples were collected from Oforikrom and Santasi in Kumasi (urban), which are 5.1 and 3.5 km from the city centre, respectively (Fig. 1), where previous studies reported high levels of PAHs in PM₁₀, soils and livers of wild rats (Bortey-Sam et al., 2013; 2014; 2015; 2015b). On the other hand, the three sites in Offinso (Twumasen Estate, Saboa and Kokote) selected for cattle urine sampling (Fig. 1) are in rural and agricultural areas where bush burning is rampant and the use and sometimes abuse of pesticides such as carbaryl (1-naphthyl-N-methylcarbamate), which could be metabolized to 1-hydroxy naphthalene, was possible

(Meeker et al., 2007; Orjuela et al., 2012). In addition, due to the lack of background urine and interferences during OH-PAHs quantification, 500 mL of cattle urine (blank stock) was collected from Hokkaido University School farm. Hokkaido University is a public university located in Sapporo, Japan, and because of the low vehicular movement and industrial activities around the farm, PAHs exposure from point sources were assumed to be negligible. However, because cattle could be exposed to PAHs through feed and/or inhalation the sample collected was measured several times to confirm levels of OH-PAHs.

From the sample sites in Ghana, spot urine (n = 95; with 30 males and 65 females) were collected, transferred into labelled corning tubes (Corning Incorporated, New York, USA) and stored at -20°C in the Department of Chemistry, Kwame Nkrumah University Science and Technology (KNUST), Ghana. Of the 5 sites, only ages of some cattle in 2 sites (Twumasen Estate and Saboa) were obtained from the herdsman. The average ages (ranges) of cattle were 2.9 ± 1.0 years (1-4.5 years) in Twumasen Estate and 4.2 ± 2.9 years (1-12 years) in Saboa, respectively. Samples were later transported to the Laboratory of Toxicology, Graduate School of Veterinary Medicine, Hokkaido University, Japan where they were stored at -30°C until analysis (quarantine number for importing is 26 douken 383).

2.2. Sample extraction and analysis

20 μL each of β -glucuronidase (bovine liver, type B-1; 1240 U/mg; Sigma Aldrich) and arylsulfatase (limpets Type V; 34 units/mg; Sigma Aldrich) enzymes, and 5 mL of 0.1 M sodium acetate buffer (pH 5.6) were added to 5 mL urine sample after spiking with three PAH internal standards ($^{13}\text{C}_6$ -2-OHFluorene, 3-OHPhenanthrene- d_9 , and $^{13}\text{C}_6$ -

1OHPyrene). The pH of sample was adjusted to 5.5 using 1 M acetic acid (Wako Pure Chemicals, Osaka, Japan) and incubated overnight at 37 °C. The sample was diluted with 4 mL of Milli-Q water and extracted twice (liquid-liquid extraction) with 10 mL each of n-pentane (Kanto Chemical Corp., Tokyo, Japan) by shaking for 1 h. To reduce the interference of sulfur metabolites, the combined extracts were washed with 2 mL of 1 N AgNO₃ solution (Wako Pure Chemicals, Osaka, Japan), concentrated to 50-100 µL, redissolved to 0.5 mL using methanol and filtered (0.20 µm DISMIC-13JP membrane filter, ADVANTEC, Toyo Roshi Kaisha Ltd., Japan) prior to instrumental analysis. All sample preparation steps were performed in darkness (by covering tubes completely with aluminum foil) to avoid possible photodegradation of target analytes. A total of 13 OH-PAHs; 2-hydroxynaphthalene (2-OHNap), 2-hydroxyfluorene (2-OHFlu), 3-hydroxyfluorene (3-OHFlu), 9-hydroxyfluorene (9-OHFlu), 1-hydroxyphenanthrene (1-OHPhe), 2-hydroxyphenanthrene (2-OHPhe), 3-hydroxyphenanthrene (3-OHPhe), 4-hydroxyphenanthrene (4-OHPhe), 9-hydroxyphenanthrene (9-OHPhe), 1-hydroxypyrene (1-OHPyr), 6-hydroxychrysene (6-OHChry), 3-hydroxybenzo(e)pyrene (3-OHBeP) and 9-hydroxybenzo(a)pyrene (9-OHBaP), were analyzed in each sample. The standards (purity ≥ 98%) were purchased from Cambridge Isotope Laboratories Inc. (Andover, MA, USA) and Toronto Research Chemical Inc. (Brisbane Road, North York, Canada). Difficulties were often associated with the separation of 1-OHPhe and 9-OHPhe, for this reason, the sum of these isomers was used as an abbreviation, 1-9-OHPhe. Similarly, 2-3-OHFlu was used as sum of 2-OHFlu and 3-OHFlu. All results were adjusted by specific gravity and expressed in ng/mL.

A Shimadzu 8030 triple quadrupole mass spectrometer, upgraded to 8040 with UF lens, (ESI MS-MS; Shimadzu, Kyoto, Japan), equipped with a Prominence UFLC system (Shimadzu, Kyoto, Japan) was used for analysis. Chromatographic separation was achieved using an Agilent Eclipse PAH column (150 mm × 2.1 mm, 3.5 μm). The mobile phases were methanol:water (2:3, v:v) (A) and methanol (B), pumped at a flow rate of 250 μL/min. The mobile phase gradient was maintained as follows: 0.0–2.0 min, 5% B; 2.0–20 min, 40% B; 20–25 min, 40% B; 25–30 min, 95% B; 30–35 min, 95% B; 35–35.01 min, 5% B. Target compounds were determined by multiple-reaction monitoring (MRM) in the negative ionization mode.

2.3. Specific gravity (SG) of cattle urine

To compensate for variations in urine dilution, urinary OH-PAH concentrations were adjusted by specific gravity (SG). Urinary SG was measured by a hand refractometer (ATAGO, PAL-095, Tokyo, Japan). Obtained mean and ranges of SG in urine of cattle in Oforikrom (1.013; [1.004-1.029]), Santasi (1.035; [1.03-1.041]), Twumasen Estate (1.035; [1.028-1.04]), Saboa (1.036; [1.026-1.049]) and Kokote (1.037; [1.029-1.042]) were used to adjust urinary OH-PAHs concentrations as illustrated by [Nermell et al \(2008\)](#). The correction formula applied to each urine concentration was as follows:

$$\text{SG\_corrected concentration} = \text{urinary OH - PAH concentration} \times \frac{(\text{SG}_{\text{target}} - 1.0)}{(\text{SG}_{\text{sample}} - 1.0)}$$

Where, $\text{SG}_{\text{target}}$ is the mean specific gravity of cattle urine per community; $\text{SG}_{\text{sample}}$ is the specific gravity of a particular sample.

2.4. *Quality control and quality assurance*

A mixture of three ¹³C-isotopically labelled OH-PAHs (¹³C6-2-OHFlu, 3-OHPhe-d9, and ¹³C6-1-OHPyr) was spiked into urine samples as internal standard prior to sample preparation and extraction. ¹³C6-2-OHFlu was used for quantification of metabolite of naphthalene and three metabolites of fluorene. 3-OHPhe-d9 was used for quantification of the five metabolites of phenanthrene and ¹³C6-1-OHPyr was used for quantification of 1OHPyr, 6-OHChry, 3-OHBeP and 9-OHBaP.

Quantitation was performed using internal standard method (five-point calibration; 1, 5, 10, 50 and 100 ng/mL), and average correlation coefficients (r^2) for the calibration curves in cattle urine were greater or equal to 0.99. The standard solutions (spiked with internal standards) for the calibration curves were prepared in urine in order to normalize differences in interferences between standards and samples. Concentrations of OH-PAHs in urine sample used for this purpose were below the limits of detection (LOD) and differences between this and sample concentration was used in this study. Analytical methods were checked for precision and accuracy. Limits of quantification (LOQs) were calculated based on $10SD/S$ (SD is the standard deviation of the response of seven replicate standard solution measurements and S is the slope of the calibration curve). LOQs (ng/mL) of OH-PAHs were, 0.29 (2-OHNap), 0.24 (2-3-OHFlu), 0.60 (9-OHFlu), 0.23 (2-OHPhe), 0.71 (1-9-OHPhe), 1.16 (3-OHPhe), 0.15 (4-OHPhe), 0.87 (1-OHPyr), 0.73 (6-OHChry), 0.54 (3-OHBeP) and 0.32 (9-OHBeP), respectively. Internal standard recoveries (¹³C6-2OHFlu, 3-OHPhe-d9, and ¹³C6-1-OHPyr) ranged from 89 ± 5.8 – $96 \pm 11.4\%$ ([Table 1](#)).

For every batch of 10 samples, a solvent blank, a spiked solvent blank (internal standards spiked into solvent), a matrix spike (internal standards spiked into blank urine), and duplicate sample were analyzed. The average recoveries in spiked solvents blanks ranged from 92 ± 4.6 – $98 \pm 8.3\%$, and that for matrix spikes was 85 ± 9.1 – $96 \pm 7.1\%$. Blanks were run periodically and contained no detectable amount of target analyte. The coefficients of variation of OH-PAH in duplicate samples were less than 15%.

1.5. Data analysis

Data analysis was performed using IBM SPSS v 20 for windows (SPSS Inc., Illinois, USA). Kolmogorov–Smirnov (K–S) and Shapiro-Wilks tests were used to determine the normality of data and were considered statistically significant if p value was less than 0.05. Concentrations of OH-PAH below their respective LOQs were replaced with a value of LOQ/2. Geometric mean concentrations were used to represent the central tendency of OHPAH in this study (Wayne, 1999). ANOVA and Tukey analyses of log transformed data were used to compare concentrations of OH-PAH in cattle urine from the study areas and differences were considered statistically significant with p value < 0.05 . Student's T-Test was also used to compare distribution of OH-PAHs between male and female cattle; and, urban and rural sites.

3. Results and discussion

3.1. Excretory levels of OH-PAH in cattle urine

As shown in Table 2, there was no significant difference ($p > 0.05$) between urinary OH-PAHs concentrations in cattle from urban and rural sites except for 2-OHPhe and 4-OHPhe. Data for 3-OHPhe was not included because concentrations from urban sites and 2

rural sites were below the LOQ (Table 3). The significant differences ($p < 0.05$) could indicate differences in cattle's exposure to PAHs from urban and rural sites. Table 3 shows the distribution of OH-PAHs in cattle urine from the 5 sample sites. From the results, 2-OHNap, 2-3-OHFlu, 1-9-OHPhe, 2-OHPhe, 3-OHPhe, 4-OHPhe and 1-OHPyr were detected. 9-OHFlu and the high molecular weight, HMW (≥ 4 rings) PAHs (6-OHChry, 3-OHB(e)P and 9-OHB(a)P) were however not detected in the cattle urine and could be due to low detection sensitivity (Campo et al., 2008) and/or because the HMW PAHs, such as BaP, are mainly excreted through feces (Burgaz et al. 1992; Li et al., 2008).

Specific gravity adjusted geometric mean concentrations (GM_{SG}) showed that 2-OHNap (2.77 ± 5.91 ng/mL; $p < 0.01$) was the most abundant OH-PAH in cattle urine from all study sites followed by 1-9-OHPhe (2.02 ± 1.16 ng/mL) > 4-OHPhe (1.74 ± 1.87 ng/mL) > 1-OHPyr (1.22 ± 0.87 ng/mL) > 2-3-OHFlu (1.08 ± 1.75 ng/mL) > 2-OHPhe (0.489 ± 0.555 ng/mL) > and 3-OHPhe (0.278 ± 0.553 ng/mL) (Fig. 2). The GM_{SG} concentration (ng/mL) of the sum of OH-PAHs (2-OHNap, 2-3-OHFlu, 1-9OHPhe, 2-OHPhe, 3-OHPhe, 4-OHPhe and 1-OHPyr) decreased in the order, Kokote (44.7 ± 10.4) > Oforikrom (7.87 ± 7.41) > Saboa (6.98 ± 3.86) > Santasi (6.68 ± 2.701) > and Twumasen Estate (5.23 ± 1.55). High urinary concentrations of 2-OHNap, 2-3-OHFlu, 2-OHPhe, 3-OHPhe and 4-OHPhe were detected in Kokote (Table 3) indicating high exposure of cattle to the parent PAHs within the sample site. Kokote is a rural area filled with many farmlands, with high agricultural and burning activities compared to the other sites, and the levels of OH-PAHs could mean that there were different or specific sources of PAHs exposure to cattle in the area.

3.2. 2-OHNaphthalene

The GM_{SG} of 2-OHNap in cattle urine in Kokote (21.9 ± 6.51 ng/mL) was significantly higher ($p < 0.05$) compared to the other sites followed by Oforikrom (4.15 ± 4.37 ng/mL) (Table 3). However, the least GM_{SG} concentration for 2-OHNap was recorded in Santasi (0.61 ± 0.23 ng/mL). Although 1-OHNap could be derived from both naphthalene and carbaryl, 2-OHNap is derived only from naphthalene (Orjuela et al., 2012). The high levels of 2-OHNap could be due to exposure through ingestion and/or inhalation, although it has been proposed as a biomarker of inhalation (Kim et al. 2000). Naphthalene is ubiquitous in ambient air with high volumes in vehicular traffic, cigarette smoke (ATSDR, 2005) and is elevated when mothballs or stoves burning biomass fuels are used (Griego et al., 2008; Riojas-Rodriguez et al., 2011). Urinary levels of 2-OHNap are markers of vehicular traffic (Li et al., 2010) and mothball exposure (Owa et al., 1993). Naphthalene is most likely the primary ingredient of mothballs in Ghana (Soghoian et al., 2012), and is frequently used in driving away insects both in and outdoors. This practice could also contribute to 2-OHNap being the most abundant metabolite in cattle urine since exposure through ambient air was also possible because of its volatile nature.

3.3. 1-OHPyrene

The highest GM_{SG} concentration of 1-OHPyr were detected in cattle in Kokote (2.29 ± 1.28 ng/mL) and Oforikrom (1.37 ± 1.18 ng/mL). Levels in Kokote were significantly higher ($p < 0.05$) compared to other sites except Oforikrom and Santasi (Table 3). GM concentrations (not adjusted by SG) of 1-OHPyr in this study (1.02 ng/mL [Twumassen Estate] to 2.41 ng/mL [Kokote]) were generally higher compared to study by Saengtienchai

et al. (2014) using cattle from Japan and Thailand (non-adjusted). A study by Ferrari et al. (2002) on determination of 1-OHPyr in bovine urine from three farms located near rural and urban areas recorded non-adjusted average concentrations of 0.66 ng/mL (urban area); 1.52 ng/mL (rural) and 5.09 ng/mL (highway). Similar to the present study, Ferrari et al. (2002) indicated higher levels of 1-OHPyr in rural areas compared to urban and suggested that other important sources besides traffic could contribute to the PAHs burden of animals. Results of non-adjusted 1-OHPyr concentrations from the present study was higher compared to study by Ferrari et al. (2002) except for levels recorded in cattle raised on farms located in the vicinity of a highway (5.09 ng/mL).

The levels in the present study could be due to vehicular activities or traffic. At high temperature combustion (that is during vehicular emissions) the HMW PAH compounds are dominant (Laflamme and Hites, 1978). Previous studies by Bortey-Sam et al. (2014; 2015) in PM10 and soils indicated pyrene as the eighth and second most abundant PAH in Kumasi, respectively, and combustion of fuel (74%) and wood/grass (23%) were the dominant sources in the region. In Ghana, some farms are generally located close to major roads with high vehicular activities or traffic (Tay and Biney, 2013), and exposure to domestic and grazing animals could be through inhalation, or picking food or water from the ground.

3.4. OHPhenanthrenes and OHFluorenes

The distribution of 2-OHPhe, 4-OHPhe and 2-3-OHFlu were significantly higher ($p < 0.05$) in Kokote than the other sites (Table 3). Urinary levels of 1-9-OHPhe in Kokote was however significantly higher ($p < 0.05$) than levels found in Oforikrom. 3-OHFlu in cattle

urine from all sites were below the LOQ except Kokote (Table 3). In this study, the most dominant OHPhe isomer from all sites was 1-9-OHPhe, which is similar to results obtained by Fan et al. (2012) in humans. However, in Kokote, 4-OHPhe was most abundant (Table 3). In human urine, Thai et al. (2016) and Levine et al. (2015) found 1-OHPhe as most dominant while Guo et al. 2013 also reported 3-OHPhe as the most dominant of four phenanthrene metabolites. These variations could be due to differences in metabolic pathway among species. The possible source of phenanthrene and fluorene exposure to cattle in Kokote could be due to inhalation during combustion at low temperatures such as wood or grass combustion since the low molecular weight, LMW (< 4 rings) PAH compounds are abundant during low temperature combustion (Lake et al., 1979). Because Kokote is mainly agricultural area with many farmlands, resident farmers frequently practice bush burning. Another possible source could be due to ingestion of soil or water since the cattle graze freely and the soil from which they pick food or water may be bound to PAHs from burning activities. Previous study by Tay and Biney (2013) indicated that agricultural soils in Accra, Ghana were dominated by LMW PAHs through which domestic animals could be exposed. PAHs tend to adsorb tightly to organic matter in soil rendering them less susceptible to biological and chemical degradation (Hatzinger et al., 1997) and in general, LMW PAHs are more water soluble than HMW PAHs (Nam et al., 2008).

3.5. Association between urinary OH-PAHs concentrations and sex

Gender differences have been used in various studies to predict differences in OH-PAHs concentrations in human (Sul et al. 2012; Levine et al., 2015; Bartolomé et al., 2015; CDC, 2015). Study by Thai et al. (2016) in human urine showed no association between sex and

urinary OH-PAHs concentrations. In this study, 2-OHNap was significantly higher ($p < 0.05$) in male cattle ($GM_{SG} = 4.43 \pm 7.16$ ng/mL) compared to females ($GM_{SG} = 2.01 \pm 5.12$ ng/mL) (Table 4). Kim et al. (2013) suggested that rates of intake, accumulation, and excretion of chemicals differ in male and female cattle, although ADME (absorption, distribution, metabolism, and excretion) data would be needed to support that assertion. Differences could also be due to different rearing systems from the various sites.

Several non pharmacogenetic factors such as age, gender, species, disease factors or exposure to environmental pollutants might contribute to the expression and regulation of hepatic P450 in man, laboratory species and domestic animals (Guengerich, 2002; Nebbia, 2001). In a study by Dacasto et al. (2005), male piedmontese cattle showed significantly higher CYP3A-dependent drug metabolizing enzymes, erythromycin *N*-demethylase (ERDEM), ethylmorphine *N*-demethylation (ETDEM) and testosterone 6 β -hydroxylation (6 β -OHT), activities compared to females, with the exception of testosterone 2 β -hydroxylase, 2 β -OHT, (whose enzymatic activity was yet lower in females). On the other hand, no gender-difference was noticed in limousin cattle.

In human, Sul et al. (2012) observed significantly higher levels of urinary 2-OHNap in men than females, and suggested that gender were predictors of urinary 2-OHNap concentrations. 1-9-OHPhe on the other hand was significantly higher in female cattle ($GM_{SG} = 2.17 \pm 1.18$ ng/mL) than males ($GM_{SG} = 1.71 \pm 1.07$ ng/mL) (Table 4), while Σ OHPhes in women were low compared to men (Bartolomé et al., 2015). These differences could be due to variations in metabolism, levels and route of exposure to PAHs. The urinary levels of 1-OHNap, 2-OHNap, Σ OHNap, 1-OHPhe, 9-OHPhe, Σ OHPhe, 1-OHPyr,

and Σ OHPAHs among women were all significantly or marginally higher than those among men workers (Guo et al. 2014). In consistency, Guo et al. (2014) found that, when exposed to similar levels of PAHs, women had significantly higher micronuclei frequencies than men. Emerging evidence also indicates that women may be at greater risk of lung cancer than men, probably because the elevated activity of CYP1A1 enzymes in women can produce higher levels of DNA adducts, and women have lower DNA repair capacity than men (Mollerup et al., 2006; Uppstad et al., 2011).

4. Conclusions

Cattle urine samples were collected from both urban (Kumasi) and rural (Offinso) sites in the Ashanti Region, Ghana, and GM_{SG} concentration of OH-PAHs indicated that, 2-OHNap was the most abundant followed by; 1-9-OHPhe > 4-OHPhe > 1-OHPyr > 2-3OHFlu > 2-OHPhe > 3-OHPhe. The results of the present study showed that cattle in Kokote (rural area) were exposed to significantly higher levels of PAHs than the other sites, and naphthalene-containing-mothballs might have contributed significantly to 2-OHNap levels detected in cattle urine. There was no significant difference between urinary OH-PAHs concentrations in cattle in urban and rural sites except for 2-OHPhe and 4-OHPhe and similar to urban areas, rural sites could also be polluted with PAHs. Levels of 2-OHNap was significantly higher in male cattle compared to females, while the opposite was for 1-9-OHPhe.

344

345

346

347

348

349

350 **Acknowledgements**

351   This work was supported by Grants-in-Aid for Scientific Research from the Ministry of
352 Education, Culture, Sports, Science and Technology of Japan awarded to M. Ishizuka (No.
353 16H0177906), Y. Ikenaka (No. 26304043, 15H0282505, 15K1221305), S. Nakayama
354 (16K16197), and the foundations of Sumitomo and JSPS Core to Core Program (AA
355 Science Platforms) and Bilateral Joint Research Project (PG36150002 and PG36150003).
356 We also acknowledge the financial support by The Mitsui & Co., Ltd. Environment Fund,
357 the Soroptimist Japan Foundation and the Nakajima Foundation. We are grateful to Mr.
358 Takahiro Ichise (Laboratory of Toxicology, Graduate School of Veterinary Medicine,
359 Hokkaido University) for technical support.

360

361

362

363

References

- Agency for Toxic Substances, Disease Registry (ATSDR), 2005. Toxicological Profile for Naphthalene, 1-Methylnaphthalene, and 2-Methylnaphthalene.
- Agency for Toxic Substances, Disease Registry (ATSDR), 2013. Detailed data for 2013 Priority List of Hazardous Substances. Public Health Service, ATSDR, Division of Toxicology and Environmental Medicine.
- Barranco, A., Alonso-Salces, R.M., Crespo, I., Burreta, L.A., Gallo, B., Vicente, F., Sarobe, M., 2004. Polycyclic aromatic hydrocarbon content in commercial Spanish fatty foods. *J. Food Prot.* 67, 2786–2971.
- Bartolomé, M., Ramos, J.J., Cutanda, F., Huetos, O., Esteban, M., Ruiz-Moraga, M., Calvo, E., Pérez-Gómez, B., González, O., Castaño, A., 2015. Urinary polycyclic aromatic hydrocarbon metabolites levels in a representative sample of the Spanish adult population: the BIOAMBIENT.ES project. *Chemosphere* 135, 436–446.
- Bortey-Sam, N., Akoto, O., Ikenaka, Y., Nakayama, S.M., Ishizuka, M., 2013. Determination of benzo[a]pyrene levels in ambient air and the source of polycyclic aromatic hydrocarbons using a diagnostic ratio method in Ghana. *Jpn. J. Vet. Res.* 61, S72-74.
- Bortey-Sam, N., Ikenaka, Y., Nakayama, S.M., Akoto, O., Yohannes, Y.B., Baidoo, E., Mizukawa, H., Ishizuka, M., 2014. Occurrence, distribution, sources and toxic potential of polycyclic aromatic hydrocarbons (PAHs) in surface soils from the Kumasi Metropolis, Ghana. *Sci. Total Environ.* 496, 471-478.
- Bortey-Sam, N., Ikenaka, Y., Akoto, O., Nakayama, S.M., Yohannes, Y.B., Baidoo, E., Mizukawa, H., Ishizuka, M., 2015. Levels, potential sources and human health risk of polycyclic aromatic hydrocarbons (PAHs) in particulate matter (PM10) in Kumasi, Ghana. *Environ. Sci. Pollut. Res.* 22, 9658–9667.
- Bortey-Sam, N., Ikenaka, Y., Akoto, O., Nakayama, S.M., Yohannes, Y.B., Baidoo, E., Saengtienchai, A., Mizukawa, H., Ishizuka, M., 2015b. Exposure levels of polycyclic

- aromatic hydrocarbons (PAHs) and heavy metals in wild rats in Kumasi, Ghana. Proceedings of the 7th International Toxicology Symposium in Africa, Johannesburg, South Africa. 31st August, 2015. Available online: [http://aa.vetmed.hokudai.ac.jp/en/uploads/2015/10/7th International Toxicology Symposium in Africa.pdf](http://aa.vetmed.hokudai.ac.jp/en/uploads/2015/10/7th_International_Toxicology_Symposium_in_Africa.pdf) pp. 67-68.
- Bouchard, M., Viau, C., 1999. Urinary 1-hydroxypyrene as a biomarker of exposure to polycyclic aromatic hydrocarbons: Biological monitoring strategies and methodology for determining biological exposure indices for various work environments. *Biomarkers* 4, 159–187.
- Burczynski, M.E., Harvey, R.G., Penning, T.M., 1998. Expression and Characterization of Four Recombinant Human Dihydrodiol Dehydrogenase Isoforms: Oxidation of trans-7,8-Dihydroxy-7,8-dihydrobenzo[a]pyrene to the Activated o-Quinone Metabolite Benzo[a]pyrene-7,8-dione. *Biochemistry* 37, 6781-6790.
- Burgaz, S., Borm, P.J.A., Jongeneelen, F.J., 1992. Evaluation of excretion of 1-hydroxypyrene and thioethers in workers exposed to bitumen fumes. *Int. Arch. Occup. Environ. Health* 63, 397–401.
- Campo, L., Rossella, F., Fustinoni, S., 2008. Development of a gas chromatography/mass spectrometry method to quantify several urinary monohydroxy metabolites of polycyclic aromatic hydrocarbons in occupationally exposed subjects, *J. Chromatogr. B* 875, 531–540.
- CDC, 2015. Fourth National Report on Human Exposure to Environmental Chemicals, Updated Tables, February 2015. <http://www.cdc.gov/exposurereport/> (last accessed 22 December 2015).
- Dacasto, M., Eeckhoutte, C., Capolongo, F., Dupuy, J., Carletti, M., Calléja, C., Nebbia, C., Alvinerie, M. and Galtier, P., 2005. Effect of breed and gender on bovine liver cytochrome P450 3A (CYP3A) expression and inter-species comparison with other domestic ruminants. *Vet. Res.* 36(2), 179-190.
- Dissanayake, A., Galloway, T.S., 2004. Evaluation of fixed wavelength fluorescence and synchronous fluorescence spectrophotometry as a biomonitoring tool of environmental contamination. *Mar. Environ. Res.* 58, 281–285.
- Fan, R., Ramage, R., Wang, D., Zhou, J., She, J., 2012. Determination of ten monohydroxylated polycyclic aromatic hydrocarbons by liquid–liquid extraction and liquid chromatography/tandem mass spectrometry. *Talanta* 93, 383-391.
- Fan, R., Wang, D., Mao, C., Ou, S., Lian, Z., Huang, S., Lin, Q., Ding, R., She, J., 2012b. Preliminary study of children's exposure to PAHs and its association with 8-hydroxy-2'-deoxyguanosine in Guangzhou, China. *Environ. Int.* 42, 53-58.

- Ferrari, S., Mandel, F., Berset, J.D., 2002. Quantitative determination of 1-hydroxypyrene in bovine urine samples using high-performance liquid chromatography with fluorescence and mass spectrometric detection. *Chemosphere* 47 (2), 173-182.
- Griego, F.Y., Bogen, K.T., Price, P.S., Weed, D.L., 2008. Exposure, epidemiology and human cancer incidence of naphthalene. *Regul. Toxicol. Pharmacol.* 51 (2), S22–26.
- Guengerich, F.P., 2002. Cytochrome P450, in: Ioannides C. (Ed.), *Enzyme systems that metabolise drugs and other xenobiotics*, John Wiley & Sons Inc., New York, pp. 33–65.
- Guo, H., Huang, K., Zhang, X., Zhang, W., Guan, L., Kuang, D., Deng, Q., Deng, H., Zhang, X., He, M. and Christiani, D., 2014. Women are more susceptible than men to oxidative stress and chromosome damage caused by polycyclic aromatic hydrocarbons exposure. *Environ. Mol. Mutagen.* 55(6), pp.472-481.
- Guo, Y., Senthilkumar, K., Alomirah, H., Moon, H.B., Minh, T.B., Mohd, M.A., Kannan, K., 2013. Concentrations and profiles of urinary polycyclic aromatic hydrocarbon metabolites (OH-PAHs) in several Asian countries. *Environ. Sci. Technol.* 47 (6), 2932-2938.
- Hatzinger, P.B., Alexander, M., 1997. Biodegradation of organic compounds sequestered in organic solids or in nanopores within silica particles. *Environ. Toxicol. Chem.* 16 (11), 2215–2221.
- Jongeneelen, F.J., 2001. Benchmark guideline for urinary 1- hydroxypyrene as biomarker of occupational exposure to polycyclic aromatic hydrocarbons. *Ann. Occup. Hyg.* 45 (1), 3–13.
- Kim, H., Cho, S.H., Kang, J.W., Kim, Y.D., Nan, H.M., Lee, C.H., Lee, H. and Kawamoto, T., 2000. Urinary 1-hydroxypyrene and 2-naphthol concentrations in male Koreans. *Int. Arch. Occ. Environ. Hea.* 74(1), 59-62.
- Kim, M., Kim, D.G., Bong, Y.H., Jang, J.H. and Son, S.W., 2013. Concentrations of PCDD/Fs, dioxin-like PCBs, PBDEs, and hexachlorobenzene in fat samples from cattle of different ages and gender in Korea. *Food chem.* 138(2), 1786-1791.
- Laflamme, R.E., Hites, R.A., 1978. The global distribution of polyaromatic hydrocarbons in recent sediments. *Geochim. Cosmochim. Acta* 42, 289– 303.
- Lake, J.L., Norwood, C., Dimock, C., Bowen, R., 1979. Origins of polycyclic aromatic hydrocarbons in estuarine sediments. *Geochim. Cosmochim. Acta* 43, 1847–1854.
- Levine, H., Berman, T., Goldsmith, R., Göen, T., Spungen, J., Novack, L., Amitai, Y., Shohat, T., Grotto, I., 2015. Urinary concentrations of polycyclic aromatic hydrocarbons in Israeli adults: demographic and life-style predictors. *Int. J. Hyg. Environ. Health* 218 (1), 123–131.
- Li, Z., Mulholland, J.A., Romanoff, L.C., Pittman, E.N., Trinidad, D.A., Lewin, M.D., et al., 2010. Assessment of non-occupational exposure to polycyclic aromatic hydrocarbons

through personal air sampling and urinary biomonitoring. *J. Environ. Monit.* 12, 1110–1118.

Li, Z., Sandau, C.D., Romanoff, L.C., Caudill, S.P., Sjodin, A., Needham, L.L., Patterson, D.G., Jr. 2008. Concentration and profile of 22 urinary polycyclic aromatic hydrocarbon metabolites in the US population. *Environ. Res.*, 107, 320–331.

Meeker, J.D., Barr, D.B., Serdar, B., Rappaport, S.M. and Hauser, R., 2007. Utility of urinary 1-naphthol and 2-naphthol levels to assess environmental carbaryl and naphthalene exposure in an epidemiology study. *J. Expo. Sci. Environ. Epidemiol.* 17 (4), 314–320.

Mollerup, S., Berge, G., Baera, R., Skaug, V., Hwer, A., Phillips, D.H., Stangeland, L., Haugen, A., 2006. Sex differences in risk of lung cancer: Expression of genes in the PAH bioactivation pathway in relation to smoking and bulky DNA adducts. *Int J Cancer* 119, 741–744.

Nam, J.J., Thomas, G.O., Jaward, F.M., Steinnes, E., Gustafsson, O., Jones, K.C., 2008. PAHs in background soils from Western Europe: Influence of atmospheric deposition and soil organic matter. *Chemosphere* 70 (9), 1596–1602.

Nebbia, C., 2001. Biotransformation enzymes as determinants of xenobiotic toxicity in domestic animals. *Vet. J.* 161, 238–252.

Nermell, B., Lindberg, A.L., Rahman, M., Berglund, M., Persson, L.A., El Arifeen, S., Vahter, M., 2008. Urinary arsenic concentration adjustment factors and malnutrition. *Environ. Res.* 106, 212–218.

National Toxicology Program, NTP, 1992. Toxicology and Carcinogenesis Study of Naphthalene in B6C3F1 Mice (Inhalation Studies). Research Triangle Park, NC.

National Toxicology Program, NTP, 2000. Toxicology and Carcinogenesis Study of Naphthalene in F344/N Rats (Inhalation Studies). Research Triangle Park, NC.

Orjuela, M.A., Liu, X., Miller, R.L., Warburton, D., Tang, D., Jobanputra, V., Perera, F.P., 2012. Urinary naphthol metabolites and chromosomal aberrations in 5-year-old children. *Cancer Epidemiol. Biomarkers Prev.* 21 (7), 1191–1202.

Owa, J.A., Izedonmwon, O.E., Ogundaini, A.O., Ogungbamila, F.O., 1993. Quantitative analysis of 1-naphthol in urine of neonates exposed to mothballs: the value in infants with unexplained anaemia. *Afr. J. Med. Med. Sci.* 22 (1), 71–76.

Rey-Salgueiro, L., Martínez-Carballo, E., García-Falcón, M.S., Simal-Gándara, J., 2008. Effects of a chemical company fire on the occurrence of polycyclic aromatic hydrocarbons in plant foods. *Food Chem.* 108, 347–353.

Riojas-Rodriguez, H., Schilman, A., Marron-Mares, A.T., Masera, O., Li, Z., Romanoff, L., et al., 2011. Impact of the improved patsari biomass stove on urinary polycyclic aromatic hydrocarbon biomarkers and carbon monoxide exposures in rural Mexican women. *Environ. Health Perspect.* 119 (9), 1301–1307.

- Saengtienchai, A., Ikenaka, Y., Nakayama, S.M., Mizukawa, H., Kakehi, M., Bortey-Sam, N., Darwish W.S., Tsubota, T., Terasaki, M., Poapolathep, A., Ishizuka, M., 2014. Identification of interspecific differences in phase II reactions: Determination of metabolites in the urine of 16 mammalian species exposed to environmental pyrene. *Environ. Toxicol. Chem.* 33 (9), 2062-2069.
- Soghoian, S., Nyadedzor, C., Ed Nignpense, B., Clarke, E.E.K., Hoffman, R.S., 2012. Health risks of using mothballs in Greater Accra, Ghana. *Trop. Med. Int. Health* 17 (1), 135-138.
- Sul, D., Ahn, R., Im, H., Oh, E., Kim, J.H., Kim, J.G., Kim, P., et al. 2012. Korea National Survey for Environmental Pollutants in the human body 2008: 1-hydroxypyrene, 2-naphthol, and cotinine in urine of the Korean population. *Environ. Res.* 118, 25-30.
- Tay, C.K., Biney, C.A., 2013. Levels and sources of polycyclic aromatic hydrocarbons (PAHs) in selected irrigated urban agricultural soils in Accra, Ghana. *Environ. Earth Sci.* 68, 1773-1782.
- Thai, P.K., Heffernan, A.L., Toms, L.M.L., Li, Z., Calafat, A.M., Hobson, P., Broomhall, S. and Mueller, J.F., 2016. Monitoring exposure to polycyclic aromatic hydrocarbons in an Australian population using pooled urine samples. *Environ. Int.* 88, 30-35.
- Uppstad, H., Osnes, G.H., Cole, K.J., Phillips, D.H., Haugen, A., Mollerup, S., 2011. Sex differences in susceptibility to PAHs is an intrinsic property of human lung adenocarcinoma cells. *Lung Cancer* 71, 264-270.
- Wayne, R.O., 1990. A physical explanation of the lognormality of pollutant concentrations. *J. Air Manag. Assoc.* 40, 1378-1383.
- Xu, X., Cook, R.L., Ilacqua, V.A., Kan, H., Talbott, E.O., Kearney, G., 2010. Studying associations between urinary metabolites of polycyclic aromatic hydrocarbons (PAHs) and cardiovascular diseases in the United States. *Sci. Total Environ.* 408, 4943-4948.

Figure captions:

Fig. 1 Map showing cattle urine sampling locations in the Ashanti Region, Ghana (yellow pins indicate sampled locations and red pin indicate city centre in Kumasi)

Fig. 2 Geometric mean concentrations (adjusted by specific gravity) of OH-PAHs in cattle urine from 5 sample sites in Kumasi and Offinso, Ghana

567

568

569

570

571 Table 1: Quality assurance and control (QA/QC) for OH-PAHs analysis in cattle urine

Compound name	LOD (ng/mL)	LOQ (ng/mL)	ISTD Recovery (%)	Spiked solvent blanks (%)	Matrix spikes (%)
2-OHNaphthelene	0.0898	0.295			
9-OHFluorene	0.181	0.603			
2-3-OHFluorene	0.0745	0.248			
<i>13C6-2-OHFluorene</i>			94	96	96
1-9-OHPhenanthrene	0.214	0.714			
2-OHPhenanthrene	0.0696	0.232			
3-OHPhenanthrene	0.348	1.16			
4-OHPhenanthrene	0.0437	0.145			
<i>3-OHPhenanthrene-d9</i>			96	98	95
1-OHPyrene	0.259	0.865			
<i>13C6-1-OHPyrene</i>			89	92	85
6-OHChrysene	0.220	0.733			
3-OHBenzo(e)Pyrene	0.162	0.542			
9-OHBenzo(a)Pyrene	0.0959	0.319			

572 Italicized compounds are internal standards (ISTD); LOD: Limit of detection; LOQ: Limit
573 of quantification

574

575

576

577

Table 2: Specific gravity adjusted urinary OH-PAHs concentrations (ng/mL) in cattle from urban and rural sites in Kumasi and Offinso, Ghana

Site	n	2-OHNap (GM _{SG} ± SD)	2-3-OHFlu (GM _{SG} ± SD)	2-OHPhe (GM _{SG} ± SD)	1-9-OHPhe (GM _{SG} ± SD)	4-OHPhe (GM _{SG} ± SD)	1-OHPyr (GM _{SG} ± SD)
Urban	17	2.29 ± 3.40 ^a	0.919 ± 0.462 ^a	0.245 ± 0.171 ^a	1.96 ± 1.14 ^a	1.09 ± 0.841 ^a	1.52 ± 0.873 ^a
Rural	78	2.91 ± 6.37 ^a	1.24 ± 1.97 ^a	0.552 ± 0.598 ^b	2.03 ± 1.17 ^a	1.88 ± 2.01 ^b	1.27 ± 0.824 ^a

n: number of samples; GM_{SG}: geometric mean concentration adjusted by specific gravity; SD: standard deviation; different letters (a and b) within a column indicate significant differences (Student's T-Test; $p < 0.05$)

Table 3: Specific gravity adjusted OH-PAHs concentrations (ng/mL) in cattle urine

Sample site	n	Location	2-OHNap (GM _{SG} ± SD)	2-3-OHFlu (GM _{SG} ± SD)	1-9-OHPhe (GM _{SG} ± SD)	2-OHPhe (GM _{SG} ± SD)	3-OHPhe (GM _{SG} ± SD)	4-OHPhe (GM _{SG} ± SD)	1-OHPyr (GM _{SG} ± SD)
Oforikrom	8	urban	4.15 ± 4.37 ^b	0.99 ± 0.62 ^b	1.67 ± 1.21 ^b	0.17 ± 0.10 ^c	nd	0.73 ± 0.41 ^c	1.37 ± 1.18 ^{ab}
Santasi	9	urban	0.61 ± 0.23 ^{cd}	0.75 ± 0.35 ^b	2.26 ± 1.14 ^{ab}	0.32 ± 0.20 ^{bc}	nd	1.14 ± 0.33 ^{bc}	1.33 ± 0.80 ^{ab}
Twumasen Estate	31	rural	0.69 ± 0.57 ^d	0.31 ± 0.13 ^c	1.73 ± 0.58 ^{ab}	0.29 ± 0.26 ^c	nd	1.50 ± 0.86 ^b	0.99 ± 0.73 ^b
Saboa	40	rural	1.24 ± 0.67 ^c	0.80 ± 0.79 ^b	2.07 ± 1.43 ^{ab}	0.41 ± 0.16 ^b	nd	1.27 ± 0.91 ^{bc}	1.16 ± 0.74 ^b
Kokote	7	rural	21.9 ± 6.51 ^a	6.74 ± 1.41 ^a	3.12 ± 0.79 ^a	2.26 ± 0.51 ^a	2.1±0.57	7.49 ± 1.73 ^a	2.29 ± 1.28 ^a

n: number of samples; nd: below limits of quantification (LOQ); different letter (a, b, c and d) within a column indicate significant difference ($p < 0.05$) among communities; GM_{SG}: geometric mean concentration adjusted by specific gravity; SD: standard deviation

Table 4: Specific gravity adjusted urinary OH-PAHs concentrations (ng/mL) in male and female cattle in Kumasi and Offinso, Ghana

Sex	2-OHNap (GM _{SG} ± SD)	2-3-OHFlu (GM _{SG} ± SD)	1-9-OHPhe (GM _{SG} ± SD)	2-OHPhe (GM _{SG} ± SD)	3-OHPhe (GM _{SG} ± SD)	4-OHPhe (GM _{SG} ± SD)	1-OHPyr (GM _{SG} ± SD)	ΣOHPAHs (GM _{SG} ± SD)
Male	4.43 ± 7.16 ^a	1.36 ± 1.94 ^a	1.71 ± 1.07 ^a	0.534 ± 0.650 ^a	0.364 ± 0.658 ^a	1.91 ± 2.22 ^a	1.19 ± 0.981 ^a	11.5 ± 14.6 ^a
Female	2.01 ± 5.11 ^b	0.950 ± 1.65 ^a	2.16 ± 1.18 ^b	0.468 ± 0.510 ^a	0.237 ± 0.499 ^a	1.65 ± 1.69 ^a	1.22 ± 0.827 ^a	8.71 ± 10.0 ^a

GM_{SG}: geometric mean concentration adjusted by specific gravity; SD: standard deviation; ΣOHPAHs: sum of OHPAHs; different letters (a and b) within a column indicate significant differences (Student's T-Test; $p < 0.05$)

Fig. 1

Fig. 2