

HOKKAIDO UNIVERSITY

Title	Chiroptical Properties of Nonanuclear Tb(III) Clusters with Chiral Champhor Derivative Ligands
Author(s)	Wada, Satoshi; Kitagawa, Yuichi; Nakanishi, Takayuki; Fushimi, Koji; Hasegawa, Yasuchika
Citation	e-Journal of Surface Science and Nanotechnology, 13, 31-34 https://doi.org/10.1380/ejssnt.2015.31
Issue Date	2015-02-14
Doc URL	http://hdl.handle.net/2115/72412
Rights	©2015 The Surface Science Society of Japan
Rights(URL)	https://creativecommons.org/licenses/by/4.0/
Туре	article
File Information	Surf.Sci.Nanotech.13.31.pdf



e-J. Surf. Sci. Nanotech. Vol. 13 (2015) 31-34

Conference - CIF'15 -

Chiroptical Properties of Nonanuclear Tb(III) Clusters with Chiral Champhor Derivative Ligands^{*}

Satoshi Wada

Graduate School of Chemical Sciences and Engineering, Hokkaido University, North 13 West 8, Kita-ku, Sapporo, Hokkaido 060-8628, Japan

Yuichi Kitagawa, Takayuki Nakanishi, Koji Fushimi, and Yasuchika Hasegawa[†] Faculty of Engineering, Hokkaido University, North 13 West 8, Kita-ku, Sapporo, Hokkaido 060-8628, Japan

(Received 29 Decemberr 2014; Accepted 26 January 2015; Published 14 February 2015)

The chiroptical properties of nonanuclear Tb(III) clusters are reported. The nonanuclear Tb(III) clusters are composed of nine Tb(III) ions, ten μ -OH parts, and sixteen chiral organic ligands, (\pm)-bornyl salicylate. Their chiroptical properties were estimated by circular dichroism (CD) and circularly polarized luminescence (CPL). Their electronic structures were calculated using TD-DFT (B3LYP/6-31G(d)) method. [DOI: 10.1380/ejssnt.2015.31]

 $Keywords: \ Lanthanides; \ Photoluminescence; \ Chirality; \ Circularly \ polarized \ luminecnece; \ Visible/ultraviolet \ absorption \ spectroscopy$

I. INTRODUCTION

Luminescence of metal complexes composed of metal ions and organic ligands has attracted wide interest for applications that include optical materials [1], organic light-emitting diodes [2, 3], and fluorescent sensors[4, 5]. The luminescent properties of these complexes strongly depend on the combination of the metal center and the organic ligands. The luminescence properties of transitionmetal complexes are derived from the d-d and metal-toligand charge-transfer (MLCT) transitions based on the outer d-orbitals of the complexes [6]. These characteristics are strongly dominated by the orbital interactions between the transition-metal ion and the organic ligands [6]. The design of the organic ligands can give controllable luminescent wavelength and intensity based on the d-d and the MLCT transitions.

On the other hand, lanthanide ions have the inner 4f orbitals, which induce different luminescence properties from those of transition-metal ions. Compared to transition-metal complexes, lanthanide complexes provide specific narrow emission band because of their inner 4f orbitals [7, 8]. They also exhibit long emission lifetime based on the 4f-4f forbidden transitions. Here, organic ligands can play an important role in the intense lanthanide luminescence. The organic ligands perform as efficient light harvesting systems for photosensitized luminescence of the lanthanide complexes. The organic ligands also enhance the radiative rate by introducing an asymmetric geometry around the lanthanide ion, effectively [9–12]. Based on their two strategies for design of the ligands, various types of strong luminescent lanthanide complexes with asymmetric structures have been reported [13, 14].

Luminescent complexes with chiral ligands show circularly polarized luminescence (CPL) [15]. This is based on the differential emission intensity of left- and righthanded circular polarized light according to the chirality



FIG. 1. Synthetic scheme of a (-)-bornyl salicylate ligand and a nonanuclear Tb(III) cluster.

of the emitter, and is often used as an indicator of chirality in a metal ion center [16–18]. The CPL property is expected to be useful for applications such as future threedimensional displays, security paints, and optical telecommunications. The magnitude of CPL is represented by the dissymmetry factor $g_{\rm CPL}$ [15]. At the present stage, a large number of scientific researches on mononuclear chiral lanthanide complexes with large $g_{\rm CPL}$ values have been performed [19–21].

Recently, polynuclear lanthanide clusters with chiral ligands have attracted attention for their CPL with characteristic structure [22, 23]. For example, Gunnlaugsson reported large CPL of dinuclear Eu(III) complexes using triple-stranded helical ligands [24]. Mazzanti reported the changeable $g_{\rm CPL}$ of polynuclear Eu(III) clusters depending on their several types of self-assembly structures [25]. In contrast, chiroptical lanthanide clusters composed of lanthanide ions and bridged oxygens, Ln-O-Ln lattices, have never been reported yet. In this study, we report the chiroptical properties of novel nonanuclear Tb(III) clusters for the first time. These clusters are constructed by nine Tb(III) ions, ten μ -OH parts, and sixteen chiral salicylic acid esters, bornyl salicylates, (\pm) -bor (Fig. 1) [26]. These ligands are champhor derivatives for enhancement of g_{CPL} according to the previous report [27]. We have demonstrated the intense CPL and electronic structures of Tb(III) clusters with bornyl salicylates, Tb- (\pm) -bor.

^{*} This paper was presented at the 15th Chitose International Forum "Nanotechnology - From Synthesis to Devices," Chitose Institute of Science and Technology, Chitose, Japan, October 2-3, 2014.

[†] Corresponding author: hasegaway@eng.hokudai.ac.jp

II. EXPERIMENTAL

A. Materials

Salicylic acid (C₆H₄(OH)COOH) and 1,4-dioxane (C₄H₈O₂) were purchased from Wako Pure Chemical Industries, Ltd. (+)-borneol (C₁₀H₁₈O, 97%) and (-)-borneol (C₁₀H₁₈O, 97%) were purchased from Aldrich. Thionyl chloride (SOCl₂) was purchased from Tokyo Chemical Industry Co., Ltd. Terbium(III) nitrate hexahydrate (Tb(NO₃)₃·6H₂O), and triethylamine ((C₂H₅)₃N) were purchased from Kanto Chemical Co., Inc.

B. Instrumental Techniques

¹H-NMR spectra were recorded in CDCl₃ on an Auto-NMR JEOL ECS 400 MHz Spectrometer; CHCl₃ ($\delta_{\rm H} =$ 7.26 ppm) was used as an internal reference. Fast atomic bombardment mass spectrometry (FAB-MS) spectra were measured on a JEOL JMS-700TZ. Electronic absorption, circular dichroism (CD), and CPL spectra were measured with a JASCO V-670 spectrophotometer, Jasco J-1500 spectropolarimeter and Jasco CPL-200 spectrofluoropolarimeter (excitation wavelength = 380 nm), respectively.

C. Synthesis of (+)-bornyl salicylate ((+)-bor, $C_{17}H_{22}O_3$) [28]

A solution of salicylic acid (1.0 g, 7.2 mmol) and (+)borneol (1.12 g, 7.2 mmol) in 30 mL of dioxane were refluxed with thionyl chloride (0.82 g, 6.9 mmol) for 1 h at 100°C. The mixture was extracted with NaHCO₃ aq. (2×20 mL) and ethyl acetate (2×20 mL). The organic layer was separated and dried with magnesium sulfate, and the solvent was evaporated. The residue was chromatographed on silica gel eluting with ethyl acetate/hexane (10/90). Yield: 0.29 g (17%). ¹H-NMR (400 MHz, CDCl₃): δ /ppm = 10.9 (s, 1H, -OH), 7.87– 7.89 (d, 1H, Ar), 7.44–7.48 (t, 1H, Ar), 6.97–6.99 (d, 1H, Ar), 6.88–6.92 (t, 1H, Ar), 5.10–5.14 (m, 1H, -CH), 2.07– 2.51 (m, 2H), 1.75–1.77 (m, 1H), 1.12–1.47 (m, 4H), 0.97 (s, 3H, -CH₃), 0.92 (s, 6H, -CH₃).

D. Synthesis of (–)-bornyl salicylate ((-)-bor, $C_{17}H_{22}O_3)$ [28]

Yield: 0.63 g (32%). ¹H-NMR (400 MHz, CDCl₃): δ /ppm = 10.9 (s, 1H, -OH), 7.87–7.89 (d, 1H, Ar), 7.44–7.48 (t, 1H, Ar), 6.97–6.99 (d, 1H, Ar), 6.88–6.92 (t, 1H, Ar), 5.10–5.14 (m, 1H, -CH), 2.07-2.51 (m, 2H), 1.75–1.77 (m, 1H), 1.12–1.47 (m, 4H), 0.97 (s, 3H, -CH₃), 0.92 (s, 6H, -CH₃).

E. Synthesis of $[Tb_9(sal-(+)-bor)_{16}(\mu-OH)_{10}]^+[NO_3]^ (Tb-(+)-bor, C_{272}H_{346}NO_{61}Tb_9)$

(+)-bornyl salicylate (0.10 g, 0.36 mmol) was dissolved in methanol, and triethylamine (0.036 g, 0.36 mmol) was added to this solution with stirring at 40°C. Then Tb(NO₃)₃·6H₂O (0.095 g, 0.21 mmol) in methanol was added dropwise to this solution with further stirring for 20 min. White powder, Tb-(+)-bor, $[Tb_9(sal-(+)-bor)_{16}(\mu-OH)_{10}]^+[NO_3]^-$, was obtained. Yield: 0.079 g (56%). MS (FAB) found: m/z = 5973.54, calcd for $C_{272}H_{346}NO_{61}Tb_9$: M⁺, 5973.75.

F. Synthesis of $[Tb_9(sal-(-)-bor)_{16}(\mu-OH)_{10}]^+[NO_3]^-$ (Tb-(-)-bor, C₂₇₂H₃₄₆NO₆₁Tb₉)

(–)-bornyl salicylate (0.10 g, 0.36 mmol) was dissolved in methanol, and triethylamine (0.036 g, 0.36 mmol) was added to this solution with stirring at 40°C. Then Tb(NO₃)₃·6H₂O (0.095 g, 0.21 mmol) in methanol was added dropwise to this solution with further stirring for 20 min. White powder, Tb-(–)-bor, [Tb₉(sal-(–)bor)₁₆(μ -OH)₁₀]⁺[NO₃]⁻, was obtained. Yield: 0.11 g (76%). MS (FAB) found: m/z = 5973.88, calcd for C₂₇₂H₃₄₆NO₆₁Tb₉: M⁺, 5973.75.

G. Computational Details

Density function theory (DFT) calculations were carried out with the Gaussian R09W B.01 employing the three-parameter hybrid functional of Becke based on the correlation functional of Lee, Yang, and Parr (B3LYP) [29, 30]. The 6-31G(d) basis set was used for all atoms.

III. RESULTS AND DISCUSSION

A. Electronic structures of the ligands and the Tb(III) clusters

The UV-vis absorption spectra of (\pm) -bor molecules and Tb- (\pm) -bor clusters are shown in Fig. 2a. The absorption bands of (\pm) -bor molecules and Tb- (\pm) -bor clusters were observed at around 307 nm and 350 nm, respectively. Their absorption bands are assigned to the π - π * transition of (\pm) -bor molecules and ligands in Tb- (\pm) -bor clusters. The absorption coefficients of ligands in Tb- (\pm) -bor clusters were approximately 60,000 M⁻¹cm⁻¹ [31], which was 200,000 times larger than that of 4f-4f transition of Tb(III) ion. The large absorption coefficients may lead to effective photosensitized luminescence of Tb- (\pm) -bor clusters.

To clarify the π -electronic properties of the (\pm) -bor ligands, the energy level and oscillator strength of the lowest singlet excited states (S1) were calculated using the TD-DFT (B3LYP/6-31G(d)) method. According to the Tb- (\pm) -bor clusters, we carried out the simplified complex model, Li-(+)-bor. The oscillator strengths of a (+)-bor molecule and a Li-(+)-bor complex were calculated to be 0.116 ($\lambda = 291.5$ nm) and 0.145 ($\lambda = 334.5$ nm), respectively. These transitions are mainly dominant to HOMO→LUMO transition (Fig. 3: transition probability $TP_{HOMO \rightarrow LUMO,(+)-bor} = 94\%$, $TP_{HOMO \rightarrow LUMO, Li-(+)-bor} = 95\%$). The energy gap between HOMO and LUMO is directly linked to the absorption wavelength. The energy gaps of (+)-bor and Li-(+)bor part are found to be 4.81 eV and 4.15 eV, respectively (Fig. 3). The magnitude of the destabilization of HOMO in Li- (\pm) -bor is larger than that of LUMO.

At the red-shifted intense absorption band, positive (or negative) CD signal was observed for Tb-(+)-bor (or Tb-(+)-bor (bard) bard) bard)



FIG. 2. UV-vis (a) and CD (b) spectra of Tb-(+)-bor (red line), Tb-(-)-bor (blue line), (+)-bor (black line), and (-)-bor (gray line) in DMF $(2.0 \times 10^{-4} \text{ M})$.



FIG. 3. Molecular orbital of (+)-bor (a) and Li-(+)-bor (b).

(-)-bor) clusters (Fig. 2b). In general, asymmetric factor of CD signals, $g_{\rm CD}$, is gained by the following equation,

$$g_{\rm CD} = 4 \frac{|m|}{|\mu|} \cos\theta \tag{1}$$

where m, μ , and θ are represented by the magnetic transition dipole moment, the electric transition dipole moment, and the angle between m and μ , respectively. Asymmetric factor $g_{\rm CD}$ of the cluster was 1.8×10^{-4} at 376 nm. The $g_{\rm CD}$ in Tb-(±)-bor clusters are the same order to those of typical chiral molecules ($g_{\rm CD} \simeq 10^{-4}$).

B. Luminescence properties of Tb(III) clusters.

Figure 4 shows photoluminescence (PL) and CPL spectra of the Tb-(\pm)-bor clusters. The spectral shapes of 4f-4f transitions in Tb-(+)-bor clusters agreed with that in Tb-(-)-bor clusters (Fig. 4a). These transition bands in PL spectra were contributed to ${}^{5}\text{D}_{4} \rightarrow {}^{7}\text{F}_{J}$ (J = 6, 5, 4, 3) transitions. We also observed the CPL spectra of Tb-(\pm)-bor clusters (Fig. 4b). The wavelengths and g_{CPL} are summarized in table 1. Asymmetric factor of CPL, gCPL,



FIG. 4. CPL (a) and PL (b) spectra of Tb-(+)-bor (red line), and Tb-(-)-bor (blue line) in DMF $(1.0^{-4} M)$.

TABLE I. Luminescent wavelengths and corresponding $|g_{CPL}|$ values of Tb(III) ions.

Transition band	Wavelength/nm	$ g_{ m gCPL} $
${}^{5}\mathrm{D}_{4} \rightarrow {}^{7}\mathrm{F}_{6} \ (\Delta J = +2)$	488	0.0056
${}^{5}\mathrm{D}_{4} \rightarrow {}^{7}\mathrm{F}_{5} \ (\Delta J = +1)$	545	0.024
${}^{5}\mathrm{D}_{4} \rightarrow {}^{7}\mathrm{F}_{4} (\Delta J = 0)$	582	0.0057
${}^{5}\mathrm{D}_{4} \rightarrow {}^{7}\mathrm{F}_{3} \ (\Delta J = -1)$	630	0.017

is expressed by the same equation to $g_{\rm CD}$ [32], and $g_{\rm CD}$ and g_{CPL} of the magnetic dipole transition band are larger than those of electric dipole transition band. According to the luminescent properties of Eu(III) complexes, transitions for $\Delta J = \pm 1$ are closely related to the magnetic dipole transition. In our experiments, g_{CPL} of $\Delta J = \pm 1$ in Tb-(\pm)-bor clusters were the larger than those of ΔJ = 0 and ± 2 in Tb-(\pm)-bor clusters. These $g_{\rm CPL}$ values $(\Delta J = 1)$ are similar to those of previous chiral Tb(III) complexes, *e.g.* previous reported mononuclear Tb(III) complex $(g_{\text{CPL}} = 0.04)$ [16]. In this study, Tb(III) cluster composed of nine Tb(III) ions also shows effective CPL properties compared with mononuclear Tb(III) complex. This result indicates that chiroptical Tb(III) cluster with Tb-O-Tb lattices is expected to be useful for future optical materials.

IV. CONCLUSIONS

We synthesized novel nonanuclear Tb(III) clusters with chiral bornyl salicylate ligands. They emitted strongly according to the 4f-4f transitions of Tb(III) ions, and showed CPL signal corresponding to these transitions. This is the first observation of CPL using Tb(III) clusters with inorganic frameworks. Novel lanthanide cluster with CPL properties is expected to open up a frontier field to coordination chemistry, photophysical chemistry and material science.

ACKNOWLEDGMENTS

This work was partly supported by Grants-in-Aid for Scientific Research on Innovative Areas of "New Polymeric Materials Based on Element-Blocks (no. 2401)" (no. 24102012) of the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan. We are

- G. Blasse and B. C. Grabmaier, *Luminescent Materials* (Springer-Verlag, New York, 1994).
- [2] X.-M. Yu, G.-J. Zhou, C.-S. Lam, W.-Y. Wong, X.-L. Zhu, J.-X. Sun, M. Wong, and H.-S. Kwok, J. Organomet. Chem. 693, 1518 (2008).
- [3] J. Kido and Y. Okamoto, Chem. Rev. 102, 2357 (2002).
- [4] R. Gao, D. G. Ho, B. Hermamdez, M. Selke, D. Murphy,
 P. I. Djurovich, and M. Thompson, J. Am. Chem. Soc. 124, 14828 (2002).
- [5] T. Gunnlaugsson, J. P. Leonard, K. Sénéchal, and A. J. Harte, J. Am. Chem. Soc. 125, 12062 (2003).
- [6] J. N. Demas and B. A. DeGraff, Anal. Chem. 63, 829 (1991).
- [7] J.-C. G. Bünzli, and C. Piguet, Chem. Rev. 102, 1897 (2002).
- [8] V. S. Sastri, J.-C. G. Bünzli, V. R. Rao, G. V. S. Rayudu, and J. R. Perumareddi, *Modern Aspects of Rare Earth and Their Complexes* (Elsevier, New York, 2003).
- [9] K. Binnemans, Chem. Rev. **109**, 4283 (2009).
- [10] A. F. Kirby and F. S. Richardson, J. Phys. Chem. 87, 2544 (1983).
- [11] J. P. Cross, M. Lauz, P. D. Badger, and S. Petoud, J. Am. Chem. Soc. **126**, 2280 (2004).
- [12] W. Liu, T. Jiao, Y. Li, Q. Liu, M. Tan, H. Wang, and L. Wang, J. Am. Chem. Soc. **126**, 16278 (2004).
- [13] K. Miyata, Y. Hasegawa, Y. Kuramochi, T. Nakagawa, T. Yokoo, and T. Kawai, Eur. J. Inorg. Chem. **32**, 4777 (2009).
- [14] P. Nockemann, B. Thijs, N. Postelmans, K. V. Heche, L. V. Meervelt, and K. Binnemans, J. Am. Chem. Soc. 128, 13658 (2006).
- [15] J. P. Riehl and F. S. Richardson, Chem. Rev. 86, 1 (1986).
- [16] M. Seitz, E. G. Moore, A. J. Ingram, G. Muller, and K. N. Raymond, J. Am. Chem. Soc. **129**, 15468 (2007).
- [17] J. Gregolinski, P. Starynowicz, K. T. Hua, J. L. Lunkley, G. Muller, and J. Lisowski, J. Am. Chem. Soc. 130, 17761

(2008).

[18] R. Carr, N. H. Evans, and D. Parker, Chem. Soc. Rev. 41, 7673 (2012).

particularly grateful for the experimental assistance by

Prof. K. Konishi, Prof. T. Ohkuma and Assoc. Prof. N. Arai of Hokkaido University, and Prof. Y. Chujo and

Assoc. Prof. Y. Morisaki of Kyoto University. We would

also thank for the support of Frontier Chemistry Center

Akira Suzuki "Laboratories for Future Creation" Project.

- [19] J. Yuasa, T. Ohno, K. Miyata, H. Tsumatori, Y. Hasegawa, and T. Kawai, J. Am. Chem. Soc. 133, 9892 (2011).
- [20] S. Petoud, G. Muller, E. G. Moore, J. Xu, J. Sokolnicki, J. P. Riehl, U. N. Le, S. M. Cohen, and K. N. Raymond, J. Am. Chem. Soc. **129**, 77 (2007).
- [21] J. A. Kitchen, D. E. Barry, L. Mercs, M. Albrecht, R. D. Peacock, and T. Gunnlaugsson, Angew. Chem. Int. Ed. 51, 704 (2012).
- [22] O. Mamula, M. Lama, S. G. Telfer, A. Nakamura, R. Kuroda, H. Stoeckli-Evans, and R. Scopelitti, Angew. Chem. Int. Ed. 44, 2527 (2005).
- [23] B. S. Murray, D. Parker, C. M. G. dos Santos, and R. D. Peacock, Eur. J. Inorg. Chem. 18, 2663 (2010).
- [24] F. Stomeo, C. Lincheneau, J. P. Leonard, J. E. O'Brien, R. D. Peacock, C. P. McCoy, and T. Gunnlaugsson, J. Am. Chem. Soc. **131**, 9636 (2009).
- [25] G. Bozoklu, C, Gateau, D. Imbert, J. Pécaut, K. Roveyns, Y. Filinchuk, F. Memon, G. Muller, and M. Mazzanti, J. Am. Chem. Soc. **134**, 8372 (2012).
- [26] T. Nakanishi, Y. Suzuki, Y. Doi, T. Seki, H. Koizumi, K. Fushimi, K. Fujita, Y. Hinatsu, H. Ito, K. Tanaka, and Y. Hasegawa, Inorg. Chem. 53, 7635 (2014).
- [27] J. L. Lunkley, D. Shirotani, K. Yamanari, S. Kaizaki, and G. Muller, Inorg. Chem. 50, 12724 (2011).
- [28] R. M. C. Vasconcelos, F. C. Leite, J. A. Leite, S. R. Mascarenhas, L. C. Rodrigues, and M. R. Piuvezam, Immunopharm. Immunot. 34, 1028 (2012).
- [29] C. T. Lee, W. T. Yang, and R. G. Parr, Phys. Rev. B 37, 785 (1988).
- [30] A. D. Becke, J. Chem. Phys. 98, 5648 (1993).
- [31] K. Manseki and S. Yanagida, Chem. Commun. 12, 1242 (2007).
- [32] F. S. Richardson, Inorg. Chem. 19, 2806 (1980).