



Title	Olefin-accelerated solid-state C-N cross-coupling reactions using mechanochemistry
Author(s)	Kubota, Koji; Seo, Tamae; Koide, Katsumasa; Hasegawa, Yasuchika; Ito, Hajime
Citation	Nature communications, 10, 111 https://doi.org/10.1038/s41467-018-08017-9
Issue Date	2019-01-10
Doc URL	http://hdl.handle.net/2115/73003
Rights(URL)	http://creativecommons.org/licenses/by/4.0/
Type	article
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Supplementary Information-1.pdf



[Instructions for use](#)

Supplementary Information

Olefin-Accelerated Solid-State C–N Cross-Coupling Reactions Using Mechanochemistry

*Koji Kubota^{*a}, Tamae Seo^a, Katsumasa Koide^a, Yasuchika Hasegawa^{a,b}, Hajime Ito^{*a,b}*

^aDivision of Applied Chemistry, Graduate School of Engineering, Hokkaido University, Sapporo, Hokkaido 060-8628, Japan.

^bInstitute for Chemical Reaction Design and Discovery (WPI-ICReDD), Hokkaido University, Sapporo, Hokkaido 060-8628, Japan.


Supplementary Methods

Chemicals and Instrumentation

The starting materials expect for 4-bromo-3',4',5',6'-tetraphenyl-1,1':2',1''-terphenyl (**1z**), 5,15-dibromo-10,20-bis(3,5-dimethylphenyl)porphyrin (**1aa**) and (10-bromoanthracen-9-yl)dimesitylborane (**1ab**) were obtained from commercial suppliers and used as received. **1z**¹, **1aa**^{2,3} and **1ab**⁴ were prepared according to the reported procedures, respectively. Solvents for substrate preparation were purchased from commercial suppliers, and further dried over molecular sieve (MS 4Å). All mechanochemical reactions were carried out using grinding vessels in a Retsch MM400 mill. Both bowls (1.5 mL or 25 mL) and balls are made of stainless. NMR spectra were recorded on JEOL JNM-ECX400P and JNM-ECS400 spectrometers (¹H: 392 or 396 or 399 or 401 MHz, ¹³C: 99 or 100 MHz). Tetramethylsilane (¹H), CDCl₃ (¹³C) was employed as external standards, respectively. Multiplicity was recorded as follows: s = singlet, brs = broad singlet, d = doublet, t = triplet, q = quartet, quint = quintet, sept = septet, o = octet, m = multiplet. Dibromomethane was used as an internal standard to determine NMR yields. GLC analyses were conducted with a Shimazu GC-2014 or GC-2025 equipped with ULBON HR-1 glass capillary column (Shinwa Chemical Industries) and a FID detector. High-resolution mass spectra were recorded at the Global Facility Center, Hokkaido University. MALDI-TOF mass spectra were recorded at the Open Facility Center, Hokkaido University. Powder diffraction data were recorded on a Rigaku SmartLab diffractometer with Cu-K_α radiation and D/teX Ultra detector covering 5-60° (2θ). Thermography was recorded with an NEC Avio Thermo GEAR G120. Transmission electron microscopy (TEM) analysis was carried out at “Joint-Use Facilities: Laboratory of Nano-Micro Material Analysis”, Hokkaido University.

General Procedure for Solid-State C–N Coupling Using a Ball Mill

Aryl halide **1** (0.5 mmol), diarylamine **2** (0.5 mmol, 1.0 equiv), Pd(OAc)₂ (0.025 mmol, 5 mol %) were placed in a ball milling vessel (stainless, 1.5 mL) loaded with one grinding ball (stainless, diameter: 3 mm). The ball milling vessel was then transferred to a glovebox. In a glovebox, *t*-Bu₃P (0.025 mmol, 5 mol %) and Na(O-*t*-Bu) (0.75 mmol, 1.5 equiv) were placed in the milling vessel. After the milling vessel was removed from glovebox, the vessel was opened under air, then 1,5-cod (0.20 μL mg⁻¹) was added via syringe. After the vessel was closed in air without the purge with inert gas, the vessel was placed in the ball mill (Retch MM400, 99 min at 30Hz). After 99 min, the mixture was passed through a short silica gel column eluting with EtOAc to remove inorganic salts. The crude mixture was then purified by flash column chromatography (SiO₂, CH₂Cl₂/hexane, typically 0-15:85) to give the corresponding arylamines **3**.




Supplementary Figure 1. How to set up the mechanochemical reactions.

Procedure for Solid-State C–N Coupling on a Gram Scale

1b (7.0 mmol, 1.970 g), **2c** (7.0 mmol, 1.381 g, 1.0 equiv), Pd(OAc)₂ (0.14 mmol, 31.4 mg, 2 mol %) were placed in a ball milling vessel (stainless, 25 mL) loaded with 4 grinding balls (stainless, diameter: 10 mm). The ball milling vessel was then transferred to a glovebox. In a glovebox, *t*-Bu₃P (0.14 mmol, 28.4 mg, 2 mol %) and Na(O-*t*-Bu) (10.5 mmol, 1.010 g, 1.5 equiv) were placed in the milling vessel. After the milling vessel was removed from glovebox, the vessel was opened under air, then 1,5-cod (840 μL, 0.20 μL mg⁻¹) was added quickly. After the vessel was closed without the purge with inert gas, the vessel was placed in the ball mill (Retch MM400, 99 min at 30Hz). After 99 min, the mixture was passed through a short silica gel column eluting with EtOAc to remove inorganic salts. The crude mixture was then purified by reprecipitation from CH₂Cl₂/MeOH to give the arylamine **3c** as a yellow solid (2.567 g, 92% yield).

Additional Results of Optimization Study





amount of 1,5-cod	conversion fo 1a (%) ^a	yield (%) ^a
1,5-cod (40 μl) (0.13 μL mg ⁻¹)	78	77
1,5-cod (40 μl) (0.13 μL mg ⁻¹) ^b	91	90
1,5-cod (60 μl) (0.20 μL mg ⁻¹)	85	81
1,5-cod (60 μl) (0.20 μL mg⁻¹)^b	>99	99
1,5-cod (80 μl) (0.26 μL mg ⁻¹)	85	83
1,5-cod (80 μl) (0.26 μL mg ⁻¹) ^b	>99	99
1,5-cod (100 μl)(0.33 μL mg ⁻¹)	81	75
1,5-cod (150 μl)(0.50 μL mg ⁻¹)	71	66



^aDetermined by NMR analysis with an internal standard.

^bReaction time was 99 min.


Supplementary Figure 2. Investigation on the amount of 1,5-cod.



Supplementary Figure 3. Scope of liquid aryl halides without olefin additives.




Supplementary Figure 4. Preliminary investigation on scope of solid aryl halides without olefin.



Supplementary Figure 5. The use of Pd(dba)₂ instead of Pd(OAc)₂.

Reaction Temperature Confirmed by Thermography


The temperature inside the milling jar after the solid-state coupling reaction was confirmed by thermography. The crude mixtures were prepared by the following conditions: 0.5 mmol of **1b**; 0.5 mmol of **2a**; 0.025 mmol of Pd(OAc)₂; 0.025 mmol of ligand; 0.75 mmol of Na(O-*t*-Bu); 1,5-cod (60 μl) in a stainless-steel ball-milling jar (1.5 mL) with a stainless-steel ball (3 mm); 30 Hz; 60 min. The obtained image showed that the temperature was around 30 °C, indicating that this reaction proceeded at round room temperature.




Supplementary Figure 6. Thermography image of the reaction mixture after the grinding with 30 Hz for 60 min.

Characterization of Palladium Nanoparticles by Transmission Electron Microscopy


The crude mixtures were prepared by the following conditions: 0.5 mmol of **1b**; 0.5 mmol of **2a**; 0.025 mmol of Pd(OAc)₂; 0.025 mmol of ligand; 0.75 mmol of Na(O-*t*-Bu); 1,5-cod (62 μl) or cyclooctane (62 μl) in a stainless-steel ball-milling jar (1.5 mL) with a stainless-steel ball (3 mm); 30 Hz; 99 min. The samples for the characterization by transmission electron microscopy (TEM) were prepared by dropping the colloidal solution of MeOH onto a copper grid covered with thin carbon film. Additional images of the reaction mixtures upon grinding for 99 min were shown in Supplementary Figure 7–9. The timescales for the aggregation of palladium particles in the reaction mixtures that contain cyclooctane or that are free of additives were shown in Supplementary Figure 10 and 11.




Supplementary Figure 7. TEM images of reaction mixture upon grinding with 1,5-cod for 99 min. Scale bars in TEM images (bottom left) indicate 10 nm.




Supplementary Figure 8. TEM images of reaction mixture upon grinding with cyclooctane for 99 min. Scale bars in TEM images (bottom left) indicate 20 nm.



Supplementary Figure 9. TEM images of reaction mixture upon grinding without additives for 99 min. Scale bars in TEM images (bottom left) indicate 20 nm.




Supplementary Figure 10. Aggregation of palladium particles as a function of time in the reaction mixture containing cyclooctane. Scale bars in TEM images (bottom left) indicate 20 nm.



Supplementary Figure 11. Aggregation of palladium particles as a function of time in the reaction mixture without additives. Scale bars in TEM images (bottom left) indicate 20 nm.


N,N-Diphenylnaphthalen-1-amine (**3a**)



The reaction was carried out with 207.1 mg (1.0 mmol) of **1a** and 169.2 mg (1.0 mmol) of **2a**. The product **3a** was obtained as a white powder (280.6 mg, 0.95 mmol, 95% yield). ¹H and ¹³C NMR were in agreement with the literature.⁵

¹H NMR (392 MHz, CDCl₃, δ): 6.92 (t, *J* = 7.2 Hz, 2H), 6.99–7.06 (m, 4H), 7.18 (t, *J* = 8.1 Hz, 4H), 7.30–7.39 (m, 2H), 7.45 (q, *J* = 7.2 Hz, 2H), 7.76 (d, *J* = 8.5 Hz, 1H), 7.88 (d, *J* = 8.1 Hz, 1H), 7.94 (d, *J* = 8.5 Hz, 1H). ¹³C NMR (99 MHz, CDCl₃, δ): 121.6 (CH), 121.8 (CH), 124.2 (CH), 126.1 (CH), 126.34 (CH), 126.36 (CH), 126.40 (CH), 127.2 (CH), 128.3 (CH), 129.0 (CH), 131.1 (C), 135.2 (C), 143.5 (C), 148.4 (C). HRMS-EI (*m/z*): [M]⁺ calcd for C₂₂H₁₇N, 295.1361; found, 295.1369.


N,N-Diphenylpyren-1-amine (**3b**)



The reaction was carried out with 140.6 mg (0.50 mmol) of **1b** and 84.6 mg (0.50 mmol) of **2a**. The product **3b** was obtained as a yellow powder (171.8 mg, 0.465 mmol, 93% yield). ¹H and ¹³C NMR were in agreement with the literature.⁶

¹H NMR (392 MHz, CDCl₃, δ): 6.95 (t, *J* = 7.4 Hz, 2H), 7.07 (d, *J* = 7.6 Hz, 4H), 7.20 (t, *J* = 7.9 Hz, 4H), 7.83 (d, *J* = 8.1 Hz, 1H), 7.93 (d, *J* = 9.4 Hz, 1H), 7.98 (t, *J* = 7.6 Hz, 1H), 8.06 (br, s, 2H), 8.09–8.19 (m, 4H). ¹³C NMR (99 MHz, CDCl₃, δ): 121.7 (CH), 122.0 (CH), 123.3 (CH), 124.8 (C), 125.1 (CH), 125.2 (CH), 126.0 (CH), 126.2 (CH), 126.3 (C), 127.1 (CH), 127.2 (CH), 127.6 (CH), 127.9 (CH), 128.2 (C), 129.1 (CH), 129.5 (C), 131.0 (C), 131.2 (C), 140.9 (C), 148.6 (C). HRMS-EI (*m/z*): [M]⁺ calcd for C₂₈H₁₉N, 369.1518; found, 369.1521.


N,N-Di-*p*-tolylpyren-1-amine (**3c**)



The reaction was carried out with 140.6 mg (0.50 mmol) of **1b** and 98.6 mg (0.50 mmol) of **2c**. The product **3c** was obtained as a yellow powder (174.9 mg, 0.44 mmol, 88% yield). ¹H and ¹³C NMR were in agreement with the literature.⁶

¹H NMR (392 MHz, CDCl₃, δ): 2.28 (s, 6H), 6.92–7.03 (m, 8H), 7.80 (d, *J* = 8.1 Hz, 1H), 7.90–7.99 (m, 2H), 8.03 (s, 2H), 8.09 (d, *J* = 7.2 Hz, 1H), 8.11–8.18 (m, 3H). ¹³C NMR (99 MHz, CDCl₃, δ): 20.7 (CH₃), 122.1 (CH), 123.5 (CH), 124.8 (C), 124.9 (CH), 125.0 (CH), 125.9 (CH), 126.1 (CH), 126.3 (C), 126.8 (CH), 127.2 (CH), 127.3 (CH), 127.6 (CH), 127.9 (C), 129.2 (C), 129.7 (CH), 130.98 (C), 131.04 (C), 131.2 (C), 141.4 (C), 146.5 (C). HRMS-EI (*m/z*): [M]⁺ calcd for C₃₀H₂₃N, 397.1831; found, 397.1829.


N,N-Bis(4-methoxyphenyl)pyren-1-amine (**3d**)



The reaction was carried out with 140.6 mg (0.50 mmol) of **1b** and 114.7 mg (0.50 mmol) of **2d**. The product **3d** was obtained as a yellow powder (146.0 mg, 0.34 mmol, 68% yield). ¹H and ¹³C NMR were in agreement with the literature.⁷

¹H NMR (392 MHz, CDCl₃, δ): 3.76 (s, 6H), 6.73–6.80 (m, 4H), 6.97 (d, *J* = 9.0 Hz, 4H), 7.75 (d, *J* = 8.1 Hz, 1H), 7.91 (d, *J* = 9.4 Hz, 1H), 7.96 (t, *J* = 7.6 Hz, 1H), 8.02 (brs, 2H), 8.05–8.18 (m, 4H). ¹³C NMR (99 MHz, CDCl₃, δ): 55.4 (CH₃), 114.5 (CH), 123.5 (CH), 123.6 (CH), 124.80 (CH), 124.84 (C), 124.9 (CH), 125.8 (CH), 126.1 (CH), 126.3 (C), 126.6 (CH), 126.7 (CH), 127.2 (CH), 127.28 (C), 127.35 (CH), 128.8 (C), 131.0 (C), 131.3 (C), 142.0 (C), 143.0 (C), 154.5 (C). HRMS-EI (*m/z*): [M]⁺ calcd for C₃₀H₂₃NO₂, 429.1729; found, 429.1715.

N,N-Di(naphthalen-2-yl)pyren-1-amine (**3e**)




3e

The reaction was carried out with 140.6 mg (0.50 mmol) of **1b** and 134.7 mg (0.50 mmol) of **2e**. The product **3e** was obtained as a yellow powder (169.1 mg, 0.36 mmol, 72% yield).

^1H NMR (392 MHz, CDCl_3 , δ): 7.29–7.41 (m, 8H), 7.45–7.50 (m, 2H), 7.69–7.78 (m, 4H), 7.88 (d, $J = 3.6$ Hz, 1H), 7.90 (d, $J = 2.7$ Hz, 1H), 7.98 (t, $J = 7.6$ Hz, 1H), 8.07–8.11 (m, 3H), 8.16–8.23 (m, 3H). ^{13}C NMR (99 MHz, CDCl_3 , δ): 118.3 (CH), 123.1 (CH), 123.2 (CH), 124.2 (CH), 124.8 (C), 125.1 (CH), 125.2 (CH), 126.0 (CH), 126.2 (CH), 126.3 (CH), 126.4 (C), 126.9 (CH), 127.2 (CH), 127.5 (CH), 127.6 (CH), 127.98 (C), 128.03 (CH), 129.0 (CH), 129.6 (C), 129.7 (C), 131.0 (C), 131.2 (C), 134.4 (C), 140.8 (C), 146.3 (C). HRMS-EI (m/z): $[\text{M}]^+$ calcd for $\text{C}_{36}\text{H}_{23}\text{N}$, 469.1831; found, 469.1821.

N-(Naphthalen-1-yl)-*N*-phenylpyren-1-amine (**3f**)




3f

The reaction was carried out with 140.6 mg (0.50 mmol) of **1b** and 109.6 mg (0.50 mmol) of **2f**. The product **3f** was obtained as a yellow powder (169.9 mg, 0.405 mmol, 81% yield).

^1H NMR (392 MHz, CDCl_3 , δ): 6.77 (d, $J = 7.6$ Hz, 2H), 6.89 (t, $J = 7.4$ Hz, 1H), 7.13 (t, $J = 7.6$ Hz, 2H), 7.23–7.27 (m, 1H), 7.29–7.39 (m, 2H), 7.43–7.49 (m, 1H), 7.71 (d, $J = 8.1$ Hz, 1H), 7.77 (d, $J = 8.1$ Hz, 1H), 7.89 (d, $J = 4.9$ Hz, 1H), 7.91 (d, $J = 5.8$ Hz, 1H), 7.97 (t, $J = 7.6$ Hz, 1H), 8.02 (brs, 2H), 8.03–8.17 (m, 4H), 8.25 (d, $J = 9.4$ Hz, 1H). ^{13}C NMR (99 MHz, CDCl_3 , δ): 120.3 (CH), 120.7 (CH), 123.4 (CH), 124.5 (CH), 124.86 (C), 124.93 (CH), 125.1 (CH), 125.5 (CH), 125.6 (CH), 125.7 (CH), 125.8 (CH), 126.0 (CH), 126.1 (CH), 126.3 (CH), 126.4 (C), 126.7 (CH), 126.8 (C), 127.2 (CH), 127.7 (CH), 128.5 (CH), 128.9 (C), 129.0 (CH), 130.2 (C), 131.0 (C), 131.3 (C), 135.2 (C), 142.4 (C), 144.9 (C), 150.7 (C). HRMS-EI (m/z): $[\text{M}]^+$ calcd for $\text{C}_{32}\text{H}_{21}\text{N}$, 419.1674; found, 419.1663.


N,N-Diphenylpyren-4-amine (**3h**)



The reaction was carried out with 140.6 mg (0.50 mmol) of **1h** and 84.6 mg (0.50 mmol) of **2a**. The product **3h** was obtained as a yellow powder (173.6 mg, 0.47 mmol, 94% yield).

^1H NMR (392 MHz, CDCl_3 , δ): 6.91–6.98 (m, 2H), 7.04–7.10 (m, 4H), 7.20 (t, $J = 8.1$ Hz, 4H), 7.83 (d, $J = 8.5$ Hz, 1H), 7.93 (d, $J = 9.0$ Hz, 1H), 7.98 (t, $J = 7.6$ Hz, 1H), 8.06 (brs, 2H), 8.09–8.19 (m, 4H). ^{13}C NMR (100 MHz, CDCl_3 , δ): 121.7 (CH), 122.0 (CH), 123.3 (CH), 124.8 (C), 125.0 (CH), 125.1 (CH), 126.0 (CH), 126.1 (CH), 126.3 (C), 127.05 (CH), 127.15 (CH), 127.6 (CH), 127.9 (CH), 128.1 (C), 129.1 (CH), 129.5 (C), 131.0 (C), 131.2 (C), 140.8 (C), 148.6 (C). HRMS-EI (m/z): $[\text{M}]^+$ calcd for $\text{C}_{28}\text{H}_{19}\text{N}$, 369.1518; found, 369.1506.


7-(*tert*-Butyl)-*N,N*-diphenylpyren-2-amine (**3i**)



The reaction was carried out with 168.2 mg (0.50 mmol) of **1i** and 85.4 mg (0.50 mmol) of **2a**. The product **3i** was obtained as a yellow powder (147.9 mg, 0.35 mmol, 70% yield).

^1H NMR (401 MHz, CDCl_3 , δ): 1.57 (s, 9H), 7.03–7.09 (m, 2H), 7.17–7.22 (m, 4H), 7.26–7.32 (m, 4H), 7.82 (d, $J = 8.4$ Hz, 2H), 7.86 (s, 2H), 7.97 (d, $J = 8.4$ Hz, 2H), 8.16 (s, 2H). ^{13}C NMR (99 MHz, CDCl_3 , δ): 32.1 (CH_3), 35.3 (C), 120.7 (CH), 121.4 (C), 122.6 (CH), 122.9 (CH), 123.0 (C), 124.4 (CH), 126.9 (CH), 128.1 (CH), 129.5 (CH), 130.5 (C), 132.3 (C), 145.7 (C), 148.4 (C), 148.5 (C). HRMS-EI (m/z): $[\text{M}]^+$ calcd for $\text{C}_{32}\text{H}_{27}\text{N}$, 425.2144; found, 425.2146.

N,N-Diphenylnaphthalen-2-amine (**3j**)




The reaction was carried out with 103.4 mg (0.50 mmol) of **1j** and 85.4 mg (0.50 mmol) of **2a**. The

product **3j** was obtained as a white powder (132.3 mg, 0.45 mmol, 90% yield). ^1H and ^{13}C NMR were in agreement with the literature.⁸

^1H NMR (401 MHz, CDCl_3 , δ): 7.01–7.07 (m, 2H), 7.11–7.16 (m, 4H), 7.23–7.30 (m, 5H), 7.31–7.41 (m, 2H), 7.42 (d, $J = 2.4$ Hz, 1H), 7.58 (d, $J = 8.0$ Hz, 1H), 7.71 (d, $J = 8.8$ Hz, 1H), 7.75 (d, $J = 8.4$ Hz, 1H). ^{13}C NMR (100 MHz, CDCl_3 , δ): 120.3 (CH), 123.0 (CH), 124.52 (CH), 124.56 (CH), 124.6 (CH), 126.4 (CH), 127.0 (CH), 127.7 (CH), 129.0 (CH), 129.4 (CH), 130.1 (C), 134.5 (C), 145.6 (C), 147.9 (C). HRMS-ESI (m/z): $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{22}\text{H}_{18}\text{N}$, 296.1434; found, 296.1437.

N,N-Diphenylphenanthren-9-amine (**3k**)




3k

The reaction was carried out with 128.6 mg (0.50 mmol) of **1k** and 84.6 mg (0.50 mmol) of **2a**. The product **3k** was obtained as a white powder (138.2 mg, 0.40 mmol, 80% yield).

^1H NMR (392 MHz, CDCl_3 , δ): 6.91–6.97 (m, 2H), 7.06–7.12 (m, 4H), 7.15–7.23 (m, 4H), 7.44–7.49 (m, 1H), 7.53–7.58 (m, 1H), 7.60 (brs, 1H), 7.61–7.66 (m, 2H), 7.75 (dd, $J = 1.3, 8.1$ Hz, 1H), 8.04 (d, $J = 8.1$ Hz, 1H), 8.69 (d, $J = 8.1$ Hz, 1H), 8.73 (d, $J = 8.1$ Hz, 1H). ^{13}C NMR (99 MHz, CDCl_3 , δ): 121.7 (CH), 121.8 (CH), 122.5 (CH), 123.0 (CH), 125.0 (CH), 126.5 (CH), 126.77 (CH), 126.81 (CH), 126.9 (CH), 127.6 (CH), 128.2 (CH), 129.1 (CH), 129.3 (C), 130.4 (C), 132.1 (C), 132.4 (C), 142.0 (C), 148.2 (C). HRMS-EI (m/z): $[\text{M}]^+$ calcd for $\text{C}_{26}\text{H}_{19}\text{N}$, 345.1518; found, 345.1501.

N,N-Diphenylanthracen-9-amine (**3l**)




3l

The reaction was carried out with 128.6 mg (0.50 mmol) of **1l** and 84.6 mg (0.50 mmol) of **2a**. The product **3l** was obtained as a yellow powder (122.6 mg, 0.355 mmol, 71% yield). ^1H and ^{13}C NMR were in agreement with the literature.⁹

^1H NMR (392 MHz, CDCl_3 , δ): 6.86 (dt, $J = 1.2, 7.3$ Hz, 2H), 7.03–7.18 (m, 8H), 7.34–7.47 (m, 4H), 8.05 (d, $J = 8.5$ Hz, 2H), 8.11 (d, $J = 8.5$ Hz, 2H), 8.50 (brs, 1H). ^{13}C NMR (100 MHz, CDCl_3 , δ): 120.2 (CH), 121.1 (CH), 124.3 (CH), 125.5 (CH), 126.7 (CH), 126.8 (CH), 128.8 (CH), 129.1 (CH), 130.6 (C), 132.7 (C), 137.2 (C), 147.7 (C). HRMS-EI (m/z): $[\text{M}]^+$ calcd for $\text{C}_{26}\text{H}_{19}\text{N}$, 345.1518; found,

45.1512.


***N,N*,10-Triphenylanthracen-9-amine (3m)**



The reaction was carried out with 166.6 mg (0.50 mmol) of **1m** and 87.3 mg (0.52 mmol) of **2a**. The product **3m** was obtained as a yellow powder (116.7 mg, 0.275 mmol, 55% yield).

^1H NMR (401 MHz, CDCl_3 , δ): 6.86–6.92 (m, 2H), 7.10–7.21 (m, 8H), 7.29–7.42 (m, 4H), 7.46–7.51 (m, 2H), 7.51–7.64 (m, 3H), 7.71 (d, $J = 8.8$ Hz, 2H), 8.19 (d, $J = 8.8$ Hz, 2H). ^{13}C NMR (99 MHz, CDCl_3 , δ): 120.4 (CH), 121.3 (CH), 124.5 (CH), 125.5 (CH), 126.6 (CH), 127.7 (CH), 128.5 (CH), 129.3 (CH), 130.5 (C), 131.4 (CH), 131.5 (C), 137.2 (C), 137.8 (C), 138.8 (C), 147.9 (C). HRMS-EI (m/z): $[\text{M}]^+$ calcd for $\text{C}_{32}\text{H}_{23}\text{N}$, 421.1831; found, 421.1813.


***N,N*,4-Triphenylnaphthalen-1-amine (3n)**



The reaction was carried out with 141.4 mg (0.50 mmol) of **1n** and 87.4 mg (0.52 mmol) of **2a**. The product **3n** was obtained as a yellow powder (177.6 mg, 0.479 mmol, 96% yield).

^1H NMR (399 MHz, CDCl_3 , δ): 6.95 (t, $J = 7.2$ Hz, 2H), 7.08 (d, $J = 7.6$ Hz, 4H), 7.21 (t, $J = 8.2$ Hz, 4H), 7.33–7.47 (m, 5H), 7.47–7.55 (m, 4H), 7.93 (d, $J = 6.4$ Hz, 1H), 8.04 (d, $J = 8.0$ Hz, 1H). ^{13}C NMR (99 MHz, CDCl_3 , δ): 121.8 (CH), 122.0 (CH), 124.6 (CH), 126.31 (CH), 126.35 (CH), 126.8 (CH), 126.9 (CH), 127.4 (CH), 127.6 (CH), 128.4 (CH), 129.2 (CH), 130.2 (CH), 131.5 (C), 133.4 (C), 138.7 (C), 140.6 (C), 143.1 (C), 148.5 (C). HRMS-ESI (m/z): $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{28}\text{H}_{22}\text{N}$, 372.1752; found, 372.1744.


N,N-Diphenyl-(1,1'-biphenyl)-4-amine (**3o**)



The reaction was carried out with 116.4 mg (0.50 mmol) of **1o** and 84.5 mg (0.50 mmol) of **2a**. The product **3o** was obtained as a white powder (142.7 mg, 0.445 mmol, 89% yield). ¹H and ¹³C NMR were in agreement with the literature.¹⁰

¹H NMR (401 MHz, CDCl₃, δ): 7.03 (td, *J* = 1.2, 7.2 Hz, 2H), 7.13 (d, *J* = 8.0 Hz, 6H), 7.23–7.27 (m, 3H), 7.28–7.33 (m, 2H), 7.42 (t, *J* = 7.6 Hz, 2H), 7.45–7.50 (m, 2H), 7.57 (d, *J* = 8.0 Hz, 2H). ¹³C NMR (99 MHz, CDCl₃, δ): 123.0 (CH), 124.0 (CH), 124.5 (CH), 126.7 (CH), 126.9 (CH), 127.9 (CH), 128.8 (CH), 129.4 (CH), 135.2 (C), 140.7 (C), 147.2 (C), 147.7 (C). HRMS-ESI (*m/z*): [M+H]⁺ calcd for C₂₄H₂₀N, 322.1596; found, 322.1581.


N,N-Diphenyl-(1,1':4',1''-terphenyl)-4-amine (**3p**)



The reaction was carried out with 154.4 mg (0.50 mmol) of **1p** and 86.2 mg (0.51 mmol) of **2a**. The product **3p** was obtained as a white powder (160.3 mg, 0.403 mmol, 81% yield).

¹H NMR (401 MHz, CDCl₃, δ): 7.01–7.07 (m, 2H), 7.12–7.18 (m, 6H), 7.24–7.31 (m, 4H), 7.33–7.38 (m, 1H), 7.43–7.48 (m, 2H), 7.50–7.55 (m, 2H), 7.62–7.67 (m, 6H). ¹³C NMR (99 MHz, CDCl₃, δ): 123.1 (CH), 124.0 (CH), 124.6 (CH), 127.1 (CH), 127.4 (CH), 127.6 (CH), 127.8 (CH), 128.9 (CH), 129.4 (CH), 134.6 (C), 139.7 (C), 140.8 (C), 147.4 (C), 147.8 (C). HRMS-EI (*m/z*): [M]⁺ calcd for C₃₀H₂₃N, 397.1831; found, 397.1818.


N,N-Diphenyl-1,2-dihydroacenaphthyl-5-amine (**3q**)



The reaction was carried out with 119.0 mg (0.51 mmol) of **1q** and 85.5 mg (0.505 mmol) of **2a**. The product **3q** was obtained as a white powder (121.5 mg, 0.378 mmol, 75% yield).

^1H NMR (399 MHz, CDCl_3 , δ): 3.33–3.50 (m, 4H), 6.91 (t, $J = 7.2$ Hz, 2H), 7.04 (d, $J = 8.4$ Hz, 4H), 7.18 (t, $J = 7.6$ Hz, 4H), 7.22–7.32 (m, 4H), 7.41 (d, $J = 7.6$ Hz, 1H). ^{13}C NMR (99 MHz, CDCl_3 , δ): 30.1 (CH_2), 30.9 (CH_2), 119.6 (CH), 119.8 (CH), 119.9 (CH), 121.5 (CH), 121.9 (CH), 128.1 (CH), 128.4 (CH), 129.1 (CH), 129.4 (C), 139.7 (C), 141.1 (C), 144.2 (C), 146.4 (C), 148.5 (C). HRMS-ESI (m/z): $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{24}\text{H}_{20}\text{N}$, 322.1590; found, 322.1596.


N,N-Diphenylfluoranthene-3-amine (**3r**)



The reaction was carried out with 141.7 mg (0.50 mmol) of **1r** and 85.3 mg (0.50 mmol) of **2a**. The product **3r** was obtained as a yellow powder (185.7 mg, 0.378 mmol, 99% yield).

^1H NMR (396 MHz, CDCl_3 , δ): 6.94–7.01 (m, 2H), 7.07–7.13 (m, 4H), 7.18–7.26 (m, 4H), 7.28–7.42 (m, 4H), 7.62 (d, $J = 8.7$ Hz, 1H), 7.80–7.88 (m, 4H). ^{13}C NMR (99 MHz, CDCl_3 , δ): 120.2 (CH), 121.2 (CH), 121.3 (CH), 121.6 (CH), 122.3 (CH), 123.0 (CH), 124.5 (CH), 127.2 (CH), 127.3 (CH), 127.7 (CH), 127.79 (CH), 127.83 (C), 129.3 (CH), 134.2 (C), 134.3 (C), 137.4 (C), 139.1 (C), 139.6 (C), 144.8 (C), 149.0 (C). HRMS-ESI (m/z): $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{28}\text{H}_{20}\text{N}$, 370.1596; found, 370.1592.

N,N-Diphenyltriphenylene-2-amine (**3s**)




The reaction was carried out with 153.6 mg (0.50 mmol) of **1s** and 84.6 mg (0.50 mmol) of **2a**. The product **3s** was obtained as a white powder (168.1 mg, 0.425 mmol, 85% yield).

^1H NMR (392 MHz, CDCl_3 , δ): 7.08 (t, $J = 7.4$ Hz, 2H), 7.18–7.24 (m, 4H), 7.27–7.34 (m, 4H), 7.39 (dd, $J = 2.3, 8.5$ Hz, 1H), 7.53 (dt, $J = 1.3, 8.6$ Hz, 1H), 7.57–7.67 (m, 3H), 8.29 (d, $J = 7.6$ Hz, 1H), 8.33 (d, $J = 2.2$ Hz, 1H), 8.48–8.56 (m, 2H), 8.59–8.64 (m, 2H). ^{13}C NMR (99 MHz, CDCl_3 , δ): 117.1 (CH), 122.9 (CH), 123.1 (CH), 123.2 (CH), 123.4 (CH), 123.8 (CH), 124.4 (CH), 125.0 (C), 126.4 (CH), 127.0 (CH), 127.2 (CH), 129.0 (C), 129.3 (C), 129.4 (CH), 129.7 (C), 130.0 (C), 130.9 (C),

146.9 (C), 147.7 (C). HRMS-EI (m/z): $[M]^+$ calcd for $C_{30}H_{21}N$, 395.1674; found, 395.1661.

9,9-Dimethyl-*N,N*-diphenyl-9*H*-fluoren-2-amine (**3t**)




3t

The reaction was carried out with 137.1 mg (0.50 mmol) of **1t** and 85.0 mg (0.50 mmol) of **2a**. The product **3t** was obtained as a white powder (163.9 mg, 0.45 mmol, 90% yield).

1H NMR (401 MHz, $CDCl_3$, δ): 1.41 (s, 6H), 7.01 (t, $J = 7.2$ Hz, 3H), 7.13 (d, $J = 8.4$ Hz, 4H), 7.19 (d, $J = 2.0$ Hz, 1H), 7.22–7.33 (m, 6H), 7.38 (d, $J = 7.2$ Hz, 1H), 7.57 (d, $J = 8.0$ Hz, 1H), 7.63 (d, $J = 7.2$ Hz, 1H). ^{13}C NMR (99 MHz, $CDCl_3$, δ): 27.2 (CH₃), 46.9 (C), 118.7 (CH), 119.5 (CH), 120.7 (CH), 122.6 (CH), 122.7 (CH), 123.4 (CH), 124.2 (CH), 126.6 (CH), 127.1 (CH), 129.3 (CH), 134.2 (C), 139.1 (C), 147.4 (C), 148.1 (C), 153.6 (C), 155.1 (C). HRMS-EI (m/z): $[M]^+$ calcd for $C_{27}H_{23}N$, 361.1831; found, 361.1825.

(*E*)-*N,N*-Diphenyl-4-styrylaniline (**3u**)




3u

The reaction was carried out with 129.6 mg (0.50 mmol) of **1u** and 84.6 mg (0.50 mmol) of **2a**. The product **3u** was obtained as a white powder (165.0 mg, 0.45 mmol, 95% yield). 1H and ^{13}C NMR were in agreement with the literature.¹¹

1H NMR (392 MHz, $CDCl_3$, δ): 7.01–7.08 (m, 5H), 7.09–7.14 (m, 4H), 7.23–7.30 (m, 6H), 7.32–7.41 (m, 4H), 7.50 (d, $J = 7.6$ Hz, 2H). ^{13}C NMR (99 MHz, $CDCl_3$, δ): 123.0 (CH), 123.6 (CH), 124.4 (CH), 126.3 (CH), 127.0 (CH), 127.2 (CH), 127.3 (CH), 128.1 (CH), 128.6 (CH), 129.2 (CH), 131.4 (C), 137.6 (C), 147.3 (C), 147.5 (C). HRMS-EI (m/z): $[M]^+$ calcd for $C_{26}H_{21}N$, 347.1674; found, 347.1670.


N,N-Diphenyl-4-(phenylethynyl)aniline (**3v**)



The reaction was carried out with 128.6 mg (0.50 mmol) of **1v** and 84.6 mg (0.50 mmol) of **2a**. The product **3v** was obtained as a white powder (158.9 mg, 0.46 mmol, 92% yield). ¹H and ¹³C NMR were in agreement with the literature.¹²

¹H NMR (392 MHz, CDCl₃, δ): 6.98–7.03 (m, 2H), 7.04–7.08 (m, 2H), 7.09–7.15 (m, 4H), 7.24–7.40 (m, 9H), 7.48–7.53 (m, 2H). ¹³C NMR (99 MHz, CDCl₃, δ): 88.6 (C), 89.6 (C), 116.0 (C), 122.3 (CH), 123.46 (CH), 123.52 (C), 124.9 (CH), 127.9 (CH), 128.2 (CH), 129.3 (CH), 131.4 (CH), 132.5 (CH), 147.1 (C), 147.8 (C). HRMS-EI (*m/z*): [M]⁺ calcd for C₂₆H₁₉N, 345.1518; found, 345.1508.


(*E*)-*N,N*-Diphenyl-4-styrylaniline (**3w**)



The reaction was carried out with 70.8 mg (0.30 mmol) of **1w** and 101.5 mg (0.60 mmol) of **2a**. The product **3w** was obtained as a white powder (112.6 mg, 0.273 mmol, 91% yield). ¹H and ¹³C NMR were in agreement with the literature.¹³

¹H NMR (392 MHz, CDCl₃, δ): 6.90–7.04 (m, 8H), 7.10 (d, *J* = 7.2 Hz, 8H), 7.19–7.31 (m, 8H). ¹³C NMR (99 MHz, CDCl₃, δ): 122.4 (CH), 123.7 (CH), 125.4 (CH), 129.1 (CH), 142.8 (C), 147.8 (C). HRMS-EI (*m/z*): [M]⁺ calcd for C₃₀H₂₄N₂, 412.1940; found, 412.1929.


N,N-Diphenyl-4-(1,2,2-triphenylvinyl)aniline (**3x**)



The reaction was carried out with 205.7 mg (0.50 mmol) of **1x** and 84.6 mg (0.50 mmol) of **2a**. The product **3x** was obtained as a white powder (167.4 mg, 0.335 mmol, 67% yield). ¹H and ¹³C NMR were in agreement with the literature.¹⁴

¹H NMR (392 MHz, CDCl₃, δ): 6.78 (d, *J* = 8.5 Hz, 2H), 6.85 (d, *J* = 8.5 Hz, 2H), 6.94–7.17 (m, 21H), 7.21 (t, *J* = 8.1 Hz, 4H). ¹³C NMR (99 MHz, CDCl₃, δ): 122.6 (CH), 122.7 (CH), 124.2 (CH), 126.25 (CH), 126.30 (CH), 126.4 (CH), 127.55 (CH), 127.57 (CH), 127.58 (CH), 129.1 (CH), 131.3 (CH), 132.1 (CH), 137.9 (C), 140.4 (C), 140.7 (C), 143.5 (C), 143.8 (C), 144.0 (C), 145.9 (C), 147.5 (C). HRMS-EI (*m/z*): [M]⁺ calcd for C₃₈H₂₉N, 499.2300; found, 499.2290.


N,N-Bis(4-methoxyphenyl)-9,10-di(naphthalen-2-yl)anthracen-2-amine (**3y**)



The reaction was carried out with 254.7 mg (0.50 mmol) of **1y** and 114.7 mg (0.50 mmol) of **2d**. The product **3y** was obtained as an orange powder (246.7 mg, 0.375 mmol, 75% yield).

¹H NMR (392 MHz, CDCl₃, δ): 3.67 (s, 6H), 6.65–6.70 (m, 4H), 6.94–7.00 (m, 5H), 7.08 (dd, *J* = 2.2, 9.4 Hz, 1H), 7.18–7.28 (m, 2H), 7.48 (d, *J* = 8.5 Hz, 1H), 7.50–7.67 (m, 7H), 7.71 (d, *J* = 8.1 Hz, 1H), 7.79–7.85 (m, 2H), 7.86–7.96 (m, 3H), 7.97–8.03 (m, 2H), 8.06 (d, *J* = 8.5 Hz, 1H). ¹³C NMR (99 MHz, CDCl₃, δ): 55.2 (CH₃), 113.0 (CH), 114.3 (CH), 122.5 (CH), 123.9 (CH), 125.2 (CH), 125.7 (CH), 125.9 (CH), 126.1 (CH), 126.3 (CH), 126.4 (CH), 126.6 (C), 127.1 (CH), 127.6 (CH), 127.78 (CH), 127.83 (CH), 127.9 (CH), 128.0 (CH), 128.8 (C), 129.4 (CH), 129.6 (CH), 130.0 (CH), 130.2 (CH), 130.4 (C), 131.2 (C), 132.4 (C), 132.6 (C), 133.2 (C), 133.3 (C), 134.1 (C), 136.3 (C), 136.6 (C), 136.7 (C), 140.1 (C), 145.3 (C), 155.8 (C). HRMS-ESI (*m/z*): [M+Na]⁺ calcd for C₄₈H₃₅O₂NNa, 680.2560; found, 680.2572.


***N,N*-Bis(4-methoxyphenyl)-3',4',5',6'-tetraphenyl-(1,1':2',1''-terphenyl)-4-amine (3z)**



The reaction was carried out with 306.8 mg (0.50 mmol) of **1z** and 114.7 mg (0.50 mmol) of **2d**. The product **3z** was obtained as a white powder (156.2 mg, 0.205 mmol, 41% yield).

^1H NMR (392 MHz, CDCl_3 , δ): 3.75 (s, 6H), 6.48 (d, $J = 8.1$ Hz, 2H), 6.61 (d, $J = 6.6$ Hz, 2H), 6.69–6.74 (m, 4H), 6.75–6.79 (m, 4H), 6.80–6.95 (m, 25H). ^{13}C NMR (99 MHz, CDCl_3 , δ): 55.4 (CH_3), 114.3 (CH), 120.9 (CH), 125.0 (CH), 125.1 (CH), 125.3 (CH), 126.5 (CH), 131.4 (CH), 131.5 (CH), 132.0 (CH), 134.0 (C), 140.05 (C), 140.13 (C), 140.3 (C), 140.4 (C), 140.6 (C), 140.8 (C), 141.4 (C), 145.6 (C), 155.0 (C). HRMS-ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{56}\text{H}_{43}\text{O}_2\text{NNa}$, 784.3186; found, 784.3196.


10,20-Bis(3,5-dimethylphenyl)-*N5,N5,N15,N15*-tetrakis(4-methoxyphenyl)porphyrin-5,15-diamine (3aa)



The reaction was carried out with 135.4 mg (0.20 mmol) of **1aa** and 114.8 mg (0.50 mmol) of **2d**. The product **3aa** was obtained as a dark purple powder (107.0 mg, 0.11 mmol, 55% yield).

^1H NMR (392 MHz, CDCl_3 , δ): 2.54 (s, 12H), 3.71 (s, 12H), 6.73 (d, $J = 8.5$ Hz, 9H), 7.16–7.27 (m, 11H), 7.71 (s, 4H), 8.66 (d, $J = 4.5$ Hz, 4H), 9.13 (d, $J = 4.9$ Hz, 4H). ^{13}C NMR (99 MHz, CDCl_3 , δ): 21.4 (CH_3), 55.4 (CH_3), 114.4 (CH), 120.3 (C), 122.6 (C), 123.2 (CH), 129.3 (CH), 132.5 (CH), 135.9 (C), 141.1 (C), 146.8 (C), 153.5 (C). HRMS-ESI (m/z): $[\text{M}]^+$ calcd for $\text{C}_{64}\text{H}_{57}\text{O}_4\text{N}_6$, 973.4436; found, 973.4460.


10-(Dimesitylboraneyl)-*N,N*-diphenylanthracen-9-amine (**3ab**)



The reaction was carried out with 252.7 mg (0.50 mmol) of **1ab** and 84.5 mg (0.50 mmol) of **2a**. The product **3ab** was obtained as a yellow powder (184.0 mg, 0.31 mmol, 62% yield).

^1H NMR (392 MHz, CDCl_3 , δ): 1.68 (brs, 6H), 2.19 (brs, 6H), 2.28 (s, 6H), 6.66 (brs, 2H), 6.87 (t, $J = 7.2$ Hz, 4H), 7.06–7.11 (m, 4H), 7.12–7.21 (m, 6H), 7.23–7.31 (m, 2H), 8.08 (d, $J = 8.5$ Hz, 2H), 8.13 (d, $J = 8.5$ Hz, 2H). ^{13}C NMR (99 MHz, CDCl_3 , δ): 21.3 (CH_3), 23.6 (br, CH_3), 120.0 (CH), 121.1 (CH), 124.9 (CH), 125.1 (CH), 126.2 (CH), 129.0 (CH), 129.2 (CH), 130.6 (C), 135.0 (C), 139.3 (C), 139.9 (C), 145.3 (br, C-B), 147.5 (C). HRMS-EI (m/z): $[\text{M}]^+$ calcd for $\text{C}_{44}\text{H}_{40}\text{BN}$, 592.3290; found, 592.3283.


N,N,9-Triphenyl-9H-carbazol-3-amine (**3ac**)



The reaction was carried out with 161.1 mg (0.50 mmol) of **1ac** and 84.6 mg (0.50 mmol) of **2a**. The product **3ac** was obtained as a yellow powder (168.3 mg, 0.41 mmol, 82% yield).

^1H NMR (392 MHz, CDCl_3 , δ): 6.95 (t, $J = 7.2$ Hz, 2H), 7.12 (d, $J = 8.1$ Hz, 4H), 7.18–7.28 (m, 6H), 7.34 (d, $J = 9.0$ Hz, 1H), 7.37–7.42 (m, 2H), 7.43–7.49 (m, 1H), 7.55–7.64 (m, 4H), 7.93 (brs, 1H), 7.99 (d, $J = 8.1$ Hz, 1H). ^{13}C NMR (99 MHz, CDCl_3 , δ): 109.9 (CH), 110.7 (CH), 118.5 (CH), 119.9 (CH), 120.5 (CH), 121.6 (CH), 122.7 (CH), 123.0 (C), 124.3 (C), 125.7 (CH), 126.1 (CH), 127.0 (CH), 127.4 (CH), 129.0 (CH), 129.9 (CH), 137.6 (C), 138.0 (C), 140.5 (C), 141.3 (C), 148.6 (C). HRMS-EI (m/z): $[\text{M}]^+$ calcd for $\text{C}_{30}\text{H}_{22}\text{N}_2$, 410.1783; found, 410.1783.

***N,N*,5-Triphenylthiophen-2-amine (3ad)**




3ad

The reaction was carried out with 119.6 mg (0.50 mmol) of **1ad** and 84.6 mg (0.50 mmol) of **2a**. The product **3ad** was obtained as a white powder (90.0 mg, 0.275 mmol, 55% yield). ¹H and ¹³C NMR were in agreement with the literature.¹⁵

¹H NMR (392 MHz, CDCl₃, δ): 6.67 (d, *J* = 4.0 Hz, 1H), 7.04 (t, *J* = 7.2 Hz, 2H), 7.11 (d, *J* = 3.6 Hz, 1H), 7.15–7.21 (m, 4H), 7.22–7.37 (m, 7H), 7.52 (d, *J* = 7.6 Hz, 2H). ¹³C NMR (99 MHz, CDCl₃, δ): 121.6 (CH), 121.8 (CH₃), 122.6 (CH), 123.0 (CH), 125.1 (CH), 127.0 (CH), 128.8 (CH), 129.2 (CH), 134.4 (C), 138.1 (C), 147.7 (C), 150.8 (C). HRMS-EI (*m/z*): [M]⁺ calcd for C₂₂H₁₇NS, 327.1082; found, 327.1078.

2-(Diphenylamino)anthracene-9,10-dione (3ae)




3ae

The reaction was carried out with 143.6 mg (0.50 mmol) of **1ae** and 84.6 mg (0.50 mmol) of **2a**. The product **3ae** was obtained as an orange powder (107.0 mg, 0.285 mmol, 57% yield).

¹H NMR (392 MHz, CDCl₃, δ): 7.16–7.23 (m, 6H), 7.26 (dd, *J* = 2.2, 8.5 Hz, 1H), 7.33–7.41 (m, 4H), 7.70–7.80 (m, 3H), 8.12 (d, *J* = 8.5 Hz, 1H), 8.22 (dd, *J* = 1.8, 7.2 Hz, 1H), 8.27–8.31 (m, 1H). ¹³C NMR (99 MHz, CDCl₃, δ): 116.5 (CH), 123.8 (CH), 125.5 (CH), 125.8 (C), 126.3 (CH), 127.0 (CH), 129.2 (CH), 129.9 (CH), 133.4 (CH), 134.6 (C), 134.0 (CH), 134.9 (C), 145.8 (C), 153.1 (C), 181.7 (C), 183.5 (C). HRMS-EI (*m/z*): [M]⁺ calcd for C₂₆H₁₇NO₂, 375.1259; found, 375.1256.


***N*1,*N*1,*N*3,*N*3,*N*6,*N*6,*N*8,*N*8-octakis(4-methoxyphenyl)pyrene-1,3,6,8-tetraamine (**3af**)**




3af

The reaction was carried out with 258.9 mg (0.50 mmol) of **1af** and 596.2 mg (2.6 mmol) of **2d**. The product **3af** was obtained as an orange powder (494.5 mg, 0.445 mmol, 89% yield). ¹H and ¹³C NMR were in agreement with the literature.¹⁶


¹H NMR (392 MHz, CDCl₃, δ): 3.737 (s, 12H), 3.740 (s, 12H), 6.66–6.75 (m, 16H), 6.84–6.94 (m, 16H), 7.45 (s, 1H), 7.46 (s, 1H), 7.90 (s, 2H), 7.91 (s, 2H). ¹³C NMR (99 MHz, CDCl₃, δ): 55.4 (CH₃), 114.4 (CH), 122.8 (CH), 123.1 (CH), 126.1 (CH), 128.4 (C), 128.9 (C), 142.4 (C), 142.5 (C), 154.4 (C). HRMS-ESI (*m/z*): [M+Na]⁺ calcd for C₇₂H₆₂O₈N₄Na, 1133.4460; found, 1133.4471.




Supplementary Figure 12. ¹H NMR spectrum of 3a.




Supplementary Figure 13. ¹³C NMR spectrum of 3a.




Supplementary Figure 14. ¹H NMR spectrum of **3b**.




Supplementary Figure 15. ¹³C NMR spectrum of **3b**.




Supplementary Figure 16. ¹H NMR spectrum of 3c.




Supplementary Figure 17. ¹³C NMR spectrum of 3c.




Supplementary Figure 18. ¹H NMR spectrum of 3d.




Supplementary Figure 19. ¹³C NMR spectrum of 3d.




Supplementary Figure 20. ¹H NMR spectrum of 3e.




Supplementary Figure 21. ¹³C NMR spectrum of 3e.




Supplementary Figure 22. ¹H NMR spectrum of 3f.




Supplementary Figure 23. ¹³C NMR spectrum of 3f.




Supplementary Figure 24. ¹H NMR spectrum of 3h.




Supplementary Figure 25. ¹³C NMR spectrum of 3h.




Supplementary Figure 26. ¹H NMR spectrum of 3i.




Supplementary Figure 27. ¹³C NMR spectrum of 3i.




Supplementary Figure 28. ^1H NMR spectrum of **3j**.




Supplementary Figure 29. ^{13}C NMR spectrum of **3j**.




Supplementary Figure 30. ¹H NMR spectrum of 3k.




Supplementary Figure 31. ¹³C NMR spectrum of 3k.




Supplementary Figure 32. ¹H NMR spectrum of 31.




Supplementary Figure 33. ¹³C NMR spectrum of 31.




Supplementary Figure 34. ¹H NMR spectrum of 3m.




Supplementary Figure 35. ¹³C NMR spectrum of 3m.




Supplementary Figure 36. ¹H NMR spectrum of **3n**.




Supplementary Figure 37. ¹³C NMR spectrum of **3n**.




Supplementary Figure 38. ¹H NMR spectrum of 3o.




Supplementary Figure 39. ¹³C NMR spectrum of 3o.




Supplementary Figure 40. ¹H NMR spectrum of 3p.




Supplementary Figure 41. ¹³C NMR spectrum of 3p.




Supplementary Figure 42. ¹H NMR spectrum of 3q.




Supplementary Figure 43. ¹³C NMR spectrum of 3q.




Supplementary Figure 46. ¹H NMR spectrum of 3s.




Supplementary Figure 47. ¹³C NMR spectrum of 3s.




Supplementary Figure 48. ¹H NMR spectrum of 3t.




Supplementary Figure 49. ¹³C NMR spectrum of 3t.




Supplementary Figure 50. ¹H NMR spectrum of 3u.




Supplementary Figure 51. ¹³C NMR spectrum of 3u.




Supplementary Figure 52. ¹H NMR spectrum of 3v.




Supplementary Figure 53. ¹³C NMR spectrum of 3v.




Supplementary Figure 54. ¹H NMR spectrum of 3w.




Supplementary Figure 55. ¹³C NMR spectrum of 3w.




Supplementary Figure S6. ¹H NMR spectrum of 3x.




Supplementary Figure S7. ¹³C NMR spectrum of 3x.




Supplementary Figure 60. ¹H NMR spectrum of 3z.




Supplementary Figure 61. ¹³C NMR spectrum of 3z.




Supplementary Figure 62. ¹H NMR spectrum of 3aa.




Supplementary Figure 63. ¹³C NMR spectrum of 3aa.




Supplementary Figure 64. ¹H NMR spectrum of 3ab.




Supplementary Figure 65. ¹³C NMR spectrum of 3ab.




Supplementary Figure 66. ¹H NMR spectrum of 3ac.




Supplementary Figure 67. ¹³C NMR spectrum of 3ac.




Supplementary Figure 68. ¹H NMR spectrum of 3ad.




Supplementary Figure 69. ¹³C NMR spectrum of 3ad.




Supplementary Figure 70. ¹H NMR spectrum of 3ae.



Supplementary Figure 71. ¹³C NMR spectrum of 3ae.



Supplementary Figure 72. ¹H NMR spectrum of 3af.



Supplementary Figure 73. ¹³C NMR spectrum of 3af.

Supplementary References

1. Kübel, C., Chen, S. -L., Müllen, K. *Macromolecules*, **31**, 6014–6021. (1998)
2. Abada, Z., Ferrié, L., Akagah, B., Lormier, A. T., Figadère, B. *Tetrahedron Lett.* **52**, 3175–3178. (2011).
3. Abada, Z., Ferrié, L., Akagah, B., Lormier, A. T., Figadère, B. *Tetrahedron Lett.* **53**, 6961–6964. (2012).
4. Jia, W. -L., Bai, D. -R., McCormick, T., Liu, Q. -D., Motala, M., Wang, R. -Y., Seward, C., Tao, Y., Wang, S. *Chem. Eur. J.* **10**, 994–1006. (2004).
5. Topchiy, M. A., Asachenko, A. F., Nechaev, M. S. *Eur. J. Org. Chem.* **16**, 994–1006. (2004).
6. Hu, J. -Y., Feng, X., Seto, N., Do, J. -H., Zeng, X., Tao, Z., Yamato, T. *J. Mol. Struct.* **1035**, 19–26. (2013).
7. Nie, H. -J., Yao, C. -J., Shao, J. -Y., Yao, J., Zhong, Y. -W. *Chem. Eur. J.* **20**, 17454–17465. (2014).
8. Tadaoka, H., Yamakawa, T. *Tetrahedron Lett.* **53**, 5531–5534. (2012).
9. Xie, C., Zhang, Y. *Org. Lett.* **9**, 781–784. (2007).
10. Monguchi, Y., Kitamoto, K., Ikawa, T., Maegawa, T., Sajiki, H. *Adv. Synth. Catal.* **350**, 2767–2777. (2008).
11. Yang, J. -S., Chiou, S. -Y., Liao, K. -L. *J. Am. Chem. Soc.* **124**, 2518–2527. (2002).
12. Deol, H., Pramanik, S., Kumar, M., Khan, I., Bhalla, V. *ACS Catal.* **6**, 3771–3783. (2016).
13. Hartwig, J. F., Kawatsura, M., Hauck, S., Sheila, I., Shaughnessy, K. H., Alcazar-Roman, L. M. *J. Org. Chem.* **64**, 5575–5580. (1999).
14. Chan, C. Y. K., Lam, J. W. Y., Zhao, Z., Chen, S., Lu, P., Sung, H. H. Y., Kwok, H. S., Ma, Y., Williams, I. D., Tang, B. Z. *J. Mater. Chem.* **2**, 4320–4327. (2014).
15. Tabet, A., Hartmann, H. *Synthesis* **4**, 610–616. (2005).
16. Jeon, N. J., Lee, J., Noh, J. H., Nazeeruddin, M. K., Grätzel, M., Seok, S. *J. Am. Chem. Soc.* **135**, 19087–19090. (2013).