Title	Incidence of anterior disc displacement without reduction of the temporomandibular joint in patients with dentofacial deformity
Author(s)	Ooi, K.; Inoue, N.; Matsushita, K.; Yamaguchi, H.; Mikoya, T.; Minowa, K.; Kawashiri, S.; Nishikata, S.; Tei, K.
Citation	International journal of oral and maxillofacial surgery, 47(4), 505-510 https://doi.org/10.1016/j.ijom.2017.11.017
Issue Date	2018-04
Doc URL	http://hdl.handle.net/2115/73362
Rights	© 2018, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	10.2 ADDwoR manuscript OoiK.pdf

[Instructions for use](#)

Incidence of anterior disc displacement without reduction of the temporomandibular joint in patients with dentofacial deformity

Kazuhiro Ooi^{1,4}, Nobuo Inoue², Kazuhiro Matsushita¹, Hiro-o Yamaguchi¹,
Tadashi Mikoya¹, Kazuyuki Minowa³, Shuichi Kawashiri⁴, Satoshi Nishikata⁵,
Kanchu Tei¹

¹ Oral and Maxillofacial Surgery, Department of Oral Patho-biological Science,
Graduate School of Dental Medicine, Hokkaido University, Kita 13 Nishi 7 kita-ku,
Sapporo, Hokkaido 060-8586, Japan

² Gerodontology, Department of Oral Health Science, Graduate School of Dental
Medicine, Hokkaido University, Kita 13 Nishi 7 kita-ku, Sapporo, Hokkaido
060-8586, Japan

³ Dental Radiology, Department of Oral Health Science, Graduate School of
Dental Medicine, Hokkaido University, Kita 13 Nishi 7 kita-ku, Sapporo,
Hokkaido 060-8586, Japan

⁴ Department of Oral and Maxillofacial Surgery, Graduate School of Medical
Science, Kanazawa University, 13-1 Takaramachi, Kanazawa, Ishikawa,
920-8641, Japan

⁵ Oral and Maxillofacial Surgery, Sapporo Higashi Tokushukai Hospital, 3-1 Kita
33 Higashi 14 higashi-ku, Sapporo, Hokkaido 060-0033, Japan

Correspondence:

Kazuhiro Ooi: Oral and Maxillofacial Surgery, Department of Oral
Patho-biological Science, Graduate School of Dental Medicine, Hokkaido
University, Kita 13 Nishi 7 kita-ku, Sapporo 060-8586, Japan.

Tel: +81-11-706 4283; FAX: +81-11-706 4283

E-mail: ooi@den.hokudai.ac.jp

Funding : Not to declare

Key words: anterior disc displacement without reduction; dentofacial deformity;
magnetic resonance imaging; anterior open bite; mandibular asymmetry

36

37 Short title : ADDwoR of TMJ in dentofacial deformity

38

39 **Abstract**

40 This study aimed to investigate the incidence of anterior disc displacement
41 without reduction (ADDwoR) of the temporomandibular joint (TMJ) in
42 patients with dentofacial deformity. Eighty-eight female patients (176 joints)
43 with Skeletal Class III and 33 female patients (66 joints) with Skeletal Class
44 II malocclusion with or without anterior open bite and asymmetry were
45 evaluated. Magnetic resonance imaging (MRI) of the TMJ was used for
46 diagnosis of ADDwoR. Statistical analysis was performed to examine the
47 relationship between ADDwoR and skeletal structure. ADDwoR was present
48 in 37 (56.1%) of the 66 joints in class II compared to 34 (19.3%) of the 176
49 joints in class III ($p < 0.05$). In class III, ADDwoR was significantly more
50 common in joints with mandibular asymmetry (24 (32.4%) of 74) than in
51 joints with open bite (9 (14.5%) of 62) and joints with open bite and without
52 mandibular asymmetry (1 (2.6%) of 38). In class II, ADDwoR was
53 significantly less common in joints with mandibular asymmetry and without
54 open bite (1 (12.5%) of 8). ADDwoR was only observed on the deviated side in

55 both class III and class II with mandibular asymmetry. The prevalence of

56 ADDwoR was quite different according to dentofacial morphology.

57

58 INTRODUCTION

59

60 Dentofacial morphology and symptoms of temporomandibular disorder are thought to
61 be related. Recent reports have shown that symptoms of temporomandibular disorder
62 are highly associated with mandibular asymmetry¹, open bite², and Skeletal Class II
63 malocclusion³, and less common with Skeletal Class III malocclusion². However, these
64 reports are only based on the relationship between subjective symptoms of
65 temporomandibular disorder and dentofacial morphology. Few objective studies using
66 MR imaging have been done.

67 Recently, we reported that anterior disc displacement without reduction (ADDwoR)
68 and bone changes of the mandibular condylar head were significantly more common in
69 patients with skeletal open bite than in volunteers without dentofacial deformity⁴. This
70 shows that skeletal open bite is one of the factors associated with the incidence of
71 ADDwoR and bone changes of the mandibular condylar head. However, bone changes
72 of the mandibular condylar head are thought to generally occur secondary to ADDwoR⁵,
73 but they can occur without ADDwoR. In the former, mandibular condylar resorption
74 occurs following ramus height decreases; if this change is unilateral, it can cause facial

asymmetry, and if it is bilateral, it can cause open bite. In the latter, bone changes occur at a stage of development of the mandibular condylar head, and disc displacement is thought to occur after mandibular condylar resorption, but the details are unknown.

In this way, ADDwoR with dentofacial deformity is a very important factor to understand the development and function of the mandible. However, the incidence of ADDwoR in dentofacial deformity has not been sufficiently investigated. Therefore, an objective imaging study was performed to determine the relationship between the incidence of ADDwoR and dentofacial structure with dentofacial deformity.

MATERIALS AND METHODS

The subjects in this study were 121 women with dentofacial deformity, including 88 with Skeletal class III malocclusion (176 joints) and 33 with Skeletal class II malocclusion (66 joints) with or without mandibular asymmetry and open bite, who underwent orthognathic surgery at Hokkaido University Hospital, Sapporo, Japan. The median age at the time of surgery was 25 years (range, 14-48 years). None of the patients had previously been diagnosed with juvenile rheumatoid arthritis. Patients with

sagittal skeletal deformities were included in this study. Mandibular asymmetry was defined as >2 mm deviation between the menton and facial midline, and open bite was defined as <0 mm overbite. No men were included in this study to avoid skewing the cephalometric measurements by sex-related differences. The subjects included some women with clinically detectable TMJ signs and symptoms (capsular pain, joint sounds, masticatory muscle tenderness) and some without symptoms. The TMJs of patients with dentofacial deformity were examined using MR imaging to assess the position of the disc before the start of orthodontic treatment. Tesla 3.0 MRI machine was used. The position of the disc was examined using sagittal and coronal slices and T1-weighted or proton-density MR imaging with the mouth closed and open. The slice thickness was 3 mm, and the slice gap was 0.5 mm on all MR images. The bilateral surface coil for TMJ was used. On PDWI, TR (repetition time) was 1300 msec, and TE (echo time) was 30 msec. On T1WI, TR (repetition time) was 700 msec, and TE (echo time) was 15 msec. Pixel size on both PDWI and T1WI was 512×192. Software processing of the MR images was not performed after the MRI had been completed. Results of MRI were classified as with or without ADDwoR by Radiologist who trained as an MRI specialist. Subjects were divided into the Skeletal class III group with mandibular asymmetry or

109 open bite and the Skeletal class II group with mandibular asymmetry or open bite.

110 ADDwoR was considered present if the disk was displaced anteriorly relative to the

111 posterior slope of the articular eminence and the head of the condyle, but without

112 reduction of the disk on mouth opening (Fig.1). The prevalence of ADDwoR was

113 examined in each group, and statistical analysis was performed using the Chi-squared

114 test. P values of less than 0.05 were considered significant.

115 This study comply with the principles stated in the Declaration of Helsinki Ethical

116 Principles for Medical Research Involving Human Subjects, adopted by the 18th World

117 Medical Assembly, Helsinki, Finland, June 1964, and as amended most recently by the

118 64th World Medical Assembly, Fontaleza, Brazil, October 2013.

119 The work has been approved by Hokkaido University Research Ethical Committee (Ref.

120 No. 010-0285).

121

122 **RESULTS**

123

124 **Prevalence of ADDwoR in patients with Skeletal Class III and Class II**

125 ADDwoR was observed in 34 (19.3%) of 176 joints in Skeletal Class III and in 37

126 (56.1%) of 66 joints in Skeletal Class II.; the difference was significant (Fig 2).

127

128 **Prevalence of ADDwoR in patients with Skeletal class III malocclusion and**

129 **Skeletal class II malocclusion according to with or without open bite and facial**

130 **asymmetry**

131 ADDwoR was significantly more common in Skeletal Class III with mandibular

132 asymmetry (24 (32.4%) of 74 joints) than in Skeletal Class III with open bite (9 (14.5%)

133 of 62 joints). ADDwoR was observed in 14 (50.0%) of 28 joints in Skeletal Class II

134 with mandibular asymmetry and in 25 (59.5%) of 42 joints in Skeletal Class II with

135 open bite; the difference was not significant (Table 1).

136

137 **Prevalence of ADDwoR in patients with Skeletal class III malocclusion according**

138 **to with or without open bite and mandibular asymmetry**

139 ADDwoR was observed in 9 (14.1%) of 64 joints without open bite and asymmetry, 1

140 (2.6%) of 38 joints with open bite and without mandibular asymmetry, 16 (32.0%) of 50

141 joints with mandibular asymmetry and without open bite, and 8 (33.0%) of 24 joints

142 with open bite and mandibular asymmetry. ADDwoR was observed only on the deviated

side in mandibular symmetry (Table 2).

Prevalence of ADDwoR in patients with Skeletal class II malocclusion according to with or without open bite and mandibular asymmetry

ADDwoR was observed in 11 (68.7%) of 16 joints both without open bite and mandibular asymmetry, 12 (54.5%) of 22 joints with open bite and mandibular asymmetry, 1 (12.5%) of 8 joints with mandibular asymmetry and without open bite, and 13 (65.0%) of 20 joints both with open bite and mandibular asymmetry. ADDwoR was only observed on the deviated side in mandibular asymmetry (Table 3).

DISCUSSION

It has been reported that symptoms of temporomandibular disorders are more frequent in patients with open bite and mandibular asymmetry, however these symptoms are less in patients with mandibular protrusion than in persons with normal occlusion⁶. Temporomandibular disorder has been found to be more common with open bite, mandibular asymmetry, and mandibular retrusion than with other dentofacial

deformities⁷. On the other hand, it has also been reported that there is no significant relationship between dentofacial deformity and temporomandibular disorder, though 62.8% of patients with dentofacial deformities had symptoms of temporomandibular disorder before orthognathic surgery⁸. One study found no relationship between cephalometric analysis and temporomandibular disorder⁹. Further, in patients with dentofacial deformities who underwent temporomandibular joint arthrography, 57% of patients had anterior disc displacement and 53% had temporomandibular joint pain, but there were no relationships among anterior disc displacement, clinical symptoms, and dentofacial morphology¹⁰. In contrast, anterior disc displacement was reported to be more common on the deviated side of temporomandibular joints with mandibular asymmetry on MR imaging¹¹. A decreased posterior facial height and backward position and rotation of the mandible are principal characteristics associated with TMJ disk displacement^{12,13}. Thus, opinions about the association between temporomandibular disorder and dentofacial deformity are divided.

Recently, we reported that anterior disc displacement was more common with open bite than in volunteers without dentofacial deformity, and bony changes were more common than with closed lock in patients with temporomandibular joint disorder⁴. We

177 want to verify prevalence of ADDwoR between Class III and Class II, because this
178 previous study was predominantly Class II. Since the incidence of temporomandibular
179 disorder is low in patients with mandibular protrusion² and high in patients with
180 mandibular retrusion³, ADDwoR was 3 times more common in patients with
181 mandibular retrusion than in patients with mandibular protrusion in this study. In the
182 present study, Skeletal Class III and II groups were divided into those with or without
183 open bite or mandibular asymmetry, because it is thought that the rate of
184 temporomandibular disorder is generally high with open bite and mandibular
185 asymmetry. The prevalence of ADDwoR was almost 2 times higher with mandibular
186 asymmetry than with open bite in Skeletal Class III, but there was no significant
187 difference in Skeletal Class II. This result indicates that the association of ADDWoR
188 with mandibular asymmetry is stronger in Skeletal Class III than in Skeletal Class II. In
189 the Class III patient, the open bite and the development of asymmetry is related to the
190 excessive growth pattern of the condyles and the facial morphology¹⁴. Open bites in
191 Class III patients are more commonly associated with accelerated condylar and
192 mandibular growth and high occlusal plane angle facial morphologies. Class III patients
193 with low occlusal plane angle facial morphologies do not commonly demonstrate an

194 open bite. Mandibular asymmetry in Class III patients is almost always related to
195 excessive or accelerated growth of the condyle and mandible on the non-deviated side,
196 that over-loads the contralateral (deviated) side that can cause the anterior disc
197 dislocation¹⁵. Thus, Class III's with open bites and asymmetries are commonly
198 associated with accelerated growth of the condyles. In the Class II patients, open bites
199 and asymmetries are commonly related to TMJ pathologies that are causing condylar
200 resorption (the opposite of what causes the open bites and asymmetries in the Class III
201 patients). The most common causes of condylar resorption in the Class II patient are:
202 1. Adolescent internal condylar resorption (AICR) where articular discs are anteriorly
203 displaced (and commonly non-reducing) by the nature of the TMJ pathology. 2.
204 Reactive arthritis with or without anteriorly disc dislocation. 3. Connective tissue
205 autoimmune diseases^{14,15}. Thus, the development of the open bite and asymmetry in
206 Class III patients has a totally different TMJ etiology as compared to the Class II
207 patients.

208 Furthermore, Skeletal Class III and Skeletal Class II were classified into 4 groups to
209 determine the incidence of ADDwoR in this study. With mandibular asymmetry, the
210 prevalence of ADDwoR was only observed at deviated side. The prevalence of

211 ADDwoR in Skeletal Class III with open bite alone was only 2.6%, and half that rate
212 was seen without both open bite and mandibular asymmetry. This incidence is
213 extremely low because the incidence of ADDwoR in volunteers without dentofacial
214 deformity was reported to be 7%¹¹. This suggests that the prevalence of ADDwoR in
215 Skeletal Class III associated only with open bite is extremely small. Given this, it is
216 desirable to investigate the details of the temporomandibular joint in dentofacial
217 morphology to determine the etiology of overgrowth of the mandible. It would be
218 expected that skeletal Class III patients with or without ADDwoR associated only with
219 anterior open bite is extremely small. Symmetric prognathic cases rarely have anteriorly
220 displaced discs; however, in the Class III asymmetry case, where the condyle on the
221 non-deviated side overgrows creating the asymmetry, increases the loading on the
222 opposite joint that contributes to the anteriorly displacement of the disc on the
223 contralateral side. However, in the Class II patient with asymmetry, the ADDwoR is
224 involved with the condyle on the deviated side that may be undergoing unilateral
225 condylar resorption causing a shift of the mandible in the Class II patient, frequently
226 related to AICR versus reactive arthritis versus connective tissue autoimmune disease¹⁶.
227 In Skeletal Class II, the prevalence of ADDwoR with mandibular asymmetry alone

228 was 12.5%, with no bilateral cases. However, it was observed in 54.5% to 68.7% of the
229 other 3 groups, with bilateral cases found in Skeletal Class III. These results indicate
230 that the process of ADDwoR expression differs between Skeletal Class III and Skeletal
231 Class II. An investigation of more detailed temporomandibular joints in Skeletal Class
232 II associated only with mandibular asymmetry would be useful in the future. Skeletal
233 Class III and Skeletal Class II had ADDwoR only on the deviated side with mandibular
234 asymmetry. These results suggest that unilateral over-development of the
235 condyle/mandible in Class III patients and unilateral condylar resorption in Class II
236 patients is strongly associated with ADDwoR.

237 It is known that condylar cartilage is important as the starting point of mandibular
238 bone development, and endochondral ossification, which was observed mainly on the
239 mandibular condyle, is important for the development of mandibular bone. Whether
240 mandibular bone formation on the displacement side decreases because a disorder
241 occurs in the cartilage of the condyle due to anterior disc displacement or mandibular
242 asymmetry occurs first is unknown, but these results mean that it is important to
243 elucidate these clinical conditions, whether they occur at the same time because anterior
244 disc displacement occurs continuously after the abnormality has occurred or in the

245 formation of the condyle. It would be desirable to follow-up the temporomandibular
246 joints of young patients with mandibular asymmetry and anterior disc displacement in
247 the future.

248 The degenerative change of the mandibular condyle is thought to have a close
249 relationship to ADDwoR¹⁷, but ADDwoR without degenerative change of the
250 mandibular condyle has also been reported³. Katzberg et al¹⁸ and Kurt et al¹⁹ reported
251 vertical shortening of ramus height by regressive change, and that growth suppression
252 of the mandibular condyle with progression of the temporomandibular disorder may
253 cause open bite. Chen et al³ reported that condylar degenerative changes may lead to
254 deformities of the jaw, in turn resulting in decreased vertical dimensions of the proximal
255 mandibular segments and open bite. We assume that probably most of these patients are
256 Class II's with either AICR, reactive arthritis, or CT/AI. We reported that bone changes
257 occurred in 79% of temporomandibular joints with ADDwoR and open bite, ADDwoR
258 appeared and increased the mandibular plane angle, bone changes developed and
259 increased the ramus plane angle, and some patients with skeletal open bite showed worn
260 facets and no protuberances on the incisal edges²⁰. These results suggest that clockwise
261 rotation occurred, resulting in mandibular condylar resorption for various reasons in

262 ADDwoR in patients who had dental articulation contact of the incisal edges in both the
263 maxilla and mandible. Vertical malocclusion develops as a result of the interaction of
264 many different etiologic factors, including thumb and finger sucking, lip and tongue
265 habits, airway obstruction, and true skeletal growth abnormalities²¹. However, this is not
266 clear, since mandibular condylar resorption with progression of temporomandibular
267 disorder is one of the causes of open bite when looking at the present and previous
268 results. We need to investigate more to pediatric patients because the onset of "vertical
269 malocclusion" in teenage and adult patients, is more likely associated in Class II
270 patients with condylar resorption. In a study using a three-dimensional finite element
271 model of mandibular bone including the temporomandibular joint, compression stress in
272 the posterior region and tensile stress in the frontal region at the temporomandibular
273 joint increased according to the mandibular plane angle or the gonial angle²². These
274 reports suggest that it can become one of the factors in which some dentofacial
275 morphologies cause temporomandibular disorders. It seems reasonable to suggest that
276 skeletal Class II profiles and hyperdivergent growth patterns are likely associated with
277 an increased frequency of TMJ disc displacement and degenerative disorders²³. In the
278 present study, it was found that the expression of ADDwoR was very different with

279 differences in dentofacial morphology. These results show that dentofacial morphology
280 is closely associated with anterior disc displacement, and that the state of the
281 temporomandibular joint is extremely important in the development of dentofacial
282 structure. In the future, it is particularly necessary to investigate the kind of change that
283 occurs over the long term in dentofacial morphology by observing the
284 temporomandibular joints of patients with open bite and mandibular asymmetry and
285 temporomandibular disorders to better determine the relationships with
286 temporomandibular disorder or the development of dentofacial structure.

287

288

289

290 **Conflict of interest**

291 We declare that we have no conflicts of interest.

292 **Role of the funding source**

293 There was no source of funding for this research.

294 **Ethical approval**

295 Approval was given by Hokkaido University Hospital Ethics Committee (Ref.

296 No 010-0285)

297 **Patient consent**

298 Not required.

299 **References**

- 300 1. Fushima K, Akimoto S, Takamoto T, Kamei T, Satoh S, Suzuki Y. Incidence of
301 Temporomandibular Joint Disorders in Patients with Malocclusion. *J Jpn Soc TMJ*
302 1989;**1**:40-50.
- 303 2. Fujisaki T, Katoh Y, Fujita Y, Sakamoto M, Kamata S, Igarashi K, Hiyama S,
304 Kuroda T. TMJ dysfunction questionnaire on patients with malocclusion. *J Jpn Soc*
305 *TMJ* 1993;**5**:320-30.
- 306 3. Chen YJ, Shih TT, Wang JS, Wang HY, Shiau YY. Magnetic resonance images
307 of the temporomandibular joints of patients with acquired open bite. *Oral Surg Oral*
308 *Med Oral Pathol Oral Radiol Endod* 2005;**99**:734-42.
- 309 4. Yura S, Ooi K, Kadowaki S, Totsuka Y, Inoue N. Magnetic resonance imaging of
310 the temporomandibular joint in patients with skeletal open bite and subjects without
311 dentofacial abnormalities. *Br J Oral Maxillofac Surg* 2010;**48**:459-61
- 312 5. Wilkes CH. Internal derangements of the temporomandibular joint. Pathological
313 variations. *Arch Otolaryngol Head Neck Surg.* 1989 ;115 : 469-77.
- 314 6. McNamara JA Jr, Seligman DA, Okeson JP. Occlusion, Orthodontic Treatment and
315 temporomandibular disorders : a review. *J Orofac Pain* 1995 ; 9: 73-90.
- 316 7. Yoshida N, Itoh R, Saitoh T, Ota F, Hirose T, Matsumoto K. Clinical evaluation of
317 incidence of temporomandibular symptoms in patient of jaw deformity *J Jpn Soc*
318 *Orthod* 2000;**1**:64-71.
- 319 8. Laskin DM, Ryan WA, Greene CS. Incidence of temporomandiblar symptoms
320 in patients with major skeletal malocculusions —A survey of oral and
321 maxillofacial surgery training programs. *Oral Surg Oral Med Oral Pathol*
322 1986;**61**:537-41.
- 323 9. Jeon DM, Jung WS, Mah SJ, Kim TW, Ahn SJ. The effects of TMJ symptoms on
324 skeletal morphology in orthodontic patients with TMJ disc displacement. *Acta*
325 *Odontol Scand* 2014;**72**:776-82.
- 326 10. Dahlberg G, Petersson A, Westesson PL, Eriksson L. Disc Displacement
327 and temporomandiblar joint symptoms in orthognathic surgery
328 patients. *Oral Surg Oral Med Oral Pathol* 1995;**76**:237-77.
- 329 11. Yura S, Inoue N, Kurihashi K, Izumiyama Y, Asada N, Mabuchi A, Oda M,
330 Kadowaki S, Yamaguchi H, Totsuka Y. Relation between disc displacement of
331 the temporomandibular joint and skeletal structure in patients with jaw deformity.

332 *J Jpn Oral Maxillo Fac Surg* 2001;**47**:105-8.

333 12 Ahn SJ, Baek SH, Kim TW, Nahm DS. Discrimination of internal derangement of
334 temporomandibular joint by lateral cephalometric analysis. *Am J Orthod*
335 *Dentofacial Orthop* 2006;**130**:331-9.

336 13 Bertram S, Moriggl A, Neunteufel N, Rudisch A, Emshoff R. Lateral cephalometric
337 analysis of mandibular morphology: discrimination among subjects with and
338 without temporomandibular joint disk displacement and osteoarthritis. *J Oral*
339 *Rehabil* 2012;**39**:93-9.

340 14 Wolford LM, Morales-Ryan CA, Garcia-Morales P, Perez D: Surgical Management
341 of Mandibular Condylar Hyperplasia Type 1. *Bayl Univ Med Cent Proc* 2009;
342 **22**:321-9.

343 15 Wolford LM, Movahed R, Perez D: A Classification System for Conditions Causing
344 Condylar Hyperplasia. *J Oral Maxillofac Surg* 2014;**72**:567-95.

345 16 Wolford LM, Goncalves JR: Condylar Resorption of the Temporomandibular Joint:
346 How Do We Treat It? *Oral and Maxillofacial Surgery Clinics of North America*
347 2015; **27**:47-67.

348 17. Gidarakou IK, Tallents RH, Stein S, Kyrkanides S, Moss ME. Comparison of
349 skeletal and dental morphology in asymptomatic volunteers and symptomatic
350 patients with unilateral disk displacement with reduction. *Angle Orthod* 2004 ; **74**:
351 212-9.

352 18. Katzberg R.W, Tallents R.H, Hayakawa K, Miller T.L, Goske M.J, Wood B.P.
353 Internal derangements of the temporomandiblar joint. Findings in the
354 pediatric age group. *Radiology* 1985;**154**:125-8.

355 19.Kurt S, Steven P, Pollei, Clyde H.Wilkes. Pediatric internal derangements of the
356 temporomandiblar joint : Effect on facial development. *Am J Orthod*
357 *Dentfac Orthp* 1993;**104**:51-9.

358 20. Ooi K, Yura S, Inoue N, Totsuka Y. Factors related to the incidence of anterior disc
359 displacement without reduction and bony changes of the temporomandibular joint in
360 patients with anterior open bite. *Oral Maxillofac Surg* 2014;**18**:397-401.

361 21. Ngan P, Fields HW. Open bite: a review of etiology and management. *Pediatr Dent*
362 1997;**19**:91-8.

363 22.Tanaka E. Stress Distributions in the TMJ during Clenching. *J Osaka Dent* 1993 ;
364 **38**:131-60.

23 Manfredini D, Segù M, Arveda N, Lombardo L, Siciliani G, Alessandro Rossi,
Guarda-Nardini L. Temporomandibular Joint Disorders in Patients With Different
Facial Morphology. A Systematic Review of the Literature *J Oral Maxillofac Surg*
2016 ;**74** :29-46.

Fig 1. MR images of Anteriorly Displaced Discs without Reduction (ADDwoR).
(A: mouth closed, B: mouth open)

ADDwoR was considered present if the disk was displaced anteriorly relative to the posterior slope of the articular eminence and the head of the condyle, but without reduction of the disk on mouth opening.

Fig. 2. Prevalence of TMJs with ADDwoR in patients with Skeletal class III and II

ADDwoR: Anteriorly Displaced Discs without Reduction

ADDwR: Anteriorly Displaced Discs with Reduction

NDD: Non-Displaced Discs

Table 1. Prevalence of joints with ADDwoR in patients with Skeletal class III and II according to with or without open bite and mandibular asymmetry.

	Class III		P value	Class II		P value
	OB (n=62)	MA (n=74)		OB (n=42)	MA (n=28)	
ADDwoR	9 (14.5)	24 (32.4)	0.0152 *	25 (59.5)	14 (50.0)	0.4319
ADDwR	41 (66.1)	42 (56.8)	0.2643	12 (28.6)	13 (46.4)	0.1266
NDD	12 (19.4)	8 (10.8)	0.1612	5(11.9)	1 (3.6)	0.2224

ADDwoR: Anteriorly Displaced Discs without Reduction
ADDwR: Anteriorly Displaced Discs with Reduction
NDD: Non-Displaced Discs
OB: open bite MA: mandibular asymmetry

* P< 0.05 chi-square test
Data are number of TMJ (%).

Table 2. Prevalence of TMJs with ADDwoR in patients with Skeletal class III according to with or without open bite and mandibular asymmetry.

	ADDwoR (n=TMJ)	bilateral			unilateral				
		ADDwoR	ADDwR	NDD	ADDwoR/ ADDwR	ADDwoR/ NDD	ADDwR/ NDD	ADDwoR DS	ADDwoR NDS
OB/MA(-/-) (n=32)	9 / 64 (14.1)	3	18	7	3	0	1		
OB/MA(+/-) (n=19)	1 / 38 (2.6) *	0	13	4	1	0	1		
OB/MA(-/+) (n=25)	16 / 50 (32.0)	4	10	2	8	0	1	8	0
OB/MA(+/+) (n=12)	8 / 24 (33.0)	2	5	1	3	1	0	4	0

ADDwoR: Anteriorly Displaced Discs without Reduction

ADDwR: Anteriorly Displaced Discs with Reduction

NDD: Non-Displaced Discs

OB: open bite MA: mandibular asymmetry

DS: deviated side NDS: non-deviated side

(+) : with (-) : without

Data are number of patients (%).

* P< 0.05

Table 3. Prevalence of TMJs with ADDwoR in patients with Skeletal class II according to with or without open bite and mandibular asymmetry.

	ADDwoR (n=TMJ)	bilateral			unilateral				
		ADDwoR	ADDwR	NDD	ADDwoR/ ADDwR	ADDwoR/ NDD	ADDwR/ NDD	ADDwoR DS NDS	
OB/MA(-/-) (n=8)	11/ 16 (68.7)	5	2	0	1	0	0		
OB/MA(+/-) (n=11)	12 / 22 (54.5)	5	2	1	2	0	0		
OB/MA(-/+) (n=4)	1/ 8 (12.5)*	0	3	0	1	0	0	1	0
OB/MA(+/+) (n=10)	13 / 20 (65.0)	3	0	0	6	1	0	7	0

ADDwoR: Anteriorly Displaced Discs without Reduction
ADDwR: Anteriorly Displaced Discs with Reduction
NDD: Non-Displaced Discs
OB: open bite MA: mandibular asymmetry
DS: deviated side NDS: non-deviated side
(+) : with (-) : without

Data are number of patients (%).

* P< 0.05