Title	Demand and supply of cultural ecosystem services: Use of geotagged photos to map the aesthetic value of landscapes in Hokkaido
Author(s)	Yoshimura, Nobuhiko; Hiura, Tsutom
Citation	Ecosystem Services, 24, 68-78 https://doi.org/10.1016/j.ecoser.2017.02.009
Issue Date	2017-04
Doc URL	http://hdl.handle.net/2115/73803
Rights	© 2017. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	Ecosystem_Services_24_68-78.pdf

[Instructions for use](#)

Title: Demand and supply of cultural ecosystem services: use of geotagged photos to map the aesthetic value of landscapes in Hokkaido

Authors: Nobuhiko Yoshimura¹ and Tsutomu Hiura²

1. Graduate School of Environmental Science, Hokkaido University. North 10 West 5, Sapporo, Hokkaido, 060-0810, Japan

2. Tomakomai Research Station, Field Science Center for Northern Biosphere, Hokkaido University. Takaoka, Tomakomai, Hokkaido 053-0035, Japan.

Email: *Corresponding Author: Email: ynobu14001@ees.hokudai.ac.jp

Phone +81-11-706-3355

T. Hiura: hiura@fsc.hokudai.ac.jp

Abstract

We proposed a mapping method for landscape aesthetic demand and potential supply area based on viewsheds, which is a direct method that provides robust results. Moreover, we mapped the aesthetic value of Hokkaido as a case study in Asia.

The Aichi Biodiversity Target refers to the importance of ecosystem service (ES) mapping methodologies. However, ES mapping in policy and practice has rarely been reported. Robust, reliable indicators are required. Recently, studies estimating aesthetic value have used geotagged photos on social networking services instead of survey results of user preferences. The methods used in these studies were cost effective and provided spatially explicit results. However, these methods used the photography positions. Using the photographed sites is a more direct method to estimate the aesthetic demand.

Therefore, we used geotagged photos on Flickr and viewsheds from each photography position to identify the photographed sites. The demand area was estimated using the viewshed. The potential supply area was estimated using MaxEnt. The demand and potential supply areas were concentrated in natural parks. Comparing the demand and potential supply areas indicates areas with potential supply despite their low demand in forest, farmland, and natural parks. This method will contribute to CES research and decision-making.

Keywords

Cultural ecosystem service, Aesthetics, Flickr, MaxEnt, GIS, Social networking service

1. Introduction

Various scales of decision-making require the measurement and visualization of the value of ecosystem services (ES), which is not included in the market price (MEA, 2005; TEEB, 2010; WBCSD, 2010). The value of cultural ecosystem services (CES) is particularly difficult to measure, both physically and monetarily, because such values are intangible and subjective (MEA, 2005). However, researchers and practitioners know that some CESs contribute to human well-being (Milcu et al., 2013). CES also motivates environmental action compared with other ESs (Hirons et al., 2016). Many CES studies have been conducted in Europe and North and Central America. In Asia, many CES studies have been carried out in China but not in Japan (Hernández-Morcillo et al., 2013; Wolff et al., 2015a). There are many CES studies on recreation, ecotourism, and the aesthetic value of landscapes because of their strong economic relevance and relative ease of estimation (Hernández-Morcillo et al., 2013; Milcu et al., 2013).

Travel and tourism accounted for 9.8% (US\$7.2 trillion) of global GDP in 2015, and this contribution is expected to increase to 10.8% (US\$11 trillion) by 2026 (WTTC, 2016). Growth will mostly come from nature-based tourism (Balmford et al., 2009); thus, the potential economic value is high. The aesthetic value of landscapes is the pleasure derived from natural beauty (TEEB, 2010). Aesthetic value is an especially important factor in recreation (Daniel et al., 2012), and is strongly related to market price. For example, residential price is affected by the amount of ocean or lake views (Benson et al., 1998; Crossman et al., 2013a). Mapping CES values provides important points of view for the development of conservation plans and for land-use management. The economic evaluation framework in the TEEB (2008) interim report states that quantification and mapping of ES values are necessary. However, only 18% of mapping studies have mapped CESs, far fewer than those that map regulating services (46%) and provisioning services (30%) (Crossman et al., 2013b). The Aichi Biodiversity Target refers to the importance of ES mapping methodologies (CBD Secretariat, 2011). Much CES mapping has reflected people's preferences and has used methods including an empirical method, a participatory approach, and a monetary valuation (Wolff et al., 2015b). The empirical method evaluates CES based on questionnaire surveys or interviews that reflect people's preferences or the value of sites (Casado-Arzuaga et al., 2013; de Vries et al., 2007; Sherrouse et al., 2011; van Zanten et al., 2016b). The participatory approach, which is based on expert knowledge or preferences of specific users (Kenter, 2016; Palomo et al., 2013; Scolozzi et al., 2014), is spatially explicit and

has high estimation accuracy in site-specific studies (Wolff et al., 2015b). Therefore, the empirical method is more appropriate than the participatory approach for evaluating CES over a large area (Wolff et al., 2015b). Monetary valuation estimates the monetary value of CES using concepts such as willingness to pay (Häyhä et al., 2015; Kenter, 2016; Nahuelhual et al., 2014). These methods have been used frequently in combination in many studies (Kenter, 2016).

Mapping the aesthetic value of landscapes reflects people's preferences in the same way as the other CESs. It has been mainly based on the results of questionnaire surveys (Casado-Arzuaga et al., 2013; Peña et al., 2015) or interviews on preferences (van Zanten et al., 2016b) gained by empirical methods, and by participatory approaches combined with environmental factors that represent attractiveness such as naturalness (Crossman et al., 2013a; de Vries et al., 2007). However, such surveys are often costly and time-consuming. One study used an Internet survey to improve the survey efficiency (Peña et al., 2015), but the preparation of the questionnaire itself remains a time-consuming process.

In recent years, the widespread use of mobile devices has led to the development of social networking services (SNS), through which text messages and photos of food, landscapes, portraits, and so on are shared on the Internet. Geotagged photos are also increasingly shared, and many of these can be collected from wide areas. SNS data uploaded by users provides user preferences and experiences, and thereby contributes to improving the empirical method. In studies related to the aesthetic value, Casalegno et al. (2013) showed the effectiveness of geotagged photos in estimating the aesthetic value on a regional scale and van Zanten et al. (2016a) demonstrated it at a continental scale. Richards and Friess (2015) demonstrated the rapidness and cost effectiveness of using geotagged photos. Martínez Pastur et al. (2016) showed that classified tags of geotagged photos and their positions can be used to map some CES values such as aesthetic and recreation values. However, many studies have also highlighted the shortcomings of SNS data. The bias of the user group was one of the big concerns (Guerrero et al., 2016; Tenerelli et al., 2016). Therefore, the points of usage of SNS data have been reported (Crampton et al., 2013), and demographic studies have also been conducted to clarify the bias of SNS data (Garcia-Palomares et al., 2015; Wood et al., 2013).

The use of ES mapping applications in policy and practice have rarely been

reported. Robust, reliable indicators and shared understanding of ES values are needed (Wolff et al., 2015b). The methods of estimating aesthetic value with geotagged photos have used the photography positions not the photographed sites. Therefore, they have usually used density of users who uploaded photos in a grid and models, such as the general linear model, to estimate the aesthetic value. However, using photographed sites is a more method direct than using photography positions and should provide robust results.

We improved the empirical method that uses geotagged photos to estimate the aesthetic value for use in policy and in practice. Mapping both the demand and supply potential of ESs also helps to understand the current state of ES. First, we developed a method to estimate aesthetic demand and potential supply area based on viewsheds and MaxEnt, which provided robust results. Second, we applied the method to Hokkaido, the northernmost island of Japan, to assess the current aesthetic value of the landscapes as a case study in Asia.

2. Materials and Methods

2.1. Study area

Our study area, which was used to develop a method for mapping aesthetic value, was Hokkaido, Japan (43° 31' N, 142° 40' E) (Fig. S1). The area is about 7.8 million ha, and its population in 2010 was 5.5 million (Statistics Bureau Ministry of Internal Affairs and Communications, 2016). The annual average temperature was about 9.8 °C in 2010 (Statistics Bureau Ministry of Internal Affairs and Communications, 2016). The area includes rich natural resources and 23 natural parks. Many tourists utilize the resort areas in Hokkaido to enjoy natural features such as wetlands, lakes, and snow. Moreover, Hokkaido is also a famous agricultural area in Japan and has rice paddies, corn fields, wheat fields, and dairy farms. The number of tourists in 2014 was 7.2 million, comprising 21.3% from overseas and 78.7% from domestic areas (Hokkaido Bureau of Tourism, Depart of Economic Affairs, 2015).

2.2. Mapping method

We developed a method for mapping the aesthetic value of landscapes in terms of demand and potential supply, by using geotagged photos on Flickr as preference data

for aesthetics. Flickr was launched in 2004 and has been operated by Yahoo, Inc. since 2005. As of August 2011, Flickr hosted approximately 6 billion uploaded photos (Flickr, 2011). Wood et al. (2013) reported that the number of Flickr users uploading photos has a positive correlation with the annual number of tourists, and thus this number can be used as a proxy for visitation. To confirm Flickr data characteristics in the studied area, we compared the number of tourists to each municipality between 2010 and 2014 (Hokkaido Bureau of Tourism, Depart of Economic Affairs, 2015) with the number of Flickr users who uploaded photos in the same municipality for the same period. This comparison revealed a positive correlation on a log–log scale ($R^2 = 0.61$, $p < 0.001$).

2.3. Developing photo datasets

Flickr provides an Application Programming Interface (API) (Flickr, 2016) that allows photos to be searched based on metadata such as photo ID, user ID, Where On Earth IDentifier (WOEID) that identifies the location, coordinates, title, tags, date, and positional accuracy. We created three datasets using this API (Table S1). Recently, photos taken with mobile phones are being increasingly uploaded to the Flickr database (Flickr, 2015). The accuracy of GPS in mobile phones is almost within 10 m (Zandbergen and Barbeau, 2011), and mobile phones use has increased since around 2010. We were able to download statistics of visitors to the study area up to 2014. We collected metadata from geotagged photos taken from 2010 to 2014 to confirm the relation between the number of Flickr photos and visitors. We created a first dataset (the Filter 1 dataset) containing geotagged photos uploaded during this time with street-level positional accuracy (provided positional accuracy level by Flickr ≥ 12 (Flickr, 2016)) and taken within 500 m of the coastline in the study area (WOEID = 7153351). This dataset has 136,023 geotagged photos including coastal area photos taken from boats. Filter 1 dataset photos included landscape photos and all other types of photos such as portraits with food and monuments. Therefore, we extracted only landscape photos using the “landscape” keyword. A sample of extracted photos is shown in Fig. S2. By this filter, portraits containing landscapes were removed.

There were 13,202 photos in the Filter 2 dataset, which included many photos taken by heavy users who had uploaded dozens of photos of a certain area or continuous-shot photos. To avoid bias, we sampled photos at random to limit the number of photos per user in each municipality to one (Filter 3). There were 2982 photos in the Filter 3 dataset, and we used this dataset to map the demand for aesthetic value of landscapes.

2.4. Mapping demand

We considered landscape sites where photos were likely to have been taken by Flickr users as sites with high landscape demand. The latitude and longitude of photography positions are stored on Flickr, but not those of the photographed sites. Since photographic orientation is rarely recorded (Shirai et al., 2013), it is hard to identify target sites. Thus, we developed an index of target probability using viewsheds calculated from photographing positions. A viewshed is a visible area from each photographing position and therefore should include the photographed target site. We assumed that the more viewsheds overlap, the higher the probability the overlapping area was a target site. However, more photos including landscape photos were likely to have been taken at crowded sites, such as popular destinations, potentially leading to the overestimation of the value of sites with high popularity and good accessibility. To adjust for this bias, we determined a viewshed score, which is the ratio of the number of landscape photos to the total number of photos taken around each photography position. We then summed scores for overlapping areas to map aesthetic demand (Fig. 1). The ratio was calculated in a 3 km radius from the photographing position considering the distribution density of Flickr photo points.

We defined and calculated each viewshed using a 50-m digital elevation model from a height of 170 cm and within a maximum radius of 10 km. Because there was no reference to the distance from the photographing position to the target site of Flickr photos, the maximum view range was set in reference to previous studies of the visible distance of wind farms (Bishop and Miller, 2007; Sullivan et al., 2013, 2012). The spatial resolution of demand mapping was 50×50 m, and the result was normalized as 0 to 1 and mapped with five ranks. The normalizing method was a Fuzzy MS Large function (ArcGIS10.3.1, ESRI Inc.). In this method, large input values were more likely to be 1. Transformation of 0 to 1 was based on mean and standard deviation (ESRI Inc., 2014).

2.5. Mapping potential supply

We can map potential supply by analyzing the relation between demand area and its environmental factors because the demand map represents the aesthetic preferences of visitors (Fig. S3). Openness, variety, and the presence of water are reported as preference factors for aesthetic value of landscapes (Uuemaa et al., 2013). In a study of Japanese preferences, Kojima et al. (1994) also reported naturalness,

uniformity or variety, openness, and the presence of water as preference factors.

We collected geospatial data as environmental factors to map the potential supply according to three categories based on the above studies: naturalness, water influence, and topography (Table S2). These data were published by the Japanese government and can be downloaded for free (Biodiversity Center of Japan, 1998; Japanese Ministry of Land Infrastructure Transport and Tourism, 2016). We also referred to previous studies of mapping CES values such as aesthetic value (Casado-Arzuaga et al., 2013; de Vries et al., 2007; Howley, 2011; Peña et al., 2015). Naturalness was based on vegetation type, and it was classified into 10 ranks according to human impact. Distances from rivers, lakes, or coastline were used as factors for the influence of water. Moreover, specific riverine landscapes, such as snaking streams, were used as geological interest for rivers. Topographic factors included 10 classes of topography, including ridges, valleys and flat plains, variety of topography, and distance from volcanic and non-volcanic topography, such as terrain caldera and cirque, as geologically interesting. Shannon's diversity index was used to calculate the variety of topography (Frank et al., 2013). None of these variables had a significant correlation with each other (all $r < 0.4$). The spatial resolution of the potential supply mapping was 50×50 m.

We used maximum entropy modeling software, MaxEnt version 3.3 (<http://www.cs.princeton.edu/~schapire/maxent/>) to estimate and map potential supply (Phillips et al., 2006, 2004). The software uses machine-learning to estimate a probability distribution by finding the probability distribution of the maximum entropy of given observed points such as species, plant, and environmental factors. The observed points are called presence data. An advantage of MaxEnt is that absence data is not required (only presence data is required). Therefore, MaxEnt is widely used to estimate species distribution (Elith et al., 2011), and it can also be used to map the social value of nature (Richards and Friess, 2015; Sherrouse et al., 2014, 2011). Flickr data is presence data, because the places where Flickr users did not upload their photos were not always the places where other social media users or people who do not use SNS did not take photos. Therefore, MaxEnt was also suitable for analyzing Flickr data.

Our demand map was the probability of distribution of demand, which was based on presence data. It had five ranks, but rank data was not used as presence data in MaxEnt. We then created new 1000 random points as presence data from the entire

demand area (rank 1 to 5). The density was about one point per 1100 ha. The Area Under the Curve (AUC) of the Receiver Operating Characteristic (ROC) plot is used to assess the MaxEnt model. ROC is the relation between sensitivity and 1-specificity. Sensitivity is the true positive rate (predicted value is true and actual value is also true) in a contingency table. 1-Specificity is the false positive rate (predicted value is true and actual value is false). Sensitivity and 1-specificity are calculated using thresholds that represent the probability of occurrence in MaxEnt. ROC can measure the efficiency of a binary classifier, such as the MaxEnt model, and AUC represents the probability of sensitivity. An AUC value of 0.5 indicates a random model, and a value of 1 indicates a model that perfectly classifies given presence data. An AUC value of 0.50 to 0.70 suggests a somewhat accurate model, a value of 0.70–0.90 suggests an accurate model, and a value exceeding 0.90 indicates a highly accurate model (Swets, 1988). In the preliminary analysis, the AUC value of a 10 km radius as maximum viewshed distance was higher than that of the 20 km radius maximum; thus, we decided the maximum radius as 10 km. We made a model of potential supply using 80% of the above presence data and validated using the others. This process was repeated 10 times for cross-validation and sensitivity and AUC was averaged as a final model.

We identified the potential supply area with similar environmental conditions to the demand area with two cutoff points. One was called the 10th percentile training presence, which is a cutoff point that includes 90% of presence data (10% threshold). Another was called equal training sensitivity and specificity, which is a cutoff point with a minimum distance between ROC and the upper left corner of the ROC plot whose sensitivity was 1. The latter is a stricter criterion than the former.

3. Results

3.1. Demand map

Figure 2 shows a demand map based on the Filter 3 dataset. This map provides an overview and detailed distribution of the aesthetic demand. For example, when enlarging high-demand areas located at the center of this study area (Fig. 3), we found that high-demand areas were located on the west side of the Tokachidake Mountain Range, and the Daisetsu Mountain Range and in the farmland at its foot. The urban area near the high-demand area also had a high demand. Comparing the photography

positions and topography, we found that demand areas depended on viewshed.

Overall, the total demand area was about 1,092,266 ha, 13.7% of the study area. Areas of high demand, especially those with ranks 4 and 5 were located in forest, farmland, river-lake, and wetland-bareland (Fig. 4). The ratio of demand area in each land use type was in the order river-lake > urban areas > sea > grassland-golf courses. The demand area corresponded well to places with tourism resources. Table 1 shows the area of demand for each natural park. Natural parks contained 24.4% of the demand area. Focusing on rank 4 and 5 areas, 40.8% of these areas were located in natural parks. The area of natural parks corresponds to 10.9% of the total study area. Therefore, rank 4 and 5 and total demand area were significantly biased toward natural parks ($\chi^2 = 190375.4, 160381.2$, both d.f. = 1, $p < 0.001$).

Areas of demand, especially those with rank 4 and 5, were located in national parks rather than in quasi-national or prefectural natural parks. However, focusing on the ratio of demanded areas in each national park, a large non-demanded area was located in the Daisetsuzan National Park; its percentile of demanded area was low, despite having the largest total area. A non-demanded area was also found in every national park, although ratios differed from park to park.

3.2. Supply map

Figure 5 shows a potential supply map and Table 2 shows the contribution of environmental factors. Environmental factors that contributed highly to the model were, in order, distance from volcanic topography (23.2%), type of topography (23.2%), variety of topography (19.6%), naturalness (13.3%), and distance from lakes (8.0%). Distance from specific riverine landscapes did not contribute to the model greatly (0.8%). In terms of the response to demand for types of topography, plains, u-shape valleys, and mountain tops-high ridges were the major contributors to the model, and low (0.0) and high (1.7) topographic variety contributed more than medium topographic variety (1.3) (Fig. S4). In the category of naturalness, 1 (urban areas), 2 (farmland, urban and residential districts with many trees), 4 (secondary short grassland), 8 (natural broad-leaved forest), and 10 (alpine heathland and wind-exposed grassland) contributed to the model. Bodies of water also contributed. Volcanic topography (23.2%) contributed more than non-volcanic topography (6.2%). Areas near and far from volcanic topography strongly contributed for volcanic topography. The AUC value of the model was 0.802.

Overall, the total potential supply area was estimated as 1,723,345 ha (equal threshold) to 2,985,729 ha (10% threshold). The area corresponded to 21.6 to 37.4% of the study area, which was wider than the demand area (13.7%). We compared the demand area of each land use type with the potential supply area (10% threshold) of that (Fig. 6). The trend of the potential supply area between land use types was similar to that of the demand area. However, the area of potential supply in each of the land use types was wider than that of demand. Many areas that had potential supply without demand (supply gap area) were located in forest and farmland. Focusing on the natural parks, few supply gap areas were located in wetland-bareland and river-lake, but there were larger supply gap areas in forest. Outside of natural parks, the large supply gap areas were located in forest and farmland. 16.6–19.0% (496,320–326,696 ha) of potential supply area were located in natural parks, and it was significantly biased to natural parks ($\chi^2 = 10\%$ threshold: 49620.5, equal threshold: 63482.6, both d.f. = 1, $p < 0.001$).

We compared the potential supply with demand in areas of each natural park (Table 1). In natural parks, areas of potential supply exceeded the demand area at both the 10% threshold and equal threshold. More areas of potential supply were located in national parks than in quasi-national or prefectural natural parks. At the 10% threshold, the area of potential supply exceeded the area of demand in the Daisetsuzan National Park and Shikotsu Toya National Park, which had low ratios of demand areas. However, areas of demand were almost same as that of potential supply in the Kushiro Shitsugen National Park.

Figure 7 shows the distribution of demand and potential supply area in forest and farmland according to the 10% threshold. Of forest areas, 70.6% did not show aesthetic value. The place that had potential supply with demand (balanced area) was 86,615.3 ha, 6.2% of total forest area. The supply gap area was 287,205.8 ha (20.6%). Many supply gap areas were located around national parks such as Daisetsuzan National Park and Shikotsu Toya National Park. Of farmland areas, 34.5% did not show aesthetic value. The balanced area was 66,321 ha, 19.1% of the total farmland area. The supply gap area was 153,257 ha (44.2%). The supply gap area was the largest in farmland. Focusing on the main agricultural areas in Hokkaido, many balanced areas were located in the Kamikawa, Sorachi, and Tokachi regions, but not many were located in the Abashiri, Nemuro, and Kushiro regions.

4. Discussion

We proposed a viewshed-based method for measuring aesthetic demand and potential supply of landscape value using Flickr photos as a preference index. Because the method for estimating the area of aesthetic demand did not use questionnaire surveys or models, such as general linear models, it was considered to be rather direct and provided robust results. MaxEnt, which is frequently used to estimate the distribution of species, was used to estimate potential supply areas. MaxEnt also provides robust results even in the spatial problem of human dimensions (Sherrouse et al., 2014). Because the photography positions of Flickr photos were presence data, the area of demand based on Flickr data also had the same characteristics. Therefore, MaxEnt was considered appropriate for estimating the potential supply area from the demand area. In contrast, because there were many copies that were shared by users or many photos posted by one user, the sample size from SNS data sometimes becomes small after filtering (Crampton et al., 2013). However, MaxEnt retains its prediction accuracy even with small sample sizes (Wisz et al., 2008). Therefore, MaxEnt is useful for CES mapping with SNS data.

Natural areas, such as forest, river-lake, and wetland-bareland, as well as farmland and urban areas, had many high-demand areas in Hokkaido. In particular, the percentage of demand area and highest demand area (rank 5) were very high for river-lake. These results are consistent with other studies that mentioned that natural areas, such as forest, lakes, and wetland, were also preferred (de Vries et al., 2007; Howley, 2011; Peña et al., 2015). Farmland was not preferred as a factor of aesthetic value (Lindemann-Matthies et al., 2010; Van Berkel and Verburg, 2014; van Zanten et al., 2016c), although some studies have reported that it was preferred (Casado-Arzuaga et al., 2013; Peña et al., 2015). Japanese people tended to prefer a single species or nature influenced by people (Kellert, 1991), and uniformness as aesthetic factors (Kojima et al., 1994). Because 70% of visitors were domestic, the fact that farmland was preferred in this study was likely to be affected by Japanese preferences for nature. The urban area had high demand based on the photos taken at viewpoints where people can enjoy city views or the photos taken near urban parks (Guerrero et al., 2016).

Demand and potential supply areas were predominantly in natural parks but were concentrated around restricted areas in many national parks. Japanese national parks are intended to protect scenery and biodiversity and to promote recreation

according to the Natural Parks Act (Japanese Ministry of the Environment, 2016, 2014). In the natural parks, many supply gap areas were located in the forest, but very few were in river-lake and wetland-bareland. There were the national parks whose supply gap area was large, such as Daisetsuzan National Park and Shikotsu Toya National Park, and those, such as Kushiro Shitsugen National park, whose supply gap area was very small. The parks with large supply gaps were considered to have poorly accessible areas in their forest, whereas Kushiro Shitsugen National Park is mainly marsh and it was visible from enough viewpoints. This information may help the park manager to understand the current conditions and manage the facility considering aesthetic aspects.

Outside the natural parks, much demand and potential supply area were located in forest and farmland (Fig. 7). Many supply gap areas in forest area were located around Daisetsuzan National Park and Shikotsu Toya National Park. Forest is expected to perform many ES, such as timber production, water purification, carbon stock, and maintaining biodiversity. Aichi Biodiversity Target 11 said that by 2020, at least 17% of terrestrial and inland water, and 10% of coastal and marine areas, especially areas of particular importance for biodiversity and ES, should be conserved having been integrated into the wider landscapes and seascapes (CBD Secretariat, 2011). Therefore, a study to prioritize conservation areas has been conducted (Kadoya et al., 2014). Although the public understanding of biodiversity is not high (CBD Secretariat, 2011), CES such as aesthetic value is well recognized and motivates public environmental awareness (Hirons et al., 2016). The total percentage contribution or importance of the variety of topography, naturalness, and distance from lakes, rivers, and coastline, which were used in this study as the environmental factors, was about 50% (Table 2). These environmental factors are related to the complexity and connectivity of landscapes that often has a strong positive relation to the biodiversity richness on various scales (Amici et al., 2015; Chisholm et al., 2011; Ishii et al., 2004; Loreau et al., 2003; Rösch et al., 2013; Takafumi and Hiura, 2009). Consequently, demand and potential supply area in this study was considered to be related to biodiversity richness. Therefore, overlaying the demand and potential supply area and prioritizing conservation areas can help to find places that are important for conservation and aesthetic enjoyment, improving the stakeholders' understanding of conservation. Farmland with potential supply has the potential for both CES and provisioning services. We found that these regions that had many supply gap areas despite having few balanced areas. These regions were considered to have good potential for developing tourism in agricultural areas. This

information could help land managers or the tourist industry identify hidden tourism resources. In addition, the distribution or value of CES varied with the seasons (Tenerelli et al., 2016; Wolff et al., 2015a). Further study related to seasonal changes in the aesthetic value will be needed.

Finally, we discuss the limitations of our proposed method and SNS data in the CES mapping study. In our method, we must consider the estimation error for the target site of each photo; an estimated target site is likely to be wider than the actual site. In this study, we fixed the maximum viewshed range as 10 km, and thus did not consider the distribution of the distance to the target site in landscape photos. Moreover, we did not fix the direction to the target sites, which may improve target site identification. In particular, urban areas near high-demand farmland also had high demand, which was overestimated because the elevation data that was used to calculate the viewsheds did not include the height of buildings or houses.

The positional accuracy of geotagged photos should also be considered. We used viewsheds to estimate locations from Flickr photos, but estimated locations can be directly affected by the positional accuracy of geotagged photos. To reduce this limitation, we used accuracy parameters provided by Flickr and filtered to street-level accuracy. We furthermore considered the data collection period (from 2010, after the spread of mobile devices). Wang et al. (2013) showed that Flickr photos have positional errors from tens to hundreds of meters. We suggest that the positional accuracy of Flickr photos will improve with an increase in the number of photos taken using mobile devices. Positional error of data also depends on the collection period, for example, those taken before the spread of mobile devices.

As of 2012, 200 million geotagged photos had been uploaded to the Flickr database, 40% of which were taken in Europe, 39% in North America, and 13% in Asia; Japan is a region that has many geotagged photos on Flickr, similar to North America and Europe (Wood et al., 2013). However, the problem of sample representativeness remains. The number of Flickr users was positively related to the number of visitors (Wood et al., 2013). Although we found this relation in this study, not all visitors used SNS (Garcia-Palomares et al., 2015). The representativeness was affected by the rate of Internet use, cameras with GPS, and mobile phones in a region (Martínez Pastur et al., 2016). These depended on age, education, and ability or motivation to use SNS (Tenerelli et al., 2016). Moreover, SNS platforms that have different user groups also

affected the representativeness (van Zanten et al., 2016a). However, methods based on questionnaire surveys or interviews also have the problem of representativeness (Tenerelli et al., 2016). Using SNS data for CES mapping is cost-effective despite having spatial coverage and spatial explicitness. Therefore, further demographic studies of SNS are needed.

5. Conclusion

We demonstrated the viewshed base method to map aesthetic demand and potential supply, which was a more direct method and provided robust results. Moreover, we also clarified the aesthetic value of Hokkaido, as a case study in Asia.

Mapping the aesthetic demand and potential supply can promote awareness of both land value and the importance of nature conservation. We hope that our results will contribute to CES research and decision-making in policy and in practice to maintain a balance between human use and nature conservation based on aesthetic demand and its potential supply.

Acknowledgments

We thank Dr. Y. Yamanaka and Dr. M. Ishihara for her valuable comments regarding this project. This study was partly supported by the Ministry of the Environment, Japan (S9-3).

Reference

- Amici, V., Rocchini, D., Filibeck, G., Bacaro, G., Santi, E., Geri, F., Landi, S., Scoppola, A., Chiarucci, A., 2015. Landscape structure effects on forest plant diversity at local scale: Exploring the role of spatial extent. *Ecol. Complex.* 21, 44–52. doi:10.1016/j.ecocom.2014.12.004
- Balmford, A., Beresford, J., Green, J., Naidoo, R., Walpole, M., Manica, A., 2009. A global perspective on trends in nature-based tourism. *PLoS Biol.* 7, 1–6. doi:10.1371/journal.pbio.1000144
- Benson, E.D., Hansen, J.L., Schwartz, A.L., Smersh, G.T., 1998. Pricing residential amenities: the value of a view. *J. Real Estate Financ. Econ.* 16, 55–73.

doi:10.1023/A:1007785315925

- Biodiversity Center of Japan, 1998. Natural Environmental Information GIS [WWW Document]. URL http://www.biodic.go.jp/trialSystem/top_en.html (accessed 4.29.16).
- Bishop, I.D., Miller, D.R., 2007. Visual assessment of off-shore wind turbines: The influence of distance, contrast, movement and social variables. *Renew. Energy* 32, 814–831. doi:10.1016/j.renene.2006.03.009
- Casado-Arzuaga, I., Onaindia, M., Madariaga, I., Verburg, P.H., 2013. Mapping recreation and aesthetic value of ecosystems in the Bilbao Metropolitan Greenbelt (northern Spain) to support landscape planning. *Landsc. Ecol.* 29, 1393–1405. doi:10.1007/s10980-013-9945-2
- Casalegno, S., Inger, R., DeSilvey, C., Gaston, K.J., 2013. Spatial Covariance between Aesthetic Value & Other Ecosystem Services. *PLoS One* 8, 6–10. doi:10.1371/journal.pone.0068437
- CBD Secretariat, 2011. Convention on Biological Diversity [WWW Document]. URL <https://www.cbd.int/sp/targets/> (accessed 1.12.17).
- Chisholm, C., Lindo, Z., Gonzalez, A., 2011. Metacommunity diversity depends on connectivity and patch arrangement in heterogeneous habitat networks. *Ecography* 34, 415–424. doi:10.1111/j.1600-0587.2010.06588.x
- Crampton, J.W., Graham, M., Poorthuis, A., Shelton, T., Stephens, M., Wilson, M.W., Zook, M., 2013. Beyond the geotag: situating “big data” and leveraging the potential of the geoweb. *Cartogr. Geogr. Inf. Sci.* 40, 130–139. doi:10.1080/15230406.2013.777137
- Crossman, N.D., Bryan, B.A., de Groot, R.S., Lin, Y.P., Minang, P.A., 2013a. Land science contributions to ecosystem services. *Curr. Opin. Environ. Sustain.* 5, 509–514. doi:10.1016/j.cosust.2013.06.003
- Crossman, N.D., Burkhard, B., Nedkov, S., Willemen, L., Petz, K., Palomo, I., Drakou, E.G., Martin-Lopez, B., McPhearson, T., Boyanova, K., Alkemade, R., Egoh, B., Dunbar, M.B., Maes, J., 2013b. A blueprint for mapping and modelling ecosystem services. *Ecosyst. Serv.* 4, 4–14. doi:10.1016/j.ecoser.2013.02.001
- Daniel, T.C., Muhar, A., Arnberger, A., Aznar, O., Boyd, J.W., Chan, K.M. a., Costanza, R., Elmqvist, T., Flint, C.G., Gobster, P.H., Gret-Regamey, A., Lave, R., Muhar, S., Penker, M., Ribe, R.G., Schauppenlehner, T., Sikor, T., Soloviy, I., Spierenburg, M., Taczanowska, K., Tam, J., von der Dunk, A., 2012. Contributions of cultural

- services to the ecosystem services agenda. *Proc. Natl. Acad. Sci.* 109, 8812–8819.
doi:10.1073/pnas.1114773109
- de Vries, S., Klein-Lankhorst, J.R., Buijs, A.E., 2007. Mapping the attractiveness of the Dutch countryside: a GIS-based landscape appreciation model. *For. Snow Landsc. Res.* 81, 43–58.
- Elith, J., Phillips, S.J., Hastie, T., Dudík, M., Chee, Y.E., Yates, C.J., 2011. A statistical explanation of MaxEnt for ecologists. *Divers. Distrib.* 17, 43–57.
doi:10.1111/j.1472-4642.2010.00725.x
- ESRI Inc., 2014. How Fuzzy Membership works—Help | ArcGIS for Desktop [WWW Document]. URL
<http://desktop.arcgis.com/en/arcmap/10.3/tools/spatial-analyst-toolbox/how-fuzzy-membership-works.htm> (accessed 1.15.17).
- Flickr, 2016. API Documentation [WWW Document]. URL
<https://www.flickr.com/services/api/flickr.photos.search.html> (accessed 4.29.16).
- Flickr, 2015. Flickr year review 2015 [WWW Document]. URL
<http://blog.flickr.net/en/2015/12/01/flickr-year-in-review-2015/> (accessed 4.29.16).
- Flickr, 2011. 6,000,000,000 | Flickr Blog [WWW Document]. URL
<http://blog.flickr.net/en/2011/08/04/6000000000/> (accessed 1.12.17).
- Frank, S., Fürst, C., Koschke, L., Witt, A., Makeschin, F., 2013. Assessment of landscape aesthetics - Validation of a landscape metrics-based assessment by visual estimation of the scenic beauty. *Ecol. Indic.* 32, 222–231.
doi:10.1016/j.ecolind.2013.03.026
- Garcia-Palomares, J.C., Gutierrez, J., Minniguez, C., 2015. Identification of tourist hot spots based on social networks: A comparative analysis of European metropolises using photo-sharing services and GIS. *Appl. Geogr.* 63, 408–417.
- Guerrero, P., Møller, M.S., Olafsson, A.S., Snizek, B., 2016. Revealing Cultural Ecosystem Services through Instagram Images: The Potential of Social Media Volunteered Geographic Information for Urban Green Infrastructure Planning and Governance. *Urban Plan.* 1, 1. doi:10.17645/up.v1i2.609
- Häyhä, T., Franzese, P.P., Paletto, A., Fath, B.D., 2015. Assessing, valuing, and mapping ecosystem services in Alpine forests. *Ecosyst. Serv.* 14, 12–23.
doi:10.1016/j.ecoser.2015.03.001
- Hernández-Morcillo, M., Plieninger, T., Bieling, C., 2013. An empirical review of cultural ecosystem service indicators. *Ecol. Indic.* 29, 434–444.

- doi:10.1016/j.ecolind.2013.01.013
- Hirons, M., Comberti, C., Dunford, R., 2016. Valuing Cultural Ecosystem Services. *Annu. Rev. Environ. Resour.* 41, 545–574.
doi:10.1146/annurev-environ-110615-085831
- Hokkaido Bureau of Tourism Depart of Economic Affairs, 2015. Tourism Statistics reports for Hokkaido [WWW Document]. URL <http://www.pref.hokkaido.lg.jp/kz/kkd/irikomi.htm> (accessed 4.29.16).
- Howley, P., 2011. Landscape aesthetics: Assessing the general publics' preferences towards rural landscapes. *Ecol. Econ.* 72, 161–169.
doi:10.1016/j.ecolecon.2011.09.026
- Ishii, H.T., Tanabe, S., Hiura, T., 2004. Exploring the relationships among canopy structure, stand productivity, and biodiversity of temperate forest ecosystems. *For. Sci.* 50, 342–355.
- Japanese Ministry of the Environment, 2016. Project to Fully Enjoy National Parks [WWW Document]. URL <https://www.env.go.jp/en/nature/enjoy-project/index.html> (accessed 1.12.17).
- Japanese Ministry of the Environment, 2014. Natural Parks Act [WWW Document]. URL <http://law.e-gov.go.jp/htmldata/S32/S32HO161.html> (accessed 4.29.16).
- Japanese Ministry of Land Infrastructure Transport and Tourism, 2016. National Land Numerical Information download service [WWW Document]. URL <http://nlftp.mlit.go.jp/ksj-e/index.html> (accessed 4.29.16).
- Jenness, J., Brost, B., Beier, P., 2013. Land Facet Corridor Designer: Extension for ArcGIS. Jenness Enterp. 110.
- Kadoya, T., Takenaka, A., Ishihama, F., Fujita, T., Ogawa, M., Katsuyama, T., Kadono, Y., Kawakubo, N., Serizawa, S., Takahashi, H., Takamiya, M., Fujii, S., Matsuda, H., Muneda, K., Yokota, M., Yonekura, K., Yahara, T., 2014. Crisis of Japanese vascular flora shown by quantifying extinction risks for 1618 taxa. *PLoS One* 9.
doi:10.1371/journal.pone.0098954
- Kellert, S.R., 1991. Japanese Perceptions of Wildlife. *Conserv. Biol.* 5, 297–308.
doi:10.1111/j.1523-1739.1991.tb00141.x
- Kenter, J.O., 2016. Integrating deliberative monetary valuation, systems modelling and participatory mapping to assess shared values of ecosystem services. *Ecosyst. Serv.* 21, 291–307. doi:10.1016/j.ecoser.2016.06.010
- Kojima, T., Furuya, K., Yui, M., 1994. A Study on the Evaluation Structure of

- Favorability in Natural Landscape. *J. Japanese Inst. Landsc. Archit.* 58, 177–180.
- Lindemann-Matthies, P., Briegel, R., Schüpbach, B., Junge, X., 2010. Aesthetic preference for a Swiss alpine landscape: The impact of different agricultural land-use with different biodiversity. *Landsc. Urban Plan.* 98, 99–109. doi:10.1016/j.landurbplan.2010.07.015
- Loreau, M., Mouquet, N., Holt, R.D., 2003. Meta-ecosystems: a theoretical framework for a spatial ecosystem ecology. *Ecol. Lett.* 6, 673–679. doi:10.1046/j.1461-0248.2003.00483.x
- Martínez Pastur, G., Peri, P.L., Lencinas, M. V., García-Llorente, M., Martín-López, B., 2016. Spatial patterns of cultural ecosystem services provision in Southern Patagonia. *Landsc. Ecol.* 31, 383–399. doi:10.1007/s10980-015-0254-9
- MEA, 2005. Millenium Ecosystem Assessment. Ecosystems and Human Well-being Synthesis [WWW Document]. URL <http://www.millenniumassessment.org/en/Synthesis.html> (accessed 4.29.16).
- Milcu, A.I., Hanspach, J., Abson, D., Fischer, J., 2013. Cultural ecosystem services: A literature review and prospects for future research. *Ecol. Soc.* 18, 44–88. doi:10.5751/ES-05790-180344
- Nahuelhual, L., Carmona, A., Laterra, P., Barrena, J., Aguayo, M., 2014. A mapping approach to assess intangible cultural ecosystem services: The case of agriculture heritage in Southern Chile. *Ecol. Indic.* 40, 90–101. doi:10.1016/j.ecolind.2014.01.005
- Palomo, I., Martín-López, B., Potschin, M., Haines-Young, R., Montes, C., 2013. National Parks, buffer zones and surrounding lands: Mapping ecosystem service flows. *Ecosyst. Serv.* 4, 104–116. doi:10.1016/j.ecoser.2012.09.001
- Peña, L., Casado-Arzuaga, I., Onaindia, M., 2015. Mapping recreation supply and demand using an ecological and a social evaluation approach. *Ecosyst. Serv.* 13, 108–118. doi:10.1016/j.ecoser.2014.12.008
- Phillips, S.B., Aneja, V.P., Kang, D., Arya, S.P., 2006. Modelling and analysis of the atmospheric nitrogen deposition in North Carolina. *Int. J. Glob. Environ. Issues* 6, 231–252. doi:10.1016/j.ecolmodel.2005.03.026
- Phillips, S.J., Dudík, M., Schapire, R.E., 2004. A Maximum Entropy Approach to Species Distribution Modeling. *Twenty-first Int. Conf. Mach. Learn. - ICML '04* 83–. doi:10.1145/1015330.1015412
- Richards, D.R., Friess, D.A., 2015. A rapid indicator of cultural ecosystem service

- usage at a fine spatial scale: Content analysis of social media photographs. *Ecol. Indic.* 53, 187–195. doi:10.1016/j.ecolind.2015.01.034
- Rösch, V., Tschardtke, T., Scherber, C., Batáry, P., 2013. Landscape composition, connectivity and fragment size drive effects of grassland fragmentation on insect communities. *J. Appl. Ecol.* 50, 387–394. doi:10.1111/1365-2664.12056
- Scolozzi, R., Schirpke, U., Detassia, C., Abdullah, S., Gretter, A., 2014. Mapping Alpine Landscape Values and Related Threats as Perceived by Tourists. *Landsc. Res. Online* (Ju, 1–17. doi:10.1080/01426397.2014.902921
- Sherrouse, B.C., Clement, J.M., Semmens, D.J., 2011. A GIS application for assessing, mapping, and quantifying the social values of ecosystem services. *Appl. Geogr.* 31, 748–760. doi:10.1016/j.apgeog.2010.08.002
- Sherrouse, B.C., Semmens, D.J., Clement, J.M., 2014. An application of Social Values for Ecosystem Services (SolVES) to three national forests in Colorado and Wyoming. *Ecol. Indic.* 36, 68–79. doi:10.1016/j.ecolind.2013.07.008
- Shirai, M., Hirota, M., Ishikawa, H., 2013. A method of Area of Interest and shooting Spot Detection using Geo-tagged photographs. *ACM*.
- Statistics Bureau Ministry of Internal Affairs and Communications, 2016. Social Indicators by Prefecture 2016 [WWW Document]. URL <http://www.stat.go.jp/english/data/shihyou/index.htm> (accessed 12.19.16).
- Sullivan, R.G., Kirchler, L.B., Cothren, J., Winters, S.L., 2013. Offshore Wind Turbine Visibility and Visual Impact Threshold Distance. *Environmental Pract.* 15, 33–49.
- Sullivan, R.G., Kirchler, L.B., Lahti, T., Roche, S., Beckman, K., Cantwell, B., Richmond, P., 2012. Wind Turbine Visibility and Visual Impact Threshold Distances in Western Landscapes. *Argonne Natl. Lab.* 1–47.
- Swets, J.A., 1988. Measuring the accuracy of diagnostic system. *Science* (80-.). 240, 1285–93. doi:10.1126/science.3287615
- T.E.Dilts, 2015. Topography Tools for ArcGIS 10.3.
- Takafumi, H., Hiura, T., 2009. Effects of disturbance history and environmental factors on the diversity and productivity of understory vegetation in a cool-temperate forest in Japan. *For. Ecol. Manage.* 257, 843–857. doi:10.1016/j.foreco.2008.10.020
- TEEB, 2010. The Economics of Ecosystems and Biodiversity: Mainstreaming the Economics of Nature provides a synthesis of the approach, conclusions and recommendations of TEEB.

- TEEB, 2008. An Interim Report. European Communities.
- Tenerelli, P., Demšar, U., Luque, S., 2016. Crowdsourcing indicators for cultural ecosystem services: A geographically weighted approach for mountain landscapes. *Ecol. Indic.* 64, 237–248. doi:10.1016/j.ecolind.2015.12.042
- Uuemaa, E., Mander, Ü., Marja, R., 2013. Trends in the use of landscape spatial metrics as landscape indicators: A review. *Ecol. Indic.* 28, 100–106. doi:10.1016/j.ecolind.2012.07.018
- Van Berkel, D.B., Verburg, P.H., 2014. Spatial quantification and valuation of cultural ecosystem services in an agricultural landscape. *Ecol. Indic.* 37. doi:10.1016/j.ecolind.2012.06.025
- van Zanten, B.T., van Berkel, D.B., Meetemeyer, R.K., Smith, J.W., Tieskens, K.F., Vergurg, P.H., 2016a. Continental scale quantification of landscape values using social media data. *Proc. Natl. Acad. Sci.* 113, 1–7. doi:10.1073/10.1073/pnas.1614158113
- van Zanten, B.T., Verburg, P.H., Scholte, S.S.K., Tieskens, K.F., 2016b. Using choice modeling to map aesthetic values at a landscape scale: Lessons from a Dutch case study. *Ecol. Econ.* 130, 221–231. doi:10.1016/j.ecolecon.2016.07.008
- van Zanten, B.T., Zasada, I., Koetse, M.J., Ungaro, F., Häfner, K., Verburg, P.H., 2016c. A comparative approach to assess the contribution of landscape features to aesthetic and recreational values in agricultural landscapes. *Ecosyst. Serv.* 17, 87–98. doi:10.1016/j.ecoser.2015.11.011
- Wang, C.-P., Wilson, K., Snavely, N., 2013. Accurate georegistration of point clouds using geographic data. *3DTV-Conference, 2013 Int. Conf.* 33–40. doi:10.1109/3DV.2013.13
- WBCSD, 2010. Vision 2050: The new agenda for business, WBCSD. doi:10.1111/j.1530-9290.2009.00117.x
- Wisz, M.S., Hijmans, R.J., Li, J., Peterson, A.T., Graham, C.H., Guisan, A., Elith, J., Dudík, M., Ferrier, S., Huettmann, F., Leathwick, J.R., Lehmann, A., Lohmann, L., Loiselle, B.A., Manion, G., Moritz, C., Nakamura, M., Nakazawa, Y., Overton, J.M., Phillips, S.J., Richardson, K.S., Scachetti-Pereira, R., Schapire, R.E., Soberón, J., Williams, S.E., Zimmermann, N.E., 2008. Effects of sample size on the performance of species distribution models. *Divers. Distrib.* 14, 763–773. doi:10.1111/j.1472-4642.2008.00482.x
- Wolff, S., Schulp, C.J.E., Verburg, P.H., 2015a. Mapping ecosystem services demand:

- A review of current research and future perspectives. *Ecol. Indic.* 55, 159–171.
doi:10.1016/j.ecolind.2015.03.016
- Wolff, S., Schulp, C.J.E., Verburg, P.H., 2015b. Mapping ecosystem services demand:
A review of current research and future perspectives. *Ecol. Indic.* 55, 159–171.
doi:10.1016/j.ecolind.2015.03.016
- Wood, S. a, Guerry, A.D., Silver, J.M., Lacayo, M., 2013. Using social media to
quantify nature-based tourism and recreation. *Sci. Rep.* 3, 2976.
doi:10.1038/srep02976
- WTTC, 2016. Travel & Tourism ECONOMIC IMPACT 2016 WORLD.
- Zandbergen, P. a., Barbeau, S.J., 2011. Positional Accuracy of Assisted GPS Data from
High-Sensitivity GPS-enabled Mobile Phones. *J. Navig.* 64, 381–399.
doi:10.1017/S0373463311000051

Fig. 1 Conceptual diagram of demand mapping

Viewshed score = number of landscape photos/total number of photos, where the number of landscape photos refers to the number of items in the Filter 2 dataset and the total number photos refers to the number of items in the Filter 1 dataset; these are counted within 3 km from every item in the Filter 3 dataset. See also Table S1.

Fig. 2 Demand map of aesthetic value of landscapes. The five ranks are based on 0–1 values normalized by a Fuzzy MS Large function (ArcGIS10.3.1, ESRI Inc.). The most highly demanded areas are ranked 5 (red), and the least demanded areas are ranked 1 (blue). Only tourism resources with at least a B grade as ranked by the Tourism Resources Evaluation Committee in Japan are shown. The grade has four ranks (Special A, A, B, C) mainly based on expert judgment. These grades are decided by the level of attractiveness. Grade Special A is the international level. Grade A is the national level. Grade B is the provincial level. Grade C is the prefectural level.

Fig. 3 Demand map around the Tokachi and Daisetsu Mountain Range (left), and the land-use map (right). Both maps show the same extent. The tourism resources with at least a B grade as ranked by the Tourism Resources Evaluation Committee in Japan are shown. The grade has four ranks (Special A, A, B, C) mainly based on expert judgment. These grades are decided by the level of attractiveness. Grade Special A is the international level. Grade A is the national level. Grade B is the provincial level. Grade C is the prefectural level.

Fig. 4 Areas of demand in each land use type. The stacked chart indicates areas of demand, which were classified by demand rank. The line charts indicate the percentage area of total demand or rank 5.

Fig. 5 Potential supply map. The potential supply area was classified by 10% threshold or equal threshold. The tourism resources with at least a B grade as ranked by the Tourism Resources Evaluation Committee in Japan are shown. The grade has four ranks

(Special A, A, B, C) mainly based on expert judgment. These grades are decided by the level of attractiveness. Grade Special A is the international level. Grade A is the national level. Grade B is the provincial level. Grade C is the prefectural level.

Fig. 6 Areas of demand and potential supply in each land use type. The stacked charts indicate areas of demand or supply potential, which are classified as inside or outside natural parks. The line charts indicate the percentage of demand or potential supply area, to the total area of each land use type.

Fig. 7 Distribution of demand and potential supply area in forest and farmland. The bar chart shows the area of gap area in farmland.

Fig. S1 Land use map study area.

Fig. S2 Sample geotagged photos extracted from Flickr database.

The left photo was taken by Kzaral. “Kushiro Marsh, Kushiro”, 2014, URL: <https://www.flickr.com/photos/32811347@N08/14963096953>, License: CC by 2.0., detail: <https://creativecommons.org/licenses/by/2.0/>.

The right photo was taken by Ryuichi Ikeda. “Biei no oka”, 2010, URL: <https://www.flickr.com/photos/8205548@N08/5007769687>, License: CC by 2.0., detail: <https://creativecommons.org/licenses/by/2.0/>.

Fig. S3 Process flow to map potential supply. MaxEnt generates potential supply based on the relation between the sampled presence data and environmental variables.

Fig. S4 Response curves of environmental variables. Red lines show mean values and blue ranges show standard deviations over 10 trials.

Fig. S5 ROC curve. The AUC value of the model was 0.802.

Table 1 Comparison of supply potential and demand in each natural park.
Percentages are shown in parentheses.

Type	Name	Area (×10 ³ ha)	Demand area (×10 ³ ha)				Potential Supply area (×10 ³ ha)			
			Rank 4 and 5		All ranks		Equal threshold		10% threshold	
National park	Akan	94.1	28.3	(30.1)	50.0	(53.1)	45.6	(48.4)	70.5	(74.9)
	Kushiro Shitsugen	26.6	12.8	(48.2)	18.7	(70.4)	13.8	(51.9)	19.1	(71.8)
	Rishiri Rebun Sarobetsu	21.4	0.6	(2.6)	10.8	(50.5)	14.1	(65.8)	17.5	(82.0)
	Shikotsu Toya	99.7	20.9	(20.9)	43.4	(43.5)	68.4	(68.7)	90.4	(90.7)
	Shiretoko	38.6	4.4	(11.3)	10.6	(27.4)	8.3	(21.6)	19.4	(50.2)
	Daisetsuzan	227.5	17.2	(7.6)	62.8	(27.6)	102.3	(44.9)	161.5	(71.0)
Subtotal		508	84.2	(16.6)	196.3	(38.6)	252.5	(49.7)	378.4	(74.5)
Quasi-national park	Abashiri	37	1.6	(4.4)	28.6	(77.4)	28.9	(78.4)	33.3	(90.2)
	Hidaka Sanmyaku Erimo	104	0.5	(0.5)	4.3	(4.1)	0.7	(0.7)	1.3	(1.2)
	Niseko Syakotan Otaru kaigan	19	1.9	(9.6)	7.5	(39.1)	11.8	(61.7)	15.9	(82.7)
	Oonuma	9	2.8	(30.0)	7.1	(76.4)	7.3	(77.9)	8.7	(93.2)
	Shokanbetsu Teuri Yagishiri	44	0.0	(0.0)	3.8	(8.7)	3.1	(7.2)	12.5	(28.7)
Subtotal		213	6.8	(3.2)	51.2	(24.1)	51.9	(24.4)	71.6	(33.6)
Prefectural natural park	Akkeshi	23	0.0	(0.0)	7.4	(32.0)	1.8	(7.8)	3.2	(13.8)
	Esan	3	0.0	(0.0)	0.4	(12.4)	0.5	(18.5)	1.5	(51.8)
	Furano Ashibetsu	36	1.1	(3.0)	5.8	(16.1)	2.0	(5.5)	8.4	(23.2)
	Hiyama	17	0.0	(0.0)	1.2	(6.7)	0.9	(5.2)	1.8	(10.4)
	Kariba Motta	24	0.0	(0.0)	0.3	(1.5)	7.2	(30.6)	16.0	(68.2)
	Matsumae yakoshi	2	0.0	(0.0)	0.1	(5.0)	0.1	(6.0)	0.3	(15.3)
	Nopporo Shinrin	2	0.0	(1.3)	0.3	(16.9)	1.4	(66.7)	2.0	(95.6)
	Notsukehuuren	13	0.4	(2.8)	1.7	(13.7)	7.6	(60.2)	9.0	(71.3)
	Kita Okhotsuku	4	0.0	(0.0)	0.2	(5.7)	0.2	(5.6)	0.9	(20.8)
	Syaridake	3	0.0	(0.0)	0.3	(8.9)	0.5	(16.6)	1.7	(58.5)
	Syumarinai	14	0.0	(0.0)	1.6	(11.3)	0.0	(0.0)	0.8	(6.0)
	Teshiodake	9	0.0	(0.0)	0.0	(0.0)	0.0	(0.2)	0.7	(7.2)
Subtotal		150	1.5	(1.0)	19.3	(12.9)	22.3	(14.9)	46.3	(30.9)
Total area of natural parks		871	92.5	(10.6)	266.8	(30.6)	326.7	(37.5)	496.3	(57.0)
(% of Hokkaido)		(10.9)	(40.8)	-	(24.4)	-	(19.0)	-	(16.6)	-
Total of Hokkaido		7984	227	(2.8)	1092	(13.7)	1723	(21.6)	2986	(37.4)

Table 2 Importance of environmental variables

Environmental variable	Contribution (%)	Importance (%)
Distance from volcanic topography	23.2	30.2
Type of topography	23.2	11.0
Variety of topography	19.6	12.3
Naturalness	13.3	7.0
Distance from lakes	8.0	9.7
Distance from non-volcanic topography	6.2	10.2
Distance from rivers	2.9	7.0
Distance from coastline	2.8	9.7
Specific riverine landscape	0.8	3.6

Table S1 Photo datasets constructed from the Flickr database

We used three filters to develop photo datasets. The Filter 2 dataset is derived from the Filter 1 dataset, and the Filter 3 dataset is derived from the Filter 2 dataset. Where On Earth Identifier (WOEID) 7153351 is assigned to the Hokkaido region in Japan.

Filter	Criteria	Number of photos	Number of total unique users
1	<p>Photos taken in Hokkaido</p> <p>(WOEID = 7153351)</p> <p>Positional accuracy level ≥ 12</p> <p>Within 500 m of the study area coastline</p>	136,023	2620
2	<p>Filter 1 +</p> <p>Photos have keyword or tag “landscape”</p>	13,202	1156
3	<p>Filter 2 +</p> <p>One photo per user in each municipality</p>	2982	1156

Table S2 Database of environmental variables

Indicator	Layer	Description	Source
Naturalness	Naturalness of vegetation	10 classes of vegetation based on degree of human impact. Bodies of water are not ranked and assigned 0 (urban = 1, high mountain vegetation = 10, etc.)	Natural Environmental Information GIS (Biodiversity Center of Japan, 1998)
Water body	River	Distance from grade 1 or 2 rivers	National Land Numerical Information download service (Japanese Ministry of Land, Infrastructure, Transport and Tourism, 2016)
	Lake	Distance from lakes	
	Coastline	Distance from coastline	
Geological interest	Special landscape with river	Distance from specific riverine landscapes (snaking stream, waterfall, etc.)	
Topography	Type of topography	Ten-class Topography Position Index	Calculated by Topography tool (T.E. Diltz, 2015) and 50-m DEM
	Variety of topography	Shannon's Diversity Index within 1 km from each cell.	Calculated by Land Facet Corridor Designer (Jenness et al., 2013) and topography type
Geological interest	Volcanic topography	Distance from volcanic topography (volcano, caldera, etc.)	National Land Numerical Information download service (Japanese Ministry of Land, Infrastructure, Transport and Tourism, 2016)
	Non-volcanic topography	Distance from mountain range, cirque, moraine etc.	

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5

Fig.6

Fig.7

Forest

		Supply potential(ha)	
		Presence	Absence
Demand (ha)	Presence	86,615 (6.2%)	36,015 (2.6%)
	Absence	287,206 (20.6%)	985,992 (70.6%)

Farmland

		Supply potential(ha)	
		Presence	Absence
Demand (ha)	Presence	66,321 (19.1%)	7,776 (2.2%)
	Absence	153,257 (44.2%)	119,628 (34.5%)

Legend

□ Natural park

□ Region

Demand vs Potential supply

□ No demand and supply potential

■ Only demand

■ Balanced area : demand and potential supply

■ Supply gap area : only potential supply

Area of demand vs potential supply

■ 63,000 ha

Fig. S1

Fig. S2

Fig. S3

Fig.S4 Response of demand to topography

1 Canyons, deeply incised streams	6 Open slopes
2 Midslope drainages, shallow valleys	7 Upper slopes, mesas
3 Upland drainages, headwaters	8 Local ridges, hills in valleys
4 U-shaped valleys	9 Midslope ridges, small hills in plains
5 Plains	10 Mountain tops, high ridges

Fig.S4 Response of demand to water body(Continued)

Fig.S4 Response of demand to naturalness (Continued)

Fig.S5 ROC curve