

HOKKAIDO UNIVERSITY

Title	Accumulation mechanisms of trace metals into Arctic sea ice [an abstract of dissertation and a summary of dissertation review]
Author(s)	Evans, La Kenya Elizabeth
Citation	北海道大学. 博士(環境科学) 甲第13542号
Issue Date	2019-03-25
Doc URL	http://hdl.handle.net/2115/73871
Rights(URL)	https://creativecommons.org/licenses/by-nc-sa/4.0/
Туре	theses (doctoral - abstract and summary of review)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	LaKenya_ElizabethEvans_abstract.pdf (論文内容の要旨)



学 位 論 文 内 容 の 要 旨

博士 (環境科学) 氏 名 Evans La Kenya Elizabeth

学位論文題名

Accumulation mechanisms of trace metals into Arctic sea ice (北極海における海氷への微量金属蓄積メカニズムの解明)

In the last 20 years the Arctic Ocean has experienced over 32% loss of summer sea ice. The loss of sea ice can influence the cycling of biogeochemical materials, affecting seawater's biology and chemistry. Sea ice is important for the supply of biogeochemical materials (trace metals, nutrients, DOM, SPM, etc) to the surface waters of the polar oceans. This thesis will focus on trace metals, as sea ice's role in the biogeochemical cycling of trace metals is not clear. To understand the geochemical behavior of trace metals and their accumulation into sea ice, dissolved (D, $<0.2 \mu$ m), and labile particulate (LP, Total Dissolvable - Dissolved) Fe, Mn, and Cd were examined in sea ice and seawater collected from the Chukchi Sea, Arctic Ocean. Samples were pre-concentrated utilizing the solid-phase extraction NOBIAS Chelate PA-1 resin (Hitachi High-Technologies Corporation) and analyzed on a Graphite Furnace Atomic Absorption Spectrometer. Chukchi seawater showed high percentages for DMn (71.5%) and DCd (66.3%) with a high percentage of LPFe (94.1%). In seawater, DCd was the only metal to correlate with phosphate (R^2 = 0.78) indicating a biogeochemical cycling source. Chukchi seawater concentrations of Fe and Mn may have been controlled through external sources. Sediments (shelf or river) supplied LPFe and LPMn. DFe and DMn were supplied by the Alaskan Coastal Current. Trace metal concentrations in Chukchi drift ice were heterogeneous. Drift ice showed high percentages for the LP fraction (99.2% Fe, 63.6% Mn and 71.2% Cd). This data indicated that, regardless of the trace metal behavior in Chukchi seawater, Chukchi seawater, to accumulate or retain the LP trace metal behavior in Chukchi seawater, Chukchi drift ice was observed to have a preference to accumulate or retain the LP trace metal fraction.

To examine possible trace metal accumulation processes utilized by Arctic sea ice, the association between trace metal concentrations and ice structure were observed in floe ice. The structure of sea ice reflects the process of ice formation, which may aid in the determination of accumulation processes. An Arctic sea ice core was examined. Using photographic analysis for the percentage of pore area and δ^{18} O analysis, sea ice structure in the core sample showed snow, granular, mixed (granular + columnar) and columnar ice. Salinity and nutrients were low, indicating brine drainage and multi-year ice. High trace metal concentrations in snow ice indicated meteoric snow as a source. High concentrations of LPFe in granular ice indicated possible particulate trace metal scavenging by frazil ice. Concentrations of LPMn and LPCd were low compared to DMn and DCd in granular ice. It is possible that reduction of LPFe and LPM after particle entrainment released DMn and DCd, indicating a chemical transformation processes. Low dissolved and labile particulate trace metal concentrations in mixed and columnar ice indicated a release due to brine drainage.

The accumulation and retention of trace metals into Arctic sea ice may be influenced by the type of sea ice. Drift sea ice can be a long-range transporter of LP metals. Floe ice can provide both D and LP metals to the local area. Therefore, sea ice type, processes of sea ice formation, chemical transformation and brine release, are important for the accumulation, retention and release of trace metals from sea ice.