Title	Historical aspects of studies on roles of the inflammasome in the pathogenesis of periodontal diseases
Author(s)	Shibata, Ken-ichiro
Citation	Molecular Oral Microbiology, 33(3), 203-221 https://doi.org/10.1111/omi.12217
Issue Date	2018-06
Doc URL	http://hdl.handle.net/2115/74519
Rights	This is the peer reviewed version of the following article: Historical aspects of studies on roles of the inflammasome in the pathogenesis of periodontal diseases, which has been published in final form at https://doi.org/10.1111/omi.12217 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving
Type	article (author version)
File Information	Mol Oral Microbiol.pdf

[Instructions for use](#)

Historical aspects of studies on roles of the inflammasome in the pathogenesis of periodontal diseases

Ken-ichiro Shibata

Department of Oral Molecular Microbiology, Faculty of Dental Medicine and Graduate School of Dental Medicine, Hokkaido University

SUMMARY

The proinflammatory cytokine interleukin (IL)-1 β is produced as inactive pro-IL-1 β and then processed by caspase-1 to become active. In 2002, it was demonstrated that the intracellular multiprotein complex known as the inflammasome functions as a molecular platform to trigger activation of caspase-1. Inflammasomes are known to function as intracellular sensors for a broad spectrum of various pathogen- and damage-associated molecular patterns.

In 1985, it was demonstrated that *Porphyromonas gingivalis*, a representative bacterium causing chronic periodontitis, induces IL-1 production by murine peritoneal macrophages. Since then, many studies have suggested that IL-1, particularly IL-1 β , plays key roles in the pathogenesis of periodontal diseases. However, the term “inflammasome” was not used until Bostanci *et al.* suggested the involvement of inflammasomes in periodontal disease in 2009. Several subsequent studies on the roles of the inflammasome in the pathogenesis of periodontal diseases have been published. Interestingly, two contradictory reports on the modulation of inflammasomes by *P. gingivalis* have been published. Some papers have described that *P. gingivalis* activates the inflammasome to produce IL-1 β , whereas some stated that *P. gingivalis* inhibits inflammasome activation to subvert immune responses. Several lines of evidence have also been accumulated that the inflammasome activation is modulated by the periodontopathic bacteria other than *P. gingivalis*.

Thus, studies on the roles of inflammasomes in the pathogenesis of periodontal diseases began only 8 years ago and many pathological roles of inflammasomes remain to be clarified.

INTRODUCTION

Interleukin-1 (IL-1), a proinflammatory cytokine, consists of IL- α and IL-1 β . IL-1 β induces inflammatory mediator production, osteoclast formation, matrix metalloproteinase expression, and matrix-producing cell death in periodontal tissues, resulting in the destruction of alveolar bone and periodontal connective tissue¹. Thus, IL-1 β plays crucial roles in the onset and progression of periodontal disease. IL-1 β is produced extracellularly after pro-IL-1 β is processed by caspase-1 to become activated by the intracellular multiprotein complex known as the inflammasome. The word “inflammasome” was first used to describe platforms for inflammatory caspase activation². IL-18 is also produced as pro-IL-18 and processed by caspase-1 to become activated by the inflammasome. The inflammasome is composed of the “nucleotide-binding domain leucine-rich repeat-containing receptor” (NLR), adaptor protein “apoptosis-associated speck-like protein containing a caspase-recruitment domain” (ASC), and procaspase-1³⁻⁵ (Fig. 1). Several types of NLRs are involved in inflammasome activation⁶. Of these inflammasomes, the NLRP3 inflammasome is the most well-studied (Fig. 1) and has been implicated in responses to a broad spectrum of pathogen- and damage-associated molecular patterns⁷⁻²³ (Table 1). We previously demonstrated that whole cells of two *Mycoplasma* species and *Streptococcus sanguinis* activate the NLRP3 inflammasome to produce IL-1 β ^{24,25}.

Periodontitis, a chronic inflammation that occurs in many adults, is a major cause of tooth loss and is characterized by chronic infection associated with Gram-negative anaerobic bacteria in the subgingival biofilm, which leads to destruction of the tissue supporting the teeth. Subgingival biofilms containing several gram-negative rods are strongly involved in the onset and progression of periodontitis. Among gram-negative

rods, *Porphyromonas gingivalis* is a representative periodontal pathogen in chronic periodontitis ²⁶.

In this review, we focus on the historical aspects of the pathological roles of inflammasomes in periodontal diseases and modulation of inflammasome activation by *P. gingivalis* as a first step for developing a new therapeutic strategy.

INVOLVEMENT OF IL-1 IN PERIODONTAL DISEASES

The reported roles of IL-1 in the pathogenesis of periodontal diseases are summarized in Fig. 2.

In 1985, the first paper regarding the functional role of IL-1 in periodontal disease was published ²⁷. However, this study only demonstrated that lipopolysaccharide (LPS) from *P. gingivalis* induces IL-1 production by murine peritoneal macrophages, but did not directly determine the roles of IL-1 in the pathogenesis of adult periodontal disease. The first report on the role of IL-1, particularly IL-1 β , in the pathogenesis of periodontal diseases was published in 1991 ²⁸. The authors demonstrated that the number of cells stained with anti-human IL-1 β was nearly 3-fold higher in periodontally diseased tissue compared to those in normal tissue, suggesting that IL-1 β produced by cells in periodontal tissues is related to the pathological processes associated with periodontal disease. In 1992, the amount of IL-1 in crevicular fluid during human experimental gingivitis was measured and shown to increase rapidly with plaque accumulation and in advance of subsequent gingival inflammation, peaking within 7 days of the start of gingivitis ²⁹. Thus, the authors suggested that IL-1 is an early marker of gingival inflammatory changes. In 1995, it was demonstrated that *in vivo* administration of recombinant IL-1 β accelerates silk ligature-induced alveolar bone resorption in rats ³⁰.

In addition, several reports showed that an IL-1 genotype associated with increased IL-1 production is a strong indicator of the susceptibility to severe periodontitis in adults ³¹⁻³³. In 1997, Ishihara *et al.* demonstrated that the total amount of IL-1 α , IL-1 β , and the total IL-1/IL-1 receptor antagonist ratio (IL-1AI), but not the total IL-1 receptor antagonist (IL-1A), were correlated with alveolar bone loss score; a similar progressive decrease in total IL-1AI was detected in gingival tissue with periodontitis ³⁴. These results suggested that the levels of both crevicular IL-1 and total IL-1AI are closely associated with periodontal disease severity. In 1997, in a *Macaca fascicularis* primate model of experimental periodontitis, IL-1 and tumor necrosis factor (TNF) antagonists, their soluble receptors, inhibited the recruitment of inflammatory cells in close proximity to bone by approximately 80% and reduced osteoclast by 67% at the experimental sites compared to the levels at control sites; bone loss was reduced by 60% ³⁵. Thus, IL-1 and TNF, representative inflammatory cytokines, play important roles in the pathological process of periodontitis. In 1998, in the same primate model, the same group demonstrated that IL-1 and TNF antagonists inhibit the progression of inflammatory cell infiltration toward alveolar bone ³⁶. This research group also demonstrated that loss of connective tissue attachment and progression of periodontal disease can be retarded by antagonists to IL-1 and TNF in the same primate model ³⁷.

Based on these reports, a previous review described the roles of IL-1 and TNF in periodontal tissue destruction ¹. These cytokines induce adhesion molecules and other mediators that facilitate and amplify the inflammatory response, stimulation of matrix metalloproteinase, and bone resorption.

INVOLVEMENT OF INFLAMMASOMES IN PERIODONTAL DISEASES

As described above, the inflammasome was first named in 2002 as a molecular platform triggering activation of caspase-1 and processing of proIL-1 β and proIL-18². Therefore, we next focus on studies of the roles of the inflammasome in the pathogenesis of periodontal diseases.

In 2009, 7 years after the discovery of the inflammasome, it was first demonstrated that NLRP3 and NLRP2, but not ASC, are expressed at significantly higher levels in gingival tissues of subjects with inflammatory periodontal disease than in healthy subjects, and a positive correlation was found between NLRP3 and IL-1 β or IL-18 expression levels in these tissues³⁸. In addition, *in vitro* data demonstrated that *P. gingivalis* upregulates the expression of NLRP3, IL-1 β , and IL-18 in a human monocytic cell line (Mono-Mac-6), but downregulates NLRP2 and ASC. Thus, the authors suggested that the NLRP3 inflammasome is involved in the pathogenesis of periodontal diseases.

In 2011, Bostanci *et al.*³⁹ demonstrated that supragingival and subgingival biofilms differentially regulate the gene expressions of NLRP3 and AIM2 inflammasomes and their down-stream IL-1 targets. Briefly, they found that the culture supernatant from supragingival biofilm containing 6-species of supragingival Zurich biofilm model⁴⁰ grown on the hydroxyapatite disc increased the expression of caspase-1, ASC, AIM2 as well as IL-1 β and IL-18, but did not upregulate NLRP3 expression. However, the culture supernatant from subgingival biofilm containing 10-species of subgingival Zurich biofilm model⁴¹ grown on the hydroxyapatite disc upregulated caspase-1, ASC, AIM2, IL-1 β and IL-18 gene expression at lower concentrations, followed by their down-regulation at higher concentrations, which was also evident for NLRP3 expression. They suggest that upregulation of transcription of inflammasome

components by supragingival biofilms correlates with early inflammatory events in periodontal disease, whereas the downregulation of the transcription by subgingival biofilms is favorable for the survival and persistence of periodontopathic bacteria including *P. gingivalis*.

In 2012, it was demonstrated that production of prostaglandin E2 does not require the inflammasome adaptor function of ASC, but was dependent on mitogen-activated protein kinase activation and that the mitogen-activated protein kinase kinase kinase CARD domain-containing protein RIPK2 was induced by *P. gingivalis* in an ASC-dependent manner⁴². Thus, they found that the inflammasome adaptor ASC-dependent RIP2 kinase regulates reduced prostaglandin E2 production in chronic periodontitis, although they did not demonstrate the direct involvement of the inflammasome in periodontal disease.

In 2014, novel findings suggested a strong correlation between the NLRP3 inflammasome and severe periodontitis⁴³. They demonstrated that blocking the functions of CD24, a negative regulator of inflammation in periodontal tissues⁴⁴⁻⁴⁷, by its antibody-enhanced expression of NLRP3 together with the co-activators ASC and caspase-1, resulting in burst release of activated IL-18, which plays important roles in inflammation. Additionally, subjects with mild chronic periodontitis showed increased titers of serum antibodies that were auto-reactive with CD24 compared with in subjects with severe periodontitis. Thus, the negative regulator CD24 upregulates NLRP3 in periodontal disease, strongly suggesting that the NLRP3 inflammasome plays pathological roles in periodontal diseases.

In 2015, it was demonstrated that expression of NLRP3 and IL-1 β in the oral gingival epithelium of patients with chronic periodontitis and/or type 2 diabetes mellitus

was significantly upregulated compared to in control individuals, and simultaneous stimulation with LPS and high glucose contributed to significant upregulation of NLRP3 expression compared to stimulation with LPS or high glucose alone⁴⁸. These results suggested that for patients with type 2 diabetes mellitus and chronic periodontitis, a hyperglycemic status may exacerbate the inflammation state of gingival tissue by activating the NLRP3 inflammasome pathway.

In 2015, it was demonstrated that the intensity of NLRP3 expression was significantly higher in chronic periodontitis or generalized aggressive periodontitis than in healthy tissue, and a more significant difference was observed in the periodontal epithelium layer⁴⁹. In addition, they showed that NLRP1 was minimally expressed in healthy and periodontitis gingival tissues, whereas absent in melanoma 2 (AIM2) was expressed at a higher level in the chronic periodontitis group than in other subjects. NLRP1 and AIM2 are also known to mediate inflammasome assembly against microbial infections^{9,10}.

In 2017, the study was carried out to compare salivary levels of NLRP3, ASC, caspase-1 and IL-1 β from patients with aggressive periodontitis (AgP) or chronic periodontitis (CP) and periodontally healthy controls (HC) as well as elucidate its association with periodontal clinical status⁵⁰. The study indicates the possibility that salivary levels of NLRP3, ASC, and IL-1 β , but not caspase-1, act as strong/independent indicators of amount and extent of periodontal breakdown in both CP and AgP.

MODULATION OF THE INFLAMMASOME BY THE PERIODONTOPATHIC BACTERIA *P. gingivalis*

There are several papers which describe modulation of inflammasomes by

periodontopathic bacteria such as *Aggregatibacter actinomycetemcomitans*⁵¹⁻⁵⁴, *Fusobacterium nucleatum*⁵⁵ other than *P. gingivalis*. In this review, we focus on involvement of the representative periodontopathic bacterium *P. gingivalis* in inflammasome modulation. There have been two contradictory reports on modulation of the inflammasomes by *P. gingivalis*, a representative periodontopathic bacteria. Some papers demonstrated that *P. gingivalis* activates the inflammasome, which plays pathological roles in periodontal diseases⁵⁶⁻⁶⁷ (Table 2). In contrast, some papers demonstrated that *P. gingivalis* inhibits inflammasomes to subvert immune responses⁶⁸⁻⁷¹ (Table 3). Olsen and Yilmaz⁷² have already published the review which discusses in-depth on several molecular mechanisms by which *P. gingivalis* modulates innate immunity by limiting the activation of the NLRP3 inflammasome. In addition, Almeida-da-Silva *et al.*⁷³ have published the review which mainly describes modulation of the ability of *P. gingivalis* to evade the immune responses by purinerigic signaling known as the second signal for the NLRP3 inflammasome activation. Therefore, this review focuses on historical flow of studies on modulation of inflammasomes by *P. gingivalis*.

1. ACTIVATION OF INFLAMMASOMES

As described above, in 1985, *P. gingivalis* LPS was shown to induce IL-1 production by using murine peritoneal macrophages²⁷. At that time, however, it was not known that the intracellular sensor inflammasome regulates the production of IL-1 β . In addition, it was demonstrated that ASC, one component of the inflammasome complex, is involved in inducing cytokines such as IL-1 β in response to *P. gingivalis* infection via a caspase-1-dependent pathway⁵⁶, although the term “inflammasome” was not used.

Activation of the inflammasome by *P. gingivalis* was reported in a study showing that treatment of gingival epithelial cells infected with *P. gingivalis* induced expression of the IL-1 β gene and intracellular accumulation of IL-1 β protein, whereas IL-1 β was not secreted unless infected cells were subsequently stimulated with ATP, as well as that knockdown of NLRP3 by siRNA attenuated the ability of ATP to induce IL-1 β secretion in infected cells⁵⁷ (Table 2). Similar findings have been reported in another study⁵⁸. They also demonstrated that activation of the NLRP3 inflammasome does not rely on *P. gingivalis* infection unless stimulated by *P. gingivalis* LPS and/or extracellular ATP. Thus, these findings suggest that the NLRP3 inflammasome is an important mediator of the inflammatory response in the gingival epithelium.

Several lines of evidence⁵⁹⁻⁶¹ support the finding that periodontitis is associated pathologically with atherosclerotic vascular disease, although the details remain unclear. It was recently found that macrophages primed with *P. gingivalis* LPS and then treated with cholesterol crystals released IL-1 β ⁶².

In 2014, it was demonstrated that expression levels of inflammasome components in gingival tissues from patients with chronic periodontitis are much higher than in those from healthy controls and that *P. gingivalis* induces the activation of NLRP3 and AIM2 inflammasome via TLR2 and TLR4 signaling, leading to IL-1 β secretion and pyroptotic cell death⁶³. In addition, studies suggested that *P. gingivalis*-induced NLRP3 inflammasome activation depends on ATP release, K⁺ efflux, and cathepsin B.

In 2015, it was reported that *P. gingivalis* can activate the NLRP3 inflammasome, in which the pathogenic factors gingipain and fimbriae play important roles, suggesting that the NLRP3 inflammasome plays a critical role in periodontal disease and atherosclerosis induced by *P. gingivalis* challenge through sustained inflammation⁶⁴. In

2017, the authors of the previous study also demonstrated that oral injection of *P. gingivalis* to wild-type mice, but not to NLRP3-deficient mice, significantly increased alveolar bone loss and gingival gene expression of pro-IL-1 β and proIL-18⁶⁵. In addition, they showed that *P. gingivalis* upregulated production of IL-1 β and IL-18 by peritoneal macrophages from wild-type mice, but not from NLRP3-deficient mice.

In 2017, Yoshida *et al.*⁷⁴ have reported that *P. gingivalis* infection activated the double-stranded RNA-dependent kinase (PKR) in osteoblasts, which in turn upregulated NLRP3 expression through activation of NF- κ B. Fleetwood *et al.*⁷⁵ have reported that treatment of murine bone marrow-derived macrophages (BMMs) with live *P. gingivalis*, heat-killed *P. gingivalis* and outer membrane vesicles (OMVs) upregulate the expression of NLRP3, procaspase-1 and pro-IL-1 β and that treatment with OMVs, and to a lesser extent heat-inactivated OMVs induces cleavage of procaspase-1 and pro-IL-1 β into their mature forms as well as inducing IL-1 β and IL-18, whereas treatment with live *P. gingivalis* and heat-killed *P. gingivalis* do not. That is, they suggest that OMVs play more important roles in the activation of the NLRP3 inflammasome than *P. gingivalis* cells. In addition, Cecil *et al.*⁷⁶ have also reported that OMVs of *P. gingivalis*, *Treponema denticola* and *Tannerella forthia* activate inflammasomes in a human monocytic THP-1 cells and its subsets to induce IL-1 β secretion and ASC speck formation and NLRP3 and/or AIM2 inflammasome(s) in murine BMMs to induce IL-1 β secretion.

There is growing evidence of a relationship between diabetes and inflammasome activation⁶⁶. Recently, infection by *P. gingivalis*, a major bacterial species in periodontal disease, was recognized as a common complication of diabetes. It has been demonstrated that human gingival fibroblasts infected with *P. gingivalis* grown in BHI

medium containing high glucose showed increased expression of IL-1 β and NLRP3⁶⁷.

Thus, these publications suggest that activation of the inflammasome by *P. gingivalis* leads to IL-1 β production and plays important roles in the pathogenicity of diabetes and atherosclerotic vascular disease as well as periodontal diseases.

2. INHIBITION OF INFLAMMASOME ACTIVATION

Several lines of evidence suggest that *P. gingivalis* inhibits inflammasome activation, leading to evasion of the host immune response (Table 3).

In 2012, it was reported that *P. gingivalis* suppresses inflammasome activation by another periodontal bacterium such as *F. nucleatum*, but not by ATP or nigericin⁶⁸. Based on the finding that *P. gingivalis* limits both the number of cells taking up beads and number of beads taken up for bead-positive cells, they speculated that the mechanism by which *P. gingivalis* inhibits inflammasome activation contributes to suppression of bacterial endocytosis.

In 2013, it was reported that 10 species from subgingival biofilms, including *P. gingivalis*, reduce NLRP3 and IL-1 β levels but do not affect AIM2 expression in human gingival fibroblasts and that exclusion of *P. gingivalis* from the biofilm partially rescued NLRP3 and IL-1 β expression⁶⁹. These results suggest that subgingival biofilms down-regulate NLRP3 and IL-1 β expression, partly because of the existence of *P. gingivalis*. Therefore, these dampened host innate immune responses may favor the survival and persistence of associated biofilm species in periodontal tissues.

A recent study found that treatment of endothelial cells with live cells, but not with heat-killed cells, of *P. gingivalis* induced proteolysis of NLRP3, but this proteolysis was not observed after ATP pre-treatment and/or *P. gingivalis*-LPS stimulation⁷⁰.

Additionally, the levels of secreted IL-1 β significantly increased after ATP pre-treatment and/or *P. gingivalis*-LPS stimulation, but not after *P. gingivalis* infection. These data demonstrate that *P. gingivalis* and its LPS differentially controlled the NLRP3 inflammasome pathway in endothelial cells, suggesting a novel potential mechanism developed by *P. gingivalis* to reduce IL-1 β secretion and escape the host immune response.

As described above, *P. gingivalis* stimulates pro-IL-1 β synthesis but not mature IL-1 β secretion, unless the P2X₇ receptor is activated by extracellular ATP⁵⁷. An additional study demonstrated that fimbriae dampens P2X₇-dependent IL-1 β secretion⁷¹. Briefly, the authors showed that when bone marrow-derived macrophages from wild-type or P2X₇-deficient mice were infected with *P. gingivalis* 381 or the isogenic fimbria-deficient (DPG3) strain with or without subsequent ATP stimulation, DPG3 induced higher IL-1 β secretion after ATP stimulation compared to 381 in wild-type bone marrow-derived macrophages, but not in P2X₇-deficient cells.

Recently, it was demonstrated that a nucleoside-diphosphate kinase of *P. gingivalis* inhibits caspase-1 activation and IL-1 β secretion in gingival epithelial cells by downregulating the ATP-P2X₇ signaling pathway and reducing the release of the high mobility group box 1 protein, a pro-inflammatory danger signal⁷⁷.

Thus, these publications suggest that *P. gingivalis* inhibits inflammasome activation to subvert host immune responses.

SUMMARY AND CONCLUSIONS

IL-1 β is produced as biologically inactive pro-IL-1 β and then processed by caspase-1,

also known as IL-1 β -converting enzyme, to active IL-1 β ⁷⁸. In 2002, it was first demonstrated that the intracellular multiprotein complex known as the inflammasome functions as a molecular platform that triggers activation of caspase-1 ². Inflammasomes are intracellular sensors that drive host immune responses. However, excessive inflammasome activity can be harmful because gain-of-function mutations in inflammasome components are associated with autoimmune and autoinflammatory disorders ^{3,4,79}.

In 1985, LPS from *P. gingivalis*, a representative bacterium in chronic periodontitis, was reported to induce IL-1 production by murine peritoneal macrophage ²⁷. After this report, many studies suggested that IL-1, particularly IL-1 β , plays key roles in the pathogenesis of periodontal diseases. Taxman *et al.* demonstrated that ASC, one component of the inflammasome complex, is involved in inducing cytokines such as IL-1 β by *P. gingivalis* via a caspase-1-dependent pathway ⁵⁶. However, they did not use the term “inflammasome,” regardless of the discovery of inflammasome. In 2009, 7 years after the discovery of inflammasomes, Bostanci *et al.* first demonstrated that inflammasome activation plays important roles in periodontal disease ³⁸. Since then, several studies on the roles of the inflammasome in the pathogenesis of periodontal diseases have been published. Of these reports, there are two contradictory reports regarding the modulation of inflammasomes by *P. gingivalis*, a representative bacterium in chronic periodontitis. Some papers described that *P. gingivalis* activates the inflammasome, whereas some papers described that *P. gingivalis* inhibits inflammasome activation. The discrepancy between studies may be attributed to the different cell types used in the studies or to differences between a live single-type bacterial infection and biofilm infection. In addition, many pathogenic bacteria such as

*Yersinia species*⁸⁰, *Legionella pneumophila*⁸¹, *Pseudomonas aeruginosa*⁸², and *Mycobacterium tuberculosis*⁸³ inhibit inflammasome activation^{3,23}, although the mechanisms differ from each other.

Inflammasomes are intracellular sensors that drive host immune responses to maintain homeostasis of host cells and tissues. Thus, the activity of *P. gingivalis* in inhibiting inflammasome activation is more strongly associated with chronic inflammation such as adult chronic periodontitis. Inhibition of inflammasome activation by *P. gingivalis* may allow other bacteria in the subgingival dental plaque as well as the bacterium to survive for longer periods of time in the gingival tissue and contribute to periodontal diseases.

As described above, studies on the roles of inflammasomes in the pathogenesis of periodontal diseases began only 8 years ago. In the oral cavity, there are teeth embedded in gingival tissue, which are mainly made of phosphate and calcium, and sometimes several artificial dental prostheses to restore intraoral defects, including resins, ceramics and metals such as titanium, cobalt and chromium, which is extremely different from nasal, stomach and gut cavities. More Recently, interestingly, it was found that titanium ions also stimulate inflammasome activation⁸⁴. Thus, many unknown pathological roles of inflammasomes in various oral diseases including periodontal diseases remain to be clarified.

CONFLICT OF INTEREST

The author has no financial conflicts of interest.

REFERENCES

1. Graves DT, Cochran D. The contribution of interleukin-1 and tumor necrosis factor to periodontal tissue destruction. *J Periodontol.* 2003;74(3):391-401.
2. Martinon F, Burns K, Tschopp J. The inflammasome: a molecular platform triggering activation of inflammatory caspases and processing of proIL-beta. *Mol Cell.* 2002;10(2):417-426.
3. Lamkanfi M, Dixit VM. Inflammasomes and their roles in health and disease. *Annu Rev Cell Dev Biol.* 2012;28:137-161.
4. Mariathasan S, Monack DM. Inflammasome adaptors and sensors: intracellular regulators of infection and inflammation. *Nat Rev Immunol.* 2007;7(1):31-40.
5. Martinon F, Mayor A, Tschopp J. The inflammasomes: guardians of the body. *Annu Rev Immunol.* 2009;27:229-265.
6. Davis BK, Wen H, Ting JP. The inflammasome NLRs in immunity, inflammation, and associated diseases. *Annu Rev Immunol.* 2011;29:707-735.
7. Duncan JA, Gao X, Huang MT, et al. . *Neisseria gonorrhoeae* activates the proteinase cathepsin B to mediate the signaling activities of the NLRP3 and ASC-containing inflammasome. *J Immunol.* 2009;182(10):6460-6469.
8. He X, Mekasha S, Mavrogiorgos N, Fitzgerald KA, Lien E, Ingalls RR. Inflammation and fibrosis during *Chlamydia pneumoniae* infection is regulated by IL-1 and the NLRP3/ASC inflammasome. *J Immunol.* 2010;184(10):5743-5754.
9. Vladimer GI, Marty-Roix R, Ghosh S, Weng D, Lien E. Inflammasomes and host defenses against bacterial infections. *Curr Opin Microbiol.*

2013;16(1):23-31.

10. Franchi L, Munoz-Planillo R, Nunez G. Sensing and reacting to microbes through the inflammasomes. *Nat Immunol.* 2012;13(4):325-332.
11. Duewell P, Kono H, Rayner KJ, et al. . NLRP3 inflammasomes are required for atherogenesis and activated by cholesterol crystals. *Nature.* 2010;464(7293):1357-1361.
12. Franchi L, Kanneganti TD, Dubyak GR, Nunez G. Differential requirement of P2X7 receptor and intracellular K⁺ for caspase-1 activation induced by intracellular and extracellular bacteria. *J Biol Chem.* 2007;282(26):18810-18818.
13. Petrilli V, Papin S, Dostert C, Mayor A, Martinon F, Tschopp J. Activation of the NALP3 inflammasome is triggered by low intracellular potassium concentration. *Cell Death Differ.* 2007;14(9):1583-1589.
14. Shimada K, Crother TR, Karlin J, et al. . Oxidized mitochondrial DNA activates the NLRP3 inflammasome during apoptosis. *Immunity.* 2012;36(3):401-414.
15. Zhou R, Yazdi AS, Menu P, Tschopp J. A role for mitochondria in NLRP3 inflammasome activation. *Nature.* 2011;469(7329):221-225.
16. Shenoy AR, Wellington DA, Kumar P, et al. . GBP5 promotes NLRP3 inflammasome assembly and immunity in mammals. *Science.* 2012;336(6080):481-485.
17. Lu B, Nakamura T, Inouye K, et al. . Novel role of PKR in inflammasome activation and HMGB1 release. *Nature.* 2012;488(7413):670-674.
18. Sander LE, Davis MJ, Boekschoten MV, et al. . Detection of prokaryotic mRNA signifies microbial viability and promotes immunity. *Nature.* 2011;474(7351):385-389.

19. Vyleta ML, Wong J, Magun BE. Suppression of ribosomal function triggers innate immune signaling through activation of the NLRP3 inflammasome. *PLoS One*. 2012;7(5):e36044.
20. Martinon F, Agostini L, Meylan E, Tschopp J. Identification of bacterial muramyl dipeptide as activator of the NALP3/cryopyrin inflammasome. *Curr Biol*. 2004;14(21):1929-1934.
21. Marina-Garcia N, Franchi L, Kim YG, et al. . Pannexin-1-mediated intracellular delivery of muramyl dipeptide induces caspase-1 activation via cryopyrin/NLRP3 independently of Nod2. *J Immunol*. 2008;180(6):4050-4057.
22. Kanneganti TD, Ozoren N, Body-Malapel M, et al. . Bacterial RNA and small antiviral compounds activate caspase-1 through cryopyrin/Nalp3. *Nature*. 2006;440(7081):233-236.
23. von Moltke J, Ayres JS, Kofoed EM, Chavarria-Smith J, Vance RE. Recognition of bacteria by inflammasomes. *Annu Rev Immunol*. 2013;31:73-106.
24. Sugiyama M, Saeki A, Hasebe A, et al. . Activation of inflammasomes in dendritic cells and macrophages by *Mycoplasma salivarium*. *Mol Oral Microbiol*. 2016;31(3):259-269.
25. Saeki A, Suzuki T, Hasebe A, et al. . Activation of nucleotide-binding domain-like receptor containing protein 3 inflammasome in dendritic cells and macrophages by *Streptococcus sanguinis*. *Cell Microbiol*. 2017;19(3).
26. Bostanci N, Belibasakis GN. *Porphyromonas gingivalis*: an invasive and evasive opportunistic oral pathogen. *FEMS Microbiol Lett*. 2012;333(1):1-9.
27. Hanazawa S, Nakada K, Ohmori Y, Miyoshi T, Amano S, Kitano S. Functional role of interleukin 1 in periodontal disease: induction of interleukin 1 production

- by *Bacteroides gingivalis* lipopolysaccharide in peritoneal macrophages from C3H/HeN and C3H/HeJ mice. *Infect Immun.* 1985;50(1):262-270.
28. Jandinski JJ, Stashenko P, Feder LS, et al. . Localization of interleukin-1 beta in human periodontal tissue. *J Periodontol.* 1991;62(1):36-43.
 29. Kinane DF, Winstanley FP, Adonogianaki E, Moughal NA. Bioassay of interleukin 1 (IL-1) in human gingival crevicular fluid during experimental gingivitis. *Arch Oral Biol.* 1992;37(2):153-156.
 30. Koide M, Suda S, Saitoh S, et al. . In vivo administration of IL-1 beta accelerates silk ligature-induced alveolar bone resorption in rats. *J Oral Pathol Med.* 1995;24(9):420-434.
 31. Thomson WM, Edwards SJ, Dobson-Le DP, et al. . IL-1 genotype and adult periodontitis among young New Zealanders. *J Dent Res.* 2001;80(8):1700-1703.
 32. Laine ML, Farre MA, Gonzalez G, et al. . Polymorphisms of the interleukin-1 gene family, oral microbial pathogens, and smoking in adult periodontitis. *J Dent Res.* 2001;80(8):1695-1699.
 33. Kornman KS, Crane A, Wang HY, et al. . The interleukin-1 genotype as a severity factor in adult periodontal disease. *J Clin Periodontol.* 1997;24(1):72-77.
 34. Ishihara Y, Nishihara T, Kuroyanagi T, et al. . Gingival crevicular interleukin-1 and interleukin-1 receptor antagonist levels in periodontally healthy and diseased sites. *J Periodontal Res.* 1997;32(6):524-529.
 35. Assuma R, Oates T, Cochran D, Amar S, Graves DT. IL-1 and TNF antagonists inhibit the inflammatory response and bone loss in experimental periodontitis. *J Immunol.* 1998;160(1):403-409.

36. Graves DT, Delima AJ, Assuma R, Amar S, Oates T, Cochran D. Interleukin-1 and tumor necrosis factor antagonists inhibit the progression of inflammatory cell infiltration toward alveolar bone in experimental periodontitis. *J Periodontol.* 1998;69(12):1419-1425.
37. Delima AJ, Oates T, Assuma R, et al. . Soluble antagonists to interleukin-1 (IL-1) and tumor necrosis factor (TNF) inhibits loss of tissue attachment in experimental periodontitis. *J Clin Periodontol.* 2001;28(3):233-240.
38. Bostanci N, Emingil G, Saygan B, et al. . Expression and regulation of the NALP3 inflammasome complex in periodontal diseases. *Clin Exp Immunol.* 2009;157(3):415-422.
39. Bostanci N, Meier A, Guggenheim B, Belibasakis GN. Regulation of NLRP3 and AIM2 inflammasome gene expression levels in gingival fibroblasts by oral biofilms. *Cell Immunol.* 2011;270(1):88-93.
40. Shapiro S, Giertsen E, Guggenheim B. An in vitro oral biofilm model for comparing the efficacy of antimicrobial mouthrinses. *Caries Res.* 2002;36(2):93-100.
41. Guggenheim B, Gmur R, Galicia JC, et al. . In vitro modeling of host-parasite interactions: the 'subgingival' biofilm challenge of primary human epithelial cells. *BMC Microbiol.* 2009;9:280.
42. Taxman DJ, Lei Y, Zhang S, Holley-Guthrie E, Offenbacher S, Ting JP. ASC-dependent RIP2 kinase regulates reduced PGE2 production in chronic periodontitis. *J Dent Res.* 2012;91(9):877-882.
43. Guo W, Ye P, Yu H, Liu Z, Yang P, Hunter N. CD24 activates the NLRP3 inflammasome through c-Src kinase activity in a model of the lining epithelium

- of inflamed periodontal tissues. *Immun Inflamm Dis*. 2014;2(4):239-253.
44. Ye P, Simonian M, Nadkarni MA, Decarlo AA, Chapple CC, Hunter N. Identification of epithelial auto-antigens associated with periodontal disease. *Clin Exp Immunol*. 2005;139(2):328-337.
 45. Chen GY, Tang J, Zheng P, Liu Y. CD24 and Siglec-10 selectively repress tissue damage-induced immune responses. *Science*. 2009;323(5922):1722-1725.
 46. Ye P, Nadkarni MA, Simonian M, Hunter N. CD24 regulated gene expression and distribution of tight junction proteins is associated with altered barrier function in oral epithelial monolayers. *BMC Cell Biol*. 2009;10:2.
 47. Ye P, Yu H, Simonian M, Hunter N. Ligation of CD24 expressed by oral epithelial cells induces kinase dependent decrease in paracellular permeability mediated by tight junction proteins. *Biochem Biophys Res Commun*. 2011;412(1):165-169.
 48. Huang X, Yang X, Ni J, et al. . Hyperglucose contributes to periodontitis: involvement of the NLRP3 pathway by engaging the innate immunity of oral gingival epithelium. *J Periodontol*. 2015;86(2):327-335.
 49. Xue F, Shu R, Xie Y. The expression of NLRP3, NLRP1 and AIM2 in the gingival tissue of periodontitis patients: RT-PCR study and immunohistochemistry. *Arch Oral Biol*. 2015;60(6):948-958.
 50. Isaza-Guzman DM, Medina-Piedrahita VM, Gutierrez-Henao C, Tobon-Arroyave SI. Salivary Levels of NLRP3 Inflammasome-Related Proteins as Potential Biomarkers of Periodontal Clinical Status. *J Periodontol*. 2017;88(12):1329-1338.
 51. Kim S, Park MH, Song YR, Na HS, Chung J. Aggregatibacter

- actinomycetemcomitans-Induced AIM2 Inflammasome Activation Is Suppressed by Xylitol in Differentiated THP-1 Macrophages. *J Periodontol.* 2016;87(6):e116-126.
52. Zhao P, Liu J, Pan C, Pan Y. NLRP3 inflammasome is required for apoptosis of *Aggregatibacter actinomycetemcomitans*-infected human osteoblastic MG63 cells. *Acta Histochem.* 2014;116(7):1119-1124.
 53. Belibasakis GN, Johansson A. *Aggregatibacter actinomycetemcomitans* targets NLRP3 and NLRP6 inflammasome expression in human mononuclear leukocytes. *Cytokine.* 2012;59(1):124-130.
 54. Shenker BJ, Ojcius DM, Walker LP, Zekavat A, Scuron MD, Boesze-Battaglia K. *Aggregatibacter actinomycetemcomitans* cytolethal distending toxin activates the NLRP3 inflammasome in human macrophages, leading to the release of proinflammatory cytokines. *Infect Immun.* 2015;83(4):1487-1496.
 55. Hung SC, Huang PR, Almeida-da-Silva CLC, Atanasova KR, Yilmaz O, Ojcius DM. NLRX1 modulates differentially NLRP3 inflammasome activation and NF-kappaB signaling during *Fusobacterium nucleatum* infection. *Microbes Infect.* 2017.
 56. Taxman DJ, Zhang J, Champagne C, et al. . Cutting edge: ASC mediates the induction of multiple cytokines by *Porphyromonas gingivalis* via caspase-1-dependent and -independent pathways. *J Immunol.* 2006;177(7):4252-4256.
 57. Yilmaz O, Sater AA, Yao L, Koutouzis T, Pettengill M, Ojcius DM. ATP-dependent activation of an inflammasome in primary gingival epithelial cells infected by *Porphyromonas gingivalis*. *Cell Microbiol.*

2010;12(2):188-198.

58. Guo W, Wang P, Liu Z, Yang P, Ye P. The activation of pyrin domain-containing-3 inflammasome depends on lipopolysaccharide from *Porphyromonas gingivalis* and extracellular adenosine triphosphate in cultured oral epithelial cells. *BMC Oral Health*. 2015;15(1):133.
59. Pussinen PJ, Tuomisto K, Jousilahti P, Havulinna AS, Sundvall J, Salomaa V. Endotoxemia, immune response to periodontal pathogens, and systemic inflammation associate with incident cardiovascular disease events. *Arterioscler Thromb Vasc Biol*. 2007;27(6):1433-1439.
60. Lockhart PB, Bolger AF, Papapanou PN, et al. . Periodontal disease and atherosclerotic vascular disease: does the evidence support an independent association?: a scientific statement from the American Heart Association. *Circulation*. 2012;125(20):2520-2544.
61. Haraszthy VI, Zambon JJ, Trevisan M, Zeid M, Genco RJ. Identification of periodontal pathogens in atheromatous plaques. *J Periodontol*. 2000;71(10):1554-1560.
62. Champaiboon C, Poolgesorn M, Wisitrasameewong W, Sa-Ard-Iam N, Rerkyen P, Mahanonda R. Differential inflammasome activation by *Porphyromonas gingivalis* and cholesterol crystals in human macrophages and coronary artery endothelial cells. *Atherosclerosis*. 2014;235(1):38-44.
63. Park E, Na HS, Song YR, Shin SY, Kim YM, Chung J. Activation of NLRP3 and AIM2 inflammasomes by *Porphyromonas gingivalis* infection. *Infect Immun*. 2014;82(1):112-123.
64. Yamaguchi Y, Kurita-Ochiai T, Kobayashi R, Suzuki T, Ando T. Activation of

- the NLRP3 inflammasome in Porphyromonas gingivalis-accelerated atherosclerosis. *Pathog Dis.* 2015;73(4).
65. Yamaguchi Y, Kurita-Ochiai T, Kobayashi R, Suzuki T, Ando T. Regulation of the NLRP3 inflammasome in Porphyromonas gingivalis-accelerated periodontal disease. *Inflamm Res.* 2017;66(1):59-65.
 66. Lee HM, Kim JJ, Kim HJ, Shong M, Ku BJ, Jo EK. Upregulated NLRP3 inflammasome activation in patients with type 2 diabetes. *Diabetes.* 2013;62(1):194-204.
 67. Kuo HC, Chang LC, Chen TC, et al. . Sterol Regulatory Element-Binding Protein-1c Regulates Inflammasome Activation in Gingival Fibroblasts Infected with High-Glucose-Treated Porphyromonas gingivalis. *Front Cell Infect Microbiol.* 2016;6:195.
 68. Taxman DJ, Swanson KV, Broglie PM, et al. . Porphyromonas gingivalis mediates inflammasome repression in polymicrobial cultures through a novel mechanism involving reduced endocytosis. *J Biol Chem.* 2012;287(39):32791-32799.
 69. Belibasakis GN, Guggenheim B, Bostanci N. Down-regulation of NLRP3 inflammasome in gingival fibroblasts by subgingival biofilms: involvement of Porphyromonas gingivalis. *Innate Immun.* 2013;19(1):3-9.
 70. Huck O, Elkaim R, Davideau JL, Tenenbaum H. Porphyromonas gingivalis-impaired innate immune response via NLRP3 proteolysis in endothelial cells. *Innate Immun.* 2015;21(1):65-72.
 71. Morandini AC, Ramos-Junior ES, Potempa J, et al. . Porphyromonas gingivalis fimbriae dampen P2X7-dependent interleukin-1beta secretion. *J Innate Immun.*

- 2014;6(6):831-845.
72. Olsen I, Yilmaz O. Modulation of inflammasome activity by *Porphyromonas gingivalis* in periodontitis and associated systemic diseases. *J Oral Microbiol.* 2016;8:30385.
 73. Almeida-da-Silva CL, Morandini AC, Ulrich H, Ojcius DM, Coutinho-Silva R. Purinergic signaling during *Porphyromonas gingivalis* infection. *Biomed J.* 2016;39(4):251-260.
 74. Yoshida K, Okamura H, Hiroshima Y, et al. . PKR induces the expression of NLRP3 by regulating the NF-kappaB pathway in *Porphyromonas gingivalis*-infected osteoblasts. *Exp Cell Res.* 2017;354(1):57-64.
 75. Fleetwood AJ, Lee MKS, Singleton W, et al. . Metabolic Remodeling, Inflammasome Activation, and Pyroptosis in Macrophages Stimulated by *Porphyromonas gingivalis* and Its Outer Membrane Vesicles. *Front Cell Infect Microbiol.* 2017;7:351.
 76. Cecil JD, O'Brien-Simpson NM, Lenzo JC, et al. . Outer Membrane Vesicles Prime and Activate Macrophage Inflammasomes and Cytokine Secretion In Vitro and In Vivo. *Front Immunol.* 2017;8:1017.
 77. Johnson L, Atanasova KR, Bui PQ, et al. . *Porphyromonas gingivalis* attenuates ATP-mediated inflammasome activation and HMGB1 release through expression of a nucleoside-diphosphate kinase. *Microbes Infect.* 2015;17(5):369-377.
 78. Thornberry NA, Bull HG, Calaycay JR, et al. . A novel heterodimeric cysteine protease is required for interleukin-1 beta processing in monocytes. *Nature.* 1992;356(6372):768-774.

79. Strowig T, Henao-Mejia J, Elinav E, Flavell R. Inflammasomes in health and disease. *Nature*. 2012;481(7381):278-286.
80. Lamkanfi M, Dixit VM. Modulation of inflammasome pathways by bacterial and viral pathogens. *J Immunol*. 2011;187(2):597-602.
81. Abdelaziz DH, Gavrilin MA, Akhter A, et al. . Apoptosis-associated speck-like protein (ASC) controls *Legionella pneumophila* infection in human monocytes. *J Biol Chem*. 2011;286(5):3203-3208.
82. Sutterwala FS, Mijares LA, Li L, Ogura Y, Kazmierczak BI, Flavell RA. Immune recognition of *Pseudomonas aeruginosa* mediated by the IPAF/NLRC4 inflammasome. *J Exp Med*. 2007;204(13):3235-3245.
83. Master SS, Rampini SK, Davis AS, et al. . *Mycobacterium tuberculosis* prevents inflammasome activation. *Cell Host Microbe*. 2008;3(4):224-232.
84. Pettersson M, Kelk P, Belibasakis GN, Bylund D, Molin Thoren M, Johansson A. Titanium ions form particles that activate and execute interleukin-1 β release from lipopolysaccharide-primed macrophages. *J Periodontal Res*. 2017;52(1):21-32.

Table 1. Activators of the NLRP3 inflammasome

Activators	References
Silica, asbestos	Davis, BK (2011)
Cholesterol crystal	Duewell, P (2010)
Various bacterial toxins	Vladimer GI (2013); Franchi L (2012)
Bacterial ATP and K ⁺ efflux	Franchi L (2007); Petrilli V (2007)
Oxidized mitochondrial DNA	Shimada K (2012)
ROS from mitochondrial	Zhou R (2011)
Guanylate-binding protein 5	Shenoy AR (2012)
dsRNA-dependent kinase	Lu B (2012)
Cathepsin B	Davis, BK (2011)
Bacterial mRNA	Sander LE (2011)
Blocking of ribosomal function	Vyleta ML (2012)
Muramyl dipeptide	Martinon F (2004); Marina-Garcia N (2008)
Bacterial RNA, PolyI:C, LPS, lipopeptide, R848	Kanneganti TD (2006)
mycoplasmas	Sugiyama M (2016)

Table 2. Activation of inflammasomes by *P. gingivalis*

Type of Inflammasome	Signal 1	Signal 2	Target cell	References
NLRP3	Pg infection	ATP	Human primary GEC	Yilmaz O (2010)
NLRP3/AIM2	Supragingival biofilm	ND	Human GF	Bostanci N (2011)
ND	Pg LPS	Cholesterol crystal	Human primary HMDM	Champaiboon C (2014)
NLRP3, AIM2	Pg infection	ND	A human monocytic cell line, THP-1	Park E (2014)
NLRP3	Pg infection	Pg LPS and/or ATP	A human GEC line, H413	Guo W (2015)
NLRP3	Pg infection	ND	Mouse PM	Yamaguchi Y (2017)
NLRP3	Pg infection	ND	Mouse osteoblast	Yoshida K (2017)
NLRP3	OMVs of Pg	ND	Mouse BMM	Fleetwood AJ (2017) & Cecil JD (2017)
NLRP3/AIM2	OMVs of Pg	ND	Mouse BMM	Cecil JD (2017)
?	OMVs of Pg, Td, Tf	ND	A human monocytic cell line, THP-1	Cecil JD (2017)

ND: not described, Pg: *P. gingivalis*, Td: *Treponema denticola*, Tf: *Tannella forthia*, HMDM: human monocyte-derived macrophage, GF: gingival fibroblasts, GEC: gingival epithelial cell, PM: peritoneal macrophages, OMVs: outer membrane vesicles, BMM: bone marrow-derived macrophage

Table 3. Inhibition of the inflammasome activation by *P. gingivalis*

Type of Inflammasome	Mechanism	Active entity	Target cell	references
NLRP3/AIM2	Subgingival biofilm	ND	Human GF	Bostanci N (2011)
NLRP3	Suppression of other bacterial endocytosis	ND	Mouse BMDM	Taxman DJ (2012)
NLRP3	ND	ND	Primary human GF	Belibasakis GN (2013)
NLRP3	Proteolysis of NLRP3	ND	Human UVEC	Huck O (2015)
NLRP3	ND	Fimbriae	Mouse BMDM	Morandini AC (2014)
ND	Downregulation of ATP/P2X ₇ -signaling and release of HMGB1	Nucleotide-diphosphate kinase	Human GEC	Johnson L (2015)

ND: not described, BMDM: bone marrow-derived macrophage, GF: gingival fibroblasts, UVEC: umbilical vein endothelial cells, GEC: gingival epithelial cell, HMGB1: high mobility group box

FIGURE LEGENDS

Fig. 1. Production of biologically active IL-1 β

IL-1 β is produced as pro-IL-1 β in the cytosol, which is biologically inactive, through activation of nuclear factor- κ B by Toll-like receptor-mediated signaling. Biologically active IL-1 β is produced after processing by caspase-1, which is also processed from pro-caspase-1 by activation of the intracellular sensor inflammasome. The inflammasome is an intracellular multiprotein complex comprising “nucleotide-binding domain leucine-rich repeat-containing receptor,” the adaptor protein “apoptosis-associated speck-like protein containing a caspase-recruitment domain,” and procaspase-1, which is formed by signals mediated by extracellular ATP, pathogen-associated molecular patterns, danger-associated molecular patterns or K⁺ efflux.

PYD: pyrin domain, CARD: caspase recruitment domain

Fig. 2. Effects of IL-1 on periodontal tissues that have been published so far.

Fig. 1

Fig. 2