Title	Heat-shock-induced cathepsin B activity during IVF and culture compromises the developmental competence of bovine embryos
Author(s)	Yamanaka, Ken-ichi; Khatun, Hafiza; Egashiraa, Junki; Balboula, Ahmed Zaky; Tatemoto, Hideki; Sakatani, Miki; Takenouchi, Naoki; Wada, Yasuhiko; Takahashi, Masashi
Citation	Theriogenology, 114, 293-300 https://doi.org/10.1016/j.theriogenology.2018.04.005
Issue Date	2018-07-01
Doc URL	http://hdl.handle.net/2115/74904
Rights	© 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	Yamanaka_et_al._manuscript.pdf

[Instructions for use](#)

Heat-shock-induced cathepsin B activity during IVF and culture compromises the developmental competence of bovine embryos

Ken-ichi Yamanaka^{a,b,*}, Hafiza Khatun^{a,b,c}, Junki Egashira^{a,b,d}, Ahmed Zaky Balboula^e, Hideki Tatemoto^{b,f}, Miki Sakatani^g, Naoki Takenouchi^g, Yasuhiko Wada^{a,b} and Masashi Takahashi^h

^a Faculty of Agriculture, Saga University, Saga, Japan

^b The United Graduate School of Agricultural Sciences, Kagoshima University, Kagoshima, Japan

^c Faculty of Animal Husbandry, Bangladesh Agricultural University, Mymensingh, Bangladesh

^d Saga Prefectural Livestock Experiment Station, Saga, Japan

^e Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt

^f Faculty of Agriculture, University of the Ryukyus, Okinawa, Japan

^g Livestock and Grassland Research Division, Kyushu Okinawa Agricultural Research Center, NARO, Kumamoto, Japan

^h Research Faculty of Agriculture, Hokkaido University, Hokkaido, Japan

*Corresponding author.

Faculty of Agriculture, Saga University, 1 Honjo-machi Saga, 840-8502, Japan.

Tel.: +81-952-28-8735

E-mail: kyama@cc.saga-u.ac.jp (K. Yamanaka).

Abstract

Heat stress can cause significant reproductive dysfunction in mammals and previous studies report that expression and activity of cathepsin B (CTSB), a lysosomal cysteine protease, is negatively correlated with the developmental competence of bovine oocytes and embryos. However, the relationship between heat shock (HS) and CTSB remains largely unknown. Here, we investigated the effects of HS during IVF and early embryonic stages of IVC on CTSB activity and developmental competence in bovine embryos. HS (40°C for 6 h during IVF and 20 h during IVC) caused a significant increase in CTSB activity irrespective of the developmental stage or duration of HS. The developmental rate to the blastocyst stage was also significantly decreased by HS. Additionally, HS during IVC significantly increased the number of apoptotic cells in blastocysts. Notably, these HS-induced changes in blastocyst development and quality were significantly improved by inhibition of CTSB activity, indicating a key role for CTSB. These results showed that CTSB activity plays an essential role in HS-induced dysfunction in bovine embryo development, and that inhibition of this activity could enhance the developmental competence of heat-shocked embryos.

Keywords: cathepsin B, heat shock, embryonic development, bovine

1. Introduction

Reproductive processes in mammals are very sensitive to hyperthermia, which can cause significant decreases in fertility in females and sperm quality in males. Hyperthermia occurs when high ambient temperatures (typically $>27^{\circ}\text{C}$) cause body temperature to rise to between 40°C and 42°C [1, 2]. Notably, exposing cows to hyperthermic heat stress reduces their conception rate and affects subsequent embryonic development [3]. *In vivo* studies indicate that the observed reduction in conception rate can occur quickly and potentially be affected by elevated body temperature on the day of insemination [1, 4, 5]. Furthermore, multiple *in vitro* studies reported that high temperatures during the period from oocyte maturation to preimplantation embryo development degrade oocyte quality and developmental competence by promoting oxidative stress and DNA damage [6-12]. Similarly, oocyte competency with regard to fertilization and blastocyst formation was also reduced by high temperature during fertilization *in vitro* [11]. Together, these data suggest that the mechanism by which heat stress affects fertility and embryo viability is multifactorial and likely involves many aspects of reproductive physiology. Because heat stress affects reproductive health, many attempts have been made to improve the developmental competence of heat-shocked oocytes and embryos via changes in media composition or supplementation. Interestingly, the addition of chemicals or antioxidants, such as insulin-like growth factor-1 [13], 2-mercaptoethanol [12, 14], astaxanthin [15], anthocyanins [16], melatonin [17, 18], and coagulansin-A [19], into the culture medium appears to enhance oocyte and embryo quality, as well as improve developmental competence during heat shock (HS).

Cathepsin B (CTSB) is a lysosomal cysteine protease that participates in the degradation of intracellular proteins in lysosomes [20]. CTSB is synthesized as an enzyme precursor in an inactive form prior to post-translational processing into its active form after passing through the endoplasmic reticulum and subsequent incorporation into the acidic environment of the lysosome [21, 22]. The active form of CTSB has disulfide-linked heavy and light-chain subunits and a molecular weight of 30 kDa [23]. CTSB is also involved in many physiological and pathological

processes, including initiation of the apoptosis pathway, remodeling of extracellular matrix, autophagy, cancer differentiation, and inflammation [24, 25]. CTSB expression is detected in many types of cells, including ovaries and cumulus cells [26, 27]. Bettegowda et al. [26] reported that higher expression of *CTSB* mRNA was observed in cumulus cells from prepubertal cows, which are a model with poor oocyte competence, relative to levels observed in adult cows. We also reported that higher CTSB activity was observed in low-quality bovine cumulus-oocyte complexes (COCs) and embryos as compared with that observed in those of better quality. Additionally, CTSB also plays a role in apoptosis, which is associated with low developmental competence in both COCs and embryos [28, 29]. Importantly, addition of a CTSB inhibitor (E-64) to the culture medium improved developmental competence in low-quality oocytes and embryos. Moreover, CTSB inhibition also improved blastocyst quality via regulation of the apoptosis pathway [28, 29]. Although bovine oocytes exposed to HS during *in vitro* maturation also exhibited increased CTSB activity, low developmental competence, and increased apoptosis, these changes could be counteracted by CTSB inhibition [30]. Despite these findings, our understanding of how CTSB inhibitors function under HS conditions during fertilization and early embryo development is limited.

In this study, we clarified the relationship between CTSB activity and low developmental competence caused by HS during fertilization and early embryonic development using *in vitro* culture techniques. Furthermore, we also investigated the effect of CTSB inhibition on the developmental competence of HS-treated embryos. To our knowledge, this is the first study investigating the role of CTSB and the benefits of its inhibition during HS-induced changes at these particular stages of embryogenesis.

2. Materials and methods

2.1. Chemicals

All chemicals were purchased from Sigma–Aldrich (St. Louis, MO, USA) unless noted otherwise.

93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119

2.2. Oocyte collection and in vitro maturation

Bovine ovaries were collected from different breeds at the local abattoir. COCs were aspirated from follicles (2–6 mm in diameter) using an 18-gauge needle attached to a 10-mL syringe and washed three times in TCM-199 (Thermo Fisher Scientific, Waltham, MA, USA, USA) containing 5% (v/v) fetal bovine serum (FBS; Thermo Fisher Scientific). A total of 50 COCs were matured in a 500-μL drop of TCM-199 supplemented with 5% FBS, FSH (0.02 IU/mL; Kyoritsu Seiyaku, Tokyo, Japan), and gentamicin (10 mg/mL; Nacalai Tesque, Kyoto, Japan), covered with liquid paraffin (Nacalai Tesque), and incubated at 38.5°C for 22 h in a humidified atmosphere of 5% CO₂ in air.

2.3. IVF and IVC

Frozen semen was thawed by immersing the straw in warm water (37°C) for 20 s. Spermatozoa were washed by centrifugation (800g for 10 min) in 90% (v/v) percoll solution (GE Healthcare Bio-Sciences AB, Stockholm, Sweden). After removing the supernatant, the pellet was diluted with IVF100 solution (Research Institute for the Functional Peptides, Yamagata, Japan) and centrifuged at 600g for 5 min. The spermatozoa pellet was then diluted with IVF100 to prepare a final sperm-cell concentration of 5.0×10^6 sperm/mL. IVF was performed at 38.5°C (control group) or at 40°C (HS group) in 5% CO₂ in air under humidified conditions for 6 h (25 oocytes per 100 μL sperm drop covered with liquid paraffin).

After fertilization, cumulus cells were removed mechanically by pipetting in CR1aa medium [31] containing 5% FBS, and putative zygotes with polar bodies were placed into micro-drops (20–25 zygotes per 50 μL drop) of CR1aa medium supplemented with 5% (v/v) FBS. The drops were then covered with liquid paraffin and cultured at 38.5°C (control group) or at 40.0°C for 20 h after fertilization, followed by incubation at 38.5°C (HS group) in a humidified atmosphere of 5% O₂, 5% CO₂, and balanced with N₂ through day 8 (day 0 represented the day of insemination).

2.4. Detection of intracellular CTSB activity

Detection of embryonic CTSB activity was performed using a Magic red CTSB detection kit (MR-RR)2 (Immunochemistry Technologies LLC, Bloomington, MN, USA) according to manufacturer instructions. Briefly, *in vitro* embryos were incubated in 500 μ L PBS along with 2 μ L of reaction mix in a humidified atmosphere of 5% CO₂ at 38.5°C for 20 min. To detect nuclei, 25 μ g/mL Hoechst 33342 was added and incubated in the same culture conditions for an additional 5 min. After rinsing with PBS containing 0.5 mg/mL polyvinylpyrrolidone (PVP; Nacalai Tesque), the stained embryos were mounted onto a glass slide and observed under a fluorescence microscope (EVOS FL; Thermo Fisher Scientific). Nuclei were observed following excitation at 375 nm, and intracellular CTSB activity was detected at 585 nm excitation. Images of CTSB activity were captured under the same exposure time and analyzed with ImageJ software (v1.55; National Institutes of Health, Bethesda, MD, USA).

2.5. Terminal deoxynucleotidyl transferase biotin-dUTP nick end labeling (TUNEL)

A TUNEL assay kit (*in situ* cell death detection kit; Roche Applied Science, Indianapolis, IN, USA) was used to assess the presence of apoptotic cells in day 8 blastocysts. Blastocysts were fixed in 4% (w/v) paraformaldehyde solution (pH 7.4) for 40 min, rinsed three times in 0.05% (w/v) PVP-PBS, and then permeabilized in 0.05% (w/v) PVP-PBS containing 0.5% TritonX-100 for 20 min, followed by three washes with 0.05% (w/v) PVP-PBS for 5 min. The fragmented DNA ends of the cells were labeled with fluorescein-dUTP for 60 min at 38.5°C. After incubation, blastocysts were washed three times in 0.05% (w/v) PVP-PBS for 5 min each, followed by mounting onto glass slides using mounting solution containing 4',6-diamidino-2-phenylindole (DAPI; Vectashield with DAPI; Vector Laboratories, Burlingame, CA, USA). The fluorescence of the fragmented DNA ends was detected using a fluorescence microscope (EVOS® FL, Thermo Fisher Scientific). The apoptotic index was calculated for each embryo as follows: apoptotic index = (number of TUNEL-positive nuclei/total number of nuclei in blastocyst) $\times$ 100.

2.6. Experimental design

We conducted three experiments described as follows and illustrated in Figure 1.

2.6.1. Experiment 1: effect of CTSB inhibitor (E-64) on the development of embryos exposed to HS during IVF or early IVC

To assess the effects of E-64 supplementation on HS-mediated changes in embryonic development, E-64 at different concentrations (0, 0.1, 0.5, and 1.0 μM) was added to the medium during HS treatment. To examine the effect of HS treatment at different times on developmental competence, HS treatment under two different conditions was conducted as follows: 1) at 40.0°C for 6 h during IVF and 2) at 40.0°C for 20 h after the beginning of IVC, during which almost all viable embryos complete the first cleavage [32]. After HS treatment, embryos were cultured at 38.5°C in E-64-free medium until day 8 (day 0 represents the day of insemination). Developmental competence was assessed by cleavage and blastocyst rates on days 2 and 8, respectively.

2.6.2. Experiment 2: effect of HS and CTSB inhibitor on CTSB activity in HS-exposed embryos

We then examined whether HS increases CTSB activity, and whether this change could be inhibited by E-64. Matured oocytes and/or zygotes were cultured with 0.5 μM E-64, followed by exposure to HS at 40.0°C for 6 h during IVF or for 20 h after the beginning of IVC. The number of fluorescent dots highlighting the active form of CTSB was analyzed immediately following HS treatment.

2.6.3. Experiment 3: effect of CTSB inhibitor on the quality of blastocysts derived from HS-exposed embryos

To examine the effects of HS and E64 inhibition on blastocyst quality, matured oocytes or zygotes were cultured with 0.5 μM E-64 and exposed to HS at 40.0°C for 6 h during IVF or for 20 h after the beginning of IVC, followed by incubation at 38.5°C until day 8 of IVC.

Developmental rate (cleavage rate on day 2 and blastocyst rate on day 8), total cell number, and apoptotic status of day 8 blastocysts were analyzed.

2.7. Statistical analysis

Each experiment was repeated at least five times. Data are expressed as the mean $\pm$ standard error. All data were analyzed using analysis of variance (ANOVA), followed by the Tukey-Kramer multiple comparison test. All percentage data were arcsine transformed prior to statistical analysis. A $P < 0.05$ was considered statistically significant.

3. Results

3.1. Inhibition of CTSB activity improves the developmental competence of bovine embryos exposed to HS during IVF or early IVC

HS during IVF decreased the developmental rate of embryos as compared with that observed in the control group ($P < 0.05$). However, embryos cultured with E-64 (0.5 μ M) under the same HS conditions increased the developmental rate of embryos to a rate similar to that observed in the control group (Table 1). As shown in Table 2, HS during early IVC decreased the developmental rate to blastocyst stage as compared with control group ($P < 0.05$); however, this rate was increased by addition of 0.5 μ M E-64 to the culture medium under HS conditions ($P < 0.05$), although the rate was lower than that in the control groups ($P < 0.05$). Based on these results, we selected the E-64 concentration that had the greatest effect on the developmental competence of HS-treated embryos. Therefore, all subsequent experiments were performed using 0.5 μ M E-64.

3.2. HS increases intracellular CTSB activity in bovine embryos

To investigate the relationship between elevated temperature during culture and CTSB activity in embryos, HS treatment was applied during IVF (6 h) or early IVC (20 h). CTSB activity was detected immediately following HS treatment. As shown in Figure 1A, the active form of CTSB

was clearly detected in the cytoplasm, with the number of fluorescent dots counted and used as an indicator of CTSB activity. Notably, the average number of fluorescent dots was higher in the HS-treated group relative to that observed in the control and HS+E-64 groups under both HS-treatment conditions (6 h during IVF: $P < 0.01$, Fig. 1B; 20 h in early IVC: $P < 0.01$, Fig. 1C). Additionally, the average number of fluorescent dots in embryos exposed to HS during early IVC was higher as compared with that in embryos exposed to HS during IVF (74.1 ± 12.2 vs. 46.6 ± 6.1 , respectively; $P < 0.05$). By contrast, addition of E-64 to the culture medium reduced the number of fluorescent dots induced by HS ($P < 0.01$; Fig. 1B and C).

3.3. Inhibition of CTSB activity improves the quality of HS-treated embryos by decreasing apoptosis in the blastocyst stage

Incubation of embryos at elevated temperatures during IVF and early IVC reduced the developmental rate in the blastocyst stage ($P < 0.05$), whereas the addition of E-64 increased the developmental rate as compared with embryos cultured without E-64 under HS conditions ($P < 0.05$; Figs. 2A and 3A). HS during IVF also reduced the cleavage rate, which was again increased by treatment with E-64 (Fig. 2A). The total cell number and apoptotic status of the blastocysts were also assessed to investigate embryo quality. HS during IVF had no significant effect on the total cell number or the percentage of apoptotic cells in the blastocyst stage (Fig. 2B and C). By contrast, although no difference was found in the total cell number among groups, the percentage of apoptotic cells in the blastocyst stage increased when embryos were exposed to HS during early IVC ($P < 0.01$; Fig. 3C). Furthermore, E-64 treatment decreased the percentage of apoptotic cells in these HS-affected blastocysts ($P < 0.01$) to a level similar to that in the control group (Fig. 3C).

4. Discussion

Multiple studies reported that HS during IVF and the early stage of IVC reduces the success of subsequent embryonic development [11, 33, 34]. The mechanism by which HS affects fertility

and embryo viability appears to be multifactorial, and various supplements have been investigated *in vitro* to determine their possible anti-HS effects. However, the efficacy and mechanisms of these and other treatments have not been fully elucidated, and HS remains an issue affecting mammalian reproductive health. CTSB is a lysosomal cysteine protease that plays an important role in the degradation of intracellular-tissue proteins in the lysosome [35]. Its activity has also been linked to changes in embryonic competency based on its high activity causing significant decreased in *in vitro* developmental rates and increases in apoptosis [28, 29]. However, the relationship between CTSB activity and low developmental competence in HS embryos remains largely unknown. To better understand the role of CTSB during HS-induced reproductive changes, we used a bovine embryo *in vitro* model. We observed that CTSB activity was elevated by HS during IVF and early IVC in bovine embryos. Moreover, supplementing the media with a CTSB inhibitor (E-64) improved the developmental competence of HS-treated embryos. These results suggested a role for CTSB in compromising the developmental competence of bovine embryos under HS conditions.

Previous study reported that higher expression of *CTSB* mRNA in cumulus cells surrounding oocytes with low developmental competence as compared with those with high developmental competence [26]. Furthermore, we also demonstrated that CTSB activity was higher in poor-quality oocytes and embryos than in good-quality ones, and that inhibition of this activity improved subsequent developmental competence [28, 29]. However, these previous studies did not specifically account for temperature-related stress. A previous study of the effects of HS investigated with regard to CTSB activity in cumulus cells and oocytes during IVM revealed that CTSB activity was elevated following HS and compromised subsequent developmental competence [30]. Moreover, administering a CTSB inhibitor neutralized these HS-mediated detrimental effects. Although the present study focused on IVF and early IVC, these data support our findings that CTSB plays a significant role in HS-induced changes during embryonic development. Similarly, we also found that addition of 0.5 μ M E-64 to the culture medium under HS conditions increased the developmental competence of treated embryos. These results are

again supported by previous studies showing that E-64 improves the subsequent development of oocytes and embryos [26, 28-30, 36].

One particularly interesting result in the present study indicated that HS during IVF reduced the cleavage rate, but that the same effect was not observed during early IVC. We previously showed that HS during IVF decreases the cleavage rate by compromising the anti-polyspermy systems of oocytes [11]. Oxidative stress also plays a role in these effects, as it can damage sperm-cell structures and decreases subsequent embryonic development when the damaged sperm are applied for IVF [37]. These findings indicate that HS during IVF could impair processes in both the oocyte and sperm, resulting in decreased cleavage. On the other hand, the inhibition of CTSB activity increased the cleavage rate of embryos fertilized under HS. Collectively, increased CTSB activity in response to any form of cellular stress might be the key to understanding the observed changes in fertilization rates and embryo competency.

The mechanism of CTSB during the observed HS-induced changes in embryo viability is likely related, at least in part, to its cellular localization in the lysosome. Lysosomes are membrane-enclosed organelles that are essential for macromolecule degradation and many other cellular functions [38]. In addition to CTSB, lysosomes also contain many other hydrolytic enzymes and play pivotal roles in endocytosis, phagocytosis, and autophagy. Previous studies demonstrated that lysosome function is significantly related to early embryonic development in mice [39-41] and cattle [42]. Furthermore, exogenous stress, such as HS and oxidative stress, in somatic cells can induce lysosomal destabilization and leakage of their components into the cytosol [43, 44]. We previously showed that lysosomal aggregation and possible release of lysosomal contents into the cytosol occur in bovine oocytes after HS treatment [30]. Therefore, leakage of lysosomal CTSB into the cytoplasm is likely the initiating step in the observed increase in CTSB activity, leading to unnecessary proteolysis and subsequent arrested embryonic development.

Leakage of lysosomal enzymes also induces apoptosis. Apoptosis is a normal physiological process often used as a quality marker in preimplantation embryo development both *in vivo* and

282 *in vitro* [45-48]. Previous studies report that HS induces apoptosis in preimplantation embryos
283 [49, 50]. Furthermore, in the present study, we also showed that HS during early IVC increased
284 the number of apoptotic cells in the blastocyst stage. The mechanism underlying these effects
285 appeared to be related to mitochondrial function, whereby HS-induced increases in
286 reactive-oxygen-species production triggers mitochondrial-permeability transition pore opening,
287 followed by release of cytochrome c to the cytosol. These changes cause apoptosis in somatic
288 cells [51, 52], as well as bovine embryos [6, 12, 14, 53]. More importantly, previous studies
289 report that CTSB can promote this process via induction of mitochondrial-membrane
290 degradation [54, 55]. These studies are supported by our present observation that CTSB
291 inhibition significantly decreased apoptosis in HS-treated embryos. Interestingly, HS during IVF
292 had no significant effect on apoptosis at the blastocyst stage, even though increased CTSB
293 activity was observed after HS treatment. This discrepancy might be related to the difference in
294 the duration of HS treatment, which was shorter during IVF (6 h) as compared with early IVC
295 (20 h). Moreover, CTSB activity in embryos exposed to HS during IVF was also lower than that
296 in embryos exposed to HS during early IVC, indicating that this treatment time was not long
297 enough to induce the level of CTSB activity needed for activation of apoptosis-signaling
298 pathways. Although we observed variable responses depending on the *in vitro* stage (IVF vs.
299 early IVC) and the duration of HS treatment (6 h vs. 20 h), the mechanism underlying the effects
300 of HS likely involves lysosome leakage leading to CTSB activation, which causes a subsequent
301 increase in apoptosis and, ultimately, dysfunctional embryo development.

302 In conclusion, our results showed that CTSB activity was elevated by HS during IVF and
303 early IVC. Moreover, inhibition of CTSB enhanced the developmental competence and quality
304 of HS-treated bovine embryos, suggesting that excess CTSB activity was indeed involved in the
305 observed HS-induced arrested development. Although further research is required to provide
306 additional details concerning the intermediate signaling cascades involved and to confirm the *in*
307 vivo development of HS embryos treated with E-64 through embryo transfer, the present study is
308 useful for elucidating the comprehensive mechanism underlying HS-induced developmental

changes in bovine embryos and provides a framework for future studies, as well as insight into developmental processes and treatment for HS-affected embryogenesis in mammals.

Acknowledgments

The authors wish to thank Sagaken-chikusan-kousha for providing the bovine ovaries. We would like to thank Editage (www.editage.jp) for English-language editing.

Funding: This research was supported by a Grant-in-Aid for Scientific Research (KAKENHI Grant Number JP25850186 and JP17K15361) from the Japan Society for the Promotion of Science (JSPS) to K.Y.

References

- [1] Nabenishi H, Ohta H, Nishimoto T, Morita T, Ashizawa K, Tsuzuki Y. Effect of the temperature-humidity index on body temperature and conception rate of lactating dairy cows in southwestern Japan. *J Reprod Dev* 2011;57:450–6.
- [2] Berman A, Folman Y, Kaim M, Mamen M, Herz Z, Wolfenson D, et al. Upper critical temperatures and forced ventilation effects for high-yielding dairy cows in a subtropical climate. *J Dairy Sci* 1985;68:1488–95.
- [3] Sakatani M. Effects of heat stress on bovine preimplantation embryos produced in vitro. *J Reprod Dev* 2017;63:347–52.
- [4] Moghaddam A, Karimi I, Pooyanmehr M. Effects of short-term cooling on pregnancy rate of dairy heifers under summer heat stress. *Vet Res Commun* 2009;33:567–75.
- [5] Badinga L, Collier RJ, Thatcher WW, Wilcox CJ. Effects of climatic and management factors on conception rate of dairy cattle in subtropical environment. *J Dairy Sci* 1985;68:78–85.
- [6] Sakatani M, Kobayashi S, Takahashi M. Effects of heat shock on in vitro development and intracellular oxidative state of bovine preimplantation embryos. *Mol Reprod Dev* 2004;67:77–82.
- [7] Edwards JL, Saxton AM, Lawrence JL, Payton RR, Dunlap JR. Exposure to a physiologically relevant elevated temperature hastens in vitro maturation in bovine oocytes. *J Dairy Sci* 2005;88:4326–33.
- [8] Nabenishi H, Takagi S, Kamata H, Nishimoto T, Morita T, Ashizawa K, et al. The role of mitochondrial transition pores on bovine oocyte competence after heat stress, as determined by effects of cyclosporin A. *Mol Reprod Dev* 2012;79:31–40.
- [9] Roth Z, Hansen PJ. Involvement of apoptosis in disruption of developmental competence of bovine oocytes by heat shock during maturation. *Biol Reprod* 2004;71:1898–906.

- [10] Roth Z, Hansen PJ. Disruption of nuclear maturation and rearrangement of cytoskeletal elements in bovine oocytes exposed to heat shock during maturation. *Reproduction* 2005;129:235–44.
- [11] Sakatani M, Yamanaka K, Balboula AZ, Takenouchi N, Takahashi M. Heat stress during in vitro fertilization decreases fertilization success by disrupting anti-polyspermy systems of the oocytes. *Mol Reprod Dev* 2015;82:36–47.
- [12] Sakatani M, Yamanaka K, Kobayashi S, Takahashi M. Heat shock-derived reactive oxygen species induce embryonic mortality in in vitro early stage bovine embryos. *J Reprod Dev* 2008;54:496–501.
- [13] Bonilla AQ, Oliveira LJ, Ozawa M, Newsom EM, Lucy MC, Hansen PJ. Developmental changes in thermoprotective actions of insulin-like growth factor-1 on the preimplantation bovine embryo. *Mol Cell Endocrinol* 2011;332:170–9.
- [14] Nabenishi H, Ohta H, Nishimoto T, Morita T, Ashizawa K, Tsuzuki Y. The effects of cysteine addition during in vitro maturation on the developmental competence, ROS, GSH and apoptosis level of bovine oocytes exposed to heat stress. *Zygote* 2012;20:249–59.
- [15] Namekawa T, Ikeda S, Sugimoto M, Kume S. Effects of astaxanthin-containing oil on development and stress-related gene expression of bovine embryos exposed to heat stress. *Reprod Domest Anim* 2010;45:e387–91.
- [16] Sakatani M, Suda I, Oki T, Kobayashi S, Kobayashi S, Takahashi M. Effects of purple sweet potato anthocyanins on development and intracellular redox status of bovine preimplantation embryos exposed to heat shock. *J Reprod Dev* 2007;53:605–14.
- [17] Wang F, Tian X, Zhang L, Tan D, Reiter RJ, Liu G. Melatonin promotes the in vitro development of pronuclear embryos and increases the efficiency of blastocyst implantation in murine. *J Pineal Res* 2013;55:267–74.
- [18] Gao C, Han HB, Tian XZ, Tan DX, Wang L, Zhou GB, et al. Melatonin promotes embryonic development and reduces reactive oxygen species in vitrified mouse 2-cell embryos. *J Pineal Res* 2012;52:305–11.

- [19] Khan I, Lee KL, Xu L, Mesalam A, Chowdhury MM, Joo MD, et al. Improvement of in vitro-produced bovine embryo treated with coagulansin-A under heat-stressed condition. *Reproduction* 2017;153:421–31.
- [20] Barrett AJ, Kirschke H. Cathepsin B, Cathepsin H, and cathepsin L. *Methods Enzymol* 1981;80 Pt C:535–61.
- [21] Buhling F, Fengler A, Brandt W, Welte T, Ansorge S, Nagler DK. Review: novel cysteine proteases of the papain family. *Adv Exp Med Biol* 2000;477:241–54.
- [22] Turk B, Turk D, Turk V. Lysosomal cysteine proteases: more than scavengers. *Biochim Biophys Acta* 2000;1477:98–111.
- [23] Klose A, Zigrino P, Dennhofer R, Mauch C, Hunzelmann N. Identification and discrimination of extracellularly active cathepsins B and L in high-invasive melanoma cells. *Anal Biochem* 2006;353:57–62.
- [24] Frlan R, Gobec S. Inhibitors of cathepsin B. *Curr Med Chem* 2006;13:2309–27.
- [25] Stoka V, Turk V, Turk B. Lysosomal cysteine cathepsins: signaling pathways in apoptosis. *Biol Chem* 2007;388:555–60.
- [26] Bettgowda A, Patel OV, Lee KB, Park KE, Salem M, Yao J, et al. Identification of novel bovine cumulus cell molecular markers predictive of oocyte competence: functional and diagnostic implications. *Biol Reprod* 2008;79:301–9.
- [27] Oksjoki S, Soderstrom M, Vuorio E, Anttila L. Differential expression patterns of cathepsins B, H, K, L and S in the mouse ovary. *Mol Hum Reprod* 2001;7:27–34.
- [28] Balboula AZ, Yamanaka K, Sakatani M, Hegab AO, Zaabel SM, Takahashi M. Intracellular cathepsin B activity is inversely correlated with the quality and developmental competence of bovine preimplantation embryos. *Mol Reprod Dev* 2010;77:1031–9.
- [29] Balboula AZ, Yamanaka K, Sakatani M, Hegab AO, Zaabel SM, Takahashi M. Cathepsin B activity is related to the quality of bovine cumulus oocyte complexes and its inhibition can improve their developmental competence. *Mol Reprod Dev* 2010;77:439–48.

- [30] Balboula AZ, Yamanaka K, Sakatani M, Kawahara M, Hegab AO, Zaabel SM, et al. Cathepsin B activity has a crucial role in the developmental competence of bovine cumulus-oocyte complexes exposed to heat shock during in vitro maturation. *Reproduction* 2013;146:407–17.
- [31] Rosenkrans CF, Jr., Zeng GQ, GT MC, Schoff PK, First NL. Development of bovine embryos in vitro as affected by energy substrates. *Biol Reprod* 1993;49:459–62.
- [32] Somfai T, Inaba Y, Aikawa Y, Ohtake M, Kobayashi S, Konishi K, et al. Relationship between the length of cell cycles, cleavage pattern and developmental competence in bovine embryos generated by in vitro fertilization or parthenogenesis. *J Reprod Dev* 2010;56:200–7.
- [33] Hendricks KE, Martins L, Hansen PJ. Consequences for the bovine embryo of being derived from a spermatozoon subjected to post-ejaculatory aging and heat shock: development to the blastocyst stage and sex ratio. *J Reprod Dev* 2009;55:69–74.
- [34] Rivera RM, Kelley KL, Erdos GW, Hansen PJ. Alterations in ultrastructural morphology of two-cell bovine embryos produced in vitro and in vivo following a physiologically relevant heat shock. *Biol Reprod* 2003;69:2068–77.
- [35] Szpaderska AM, Frankfater A. An intracellular form of cathepsin B contributes to invasiveness in cancer. *Cancer Res* 2001;61:3493–500.
- [36] Min SH, Song BS, Yeon JY, Kim JW, Bae JH, Park SY, et al. A cathepsin B inhibitor, E-64, improves the preimplantation development of bovine somatic cell nuclear transfer embryos. *J Reprod Dev* 2014;60:21–7.
- [37] Silva PF, Gadella BM, Colenbrander B, Roelen BA. Exposure of bovine sperm to pro-oxidants impairs the developmental competence of the embryo after the first cleavage. *Theriogenology* 2007;67:609–19.
- [38] Eskelinen EL, Tanaka Y, Saftig P. At the acidic edge: emerging functions for lysosomal membrane proteins. *Trends Cell Biol* 2003;13:137–45.

- [39] Tsukamoto S, Hara T, Yamamoto A, Kito S, Minami N, Kubota T, et al. Fluorescence-based visualization of autophagic activity predicts mouse embryo viability. *Sci Rep* 2014;4:4533.
- [40] Tsukamoto S, Hara T, Yamamoto A, Ohta Y, Wada A, Ishida Y, et al. Functional analysis of lysosomes during mouse preimplantation embryo development. *J Reprod Dev* 2013;59:33–9.
- [41] Tsukamoto S, Kuma A, Murakami M, Kishi C, Yamamoto A, Mizushima N. Autophagy is essential for preimplantation development of mouse embryos. *Science* 2008;321:117–20.
- [42] Song B-SS, Yoon S-BB, Kim J-SS, Sim B-WW, Kim Y-HH, Cha J-JJ, et al. Induction of autophagy promotes preattachment development of bovine embryos by reducing endoplasmic reticulum stress. *Biology of reproduction* 2012;87:11.
- [43] Melo FR, Lundequist A, Calounova G, Wernersson S, Pejler G. Lysosomal membrane permeabilization induces cell death in human mast cells. *Scand J Immunol* 2011;74:354–62.
- [44] Brunk UT, Zhang H, Dalen H, Ollinger K. Exposure of cells to nonlethal concentrations of hydrogen peroxide induces degeneration-repair mechanisms involving lysosomal destabilization. *Free Radic Biol Med* 1995;19:813–22.
- [45] Gjørret JO, Knijn HM, Dieleman SJ, Avery B, Larsson LI, Maddox-Hyttel P. Chronology of apoptosis in bovine embryos produced in vivo and in vitro. *Biol Reprod* 2003;69:1193–200.
- [46] Yu RA, He LF, Chen XM. Effects of cadmium on hepatocellular DNA damage, proto-oncogene expression and apoptosis in rats. *Biomed Environ Sci* 2007;20:146–53.
- [47] Yu F, Megyesi J, Safirstein RL, Price PM. Involvement of the CDK2-E2F1 pathway in cisplatin cytotoxicity in vitro and in vivo. *Am J Physiol Renal Physiol* 2007;293:F52–9.
- [48] Su J, Wang Y, Li Y, Li R, Li Q, Wu Y, et al. Oxamflatin significantly improves nuclear reprogramming, blastocyst quality, and in vitro development of bovine SCNT embryos. *PLoS One* 2011;6:e23805.

- 447 [49] Krininger CE, 3rd, Stephens SH, Hansen PJ. Developmental changes in inhibitory effects of
448 arsenic and heat shock on growth of pre-implantation bovine embryos. *Mol Reprod Dev*
449 2002;63:335–40.
- 450 [50] Paula-Lopes FF, Hansen PJ. Heat shock-induced apoptosis in preimplantation bovine
451 embryos is a developmentally regulated phenomenon. *Biol Reprod* 2002;66:1169–77.
- 452 [51] Elmore S. Apoptosis: a review of programmed cell death. *Toxicol Pathol* 2007;35:495–516.
- 453 [52] Taylor RC, Cullen SP, Martin SJ. Apoptosis: controlled demolition at the cellular level. *Nat*
454 *Rev Mol Cell Biol* 2008;9:231–41.
- 455 [53] Hansen PJ. Effects of heat stress on mammalian reproduction. *Philos Trans R Soc Lond B*
456 *Biol Sci* 2009;364:3341–50.
- 457 [54] Bossy-Wetzel E, Newmeyer DD, Green DR. Mitochondrial cytochrome c release in
458 apoptosis occurs upstream of DEVD-specific caspase activation and independently of
459 mitochondrial transmembrane depolarization. *EMBO J* 1998;17:37–49.
- 460 [55] Boya P, Andreau K, Poncet D, Zamzami N, Perfettini JL, Metivier D, et al. Lysosomal
461 membrane permeabilization induces cell death in a mitochondrion-dependent fashion. *J Exp*
462 *Med* 2003;197:1323–34.

Figure legends

Fig. 1. Schematic diagrams of the experimental design. In all experiments, two types of HS treatment with a different condition was conducted as follows. HS during IVF: mature oocytes were cultured at 40.0°C for 6 h during IVF; HS during early IVC: putative zygotes were cultured at 40.0°C for 20 h after the beginning of IVC. Experiment 1: CTSB inhibitor (E-64) with different concentrations (0, 0.1, 0.5, and 1.0 μ M) was added to the medium during HS treatment. After HS treatment, zygotes/oocytes were cultured at 38.5°C in E-64-free medium until day 8. Developmental competence was assessed by cleavage and BL rates on days 2 and 8, respectively. Experiment 2: mature oocytes or putative zygotes were cultured with or without E-64 (0.5 μ M) during each HS treatment. After HS treatment, CTSB activity in their respective cytoplasm was analyzed immediately. Experiment 3: mature oocytes or putative zygotes were cultured with or without E-64 (0.5 μ M) during each HS treatment. After HS treatment, they were cultured at 38.5°C in E-64-free medium until day 8. BL quality was evaluated by counting total cell number and TUNEL-positive cells in BLs.

BL, blastocyst; CTSB, cathepsin B; E-64, cathepsin inhibitor; HS, heat shock; TUNEL, terminal deoxynucleotidyl transferase biotin-dUTP nick end labeling.

Fig. 2. Effect of HS on intracellular CTSB activity. Intracellular localization of the active form of CTSB was detected (red fluorescent dots) (A). Average number of dots showing the active form of CTSB in embryos exposed to HS (40.0°C) for 6 h during IVF (B) and for 20 h during early IVC (C). HS and HS+E-64 groups were supplemented with and without 0.5 μ M E-64 during HS treatment, respectively. In the control group, IVF and IVC were performed without HS and E-64 supplementation. Data represent the mean $\pm$ standard error. Bars labeled with different letters represent significant differences ($P < 0.01$).

CTSB, cathepsin B; E-64, cathepsin inhibitor; HS, heat shock.

Fig. 3. Effect of HS and E-64 administration during IVF on subsequent embryonic development, total cell number, and apoptotic index of blastocysts. Mature oocytes were exposed to HS (40.0°C) for 6 h during IVF with or without the addition of E-64. Developmental rates (A), total cell number (B), and apoptotic index (C) of the blastocysts were analyzed. HS and HS+E-64 groups were supplemented with and without 0.5 µM E-64 during HS treatment, respectively. In the control group, IVF and IVC were performed without HS and E-64 supplementation. Data represent the mean ± standard error. Bars labeled with different letters indicate significant differences ($P < 0.01$).

CTSB, cathepsin B; E-64, cathepsin inhibitor; HS, heat shock.

Fig. 4. Effect of HS and CTSB administered during early IVC on subsequent embryonic development, total cell number, and apoptotic index of blastocysts. Putative zygotes were exposed to HS (40.0°C) for 20 h during early IVC with or without the addition of E-64. Developmental rates (A), total cell number (B), and apoptotic index (C) of the blastocysts were analyzed. HS and HS+E-64 groups were supplemented with and without 0.5 µM E-64 during HS treatment, respectively. In the control group, IVF and IVC were performed without HS and E-64 supplementation. Data represent the mean ± standard error. Bars labeled with different letters indicate significant differences ($P < 0.01$).

CTSB, cathepsin B; E-64, cathepsin inhibitor; HS, heat shock.

514 **Table 1** Effect of E-64 on *in vitro* development of bovine embryos exposed to HS (40.0°C) for 6
515 h during IVF.

Treatment	No. embryos	No. (%) cleaved embryos	No. (%) blastocysts
Control	148	126 (85.1 ± 2.3) ^a	46 (31.1 ± 3.0) ^a
HS	155	104 (67.1 ± 1.5) ^b	17 (11.0 ± 1.1) ^b
HS+E-64 (0.1 µM)	142	118 (83.1 ± 3.0) ^a	26 (18.3 ± 1.7) ^{bc}
HS+E-64 (0.5 µM)	138	112 (81.8 ± 1.0) ^{ac}	33 (23.9 ± 2.0) ^{ac}
HS+E-64 (1.0 µM)	141	103 (73.0 ± 1.7) ^{bc}	24 (17.0 ± 1.0) ^{bc}

516 HS and HS+E-64 groups were supplemented with and without 0.5 µM E-64 during HS treatment,
517 respectively. In the control group, IVF and IVC were performed without HS and E-64
518 supplementation. Data represent the mean ± standard error from five replicates. The data are
519 presented as the means ± standard error from 5 replicates. ^{a-c} Values with different superscripts
520 within the same column are significantly different ($P < 0.05$).

521 E-64, cathepsin inhibitor; HS, heat shock.

522 **Table 2** Effect of E-64 on *in vitro* development of bovine embryos exposed to HS (40.0°C) for
523 20 h during early IVC.

Treatment	No. embryos	No. (%) cleaved embryos	No. (%) blastocysts
Control	153	123 (80.4 ± 2.0)	47 (30.7 ± 2.3) ^a
HS	164	135 (82.3 ± 1.4)	19 (11.6 ± 0.8) ^b
HS+E-64 (0.1 µM)	152	120 (78.9 ± 3.1)	23 (15.1 ± 1.4) ^{bc}
HS+E-64 (0.5 µM)	159	127 (79.9 ± 2.8)	35 (22.0 ± 1.5) ^c
HS+E-64 (1.0 µM)	139	114 (82.0 ± 2.1)	24 (17.3 ± 1.3) ^{bc}

524 HS and HS+E-64 groups were supplemented with and without 0.5 µM E-64 during HS treatment,
525 respectively. In the control group, IVF and IVC were performed without HS and E-64
526 supplementation. Data represent the mean ± standard error from five replicates.

527 ^{a-c} Values with different superscripts within the same column are significantly different (*P* <
528 0.05).

529 E-64, cathepsin inhibitor; HS, heat shock.

Fig.1

HS during IVF

HS during early IVC

Fig. 2

Fig. 3

Fig. 4