Title	Secondary formation of oxalic acid and related organic species from biogenic sources in a larch forest at the northern slope of Mt. Fuji
Author(s)	Mochizuki, Tomoki; Kawamura, Kimitaka; Miyazaki, Yuzo; Wada, Ryuichi; Takahashi, Yoshiyuki; Saigusa, Nobuko; Tani, Akira
Citation	Atmospheric environment, 166, 255-262 https://doi.org/10.1016/j.atmosenv.2017.07.028
Issue Date	2017-07-18
Doc URL	http://hdl.handle.net/2115/74956
Rights	© 2017. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	Accepted Manuscript AE_kk.pdf

[Instructions for use](#)

Accepted by Atmospheric Environment

**Secondary formation of oxalic acid and related organic species from biogenic sources in
a larch forest at the northern slope of Mt. Fuji**

Tomoki Mochizuki^{1,4}, Kimitaka Kawamura^{*,1,5}, Yuzo Miyazaki¹, Ryuichi Wada², Yoshiyuki
Takahashi³, Nobuko Saigusa³, and Akira Tani⁴

¹Institute of Low Temperature Science, Hokkaido University, Sapporo 060-0819, Japan

²Department of Natural and Environmental Science, Teikyo University of Science, Uenohara
409-0193, Japan

³National Institute for Environmental Studies, Tsukuba 305-8506, Japan

⁴Graduate Division of Nutritional and Environmental Sciences, University of Shizuoka,
Shizuoka 422-8526, Japan

⁵Now at Chubu Institute for Advanced Studies, Chubu University, Kasugai 487-8501, Japan

*Correspondence to: K. Kawamura (kkawamura@isc.chubu.ac.jp)

Highlights:

We analyzed water-soluble organic aerosols from a larch forest

Oxalic acid was the dominant dicarboxylic acid species

Terpenes and unsaturated fatty acids are important precursors of dicarboxylic acids

Inflow of O₃ and other oxidants enhanced the formation of dicarboxylic acids

Keywords: Water-soluble organic aerosol, Oxalic acid, Unsaturated fatty acids, *Larix
kaempferi* forest, Terpenes, Ozone

Abstract

To better understand the formation of water-soluble organic aerosols in the forest atmosphere, we measured low molecular weight (LMW) dicarboxylic acids, oxocarboxylic acids, α -dicarbonyls, unsaturated fatty acids (UFAs), and water-soluble organic carbon (WSOC) in aerosols from a *Larix kaempferi* forest located at the northern slope of Mt. Fuji, Japan, in summer 2012. Concentrations of dicarboxylic acids, oxocarboxylic acids, α -dicarbonyls, and WSOC showed maxima in daytime. Relative abundance of oxalic acid in LMW dicarboxylic acids was on average 52% and its average concentration was 214 ng m^{-3} . We found that diurnal and temporal variations of oxalic acid are different from those of isoprene and α -pinene, whereas biogenic secondary organic aerosols (BSOAs) derived from isoprene and α -pinene showed similar variations with oxalic acid. The mass concentration ratios of oxalic acid/BSOAs were relatively constant, although a large variation in the concentrations of toluene that is an anthropogenic volatile organic compound was observed. These results suggest that formation of oxalic acid is associated with the oxidation of isoprene and α -pinene with O_3 and other oxidants in the forest atmosphere. In addition, concentrations of UFAs were observed, for the first time, to decrease dramatically during daytime in the forest. Mass concentration ratios of azelaic acid to UFAs showed a positive correlation with O_3 , suggesting that UFAs are oxidized to yield azelaic acid, which may be further decomposed to oxalic acid in the forest atmosphere. We found that contributions of oxalic acid to WSOC are significantly high ranging from 3.7 to 9.7% (average 6.0%). This study demonstrates that forest ecosystem is an important source of oxalic acid and other dicarboxylic acids in the atmosphere.

1. Introduction

Low molecular weight (LMW) dicarboxylic acids (diacids) have been detected in atmospheric particles from urban, marine, forest, and polar regions (e.g., Li and Winchester, 1989; Kawamura and Bikkina, 2016; Kourtchev et al., 2009). Oxalic acid is the most abundant species of organic aerosols and major portion of oxalic acid (90%) is present in particle phase (Lim et al., 2005). LMW diacids are directly emitted from motor exhausts (Kawamura and Kaplan, 1987), biomass burning (Kawamura et al., 2013), domestic cooking (Schauer et al., 2002) and marine sources (Rinaldi et al., 2011). Furthermore, photochemical oxidations of natural organic precursors (hydrocarbons) are important sources of LMW diacids (e.g., Huang et al., 2011). Diacids are highly water-soluble (Giebl et al., 2002), and thus they can act as cloud condensation nuclei (CCN) to result in the formation of clouds having an influence on the radiative forcing and climate changes (Kanakidou et al., 2005).

Biogenic volatile organic compounds (BVOCs), such as isoprene and monoterpenes, are emitted from vegetation. In particular, plant leaves, leaf litter, and roots are important sources of BVOCs (Lin et al., 2007; Tani and Kawawata, 2008; Mochizuki et al., 2011). The annual global emissions of BVOCs were estimated to be 1000 Tg year⁻¹ (Guenther et al., 2012), which is several times higher than that of anthropogenic VOCs (AVOCs) including toluene (Boucher et al., 2013). Because BVOCs are highly reactive with OH radicals and O₃ (Atkinson and Arey, 2003), they are closely involved with the formation of secondary organic aerosols (SOAs). In model experiment, the amount of biogenic SOAs (BSOAs), including 2-methyltetrols as isoprene SOA tracers and pinonic acid as α -pinene SOA tracer, is several times higher than that of anthropogenic SOA on a global scale (Hallquist et al., 2009). However, BSOAs loadings from BVOC oxidation are highly uncertain (Hallquist et al., 2009).

In our previous study, we conducted simultaneous measurements of BVOCs such as isoprene and α -pinene and their oxidation products such as isoprene- and α -pinene-SOA

tracers as well as oxidants (O_3) in a *Larix kaempferi* forest (Mochizuki et al., 2015). We reported that isoprene and α -pinene emitted from the forest ecosystem are important precursors of BSOAs and that O_3 and other oxidants promote the formation of BSOAs from BVOCs just below the forest canopy. Based on laboratory experiments, BVOCs are known to contribute to the formation of LMW dicarboxylic acids (Huang et al., 2011; Kawamura et al., 2014). However, there is no study on the simultaneous measurements of BVOCs and dicarboxylic acids such as oxalic acid, other organic acids and intermediate compounds that may be dominant SOA components in the forest atmosphere.

Unsaturated fatty acids (UFAs) are emitted from terrestrial higher plants and marine algae (Yokouchi and Ambe, 1986; Kawamura and Gagosian, 1987). UFAs such as oleic and linoleic acids are important sources for the formation of azelaic acid (C_9 diacid) (Yokouchi and Ambe, 1986), although simultaneous observations have never been conducted for the source and product. Longer-chain diacids can produce shorter-chain diacids via photochemical breakdown of UFAs with OH radicals (Kawamura et al., 1996; Pavuluri et al., 2015; Enami et al., 2015). However, there is no study on time-resolved diurnal variations of UFAs and their oxidation products (mid- and short-chain diacids) and their controlling factors in the forest atmosphere.

In the present study, to better understand the formation of oxalic acid and controlling factors that control the formation of oxalic acid in the forest atmosphere, we conducted simultaneous measurements of LMW diacids, oxocarboxylic acids (oxoacids), α -dicarbonyls, UFAs, and water soluble organic carbon (WSOC) in aerosols collected at the site of a *L. kaempferi* forest on the northern foothill of Mt. Fuji. The results of BVOCs, AVOCs, biogenic SOA tracers, and O_3 concentrations at the same time were reported previously (Mochizuki et al., 2015). Here, we report diurnal and temporal variations of diacids and related compounds and compositions of diacids. We discuss sources of oxalic acid and major contributors to the formation of oxalic acid over a Japanese larch forest.

2. Experimental

The Fuji-Hokuroku Flux Research site (35°26' N, 138°45' E) is located in Fujiyoshida city, Yamanashi, Japan on the northern foothill of Mt. Fuji (Fig. A1). *L. kaempferi* was planted in the area of over 150 ha. Almost all trees were 55 years old. The tree canopy height was approximately 20–25 m. The major understory is *Dryopteris crassirhizoma* (fern species) in June to August. Continuous measurements of ambient temperature, solar radiation, wind direction, and wind speed were conducted simultaneously on a meteorological tower (32 m). Details of the site information have been described in Urakawa et al. (2015).

Total suspended particulate matter (TSP) was collected using a high-volume air sampler (AS-810B, Kimoto Electric, Japan) and precombusted (450 °C, 6 h) quartz-fiber filters (8 inch × 10 inch) (Tissuquartz 2500QAT-UP, Pallflex, USA) at a flow rate of 1200 L min⁻¹ on the steel scaffold for tree survey at a height level of 16 m within the Japanese larch canopy. The samples were collected every 3 hours (06:00–09:00, 09:00–12:00, 12:00–15:00 and 15:00–18:00 LT) in daytime (n = 20) and 12 hours (18:00–06:00 LT) in nighttime (n = 4). Before and after the sampling, each filter was placed in a clean glass bottle with a Teflon-lined screw cap. After the sampling, filters were stored in a freezer at -20°C prior to analysis. Samplings were carried out on 5, 10, and 15–17 July 2012. We focused on 5 days to investigate the formation of oxalic acid and their controlling factors. Large variations of concentration of BVOCs, AVOCs, and O₃ and ambient temperature, solar radiation, wind speed, and wind direction were observed in the specific 5 days (Fig. A2) (Mochizuki et al., 2015).

Details of the analytical procedure of diacids, oxoacids, α -dicarbonyls and unsaturated fatty acids were described in Kawamura et al. (1996) and Mochida et al. (2003a). Briefly, water-soluble organic compounds were extracted from the filter portion (12.6 cm²) with ultrapure water (resistivity of > 18.2 M Ω cm). To remove particles, the extracts were passed

through quartz wool packed in a Pasteur pipette. After adjustment of pH to 8.5–9.0 with 0.05 M KOH solution, the filtrates were dried using a rotary evaporator under vacuum at 50 °C. Organic acids and α -dicarbonyls were derivatized to dibutyl esters and dibutoxy acetals with 14% BF_3 /n-butanol at 100 °C for 1 h. The derivatives were identified and quantified using a capillary gas chromatograph (Agilent GC7890A, Agilent, USA) and GC-mass spectrometer (Agilent GC7890A and 5975C MSD, Agilent, USA). Recoveries of authentic dicarboxylic acids (C_2 – C_6) spiked to a quartz filter were 78–91%. Analytical errors were within 9% based on replicate analyses using same sample filter (different parts). We confirmed that levels of field blanks were less than ~5% (diacids), ~7% (oxoacids), and ~9% (α -dicarbonyls) of those of ambient aerosols.

For the measurements of water-soluble organic carbon (WSOC), a portion of filter was extracted with ultrapure water under ultrasonication. The water extracts were filtered through a membrane disk filter (0.22 μm , Millipore Millex-GV, Merck, USA) and then analysed using a total organic carbon analyser (Model TOC-Vcsh, Shimadzu) (Miyazaki et al., 2011).

Concentrations of isoprene and α -pinene and their oxidation products (isoprene-SOA tracers: 2-methylglyceric acid and 2-methyltetrols, α -pinene-SOA tracers: pinonic acid and 3-methyl-1,2,3-butanetricarboxylic acid) and inorganic ions (SO_4^{2-}) and pH values of the water extracted samples were reported in Mochizuki et al. (2015) (Fig. A2).

3. Results and discussion

3.1 Diurnal and temporal variations of dicarboxylic acids, oxocarboxylic acids, and α -dicarbonyls

We identified straight chain saturated (C_2 – C_{10}) diacids, oxoacids (ωC_2 – ωC_9 and pyruvic: Pyr), and α -dicarbonyls (glyoxal: Gly and methylglyoxal: MeGly) in the ambient aerosols collected from the *L. kaempferi* forest (Table 1). Total concentrations of diacids, oxoacids and α -dicarbonyls started to increase in the morning with a maximum around at noontime or in the

afternoon (Fig. 1). Average concentration of total diacids in daytime was 424 ng m^{-3} , which is several times or one order of magnitude higher than those of total oxoacids (72 ng m^{-3}) and α -dicarbonyls (13 ng m^{-3}). Oxalic acid accounts for 52% of total straight chain diacids ($\text{C}_2\text{-C}_{10}$), followed by malonic acid (23%).

Diurnal and temporal variations of oxalic acid in *L. kaempferi* forest aerosols are provided in Fig. 2. Oxalic acid concentrations started to increase in the morning with a maximum around at noontime or in the afternoon, and then decreased to minimize in nighttime. Concentration of oxalic acid peaked ca. 3 hours behind the peaks of isoprene and α -pinene (Fig. 1 and Fig. A2). Concentrations of oxalic acid ($76 - 458 \text{ ng m}^{-3}$) as well as other diacids ($61 - 482 \text{ ng m}^{-3}$), oxoacids ($19 - 180 \text{ ng m}^{-3}$), and α -dicarbonyls ($5 - 33 \text{ ng m}^{-3}$) on 10 and 17 July were higher than those (oxalic acid: $35 - 202 \text{ ng m}^{-3}$, other diacids: $18 - 247 \text{ ng m}^{-3}$, oxoacids: $8 - 55 \text{ ng m}^{-3}$, and α -dicarbonyls: $2 - 15 \text{ ng m}^{-3}$) on 15 and 16 July (Fig. 2 and Fig. A3). Such diurnal and temporal trends of oxalic acid are similar to those of O_3 , toluene, and BSOAs (Fig. 2 and Fig. A2).

This site is located approximately 8 km northwest of the urban area of Fuji-Yoshida city. Particles containing oxalic acid of anthropogenic origin are likely transported from the urban area. To differentiate the local formation of oxalic acid in this site and the atmospheric transport from the surrounding urban areas, we show diurnal and temporal variations in the mass concentration ratios of oxalic acid to BSOAs (C_2/BSOAs) in Fig. 2. In our previous study, production rate of BSOA tracers from isoprene and α -pinene is around 3 hours in this forest. This time scale is comparable to the production rate of oxalic acid from isoprene via O_3 and OH radical oxidation in laboratory experiment (Huang et al., 2011; Kawamura et al., 2014). The wind direction in daytime on 10 July was mainly northeast, indicating an inflow of urban air from the surrounding urban area (Fig. A2e). Conversely, the wind direction in daytime on 17 July was mainly northwest, indicating an inflow of air mass from the surrounding forest area. However, the ratio of C_2/BSOAs did not differ between these days (3

on 10 July, 2 on 17 July) (Fig. 2). These results suggest that transport of oxalic acid from the surrounding urban area is rather minor in this forest. On the other hand, high ratio of C_2 /BSOAs (average: 11) was observed on 15 July. Because wind speed on 15 July (average 4.2 m s^{-1}) was higher than that on 5, 10, 16 and 17 July (average: 1.9 m s^{-1}), oxalic acid may have been transported from the outside of this forest on 15 July. In order to evaluate the factors controlling the production of oxalic acid in a *L. kaempferi* forest, the data on 15 July were excluded from analysis. The factors controlling the production of oxalic acid from isoprene and α -pinene via photochemical reactions with O_3 and other oxidants will be discussed in section 3.2, except for the date of 15 July.

3.2 Production of oxalic acid from isoprene and α -pinene

In our present study, a positive correlation was observed between methylglyoxal and oxalic acid ($r^2 = 0.66$) and glyoxal and oxalic acid ($r^2 = 0.80$) in a *L. kaempferi* forest (Fig. 3). Oxalic acid is a product of the chain reactions of α -dicarbonyls as suggested by a strong relation between oxalic acid and α -dicarbonyls. During the same sampling time, isoprene and α -pinene accounted for 23% and 44% of total terpenoids, respectively (Mochizuki et al., 2015); the former is mainly emitted from fern species (*D. crassirhizoma*) whereas the latter is from *L. kaempferi* (Mochizuki et al., 2014). Laboratory experiments demonstrated that aqueous or gas phase oxidation of isoprene (Paulot et al., 2009; Carlton et al., 2009; Huang et al., 2011; Kawamura et al., 2014) and α -pinene (Fick et al., 2003; Kawamura et al., unpublished data, 2016) with OH radical and/or O_3 produce methylglyoxal and glyoxal. OH radical reactions of methylglyoxal and glyoxal lead to the production of oxalic acid (Carlton et al., 2007; Tan et al., 2009; Tan et al., 2010).

In addition, oxalic acid showed a significant correlation with O_3 ($r^2 = 0.65$) (Fig. 4a). In order to examine the effect of gaseous oxidants on the formation of oxalic acid from isoprene and α -pinene, we plotted the mass concentration ratios of oxalic acid to isoprene plus

α -pinene (oxalic acid/(isoprene+ α -pinene)) against O_3 concentrations (Fig. 4b). Because oxidants such as OH radicals were not measured in this study, O_3 is used here as an indicator of the oxidant concentrations. A strong positive correlation was found between the mass concentration ratios of oxalic acid to isoprene plus α -pinene (oxalic acid/(isoprene+ α -pinene)) and O_3 concentrations ($r^2 = 0.69$) (Fig. 4b). These results suggest that O_3 and other oxidants promote the formation of oxalic acid from isoprene and α -pinene in the forest of *L. kaempferi*.

Aerosol acidity contributes to the formation of SOA from aldehydes such as glyoxal and isoprene through acid-catalyzed heterogeneous reactions (e.g., Jang et al., 2002). Sorooshian et al. (2006) and Ervens et al. (2008) reported that aerosol aqueous phase is important for the formation of oxalic acid from isoprene. In this study, we found that oxalic acid negatively correlated with pH (range: 4.2 – 6.4) of the water extracts from aerosol samples ($r^2 = 0.75$), but positively correlated with SO_4^{2-} ($r^2 = 0.38$). Further, key intermediates such as methylglyoxal plus glyoxal weakly correlated with SO_4^{2-} ($r^2 = 0.14$). Based on the field measurement, Yu et al. (2005) reported that oxalic acid in urban aerosols correlate with sulfate. Conversely, Tan et al. (2009, 2010) reported that acidity is not needed for oxalic acid formation from glyoxal and related compounds in aqueous phase. Production of oxalic acid and sulfate may simultaneously occur in the aqueous phase. In addition, oxalic acid did not show a correlation with relative humidity (RH). Zhang et al. (2011) reported that RH does not affect SOA yields via the oxidation of isoprene. In contrast, Tillmann et al. (2010) reported that high humidity enhances the SOA yields from α -pinene via ozonolysis. Higher RH may promote the partitioning of gaseous precursors of oxalic acid to aerosol phase and may enhance the acid-catalyzed heterogeneous reactions. Our results suggest that aerosol acidity in aqueous phase may play an important role in the formation of oxalic acid from isoprene and α -pinene in the forest atmosphere.

Malonic/succinic acid (C_3/C_4) ratio has been often used as an indicator of photochemical processing (aging) of organic aerosols, during which the C_3/C_4 ratio increases (e.g.,

Kawamura and Ikushima, 1993). The C_3/C_4 ratios ranged from 1.2 to 1.8 (average: 1.4) in this study. However, we did not find any diurnal and temporal trends of C_3/C_4 ratio during the measurement period, suggesting that photochemical aging of organic aerosols is not significant during an early stage of photochemical processing of aerosols although the formation of oxalic acid is important in the forest atmosphere. In addition, concentrations of oxalic acid did not show any correlations with ambient temperature ($r^2 = 0.0013$) and solar radiation ($r^2 = 0.0001$), suggesting that these parameters are not important for the formation of oxalic acid in the forest atmosphere. Our results suggest that O_3 is one of major contributors to the formation of oxalic acid from isoprene and α -pinene.

3.3 Possible formation of oxalic acid from unsaturated fatty acids

We identified unsaturated fatty acids (myristoleic: $FA_{14:1}$, palmitoleic: $FA_{16:1}$, oleic: $FA_{18:1}$, and linoleic: $FA_{18:2}$) in the ambient aerosols collected under the canopy of the *L. kaempferi* forest (Table 1). UFAs are dominant lipid class compounds that are generally present in terrestrial higher plants and algae. UFAs in TSP samples showed higher concentrations in nighttime with a drastic decline in daytime (Fig. 5). Because of a double bond, UFAs can be easily oxidized by O_3 and OH radicals in the atmosphere in daytime (Yokouchi and Ambe, 1986; Kawamura and Gagosian, 1987).

Azelaic acid (C_9) in the *L. kaempferi* forest is a dominant species in the range of C_6 - C_{10} diacids (Table 1). UFAs such as $FA_{18:1}$ and $FA_{18:2}$ have a double bond at C9-position. Azelaic and 9-oxononanoic acids are produced by the oxidation of oleic and linoleic acids via the cleavage of a double bond at C9-position. ω -Oxidation of 9-oxononanoic acid results in azelaic acid. Concentration of azelaic acid had the highest value around noon (Fig. 5). Fig. 5 shows diurnal and temporal variations of the mass concentration ratios of azelaic acid to oleic plus linoleic acids ($C_9/(FA_{18:1}+FA_{18:2})$). The ratios started to increase in the morning with a maximum around noontime or in the afternoon. The ratios of $C_9/(FA_{18:1}+FA_{18:2})$ showed a

variation pattern similar to that of O₃ concentrations with a good correlation ($r^2 = 0.56$) (Fig. 6). Azelaic acid can be derived from oleic and linoleic acids via oxidation with oxidants (Yokouchi and Ambe, 1986; Kawamura and Gagosian, 1987; Nouredдини and Kanabur, 1999). We suggest that C₉ is mainly produced by the oxidation of oleic and linoleic acids emitted from the forest ecosystem via the oxidation with O₃ and other oxidants. Because longer-chain diacids (C₃–C₉) can produce oxalic acid via photochemical breakdown with OH radicals (Kawamura et al., 1996; Pavuluri et al., 2015; Enami et al., 2015), UFAs also contribute to the formation of oxalic acid in the *L. kaempferi* forest.

3.4 Comparison with previous studies

Average contributions of oxalic acid (C₂) to total straight chain saturated diacids (C₂–C₁₀) in daytime was 50% (range: 41–65%). Contributions of C₂ to C₂–C₁₀ is lower than those of the aged organic aerosols from Chichi-jima Island (81%) in the remote western North Pacific (Mochida et al., 2003b) and Okinawa Island (74%) in the western North Pacific Rim (Kunwar and Kawamura, 2014a). On the other hand, this level is comparable to those reported from the western North Pacific during the period of high biological activity in summer (Bikkina et al., 2014) and from Tokyo in summer (55%) via photo-oxidation of anthropogenic VOCs (Kawamura and Yasui, 2005). Our result suggests that oxalic acid in this site is less aged and mainly produced via the oxidation of local BVOCs and UFAs.

Average concentration of oxalic acid in daytime was 214 ng m⁻³. The mean concentration level of oxalic acid in TSP in the *L. kaempferi* forest is comparable to or higher than those reported from mixed coniferous/deciduous forest in Hungary (196 ng m⁻³ in PM_{2.5}) (Kourtchev et al., 2009) and broad-leaved forest in Taiwan (161 ng m⁻³ in PM_{2.5}) (Tsai and Kuo, 2013). On the other hand, this level is one order of magnitude higher than those from the remote site in high Arctic Alert, Canada (17 ng m⁻³ in TSP) (Kawamura et al., 2010), but lower than those from the urban areas in 14 cities of China (513 ng m⁻³ in PM_{2.5}) (Ho et al.,

2007) and from Tokyo (270 ng m^{-3} in TSP) (Kawamura and Ikushima, 1993). These comparisons demonstrate that larch forest is an important source of oxalic acid in the atmosphere.

Monocarboxylic acids such as acetic and formic acids are produced in the atmosphere by photo-oxidation of BVOCs such as isoprene, in which acetic acid is key intermediate in the oxidation of isoprene (Ervens et al., 2008). These organic acids exist in the atmosphere as both gases (Kawamura et al., 2000; Khwaja et al., 1995; Liu et al., 2012; Yuan et al., 2015) and aerosols (Khwaja et al., 1995; Kawamura et al., 2000; Li and Winchester, 1989; Liu et al., 2012; Mochizuki et al., 2016). Although we did not measure acetic and formic acids in this study, we presume that acetic and formic acids are abundantly present in the forest atmosphere in both gas and particle phases.

Fig. 7 shows diurnal and temporal variations of concentrations of WSOC and mass concentration ratio of oxalic acid to WSOC (C_2/WSOC) for the aerosol samples collected at the *L. kaempferi* forest. WSOC concentrations started to increase in the morning with a maximum around at noontime or in the afternoon. Higher concentrations of WSOC were observed on 10 and 17 July. The variation pattern of WSOC concentrations was similar to that of oxalic acid ($r^2 = 0.64$). Overall, the peak ratios of C_2/WSOC were observed in daytime, which is consistent with O_3 concentration. Although a weak correlation was observed between the concentrations of WSOC and O_3 ($r^2 = 0.15$) (Fig. 8a), we found a good positive correlation between the ratios of C_2/WSOC and O_3 ($r^2 = 0.41$) (Fig. 8b). We suggest that secondary formation of oxalic acid from BVOCs and UFAs emitted from the forest ecosystem enhances its contribution to WSOC in the forest aerosols. The mean ratio of C_2/WSOC was 6% (range: 3.7 – 9.7%). This level is higher than that reported in aerosol samples from the western North Pacific (4.4% in TSP) (Bikkina et al., 2015), but lower than those from Okinawa Island (14% in TSP) (Kunwar and Kawamura, 2014a; Kunwar et al., 2014b) and 14 cities of China (7% in $\text{PM}_{2.5}$) (Ho et al., 2007).

4. Conclusions

Dicarboxylic acids, oxocarboxylic acids, α -dicarbonyls, and UFAs were detected in organic aerosols in a larch (*L. kaempferi*) forest. Oxalic acid was dominant organic acid followed by malonic acid. Higher concentrations of oxalic acid and key intermediate compounds were observed in daytime. The peak concentration of oxalic acid appeared after the peaks of isoprene and α -pinene, whereas BSOAs such as isoprene and α -pinene oxidation products and O_3 showed strong positive correlations with oxalic acid. Although oxalic acid is not only derived from biogenic VOCs but also from anthropogenic VOCs, isoprene and α -pinene were dominant sources of oxalic acid in the forest aerosols. O_3 and other oxidants enhanced the formation of oxalic acid in a *L. kaempferi* forest. In addition, oxalic acid is negatively correlated with pH of the water extracts from aerosol samples, and is positively correlated with SO_4^{2-} . Oxalic acid did not show any correlations with temperature, solar radiation, and relative humidity. Our results suggest that aerosol acidity of aqueous phase particle plays an important role in the formation of oxalic acid from isoprene and α -pinene in the forest atmosphere. In addition, lower concentration of UFAs was observed in daytime, whereas concentrations of azelaic acid that is a specific oxidation product of oleic and linoleic acids were higher in daytime. The mass concentration ratio of azelaic acid to UFAs showed a positive correlation with O_3 , suggesting that azelaic acid is derived from UFAs. Oxalic acid is a product of the chain reactions of azelaic acid. Oxalic acid accounted for 6% of WSOC in a *L. kaempferi* forest. Our results suggest that forest ecosystem is an important source of oxalic acid and other dicarboxylic acids in the atmosphere.

Acknowledgements

This work was supported by Japan Society for the Promotion of Science (Grant-in-Aid for JSPS Fellows 2410958, and 24221001) and Grant-in-Aid for Scientific Research (B)

(25281002 and 25281010) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan. The data of this paper are available upon request to K. Kawamura (kkawamura@isc.chubu.ac.jp).

References

- Atkinson, R., Arey, J., 2003. Gas-phase tropospheric chemistry of biogenic volatile organic compounds: a review. *Atmos. Environ.* 37 (Supplement 2), S197-S219.
- Bikkina, S., Kawamura, K., Miyazaki, Y., Fu, P.Q., 2014. High abundances of oxalic, azelaic, and glyoxylic acids and methylglyoxal in the open ocean with high biological activity: Implication for secondary SOA formation from isoprene. *Geophys. Res. Lett.* 41.
- Bikkina, S., Kawamura, K., Miyazaki, Y., 2015. Latitudinal distributions of atmospheric dicarboxylic acids, oxocarboxylic acids, and α -dicarbonyls over the western North Pacific: Sources and formation pathways. *J. Geophys. Res. Atmos.* 120, 5010-5035.
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S.K., Sherwood, S., Stevens, B., Zhang, X.Y., 2013. Clouds and Aerosols. In: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (eds.), *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Carlton, A.G., Turpin, B.J., Altieri, K.E., Seitzinger, S., Reff, A., Lim, H.J., Ervens, B., 2007. Atmospheric oxalic acid and SOA production from glyoxal: Results of aqueous photooxidation experiments. *Atmos. Environ.* 41, 7588-7602.
- Carlton, A.G., Wiedinmyer, C., Kroll, J.H., 2009. A review of secondary organic aerosol (SOA) formation from isoprene. *Atmos. Chem. Phys.* 9, 4987-5005.
- Enami, S., Hoffmann, M.R., Colussi, A.J., 2015. Stepwise oxidation of aqueous dicarboxylic acids by gas-phase OH radicals. *J. Phys. Chem. Lett.* 6, 527-534.
- Ervens, B., Carlton, A.G., Turpin, B.J., Altieri, K.E., Kreidenweis, S.M., Feingold, G., 2008. Secondary organic aerosol yields from cloud-processing of isoprene oxidation products. *Geophys. Res. Lett.* 35, L02816.
- Fick, J., Pommer, L., Nilsson, C., Andersson, B., 2003. Effect of OH radicals, relative humidity, and time on the composition of the products formed in the ozonolysis of α -pinene. *Atmos. Environ.* 37, 4087-4096.
- Giebl, H., Berner, A., Reischl, G., Puxbaum, H., Kasper-Giebl, A., Hitzemberger, R., 2002. CCN activation of oxalic and malonic acid test aerosols with the University of Vienna Cloud Condensation Nuclei Counter. *J. Aerosol. Sci.* 33, 1623-1634.
- Guenther, A.B., Jiang, X., Heald, C.L., Sakulyanontvittaya, T., Duhl, T., Emmons, L.K., Wang,

- X., 2012. The model of emissions of gases and aerosols from nature version 2.1 (MEGAN2.1): an extended and updated framework for modelling biogenic emissions. *Geosci. Model Dev.* 5, 1471-1492.
- Hallquist, M., Wenger, J.C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N.M., George, C., Goldstein, A.H., Hamilton, J.F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M.E., Jimenez, J.L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th.F., Monod, A., Prévôt, A.S.H., Seinfeld, J.H., Surratt, J.D., Szmigielski, R., Wildt, J., 2009. The formation, properties and impact of secondary organic aerosol: current and emerging issues. *Atmos. Chem. Phys.* 9, 5155-5236.
- Ho, K.F., Cao, J.J., Lee, S.C., Kawamura, K., Zhang, R.J., Chow, J.C., Watson, J.G., 2007. Dicarboxylic acids, ketocarboxylic acids, and dicarbonyls in the urban atmosphere of China. *J. Geophys. Res.* 112, D22S27.
- Huang, D., Zhang, X., Chen, Z.M., Zhao, Y., Shen, X.L., 2011. The kinetics and mechanism of an aqueous phase isoprene reaction with hydroxyl radical. *Atmos. Chem. Phys.* 11, 7399-7415.
- Jang, M., Czoschke, N.M., Lee, S., Kamens, R.M., 2002. Heterogeneous atmospheric aerosol production by acid-catalyzed particle-phase reactions. *Science*, 298, 814-817.
- Kanakidou, M., Seinfeld, J.H., Pandis, S.N., Barnes, I., Dentener, F.J., Facchini, M.C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C.J., Moortgat, G.K., Winterhalter, R., Myhre, C.E.L., Tsigaridis, K., Vignati, E., Stephanou, E.G., Wilson, J., 2005. Organic aerosol and global climate modeling: a review. *Atmos. Chem. Phys.* 5, 1053-1123.
- Kawamura, K., Kaplan, I.R., 1987. Motor exhaust emissions as a primary source for dicarboxylic-acids in Los-Angeles ambient air. *Environ. Sci. Technol.* 21, 105-110.
- Kawamura, K., Gagosian, R.B., 1987. Implication of ω -oxocarboxylic acids in the remote marine atmosphere for photooxidation of unsaturated fatty acids. *Nature* 325, 330-332.
- Kawamura, K., and Ikushima, K., 1993, Seasonal changes in the distribution of dicarboxylic acids in the urban atmosphere. *Environ. Sci. Technol.* 27, 2227-2235.
- Kawamura, K., Kasukabe, H., Barrie, L.A., 1996. Source and reaction pathways of dicarboxylic acids, ketoacids and dicarbonyls in Arctic aerosols: One year of observations. *Atmos. Environ.* 30, 1709-1722.
- Kawamura, K., Steinberg, S., Kaplan, I.R., 2000. Homologous series of C₁-C₁₀ monocarboxylic acids and C₁-C₆ carbonyls in Los Angeles and motor vehicle exhausts. *Atmos. Environ.* 34, 4175-4191.
- Kawamura K., and Yasui, O., 2005, Diurnal changes in the distribution of dicarboxylic acids, ketocarboxylic acids and dicarbonyls in the urban Tokyo atmosphere. *Atmos. Environ.* 39, 1945-1960.
- Kawamura, K., Kasukabe, H., Barrie, L.A., 2010. Secondary formation of water-soluble organic acids and α -dicarbonyls and their contributions to total carbon and water-soluble organic carbon: Photochemical aging of organic aerosols in the Arctic spring. *J. Geophys.*

- 412 Res. Atmos. 115, D21306.
- 413 Kawamura, K., Tachibana, E., Okuzawa, K., Aggarwal, S.G., Kanaya, Y., Wang, Z.F., 2013.
 414 High abundances of water-soluble dicarboxylic acids, ketocarboxylic acids and
 415 α -carbonyls in the mountaintop aerosols over the North China Plain during wheat burning
 416 season. Atmos. Chem. Phys. 13, 8285-8302.
- 417 Kawamura, K., Tachibana, E., Sakamoto, Y., Hirokawa, J., 2014. GC/MS analyses of
 418 dicarboxylic acids, oxocarboxylic acids, glyoxal and methylglyoxal in the ozone
 419 oxidation products of isoprene. Res. Org. Geochem. 30, 37-41.
- 420 Kawamura, K., Bikkina, S., 2016. A review of dicarboxylic acids and related compounds in
 421 atmospheric aerosols: Molecular distributions, sources and transformation. Atmos. Res.
 422 170, 140-160.
- 423 Khwaja, H.A., 1995. Atmospheric concentrations of carboxylic acids and related compounds
 424 at a semiurban site. Atmos. Environ. 29(1), 127-139.
- 425 Kourtchev, I., Copolovici, L., Claeys, M., Maenhaut, W., 2009. Characterization of
 426 atmospheric aerosols at a forested site in central Europe. Environ. Sci. Technol. 43,
 427 4665-4671.
- 428 Kunwar, B., and Kawamura, K., 2014a. Seasonal distributions and sources of low molecular
 429 weight dicarboxylic acids, ω -oxocarboxylic acids, pyruvic acid, α -dicarbonyls and fatty
 430 acids in ambient aerosols from subtropical Okinawa in the western Pacific Rim. Environ.
 431 Chem. 11, 673-689.
- 432 Kunwar, B., and Kawamura, K., 2014b. One-year observations of carbonaceous and
 433 nitrogenous components and major ions in the aerosols from subtropical Okinawa Island,
 434 an outflow region of Asian dusts. Atmos. Chem. Phys. 14, 1819-1836.
- 435 Li, S.M., Winchester, J.W., 1989. Geochemistry of organic and inorganic ions of late winter
 436 arctic aerosols. Atmos. Environ. 33(11), 2401-2415.
- 437 Lim, H.J., Carlton, A.G., Turpin, B.J., 2005. Isoprene forms secondary organic aerosol
 438 through cloud processing: Model simulations. Environ. Sci. Technol. 39, 4441-4446.
- 439 Lin, C., Owen, S.M., Peñuelas, J., 2007. Volatile organic compounds in the roots and
 440 rhizosphere of *Pinus* spp.. Soil Biol. Biochem. 39 (4), 951-960.
- 441 Liu, J., Zhang, X., Parker, E.T., Veres, P.R., Roberts, J.M., de Gouw, J.A., Hayes, P.L.,
 442 Jimenez, J.L., Murphy, J.G., Ellis, R.A., Huey, L.G., Weber, R.J., 2012. On the
 443 gas-particle partitioning of soluble organic aerosol in two urban atmospheres with
 444 contrasting emissions: 2. Gas and particle phase formic acid. J. Geophys. Res. 117,
 445 D00V21.
- 446 Mochida, M., Kawamura, K., Umemoto, N., Kobayashi, M., Matsunaga, S., Lim, H., Turpin,
 447 B.J., Bates, T.S., Simoneit, B.R.T., 2003a. Spatial distribution of oxygenated organic
 448 compounds (dicarboxylic acids, fatty acids and levoglucosan) in marine aerosols over the
 449 western Pacific and off coasts of East Asia: Asian outflow of organic aerosols during the
 450 ACE-Asia campaign. J. Geophys. Res. 108 (D23), 8638.

- 451 Mochida, M., Kawabata, A., Kawamura, K., 2003b. Seasonal variation and origins of
 452 dicarboxylic acids in the marine atmosphere over the western North Pacific. J. Geophys.
 453 Res. 108(D6), 4193.
- 454 Mochizuki, T., Endo, Y., Matsunaga, S., Chang, J., Ge, Y., Huang, C., Tani, A., 2011. Factors
 455 affecting monoterpene emission from *Chamaecyparis obtusa*. Geochem. J. 45, e15-e22.
- 456 Mochizuki, T., Tani, A., Takahashi, Y., Saigusa, N., Ueyama, M., 2014. Long-term
 457 measurement of terpenoid flux above a *Larix kaempferi* forest using a relaxed eddy
 458 accumulation method. Atmos. Environ. 83, 53-61.
- 459 Mochizuki, T., Miyazaki, Y., Ono, K., Wada, R., Takahashi, Y., Saigusa, N., Kawamura, K.,
 460 Tani, A., 2015. Emissions of biogenic volatile organic compounds and subsequent
 461 formation of secondary organic aerosols in a *Larix kaempferi* forest. Atmos. Chem. Phys.
 462 15, 12029-12041.
- 463 Mochizuki, T., Kawamura, K., Aoki, K., Sugimoto, N., 2016. Long-range atmospheric
 464 transport of volatile monocarboxylic acids with Asian dust over high mountain snow site,
 465 central Japan. Atmos. Chem. Phys. 16, 14621-14633.
- 466 Miyazaki, Y., Kawamura, K., Jung, J., Furutani, H., Uematsu, M., 2011. Latitudinal
 467 distributions of organic nitrogen and organic carbon in marine aerosols over the western
 468 North Pacific. Atmos. Chem. Phys. 11, 3037-3049.
- 469 Nouredдини, H., and Kanabur, M., 1999. Liquid-phase catalytic oxidation of unsaturated fatty
 470 acids. J. Amer. Oil Chem. Soc. 73(3), 305-312.
- 471 Paulot, F., Crounse, J.D., Kjaergaard, H.G., Kroll, J.H., Seinfeld, J.H., Wennberg, P.O., 2009,
 472 Isoprene photooxidation: new insights into the production of acids and organic nitrates.
 473 Atmos. Chem. Phys. 9, 1479-1501.
- 474 Pavuluri, C.M., Kawamura, K., Mihalopoulos, N., Swaminathan, T., 2015. Laboratory
 475 photochemical processing of aqueous aerosols: formation and degradation of dicarboxylic
 476 acids, oxocarboxylic acids and α -dicarbonyls. Atmos. Chem. Phys. 15, 7999-8012.
- 477 Rinaldi, M., Decesari, S., Carbone, C., Finessi, E., Fuzzi, S., Ceburnis, D., O'Dowd, C.D.,
 478 Sciare, J., Burrows, J.P., Vrekoussis, M., Ervens, B., Tsigaridis, K., 2011. Evidence of a
 479 natural marine source of oxalic acid and a possible link to glyoxal. J. Geophys. Res. 116,
 480 D16204.
- 481 Schauer, J.J., Kleeman, M.J., Cass, G.R., Simoneit, B.R.T., 2002. Measurement of emissions
 482 from air pollution sources. 4. C1-C27 organic compounds from cooking with seed oils.
 483 Environ. Sci. Technol. 36, 567-575.
- 484 Sorroshian, A., Varutbangkul, V., Brechtel, F.J., Ervens, B., Feingold, G., Bahreini, R.,
 485 Murphy, S.M., Holloway, J.S., Atlas, E.L., Buzorius, G., Jonsson H., Flagan, R.C.,
 486 Seinfeld, J.H., 2006. Oxalic acid in clear and cloudy atmospheres: Analysis of data from
 487 International Consortium for Atmospheric Research on Transport and Transformation
 488 2004. J. Geophys. Res. 111, D23S45.
- 489 Tan, Y., Perri, M.J., Seitzinger, S.P., Turpin, B.J., 2009. Effects of precursor concentration and

- acidic sulfate in aqueous glyoxal –OH radical oxidation and implications for secondary organic aerosol. *Environ. Sci. Technol.* 43, 8105-8112.
- Tan, Y., Carlton, A.G., Seitzinger, S.P., Turpin, B.J., 2010. SOA from methylglyoxal in clouds and wet aerosols: Measurement and prediction of key products. *Atmos. Environ.* 44, 5218-5226.
- Tani, A., Kawawata, Y., 2008. Isoprene emission from native deciduous *Quercus* spp. in Japan. *Atmos. Environ.* 42, 4540-4550.
- Tillmann, R., Hallquist, M., Jonsson Å.M., Kiendler-Scharr, A., Saathoff, H., Iimura, Y., Mentel, Th.F., 2010. Influence of relative humidity and temperature on the production of pinonaldehyde and OH radicals from the ozonolysis of α -pinene. *Atmos. Chem. Phys.* 10, 7057-7072.
- Tsai, Y.I., Kuo, S.C., 2013. Contributions of low molecular weight carboxylic acids to aerosols and wet deposition in a natural subtropical broad-leaved forest environment. *Atmos. Environ.* 81, 270-279.
- Urakawa, R., Ohte, N., Shibata, H., Tateno, R., Hishi, T., Fukushima, K., Inagaki, Y., Hirai, K., Oda, T., Oyanagi, N., Nakata, M., Toda, H., Tanaka, K., Fukuzawa, K., Watanabe, T., Tokuchi, N., Nakaji, T., Saigusa, N., Yamao, Y., Nakanishi, A., Enoki, T., Ugawa, S., Hayakawa, A., Kotani, A., Kuroiwa, M., Isobe, K., 2015. Biogeochemical nitrogen properties of forest soils in the Japanese archipelago. *Ecol. Res.* 30, 1-2.
- Yokouchi, Y., Ambe, Y., 1986. Characterization of polar organics in airborne particulate matter. *Atmos. Environ.* 20 (9), 1727-1734.
- Yu, J.Z., Huang, X.F., Xu, J., Hu M., 2005. When aerosol sulfate goes up, so does oxalate: Implication for the formation mechanisms of oxalate. *Environ. Sci. Technol.* 39, 128-133.
- Yuan, B., Veres, P.R., Warneke, C., Roberts, J.M., Gilman, J.B., Koss, A., Edwards, P.M., Graus, M., Kuster, W.C., Li, S.-M., Wild, R.J., Brown, S.S., Dubé, W.P., Lerner, B.M., Williams, E.J., Johnson, J.E., Quinn, P.K., Bates, T.S., Lefer, B., Hayes, P.L., Jimenez, J.L., Weber, R.J., Zamora, R., Ervens, B., Millet, D.B., Rappenglück, B., de Gouw, J.A., 2015. Investigation of secondary formation of formic acid: urban environment vs. oil and gas producing region. *Atmos. Chem. Phys.* 15, 1975-1993.
- Zhang, H., Surratt, J.D., Lin, Y.H., Bapat, J., Kamens, R.M., 2011. Effect of relative humidity on SOA formation from isoprene/NO photooxidation: enhancement of 2-methylglyceric acid and its corresponding oligoesters under dry conditions. *Atmos. Chem. Phys.* 11, 6411-6424.

Fig. 1. Diurnal and temporal variations in the concentrations of (a) total dicarboxylic acids, (b) oxocarboxylic acids, and (c) α -dicarbonyls in a *L. kaempferi* forest. Shaded areas indicate nighttime.

Fig. 2. Diurnal and temporal variations in the concentrations of (a) oxalic acid and (b) O_3 , and (c) mass concentration ratios of oxalic acid/BSOAs in a *L. kaempferi* forest. Shaded areas indicate nighttime.

Fig. 3. Correlation plot of oxalic acid and α -dicarbonyls.

Fig. 4. Correlation plots of (a) oxalic acid with O_3 and (b) mass concentration ratio of oxalic acid to isoprene plus α -pinene (oxalic acid/(isoprene+ α -pinene)) with O_3 .

Fig. 5. Diurnal and temporal variations in (a) concentrations of unsaturated fatty acids (UFAs: oleic plus linoleic acids) and azelaic acid (C₉), (b) mass concentration ratios of azelaic acid/UFAs (C₉/UFAs), and O₃ concentrations in a *L. kaempferi* forest. Shaded areas indicate nighttime.

Fig. 6. Correlation plots of concentration ratios of azelaic acid (C₉) to oleic plus linoleic acids ($C_9 / (FA_{18:1} + FA_{18:2})$) as a function of O₃.

Fig. 7. Diurnal and temporal variations in (a) concentrations of water-soluble organic carbon (WSOC) and oxalic acid, (b) mass concentration ratios of oxalic acid to WSOC ($C_2/WSOC$) and (c) O_3 concentrations in a *L. kaempferi* forest. Shaded areas indicate nighttime.

Fig. 8. Correlation plots of (a) water soluble organic carbon (WSOC) and (b) mass concentration ratios of oxalic acid to WSOC ($C_2/WSOC$) as a function of O_3 .

Table 1. Average concentrations (ng m^{-3}) and standard deviation (S.D.) of low molecular weight dicarboxylic acids, oxocarboxylic acids, α -dicarbonyls, and unsaturated fatty acids in a *L. kaempferi* forest.

Compounds	Daytime		Nighttime	
	Average	S.D.	Average	S.D.
Dicarboxylic acids				
Oxalic, C ₂	214	111	162	124
Malonic, C ₃	100	63.9	52.6	42.0
Succinic, C ₄	75.0	44.5	32.9	24.1
Glutaric, C ₅	11.4	7.9	5.2	3.8
Adipic, C ₆	4.4	3.7	2.0	1.5
Pimelic, C ₇	1.7	2.2	1.4	1.7
Suberic, C ₈	4.4	2.9	1.5	1.2
Azelaic, C ₉	9.9	6.1	7.7	5.8
Sebacic, C ₁₀	3.3	5.5	0.5	0.2
Total C ₂ -C ₁₀	424		266	
Oxocarboxylic acids				
Glyoxylic, ω C ₂	18.8	11.2	12.1	10.0
3-Oxopropanoic, ω C ₃	7.5	5.2	5.1	4.4
4-Oxobutanoic, ω C ₄	13.0	9.1	7.5	7.4
5-Oxopentanoic, ω C ₅	2.6	1.6	1.6	1.3
6-Oxohexanoic, ω C ₆	2.2	1.4	2.6	2.1
7-Oxoheptanoic, ω C ₇	7.7	4.3	5.5	4.7
8-Oxooctanoic, ω C ₈	6.3	4.1	4.7	4.4
9-Oxononanoic, ω C ₉	2.6	1.7	2.8	2.0
Pyruvic, Pyr	11.8	8.7	7.0	6.5
Total oxoacids	72		49	
α-Dicarbonyls				
Glyoxal, Gly	5.9	3.8	3.7	2.6
Methylglyoxal, MeGly	7.4	4.0	4.6	3.3
Total α -dicarbonyls	13		8	
Unsaturated fatty acids				
Myristoleic, FA _{14:1}	0.8	0.8	1.9	0.9
Palmitoleic, FA _{16:1}	5.4	4.8	8.9	5.5
Linoleic, FA _{18:2}	5.9	4.4	12.8	5.5
Oleic, FA _{18:1}	12.4	6.5	31.1	17.9
Total unsaturated fatty acids	25		55	