

# HOKKAIDO UNIVERSITY

| Title                  | Staufen1, Kinesin1 and microtubule function in cyclin B1 mRNA transport to the animal polar cytoplasm of zebrafish oocytes           |
|------------------------|--|
| Author(s)              | Takahashi, Kazuki; Ishii, Kana; Yamashita, Masakane  |
| Citation               | Biochemical and biophysical research communications, 503(4), 2778-2783<br>https://doi.org/10.1016/j.bbrc.2018.08.039                 |
| Issue Date             | 2018-09-18   |
| Doc URL                | http://hdl.handle.net/2115/75463   |
| Rights                 | ©2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/ |
| Rights(URL)            | http://creativecommons.org/licenses/by-nc-nd/4.0/  |
| Туре                   | article (author version)   |
| Additional Information | There are other files related to this item in HUSCAP. Check the above URL.   |
| File Information       | Takahashi_Text_Figs.pdf  |



| 1  | Staufen1, Kinesin1 and microtubule function in cyclin B1 mRNA transport to the animal polar                      |
|----|--|
| 2  | cytoplasm of zebrafish oocytes   |
| 3  |  |
| 4  | Kazuki Takahashi <sup>a</sup> , Kana Ishii <sup>a</sup> and Masakane Yamashita <sup>b,*</sup>                    |
| 5  |  |
| 6  | <sup>a</sup> Laboratory of Reproductive and Developmental Biology, Graduate School of Life Science, Hokkaido     |
| 7  | University, Sapporo 060-0810, Japan.   |
| 8  | <sup>b</sup> Laboratory of Reproductive and Developmental Biology, Department of Biological Sciences, Faculty of |
| 9  | Science, Hokkaido University, Sapporo 060-0810, Japan  |
| 10 |  |
| 11 | *Corresponding author.   |
| 12 | Address: Laboratory of Reproductive and Developmental Biology, Department of Biological Sciences,                |
| 13 | Faculty of Science, Hokkaido University, North 10, West 8, Sapporo 060-0810, Japan.                              |
| 14 | Phone: +81-11-706-4454; Fax: +81-11-706-4456   |
| 15 | E-mail address: myama@sci.hokudai.ac.jp (M. Yamashita)   |
| 16 |  |

# **1 ABSTRACT**

- 2 In zebrafish oocytes, cyclin B1 mRNAs are transported to the animal polar cytoplasm. To elucidate the
- 3 molecular basis of *cyclin B1* mRNA transport, we analyzed zebrafish Staufen1, a protein known to play a
- 4 central role in mRNA transport to the vegetal pole of *Xenopus* oocytes. Zebrafish Staufen1 interacts with
- 5 cyclin B1 mRNA throughout oocyte growth. Both cyclin B1 mRNA and Staufen1 are evenly distributed in
- 6 the cytoplasm of young oocytes but are co-localized to the animal polar cytoplasm in later stages.
- 7 Real-time imaging showed that the plus ends of oocyte microtubules are free in the cytoplasm in early
- 8 stages but anchored to the animal polar cytoplasm in later stages. Transport of *cyclin B1* reporter mRNA
- 9 to the animal polar cytoplasm was inhibited by disruption of microtubules and injection of antibodies
- against Staufen1 or Kinesin1, a plus-end-directed microtubule motor that interacts with Staufen1,
- 11 indicating that the transport depends on movement along microtubules toward the plus ends. Reporter
- 12 mRNAs with an element required for the vegetal localization of vg1 mRNA in Xenopus oocytes were
- 13 localized to the animal polar cytoplasm in zebrafish oocytes, indicating that the element is functional for
- 14 animal polar localization in zebrafish oocytes. Our findings suggest that *cyclin B1* mRNA-Staufen1
- 15 protein complexes are transported toward the animal pole of zebrafish oocytes by the plus-end-directed
- 16 motor protein Kinesin1 along microtubules and that a common mRNA transport machinery functions in
- 17 zebrafish and *Xenopus* oocytes, although its transport direction is opposite due to different organizations
- 18 of microtubules.
- 19
- 20 Keywords: Kinesin1, microtubule, mRNA transport, Staufen1, zebrafish oocyte
- 21
- 22

### **1 1. Introduction**

2 Specific subcellular localization of mRNAs enables spatio-temporal control of gene expression, which

- 3 assures normal progression of various biological events. Cyclin B1 protein, a regulatory subunit of
- 4 maturation/M-phase-promoting factor, is absent in immature zebrafish oocytes [1,2], because its mRNA is
- 5 translationally repressed and localized as aggregates to the animal polar cytoplasm until the oocytes are
- 6 stimulated by maturation-inducing hormone [3,4]. We previously reported that cyclin B1 mRNA
- 7 localization in zebrafish oocytes and its translational timing after hormonal stimulation are regulated by at
- 8 least two components, a *cis*-acting element situated in the open-reading frame (ORF) and Pumilio1
- 9 (Pum1) bound to the Pumilio-binding element (PBE) in the 3' UTR of *cyclin B1* mRNA, based on the
- 10 findings that *cyclin B1* reporter mRNA with mutations in the ORF element and PBE failed to stay in the
- animal polar cytoplasm and underwent precocious translational activation [5,6,7]. However, the mutant
- 12 mRNAs were transported from the nucleus to the animal polar cytoplasm, although they were not
- 13 anchored there [6]. These findings indicate that the mechanisms of cyclin B1 mRNA transport is
- 14 different from those of its anchoring to the animal polar cytoplasm. The transport mechanism of *cyclin B1*
- 15 mRNA in zebrafish oocytes is largely unknown.
- 16 We previously identified Insulin-like growth factor 2 mRNA-binding protein 3 (IMP3) as a *cyclin B1*
- 17 mRNA-binding protein and suggested its involvement in translational regulation [8]. In *Xenopus* oocytes,
- 18 IMP3 plays a crucial role in the localization of *vg1* mRNA to the vegetal cortex [9,10]. In zebrafish
- 19 oocytes, however, IMP3 and its binding targets, *vg1* mRNA and *cyclin B1* mRNA, are localized to the
- 20 animal polar cytoplasm but not to the vegetal polar cytoplasm [8]. Therefore, it remains elusive whether a
- 21 pathway similar to that in *Xenopus* oocytes exists for the animal polar localization of *vg1* and *cyclin B1*
- 22 mRNAs in zebrafish oocytes.
- Staufen1 (Stau1) is thought to play an important role in mRNA transport to the vegetal pole of
   *Xenopus* oocytes, in collaboration with microtubules and Kinesin1 [11]. In this study, we investigated the
- 25 involvement of Stau1, Kinesin1 and microtubule in cyclin B1 mRNA transport to the animal polar
- 26 cytoplasm in zebrafish oocytes, with reference to the generality and diversity of mRNA transport
- 27 mechanisms in fish and amphibian oocytes.
- $\mathbf{28}$

#### **29 2.** Materials and methods

30 2.1. Oocyte culture and extraction

All animal experiments in this study were approved by the Committee on Animal Experimentation,
Hokkaido University (permission No. 13-0099). Oocyte culture and extraction were carried out as
described previously [8]. For immunoprecipitation analysis, Triton X-100 (1% at the final concentration)
was added to the extracts.

35

#### 1 2.2. Production of recombinant proteins and antibodies

- 2 To produce N-terminally Flag-tagged Stau1 (Flag-Stau1), an ORF of zebrafish Stau1 was amplified 3 with a primer set of Stau1 F and Stau1 R (Supplementary Table 1), which were designed according to a 4 sequence, NM\_205561.1, deposited in the NCBI nucleotide database. The PCR product was inserted into 5 the *EcoR*I-digested pCS2+FT-N vector [12] using an In-Fusion HD Cloning Kit (TaKaRa). 6 A cDNA fragment of zebrafish Stau1 was amplified with a primer set of Stau1 exF and Stau1 exR 7 (Supplementary Table 1) and inserted into the pENTR/D-TOPO Gateway vector (Invitrogen). The 8 resulting plasmids were recombined with the destination vector pET161-DEST using a Gateway cloning 9 system (Invitrogen) to produce Stau1 tagged with a polyhistidine (His) at the C terminus (Stau1-His). 10 Stau1-His was expressed in E. coli BL21 (DE3) and purified according to the method described 11 previously [13]. Polyclonal antibodies against Stau1-His were produced in mice (anti-Stau1 (M)) and 12 guinea pigs (anti-Stau1 (GP)) and affinity-purified with antigenic Stau1-His protein as described 13 previously [8]. Similarly, a mouse polyclonal antibody against End-binding protein 1 (EB1) was raised 14 against zebrafish EB1 (NM 213640.1). In brief, a cDNA fragment of zebrafish EB1 was amplified with a 15primer set of EB1 exF and EB1 exR (Supplementary Table 1). E. coli-produced EB1-His proteins were
- 16 17

### 18 2.3. Immunoblotting and immunoprecipitation

injected into mice.

Immunoblotting was performed as described previously [14], using anti-Stau1 (M), anti-Stau1 (GP),
anti-EB1, anti-IMP3 [8], anti-α-Tubulin (T9026; Sigma-Aldrich), anti-Pum1 [5] or anti-Flag M2 (F1804;

21 Sigma-Aldrich) antibodies. All immunoprecipitation assays in this study were performed using crude

extracts equivalent to 10-30 full-grown immature (stage IV) oocytes (100-300 µg proteins). Interaction of

23 Stau1 with Pum1 or IMP3 was examined by co-immunoprecipitation assays, as described previously [8].

24

25 2.4. RT-PCR and quantitative real-time RT–PCR (qRT-PCR)

Oocyte extracts (100 µg protein) were subjected to RT-PCR [8], using the primer sequences shown in
Supplementary Table 1. For qRT-PCR analysis, cDNA was synthesized with a PrimeScript RT Master
Mix (Perfect Real Time) (TaKaRa) according to the manufacturer's instructions. Real-time PCR
amplification was performed with a SYBR Premix Ex Taq II (Tli RNaseH Plus) (TaKaRa) on Thermal
Cycler Dice Real Time System II (TaKaRa).

31

32 2.5. Section in situ hybridization, immunostaining and whole-mount in situ hybridization

Histological sections were simultaneously analyzed by *in situ* hybridization and immunostaining [5].

- 34 The sections were observed on an LSM 5 LIVE microscope equipped with Plan-Apochromat 10x/0.45
- 35 M27 and Plan-Apochromat 20x/0.8 M27 objective lenses (Carl Zeiss). Images were acquired with LSM 5

DUO 4.2 software (Carl Zeiss) and processed with Photoshop and Illustrator (Adobe). Whole-mount *in situ* hybridization was performed according to the procedure reported previously [15].

3

#### 4 2.6. Production of Reporter mRNAs

5 A construct containing vg1/vegetal localization element (VLE), a *cis*-acting element responsible for 6 the transport of *vg1* mRNA to the vegetal pole of *Xenopus* oocytes [11], was produced by replacing the 7 cyclin B1 3' UTR of tgo3' [6] with the Xenopus VLE sequence [11,16]. Using total cDNA from Xenopus 8 oocytes as a template, *Xenopus* VLE was amplified with a primer set of xVg1VLE\_F and xVg1VLE\_R 9 and cloned into the tgo3' vector amplified with a primer set of cyclinB1\_ORF\_R and 10 pT2KXIG∆in BgIII F (Supplementary Table 1) using an In-Fusion HD Cloning Kit (TaKaRa). The 11 SV40 reporter construct was produced as described previously [7]. 12 For an in vivo reporter mRNA localization assay, full-grown immature (stage IV) oocytes were 13 co-injected with distilled water containing 250 pg of reporter vectors that encode gfp-cyclin B1 mRNA 14 and either 10 ng/µl of nocodazole, 100 ng/µl of anti-GST [14], anti-Kinesin1 (SUK4; DSHB; reported to

15 inhibit the Kinesin1 function in *Xenopus* oocytes [11,17]) or anti-Stau1 (M) antibodies. After injection,

16 oocytes were incubated in culture medium for 4 hr at room temperature. The oocytes were fixed with 4%

17 paraformaldehyde and subjected to whole-mount *in situ* hybridization. To quantify reporter mRNAs

18 produced from the injected vectors, total RNA extracted from 10 injected oocytes was subjected to

19 qRT-PCR using primer sets for *gfp* and  $\beta$ -*actin* (Supplementary Table 1). The content of *gfp-cyclin B1* 

20 reporter mRNA was normalized to that of  $\beta$ -actin mRNA.

 $\mathbf{21}$ 

22 2.7. Real-time imaging of the plus end of microtubules in oocytes

A transgenic zebrafish that expresses EB1-GFP [Tg(-0.5zp3b:Mmu.Mapre1-GFP)/nub1Tg] [18] was
provided from National BioResource Project (NBRP) Zebrafish, Japan. The movement of EB1-GFP
signals in stage I and IV oocytes was traced under a confocal laser microscope LSM 880 with Airyscan
(Carl Zeiss).

27

# **28 3. Results**

29 3.1. Behavior of zebrafish Staul during oocyte growth

30 Newly produced anti-Stau1antibodies recognized a single band of ca. 70 kDa, the size of which is

31 comparable to that of Stau1 consisting of 702 amino acids, in oocyte extracts and immunoprecipitates

32 (Supplementary Fig. 1A and B). Since Flag-Stau1 overexpressed in zebrafish oocytes also showed a

33 similar size (cf. Fig. 1B), we conclude that the 70-kDa band is Stau1 in zebrafish oocytes.

The expression levels of Stau1 during oocyte growth were examined by immunoblotting using oocytes classified into 4 stages. Stages I to III are growing and stage IV is full-grown immature oocytes [19]. 1 Zebrafish Stau1 was detected in stage I oocytes, although its level was very low. The expression levels of

2 Staul increased rapidly in stage II oocytes but decreased gradually as the oocyte increased in size (Fig.

3 1A). The expression levels of  $\alpha$ -Tubulin were low in stage I oocytes and increased rapidly in stage II. In

4 contrast to Stau1, however, the expression levels of  $\alpha$ -Tubulin increased during the oocyte growth (Fig.

- 5 1A).
- 6 7

## 3.2. Interaction of Stau1 with cyclin B1 mRNA in oocytes

8 Anti-Flag immunoprecipitates from Flag-Stau1-overexpressing oocytes but not control precipitates 9 contained cyclin B1 mRNA (Fig. 1B), indicating that Flag-Stau1 binds to cyclin B1 mRNA in oocytes. 10 Since Pum1 [5] and IMP3 [8] are cyclin B1 mRNA-binding proteins in zebrafish oocytes, we examined 11 the interaction of endogenous proteins, between Stau1and Pum1 (Fig. 1C) or IMP3 (Fig. 1D) in oocytes 12by a co-immunoprecipitation assay. The results clearly showed that the interaction of Stau1 with Pum1 or 13IMP3. The interaction disappeared by RNase treatment, indicating that Stau1 does not bind directly to 14 Pum1 or IMP3 but binds via cyclin B1 or other mRNAs. Interaction of endogenous Stau1 with cyclin B1 15 mRNA during oocyte growth (stages I to IV) was analyzed by immunoprecipitation followed by RT-PCR. 16 In all of the stages, the interaction was detected (Fig. 1E).

17

## 18 *3.3 Subcellular distribution of Stau1 and cyclin B1 mRNA during oocyte growth*

The subcellular distribution of *cyclin B1* mRNA changes during oocyte growth in zebrafish [3]. Since
Stau1 continues to interact with *cyclin B1* mRNA (Fig. 1E), we investigated the subcellular distribution of
Stau1 in oocytes from stages I to IV by immunostaining with anti-Stau1 (M) antibody. Control

22 experiments without anti-Stau1 (M) antibody showed no clear signals. Control hybridization with the

23 DIG-labeled sense *gfp* probe also showed no specific signals.

Anti-Stau1 (M) immunostaining revealed changes in subcellular distribution of Stau1 during oogenesis (Fig. 2). In stage I oocytes, Stau1 was distributed throughout the cytoplasm like *cyclin B1* mRNA, with strong signals at the cortical cytoplasm of oocytes (Fig. 2A1-3). In stage II oocytes, some signals of both *cyclin B1* mRNA and Stau1 were localized to the animal polar cytoplasm (Fig. 2B1-3). In stage III oocytes, the vast majority of *cyclin B1* mRNA was localized to the animal polar cytoplasm, while Stau1 was distributed around the entire oocyte cortex and the periphery of the nucleus (Fig. 2C1-6). In stage IV oocytes, the animal polar distributions of both *cyclin B1* mRNA and Stau1 became tighter than those in

31 stage III oocytes (Fig. 2D1-6).

32

- 34 cytoplasm
- 35 When a DNA construct that expresses *gfp-cyclin B1* reporter mRNA (Fig. 3A, 3' UTR) was injected

<sup>33</sup> *3.4. Involvement of microtubule, Kinesin1 and Stau1 in cyclin B1 mRNA transport to the animal polar* 

1 into the nucleus of full-grown zebrafish oocytes, reporter mRNAs transcribed in the nucleus were

2 transported and anchored to the animal polar cytoplasm and formed aggregates, as in the case of

- 3 endogenous cyclin B1 mRNAs (Fig. 3A and B) [6,7]. We therefore used this system to examine the
- 4 involvement of microtubule, Stau1 and Kinesin1 in cyclin B1 mRNA transport in zebrafish oocytes, since
- 5 these molecules are known to function in mRNA transport in *Xenopus* oocytes [11].

6 Nocodazole abolished the transport of reporter mRNAs to the animal polar cytoplasm (Supplementary

7 Fig. 2A), suggesting that the transport depends on microtubules, in contrast to anchoring to the animal

8 polar cytoplasm which depends on microfilaments but not on microtubules [3,5,7]. In comparison with

9 water or anti-GST antibody as controls, anti-Kinesin1 or anti-Stau1 antibodies inhibited the transport of

- 10 reporter mRNAs (Supplementary Fig. 2A). In these experiments, expression levels of reporter mRNAs
- 11 showed no statistically significant differences (Supplementary Fig. 2B). These results suggest the

12 involvement of microtubule, Kinesin1 and Stau1 in mRNA transport in zebrafish oocytes, as in the case of

- 13 Xenopus.
- 14

### 15 *3.5. VLE-directed mRNA transport in zebrafish oocytes*

16 In Xenopus oocytes, Stau1 transports vg1 mRNA to the vegetal pole via a cis-acting element of the 17 mRNA called VLE [11]. We analyzed the behavior of *gfp-cyclin B1* reporter mRNA harboring VLE 18 instead of cyclin B1 3' UTR (Fig. 3A-C). Consistent with results previously reported [5,7], gfp-cyclin B1 19 reporter mRNAs with the 3' UTR (a positive control) showed aggregated signals localized to the animal 20 polar cytoplasm, whereas those harboring SV40 polyadenylation signal (a negative control) were not  $\mathbf{21}$ localized to any regions. The reporter mRNAs harboring VLE were apparently localized to the animal 22 polar cytoplasm. Although the localization rate was lower than that of the reporter mRNAs harboring 23cyclin B1 3' UTR, this finding suggests that the VLE sequence is functional in zebrafish oocytes as a  $\mathbf{24}$ cis-acting element for animal polar localization, different from its original function as an element for 25 vegetal polar localization in Xenopus oocytes.

We checked interaction between Stau1 and reporter mRNAs by anti-Stau1 immunoprecipitation followed by RT-PCR. Both VLE and 3' UTR reporter mRNAs, but not SV40 reporter mRNA as a control, were detected in anti-Stau1 immunoprecipitates (Fig. 3D), indicating that Stau1 binds to VLE reporter mRNA. Thus, it is plausible that VLE is recognized by Stau1 that is a component of the mRNA transport machinery to the animal pole of zebrafish oocytes.

31

# 32 *3.6. Microtubule orientation in zebrafish oocytes*

Since our results suggested that Kinesin1, a motor protein that drives cargos toward the plus ends of
 microtubules, is involved in *cyclin B1* mRNA transport (Supplementary Fig. 2A), we investigated the
 orientation of microtubules in zebrafish oocytes by immunostaining of EB1, a microtubule plus-end

1 tracking protein [20]. A newly produced antibody against zebrafish EB1 (Supplementary Fig. 1C)

2 revealed that EB1 was accumulated in the animal polar cytoplasm of full-grown immature (stage IV)

3 oocytes, whereas it showed no clear accumulation in other regions including the vegetal polar cytoplasm
4 (Fig. 4).

5 We also observed the behavior of plus end of microtubules by real-time imaging of transgenic 6 zebrafish oocytes that express EB1-GFP [18]. EB1-GFP signals were evenly distributed in stage I oocytes 7 (Supplementary Fig. 3A). In stage IV oocytes, however, some of them were accumulated to the animal 8 polar cytoplasm (Supplementary Fig. 3B). We also found that EB1-GFP signals move randomly in the 9 whole cytoplasm of stage I oocytes (Supplementary Fig. 3C, Supplementary Movie 1), while they are 10 almost immovable in the animal polar cytoplasm in stage IV oocytes (Supplementary Fig. 3D, 11 Supplementary Movie 2). These findings indicate that the orientation of microtubules is not fixed in 12young oocytes because their plus ends can move freely in the oocytes. In later stages, the plus ends 13become anchored to the animal polar cytoplasm, which allows microtubules to organize their plus ends 14 toward the animal pole.

15

#### **16 4. Discussion**

17 We showed that *gfp-cyclin B1* reporter mRNA harboring VLE, a sequence essential for vegetal polar 18 transport of vg1 mRNA in Xenopus oocytes, was transported to the animal polar cytoplasm in zebrafish 19 oocytes (Fig. 3C). These results suggest that a conserved mechanism functions in the intracellular 20 transport of zebrafish cyclin B1 mRNA and Xenopus vg1 mRNA, although the directions of transport are 21 opposite (animal pole in zebrafish and vegetal pole in *Xenopus*). Concerning the difference in direction, it 22 is noteworthy that the microtubule plus end-tracking protein EB1 accumulates in the animal polar 23cytoplasm of full-grown immature zebrafish oocytes (Fig. 4). In full-grown immature Xenopus oocytes,  $\mathbf{24}$ the majority of microtubules are oriented with the minus ends towards the cortex and a subpopulation of 25 microtubules with plus ends directs toward the vegetal pole [17,21]. Although the organization of 26 microtubules in zebrafish oocytes remains to be examined in detail, differences in the direction of mRNA 27 transport might be simply explainable by differences in the microtubule organization between zebrafish 28 and Xenopus oocytes.

At least two types of Stau1 can be recognized in zebrafish oocytes: one is co-localized to *cyclin B1* mRNA and the other is not (Fig. 2). Stau1 should interact with various mRNAs in oocytes, besides *cyclin B1* mRNA. Indeed, we found that Stau1 interacts with *zorba*, *mos*, *vg1* and *buc* mRNAs but not with *dazl*, *brul* and *vasa* mRNAs in zebrafish oocytes (data not shown). According to the localization of these mRNAs in oocytes [22,23,24], it is likely that Stau1 preferentially binds to mRNAs that are finally localized to the animal polar cytoplasm of full-grown immature oocytes, although molecular mechanism

that allows Stau1 to bind to specific mRNAs remains a mystery. Our findings suggest the involvement of

1 Stau1 in cyclin B1 mRNA transport. However, it is also plausible that other molecules including IMP3

- 2 contribute to cyclin B1 mRNA transport, since IMP3 is bound to Stau1 via cyclin B1 mRNA (Fig. 1D)
- 3 and localized to the animal polar cytoplasm [8]. In addition, our preliminary experiment showed that an
- 4 anti-IMP3 antibody inhibits the animal polar localization of *cyclin B1* mRNA. A better understanding of
- 5 the molecular composition of *cyclin B1* mRNA transport machinery is necessary.

6 We found that cyclin B1 mRNA and Stau1 are bound to each other (Fig. 1E) but distributed throughout 7 the cytoplasm of stage I oocytes (Fig. 2A), suggesting that the binding of Stau1 to cyclin B1 mRNA is 8 insufficient to initiate mRNA transport. What is required to initiate cyclin B1 mRNA transport? The finding 9 that protein levels of α-Tubulin are low in stage I oocytes and increase with oocyte growth (Fig. 1A), 10 raising the possibility that microtubules are not fully organized to a sufficient level for transporting the 11 Stau1-cyclin B1 mRNA complex to the animal pole in stage I oocytes. Real-time imaging of the plus-ends 12 of microtubules revealed that they move randomly in stage I oocytes (Supplementary Fig. 3C and 13Supplementary Movie 1), indicating that the direction of microtubules is not fixed in this stage. Further 14 studies are required to understand what organization of microtubules is required for cyclin B1 mRNA 15 transport and what molecules are involved in it. Especially, the findings in this study suggest that anchoring 16 of the plus ends of microtubules to the animal polar cytoplasm is a key event for the initiation of cyclin B1 17 mRNA transport (Fig. 4). Our next goal is to determine the anchoring mechanism.

18

#### **19** Acknowledgements

We are grateful to Prof. Masahiko Hibi, Nagoya University, for providing EB1-GFP zebrafish. We also
thank Drs. Yoshinao Katsu, Tomoya Kotani and Katsueki Ogiwara, Hokkaido University, for helpful

suggestions. This work was supported by a Grant-in-Aid for JSPS Fellows (15J01844 to KT) and a

23 Grant-in-Aid for Scientific Research (23370027 to MY) from the Ministry of Education, Culture, Sports,

- 24 Science and Technology, Japan.
- $\mathbf{25}$

### **26** References

- [1] Y. Nagahama, M. Yamashita, Regulation of oocyte maturation in fish., Dev Growth Differ. 50 Suppl 1
  (2008) S195-219.
- [2] M. Yamashita, Toward modeling of a general mechanism of MPF formation during oocyte maturation
   in vertebrates, Zoological Science 17 (2000) 841-851.
- [3] T. Kondo, T. Kotani, M. Yamashita, Dispersion of cyclin B mRNA aggregation is coupled with
   translational activation of the mRNA during zebrafish oocyte maturation., Dev Biol. 229 (2001)
   421-431.
- [4] T. Kondo, T. Yanagawa, N. Yoshida, M. Yamashita, Introduction of cyclin B induces activation of the
   maturation-promoting factor and breakdown of germinal vesicle in growing zebrafish oocytes

unresponsive to the maturation-inducing hormone., Dev Biol. 190 (1997) 142-152.

- [5] T. Kotani, K. Yasuda, R. Ota, M. Yamashita, Cyclin B1 mRNA translation is temporally controlled
   through formation and disassembly of RNA granules., J Cell Biol. 202 (2013) 1041-1055.
- [6] K. Yasuda, T. Kotani, R. Ota, M. Yamashita, Transgenic zebrafish reveals novel mechanisms of
  translational control of cyclin B1 mRNA in oocytes., Dev Biol. 348 (2010) 76-86.
- [7] K. Yasuda, T. Kotani, M. Yamashita, A cis-acting element in the coding region of cyclin B1 mRNA
  couples subcellular localization to translational timing., Dev Biol. 382 (2013) 517-529.
- [8] K. Takahashi, T. Kotani, Y. Katsu, M. Yamashita, Possible involvement of insulin-like growth factor 2
  mRNA-binding protein 3 in zebrafish oocyte maturation as a novel cyclin B1 mRNA-binding protein
  that represses the translation in immature oocytes., Biochem Biophys Res commun. 448 (2014) 22-27.
- 11 [9] J.O. Deshler, M.I. Highett, T. Abramson, B.J. Schnapp, A highly conserved RNA-binding protein for
- 12 cytoplasmic mRNA localization in vertebrates., Curr Biol. 8 (1998) 489-496.
- 13 [10] A. Git, R. Allison, E. Perdiguero, A.R. Nebreda, E. Houliston, N. Standart, Vg1RBP phosphorylation
- by Erk2 MAP kinase correlates with the cortical release of Vg1 mRNA during meiotic maturation of
   Xenopus oocytes., RNA 15 (2009) 1121-1133.
- [11] Y.J. Yoon, K.L. Mowry, Xenopus Staufen is a component of a ribonucleoprotein complex containing
   Vg1 RNA and kinesin., Development 131 (2004) 3035-3045.
- [12] R. Ota, T. Kotani, M. Yamashita, Possible involvement of Nemo-like kinase 1 in Xenopus oocyte
   maturation as a kinase responsible for Pumilio1, Pumilio2, and CPEB phosphorylation., Biochemistry
   50 (2011) 5648-5659.
- [13] R. Ota, K. Suwa, T. Kotani, K. Mita, M. Yamashita, Possible involvement of phosphatidylinositol
  3-kinase, but not protein kinase B or glycogen synthase kinase 3 β, in progesterone-induced oocyte
  maturation in the Japanese brown frog, Rana japonica., Zool Sci. 25 (2008) 773-781.
- [14] R. Ota, T. Kotani, M. Yamashita, Biochemical characterization of Pumilio1 and Pumilio2 in Xenopus
   oocytes., J Biol Chem. 286 (2011) 2853-2863.
- [15] S. Schulte-Merker, R.K. Ho, B.G. Herrmann, C. Nüsslein-Volhard, The protein product of the
  zebrafish homologue of the mouse T gene is expressed in nuclei of the germ ring and the notochord of
  the early embryo., Development 116 (1992) 1021-1032.
- [16] K.L. Mowry, D.A. Melton, Vegetal messenger RNA localization directed by a 340-nt RNA sequence
   element in Xenopus oocytes., Science 255 (1992) 991-994.
- [17] T.J. Messitt, J.A. Gagnon, J.A. Kreiling, C.A. Pratt, Y.J. Yoon, K.L. Mowry, Multiple kinesin motors
   coordinate cytoplasmic RNA transport on a subpopulation of microtubules in Xenopus oocytes., Dev
   Cell 15 (2008) 426-436.
- [18] L.D. Tran, H. Hino, H. Quach, S. Lim, A. Shindo, Y. Mimori-Kiyosue, M. Mione, N. Ueno, C.
  Winkler, M. Hibi, K. Sampath, Dynamic microtubules at the vegetal cortex predict the embryonic axis

<sup>1</sup> 

- 1 in zebrafish, Development 139 (2012) 3644-3652.
- 2 [19] K. Selman, R.A. Wallace, A. Sarka, X. Qi, Stages of oocyte development in the zebrafish,
- Brachydanio rerio, J Morphol. 218 (1993) 203-224.
- 4 [20] Y. Mimori-Kiyosue, N. Shiina, S. Tsukita, The dynamic behavior of the APC-binding protein EB1 on
  5 the distal ends of microtubules., Curr Biol. 10 (2000) 865-868.
- 6 [21] D.C. Pfeiffer, D.L. Gard, Microtubules in Xenopus oocytes are oriented with their minus-ends
  7 towards the cortex., Cell Motil Cytoskeleton 44 (1999) 34-43.
- 8 [22] C. Howley, R.K. Ho, mRNA localization patterns in zebrafish oocytes., Mech Dev. 92 (2000)
  9 305-309.
- [23] K. Kosaka, K. Kawakami, H. Sakamoto, K. Inoue, Spatiotemporal localization of germ plasm RNAs
   during zebrafish oogenesis., Mech Dev. 124 (2007) 279-289.
- 12 [24] H. Suzuki, T. Tsukahara, K. Inoue, Localization of c-mos mRNA around the animal pole in the
- 13 zebrafish oocyte with Zor-1/Zorba., BioSci Trends 3 (2009) 96-104.
- 14

15

### **1** Figure legends

2

3 Fig. 1. Staufen1 protein (Stau1) during zebrafish oocyte growth. (A) Oocyte extracts from stages I to IV

4 were analyzed by immunoblotting with anti-Stau1 (GP) and anti-α-Tubulin antibodies. (B) Stage IV

- 5 oocytes overexpressed with (+) or without (-) Flag-Stau1 were immunoprecipitated with (+) or without (-)
- 6 anti-Flag M2 antibody (Flag IP). Flag-Stau1 protein and cyclin B1 mRNA in the immunoprecipitates were
- 7 detected by immunoblotting with anti-Flag M2 antibody and RT-PCR, respectively. (C) Extracts from
- 8 stage IV oocytes were treated with (+) or without (-) RNase A and immunoprecipitated with anti-Pum1,
- 9 anti-Stau1 or control antibodies (-). Pum1 and Stau1 in the immunoprecipitates were detected by
- 10 immunoblotting. (D) Experiments similar to those in (C) using anti-IMP3 and anti-Stau1 antibodies. (E)
- 11 Extracts from stage I to IV oocytes were immunoprecipitated (Stau1 IP) with anti-Stau1 (+) or control
- 12 antibodies (-) and *cyclin B1* mRNA in the immunoprecipitates was detected by RT-PCR. Extracts from
- 13 stage IV oocytes before immunoprecipitation were also subjected to RT-PCR to confirm the presence of
- 14 *cyclin B1* mRNA in the original extracts (initial).
- 15

16 **Fig. 2.** Distribution of *cyclin B1* mRNA and Stau1 protein during zebrafish oocyte growth. Paraffin

17 sections of zebrafish oocytes were simultaneously analyzed by *in situ* hybridization and immunostaining.

18 Hoechst 33258 staining of DNA (C, D), *in situ* hybridization of *cyclin B1* mRNA (A1; B1; C1, 4; D1, 4),

19 immunostaining of Stau1 (A2; B2; C2, 5; D2, 5) and their merged and magnified images (A3; B3; C3, 6;

20 D3, 6) are shown. CA, cortical alveolus; EC, egg chorion; FC, follicle cell; GV, germinal vesicle; MP,

- 21 micropyle; YG, yolk granule. Scale bars, 100 µm.
- $\mathbf{22}$
- **Fig. 3.** Effects of 3' UTR sequences on the *gfp-cyclin B1* reporter mRNA localization to the animal polar
- 24 cytoplasm. (A) Structures of reporter mRNAs. The cyclin B1 3' UTR sequence was replaced with either
- 25 *Xenopus* vg1/vegetal localization element (VLE) or SV40 polyadenylation signal (SV40) sequences. (B)
- 26 Reporter mRNAs were detected by whole-mount *in situ* hybridization using a *gfp* probe, and the resulting
- signals were classified into 5 groups (Aggregated, Class I, II, III and Not localized: Class I, signal
- restricted to less than 1/4 of the oocyte hemisphere; class II, signal ranging from 1/4 to 1/2 of the
- 29 hemisphere; and class III, signal expanding over 1/2 of the hemisphere [6]). Localization of endogenous
- 30 *cyclin B1* mRNAs is also shown (Wild type). Arrows indicate localized signals. Scale bars, 100 µm. (C)
- 31 The numbers of oocytes showing each localized signal were counted. Data were collected from three
- 32 independent experiments. The numbers in parentheses indicate the total numbers of oocytes analyzed. (D)
- 33 Extracts from oocytes injected with reporter constructs (3' UTR, VLE, SV40) were subjected to
- immunoprecipitation with (+) or without (-) anti-Stau1 antibody. The reporter mRNAs in the
- 35 immunoprecipitates (Stau1 IP) and extracts before immunoprecipitation (initial) were detected by

- 1 RT-PCR using the *gfp* sequence.
- 2
- 3 Fig. 4. Distribution of EB1 in zebrafish stage IV oocytes revealed by *in situ* hybridization of *cyclin B1*
- 4 mRNA, immunostaining of EB1 and merged images. The animal (upper box) and vegetal (lower box)
- 5 polar regions in (A) are magnified in (B) and (C), respectively. Hoechst 33258 staining of DNA is shown
- 6 in blue. Broken lines indicate the boundary between the oocyte and follicle layer (B, C). Scale bars, 100
- **7** μm (A), 50 μm (B, C).
- 8







