Title	Otolith microstructure of Arabesque greenling <i>Pleurogrammus azonus</i> : A species with a long embryonic period
Author(s)	Marannu, Savitri; Nakaya, Mitsuhiro; Takatsu, Tetsuya; Takabatake, Shin-ichi; Joh, Mikimasa; Suzuki, Yutaro
Citation	Fisheries research, 194, 129-134 https://doi.org/10.1016/j.fishres.2017.06.002
Issue Date	2017-10
Doc URL	http://hdl.handle.net/2115/75599
Rights	© 2017. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	nakaya.pdf

[Instructions for use](#)

**Otolith microstructure of Arabesque greenling *Pleurogrammus azonus* a species
with a long embryonic period**

Savitri Marannu¹, Mitsuhiro Nakaya^{2*}, Tetsuya Takatsu¹, Shin-ichi Takabatake³,
Mikimasa Joh³, Yutaro Suzuki⁴

¹ Faculty of Fisheries Sciences, Hokkaido University, 3-1-1 Minato-cho, Hakodate,
Hokkaido, 041-8611, Japan

² Faculty of Fisheries, Hokkaido University, 3-1-1 Minato-cho, Hakodate, Hokkaido,
041-8611, Japan

³ Mariculture Fisheries Research Institute, Hokkaido Research Organization, 1-156-3
Funami-cho, Muroran, Hokkaido 051-0013, Japan

⁴ Wakkanai Fisheries Research Institute, Hokkaido Research Organization, 4-5-15
Suehiro-cho, Wakkanai, Hokkaido 097-0001, Japan

* corresponding author, e-mail: mnakaya@fish.hokudai.ac.jp

Abstract:

We investigated the relationship between morphological development in the embryonic stage and the formation of otolith microstructure in Arabesque greenling *Pleurogrammus azonus*, and then validated that the otolith increments were formed on a daily basis. Increments were more consistent to count and measure in sagittal otoliths from wild caught larvae compared to lapillar otoliths, and we suggest that sagittal otoliths as suitable for otolith microstructure analysis. This study clarified the formation time of three prominent increments on the sagittal otolith. First prominent increment (approx. 17 μm in otolith radius (OR)) was coincident with the time of eye-pigmentation in the embryonic stage. Second (approx. 36 μm OR) was considered to be the hatch increment and the third (approx. 38 μm OR) was at the time of transition from endogenous to exogenous nutrition. The third prominent increment which was the clearest radius (termed below as the check) was used as the starting point for validating the daily increment formation in sagittal otolith, and also there was no significant difference in the check radius among reared (6, 8, and 10 °C) and wild caught larvae. The relationship between number of days after hatching and the number of increments formed after check was significant and the slope of the regression line was not different from 1, validating the assumption that growth increments are formed on a daily basis in sagittae of *P. azonus*. The check was formed on the otolith at the transition from endogenous to exogenous nutrition, and the number of days required prior to its formation varied with water temperature. For the species in which the check is formed at the nutritional transition from endogenous to exogenous feeding, the relationship between number of days to check formation and water temperature is essential to estimate the hatch date of wild caught individuals.

39

40 *Keywords:* Otolith microstructure, Arabesque greenling, *Pleurogrammus azonus*, Long

41 embryonic period, Validation

42

1. INTRODUCTION

Year-class strength is affected by survival during the early life history of fishes (Houde, 1987). Otolith microstructure is useful for determining the early life histories of fishes, since many life history events, such as completion of eye pigmentation, mouth opening, hatching, transition from endogenous to exogenous nutrition, and metamorphosis, are reflected by changes in the width of increments and in the elemental composition of otoliths (Jones, 1992). The formation of otolith growth increments theoretically enables age determination with a high accuracy, often to the daily level and such studies are perhaps the most common application of microstructure examinations (Campana and Neilson, 1985). In order to estimate age of specimens from field samples, validation needs to be undertaken to determine the age when the first increment is formed and the accuracy of the increment counts. The age at the first increment deposition is best determined through laboratory rearing (Jones, 1992).

In general, fish with long embryonic periods start forming increments in the otoliths before hatching (Geffen, 1983; Tsukamoto and Shima, 1990; Narimatsu and Munehara, 1997; Morales-Nin, 2000; Joh *et al.*, 2008; Nakaya *et al.*, 2008). However there is no detailed information on the relationship between morphological development during the embryonic period and the formation of the otolith microstructure.

Arabesque greenling *Pleurogrammus azonus* is one of the most important fishery resources in northern Japan, and since 2010 its population size has decreased markedly around Hokkaido (Takashima *et al.*, 2016). In the Sea of Japan and Okhotsk Sea around Hokkaido, Arabesque greenling recruit to the fishery stock before becoming 1-year-old and a high proportion of the fishery catch consists of 0- and 1-year-old fishes (Hoshino *et al.*, 2009; Takashima *et al.*, 2013). Therefore, the early determination of

year class strength is indispensable for the fisheries management of this species. A number of early life studies enabling the determination of year class strength have already been carried out: main spawning grounds are located in coastal reefs in the Sea of Japan around Hokkaido (Irie, 1983; Fig. 1) and they spawn in water temperatures ranging from 10–13 °C (Kambara, 1957: from early October to early November in the Sea of Japan off northern Hokkaido, and from middle November to middle December in the Sea of Japan off southern Hokkaido). The eggs (2.3–2.7 mm in diameter) are demersal and adhere strongly to one another to form egg masses, and the incubation period is about two months at a water temperature of approximately 10 °C (Yusa, 1957). The larvae have been collected by plankton net in the surface layer in the Sea of Japan around Hokkaido and Tsugaru Strait from January to February (Suzuki *et al.*, unpublished data; Toyonaga *et al.*, unpublished data, Fig. 1). However, little information is available on age and growth in the early life stage of this species.

This study investigated the relationships between morphological development and the formation of otolith microstructure especially in the embryonic stage and validated that otolith increments were formed daily.

2. MATERIALS AND METHODS

2.1. Sampling of the wild larvae

Pleurogrammus azonus larvae were collected with a plankton net (mouth diameter: 80 cm; mesh aperture: 330 µm) at ca. 1 m depth from the sea surface in the sampling area off Matsumae (Fig. 1) by the T/S *Ushio-maru*, Hokkaido University during 14–15 January and 15–16 February, 2013. Theilacker (1980) and Fox (1996) reported that lengths of fish larvae shrunk more when subjected to abrasion in capture nets compared

with fixing or preservation of undamaged larvae. In 80–90% ethanol solution, there is no typical shrinkage (Theilacker, 1980; Oozeki *et al.*, 1991), and Joh *et al.* (2003) suggested that 90% ethanol is a better overall preservative, particularly for > 5mm NL fish larvae. In this study all larvae collected were preserved in 90% ethanol solution (ethanol : distilled water = 9 : 1) and we used individuals that during collection were only minimally damaged. Water temperature was measured with a CTD (SBE-19 plus, Sea-Bird Electronics Inc., Bellevue, USA) at some stations in the sampling area.

2.2. Laboratory experiments

Exp. 1: Observations of embryonic development and otolith growth

We firstly investigated in otoliths of pre-hatch embryos, the relationship between the occurrence and timing of prominent increments with the morphological development and otolith size. Broodstock of *P. azonus* were caught with trap net and angling in the sampling area off Matsumae (Fig. 1). Eggs were obtained by stripping the females. Sperm were obtained from the testis by dissecting the males. The eggs (diameter: 2.3 ± 0.11 mm (mean $\pm$ standard deviation (SD))) and sperm were mixed in a bowl, and then transferred to a hatching jar. Temperature has been considered as one of the major factors controlling development rate of many species (Heath, 1992). Rearing temperatures that were representative of the conditions during the spawning period in the Sea of Japan, coast of Hokkaido (Kambara, 1957; Irie, 1983) were set as constant at 8, 10, and 12 °C. The eggs were incubated in 30 L tanks at each water condition. The rearing water temperature and salinity were monitored (Table 1) by making daily measurements using a conductivity meter (YSI Pro30, YSI/Nanotech Ltd.). The light condition was 12 hours light (08:00–17:59) and 12 hours dark (18:00–07:59).

Embryonic development stages and growth patterns (otolith formation, eye pigmentation, and mouth opening) were identified based on Yusa (1957).

Over 20 eggs per daily sampling for each rearing condition were observed for the embryonic development stages and growth patterns (otolith formation, eye pigmentation, and mouth opening) until hatching.

Exp. 2: Observations of larval and otolith development, yolk-sac absorption, and feeding activity

Secondly, we investigated in otoliths of hatched larvae the relationship between the occurrence and timing of prominent increments with the morphological development and otolith size. Eggs were incubated at 8 °C until hatching. After hatching, larvae were reared in 10 L cylindrical tanks with set temperature conditions (6, 8, and 10 °C: representative of the conditions during larval period in the Sea of Japan, coast of Hokkaido (Suzuki *et al.*, unpublished)) in 150 L water baths. The light conditions were the same as in Exp. 1. The larvae were fed rotifers which had been enriched with Super Capsule Powder (Chlorella Industry Co. Ltd., Fukuoka, Japan), two times a day at 08:00 and 15:00 through the rearing period. Three to five larvae were sacrificed every day up to the stage when they had consumed their yolk-sac in order to measure the yolk-sac volume (see section 2.3). Furthermore, we observed the frequency of occurrence of feeding at each rearing temperature up until almost all of the larvae had started feeding.

Exp. 3: Somatic growth and otolith increment validation

Thirdly, we investigated whether otolith micro-increments are deposited daily after hatching by rearing larvae in a 200 L circular tank at 10.2 ± 1.12 °C (mean $\pm$ SD). The fish were fed rotifers cultured with Marine Glos EX (Marine Tech, Co. Ltd., Aichi,

Japan; 1–25 days after hatching; DAH), *Artemia* sp. nauplii enriched with Super Capsule Powder (20–70 DAH), and frozen mysid (66–70 DAH). The light conditions were the same as Exp. 1. Five to twenty fish were randomly collected at 10 day intervals until 70 DAH.

2.3 Measurements of otolith microstructure and yolk-sac volume estimation

Eggs and larvae preserved in 90% ethanol solution obtained from Exps. 1–3 were used. We measured the notochord length (NL) of preflexion and flexion larvae of *P. azonus* and the standard length (SL) of postflexion larvae to the nearest 0.1 mm using an electric slide caliper under a stereo microscope, with both measures being referred to as body length (BL) in this report. Otoliths were extracted and prepared for the daily increment validation process. Notochord length (NL), yolk-sac sizes ($Volume = 3/4 \pi (L/2 \times H/2 \times W/2)$, where L , H and W are yolk-sac length, height and width, respectively) were measured to the nearest 0.1 mm under a stereo-microscope. Lapillar and sagittal otoliths were removed from each larva and mounted on a glass slide with epoxy resin. All counts, measurements, and observations of otoliths were conducted with a binocular microscope connected to a video monitor. Otoliths were observed with translucent light at 1000 x magnification (100 x objective lens with 10 x eye-piece lens). Otolith radius (the longest distance from the center of the core to the edge of the sagitta) and daily increments were counted and measured using otolith microstructure analysis software (ARP Ver. 5.27, Ratoc System Engineering Co. Ltd., Tokyo, Japan). These measurements were repeated three times by different researchers and adopted when two or more counts of the number of increments agreed. The mean increment width was used as the data point for the individual.

3. RESULTS

3.1. Otolith features of wild larvae of *P. azonus*

Mean larval size of wild *P. azonus* was 8.5 ± 0.96 mm NL ($\pm$ standard deviation (SD)) collected from the sea surface with 9.4–9.7 °C and salinity 33.9–34.0 during 14–15 January, 2013. During 15–16 February, 2013, mean larval size was 11.4 ± 1.88 mm NL with 8.0–9.0 °C and 34.0–34.2. A pair of lapillar and sagittal otoliths were visible in each larva but the asteriscus could not be located. Multiple primordia (Fig. 2a,b) were observed in lapilli (23%) and sagittae (9%) from 90 pairs of otoliths. These multiple primordia tended to lead to some abnormalities in the increment structure, however their occurrence was low in sagittae. Therefore, sagittal otoliths were used for further observations of daily increment analysis.

There were three prominent increments on the sagittal otoliths. The first prominent increment was 17 ± 2.6 μ m (mean $\pm$ SD in otolith radius (OR)), second was 36 ± 1.9 μ m OR, and third was 38 ± 1.5 μ m OR (Fig. 2c). The second prominent increment was harder to differentiate in comparison with the first and third, therefore the increments which are outside of the third prominent were counted and measured. Mean increment width that formed the 2–7 rings outside of the third prominent increment was 0.8 ± 0.16 μ m ($\pm$ SD).

3.2. Laboratory experiment

Exp. 1: Embryonic development, and otolith growth

The number of days required to complete morphological developments was different among the rearing water temperatures (Table 2). Otolith formation in 8, 10, and 12 °C

was first completed from 23, 13, and 11 DAF, respectively. The first eye pigmentation day occurred on 32, 21, and 20 DAF, mouth opening was observed on 49, 36, and 35 DAF, and newly hatched larvae were first observed on 58, 46, and then 39 DAF in 8, 10, and 12 °C, respectively. Newly hatched larvae had eyes with guanine pigmentation and open mouths. Notochord lengths (mean $\pm$ SD) of newly hatched larvae were 9.6 $\pm$ 0.33 mm for larvae reared at 8 °C, 9.6 $\pm$ 0.16 mm for larvae reared at 10 °C, and 9.5 $\pm$ 0.48 mm for larvae reared at 12 °C. There was no significant difference in the NL among the three rearing temperatures (8, 10, and 12 °C: one way-ANOVA, $p = 0.42$).

The equations between DAF and sagittal otolith radius (OR) were as follows (8 °C: $OR = 0.65DAF - 1.50$ ($r^2 = 0.93$, $p < 0.001$, $N = 21$); 10 °C: $OR = 0.72DAF + 1.86$ ($r^2 = 0.95$, $p < 0.001$, $N = 16$), and 12 °C: $OR = 1.01DAF - 3.78$ ($r^2 = 0.93$, $p < 0.001$, $N = 30$); Fig. 3). Otoliths grew more rapidly at higher water temperature (ANCOVA, $p < 0.001$).

The first prominent increment at approx. 17 μ m OR corresponded to the start of the eye-pigmentation process (8–12 °C, 20–32 DAH, 17–19 μ m OR). The second prominent increment (approx. 36 μ m OR) did not correspond to mouth opening (8–12 °C, 35–49 DAH, 28–32 μ m OR) but it corresponded to hatching (8–12 °C, 39–58 DAH, 35–36 μ m OR).

Exp. 2: Larval and otolith development, yolk-sac absorption, and feeding activity

Under 6, 8, and 10 °C conditions, the larvae had already absorbed the yolk (with yolk volume < 0.01 mm³) at 9, 8, and 7 DAH, respectively. The first feeding was initiated 8, 7, and 6 DAH in 6, 8, and 10 °C, respectively. The third prominent increment was observed on 9, 8, and 6–7 DAH in 6, 8, and 10 °C, respectively (Fig. 4).

Their sagittal otolith sizes were $38 \pm 1.0 \mu\text{m}$ OR for 6°C ($N = 16$), $39 \pm 1.0 \mu\text{m}$ OR for 8°C ($N = 20$), and $38 \pm 1.0 \mu\text{m}$ OR for 10°C ($N = 21$), and hereafter called the “check”. There was no significant difference in the check radius among the three rearing temperatures (6, 8, and 10°C) and wild caught larvae ($38 \pm 1.5 \mu\text{m}$ OR; one way-ANOVA, $p = 0.31$).

Exp. 3: Somatic growth and otolith increment validation

The relationship between BL and DAH is shown in Fig. 5a. Mean BL at 0 DAH larvae was $10.3 \pm 0.74 \text{ mm}$ ($\pm \text{SD}$), and $16.5 \pm 0.35 \text{ mm}$ at 40 DAH. Mean increment width in sagittal otoliths was $0.8 \pm 0.24 \mu\text{m}$ from 6–7 DAH to 40 DAH (Fig. 5b). The minimum width was $0.3 \mu\text{m}$, and $< 0.35 \mu\text{m}$ was observed in only 0.3% of increments. After 50 DAH ($18.1 \pm 0.41 \text{ mm BL}$), the growth increased drastically and all individuals collected at 70 DAH ($28.3 \pm 1.60 \text{ mm BL}$) had already reached the juvenile stage. Their increment widths increased drastically from 50 DAH ($2.0 \pm 0.53 \mu\text{m}$; mean $\pm \text{SD}$) in flexion stage larvae to 70 DAH ($3.3 \pm 0.84 \mu\text{m}$) in juveniles. The third prominent increments (check; $37 \pm 1.9 \mu\text{m}$ OR), which was used as the starting point of otolith increments count, were observed in 6–7 DAH. The relationship between DAH and the number of increments formed after check (NIAC) is $NIAC = 0.99DAH - 6.18$ ($r^2 = 0.99$, $p < 0.001$, $N = 49$; Fig. 6). The slope of the regression line was not different from 1 (t -test, $p = 0.09$).

4. DISCUSSION

4.1. Relationship between the morphological development in embryonic-larval stage and the otolith microstructure

Species with long embryonic periods have been reported to start forming increments

in the otoliths before hatching (Geffen, 1983; Tsukamoto and Shima, 1990; Narimatsu and Munehara, 1997; Morales-Nin, 2000; Joh *et al.*, 2008; Nakaya *et al.*, 2008). Most of these workers reported that the prominent increment (check) was associated with eye-pigmentation, but has been shown to form at the time of hatching in fat greenling *Hexagrammos otakii* (Joh *et al.*, 2008), which is a closely-related species of Arabesque greenling *P. azonus*. Otolith microstructure observations in *P. azonus* showed that some increments were already formed before hatching. This study clarified the timing of the formation of three prominent increments on sagittal otoliths. The timing of formation of the first prominent increment (approx. 17 μm OR) was coincident with the time when eye-pigmentation occurred in the embryonic stage. The second prominent increment (approx. 36 μm OR) agreed with the day of hatching, and was considered to be the hatch increment, though the second prominent increment was harder to differentiate in comparison with the first and third, while the third (approx. 38 μm OR) occurred at the time of transition from endogenous to exogenous nutrition.

Microstructural features formed during the early life stages may be the result of changes in physiological metabolism, stress, and/or growth cycles that are generally associated with these transitions (Campana and Neilson, 1985). Especially the check formation at the first feeding stage might be related to a shift from endogenous nutrition to exogenous feeding and related to circadian activity patterns. It was noted by Campana (1983) that stress can cause an inhibition of calcium uptake from the water which significantly reduces the calcium deposition in the otolith, resulting in check formation. The third prominent increment (approx. 38 μm OR) in sagittal otolith was the most prominent increment and was used as the starting point (check) for validating the daily increment formation for *P. azonus*. This check radius coincided with the

transition from endogenous to exogenous nutrition and relates to a life history event as found in other species (Hayashi *et al.*, 1989; Maillet and Checkley, 1990; Joh *et al.*, 2005).

4.2. Validation of daily increment formation

To estimate age in wild caught larvae and juveniles, two pieces of information must be known: the age at first increment formation (check) and the accuracy of the increment deposition rate (Jones, 1992). In this study for *P. azonus*, the third prominent increment was used as the starting point for validating the daily increment formation in sagittal otoliths. There was no significant difference in the check radius among reared (6, 8, and 10 °C), and wild caught larvae, though the timing of the check formation was different between rearing temperature conditions. Temperature is a major determinant of somatic growth and metabolic rate where higher temperature induces faster somatic growth and metabolic rate up to a certain limit (Hare and Cowen, 1995). In the higher temperatures, the check was deposited earlier which seems to be due to the elevated metabolic rate. Life history events such as hatching and first feeding are also regulated by temperature, since it plays an important role in embryonic and larval development stage. The mean increment width of larval *P. azonus* was $0.8 \pm 0.24 \mu\text{m}$ ($\pm$ SD) and also 99.7% of increment measurements were above the resolution limits of light microscopes ($0.3 \mu\text{m}$, Fox *et al.*, 2003; $0.35 \mu\text{m}$, Campana *et al.*, 1987). Therefore, it is possible to count the number of increments.

In this study, we investigated the relationship between the morphological development and otolith formation in the range of water temperatures experienced by eggs in the field. Life history events (i.e. completion of eye pigmentation, mouth

283 opening, hatching, transition from endogenous to exogenous nutrition) corresponded to
284 prominent increments, and the otolith size did not vary even if water temperature
285 changed. The present study clarified the duration under 3 different temperature
286 conditions and established the essential basis of growth analysis and hatch date
287 estimation of wild caught *P. azonus* larvae and juveniles. For species with a check
288 formation at a transition from endogenous to exogenous nutrition, understanding of the
289 duration from hatching to the check formation and the effect of the water temperature
290 on its formation is requisite data for the hatch date estimation of wild caught individuals.

Acknowledgements

The authors thank Y. Sakurai and T. Nakatani of the Faculty of Fisheries Sciences, Hokkaido University for valuable comments on the manuscript. We are also grateful to T. Terada, M. Sato, and H. Horikawa of the Matsumae-Sakura Fisheries Cooperative Association who provided the broodstock, and crew of T/S *Ushio-maru* of Faculty of Fisheries, Hokkaido University, who helped with the outdoor sampling. Finally, we thank the editor and two anonymous reviewers for their critical suggestions.

References

- Campana SE. (1983) Calcium deposition and otolith check formation during periods of stress in coho salmon, *Oncorhynchus kisutch*. *Comp Biochem Physiol* **75**: 215–220.
- Campana SE, Neilson JD. (1985) Microstructure of fish otoliths. *Can J Fish Aquat Sci* **42**: 1014–1032.
- Campana SE, Gagné JA, Munro J. (1987) Otolith microstructure of larval herring (*Clupea harengus*): image or reality? *Can J Zool* **67**: 1401–1410.
- Fox CJ. (1996) Length changes in herring (*Clupea harengus*) larvae: effects of capture and storage in formaldehyde and alcohol. *J Plankton Res* **18**: 483–493.
- Fox CJ, Folkvord A, Geffen AJ. (2003) Otolith micro-increment formation in herring *Clupea harengus* larvae in relation to growth rate. *Mar Ecol Prog Ser* **264**: 83–94.
- Geffen AJ. (1983) The deposition of otolith rings in Atlantic salmon, *Salmo salar* L., embryos. *J Fish Biol* **23**: 467–474.
- Hare JA, Cowen RK. (1995) Effect of age, growth rate, and ontogeny on the otolith size-fish size relationship in bluefish, *Pomatomus saltatrix*, and the implications for back calculation of size in fish early life history stages. *Can J Fish Aquat Sci* **52**: 1909–1922.
- Hayashi A, Yamashita Y, Kawaguchi K. (1989) Rearing method and daily otolith ring of Japanese sardine larvae. *Nippon Suisan Gakkaishi* **55**: 997–1000.

- 319 Heath MR. (1992) Field investigations of the early life stages of marine fish. Adv
320 Mar Biol **28**: 1–174.
- 321 Hoshino N, Takashima T, Watanobe M, Fujioka T. (2009) Age structures and catch
322 fluctuations of Arabesque greenling (*Pleurogrammus azonus*) in coastal area of
323 southern Hokkaido, Japan. Sci Rep Hokkaido Fish Exp Stn **76**: 1–11.
- 324 Houde ED. (1987) Fish early life dynamics and recruitment variability. Am Fish Soc
325 Symp **2**: 17–29.
- 326 Irie T. (1983) Spawning grounds for greenling *Pleurogrammus azonus* in Hidaka, off
327 the Pacific coast of Hokkaido. Bull Hokkaido Reg Fish Res Lab **48**: 1–9 (in
328 Japanese with English abstract).
- 329 Joh M, Takatsu T, Nakaya M, Imura K, Higashitani T. (2003) Body length shrinkage
330 of marbled sole *Pseudopleuronectes yokohamae* larvae preserved in formalin and
331 ethanol solutions. Aquaculture Sci **51**: 227–228.
- 332 Joh M, Takatsu T, Nakaya M, Higashitani T, Takahashi T. (2005) Otolith
333 microstructure and daily increment validation of marbled sole (*Pseudopleuronectes*
334 *yokohamae*). Mar Biol **147**: 59–69.
- 335 Joh M, Joh T, Matsu-ura T, Takatsu T. (2008) Validation of otolith increment
336 formation and the growth rate of fat greenling larvae. Aquaculture Sci **56**:
337 157–166.
- 338 Jones CM. (1992) Development and application of the otolith increment technique. In:
339 Stevenson DK, Campana SE (Eds). Otolith microstructure examination and analysis.
340 Can Spec Publ Fish Aquat Sci **117**: 1–11.

- 341 Kambara H. (1957) Studies on “Hokke” *Pleurogrammus azonus* Jordan et Metz (II)
 342 —Ecological study of spawning—. Hokusuisi-Geppo **14**: 3–13 (in Japanese).
- 343 Maillet GL, Checkley DM. (1990) Effects of starvation on the frequency of formation
 344 and width of growth increments in sagittae of laboratory-reared Atlantic menhaden
 345 *Brevoortia tyrannus* larvae. Fish Bull **88**: 155–165.
- 346 Morales-Nin B. (2000) Review of the growth regulation processes of otolith daily
 347 increment formation. Fish Res **46**: 53–67.
- 348 Nakaya M, Morioka T, Fukunaga K, Sekiya S, Jinbo T, Nagashima H, Ueno Y. (2008)
 349 Validation of otolith daily increments for larval and juvenile Japanese halfbeak
 350 *Hyporhamphus sajori*. Fish Res **93**: 186–189.
- 351 Narimatsu Y, Munehara H. (1997) Age determination and growth from otolith daily
 352 growth increments of *Hypoptychus dybowskii* (Gasterosteiformes). Fish Sci **63**:
 353 503–508.
- 354 Oozeki Y, Watanabe Y, Kuji Y, Takahashi S. (1991) Effects of various preservatives
 355 on the body length of saury larvae. Bull Tohoku Natl Fish Res Inst **53**: 15–21 (in
 356 Japanese with English abstract).
- 357 Takashima T, Hoshino N, Itaya K, Maeda K, Miyashita K. (2013) Age variation using
 358 sectioned otoliths and age-size relationship for Northern Hokkaido stock of the
 359 Arabesque greenling *Pleurogrammus azonus*. Nippon Suisan Gakkaishi **79**:
 360 383–393.
- 361 Takashima T, Okada N, Asami H, Hoshino N, Shida O, Miyashita K. (2016)
 362 Maturation process and reproductive biology of female Arabesque greenling

363 *Pleurogrammus azonus* in the Sea of Japan, off the west coast of Hokkaido. Fish
 364 Sci **82**: 225–240.

 365 Theilacker GH. (1980) Changes in body measurements of larval northern anchovy,
 366 *Engraulis mordax*, and other fishes due to handling and preservation. Fish Bull US
 367 **78**: 685–692.

 368 Tsukamoto K, Shima Y. (1990) Otolith daily increment in sandfish. Nippon Suisan
 369 Gakkaishi **56**: 1083–1087.

 370 Yusa T. (1957) Studies on “hokke” *Pleurogrammus azonus* Jordan et Mets. (III)
 371 —Embryonic and larval development—. J Hokkaido Fish Sci Inst **14**: 3–18.
 372

Fig. 1. Major spawning grounds (Irie, 1983) and horizontal distribution of larvae and juveniles of *P. azonus* in the Sea of Japan and Tsugaru Strait (Suzuki *et al.*, unpublished data; Toyonaga *et al.*, unpublished data). Location of the sampling area of *P. azonus* larvae by T/S *Ushio-maru* off Matsumae on January 14–15 and February 15–16 in 2013.

Fig. 2. Otoliths of wild caught *P. azonus* larvae that were collected off Matsumae from January to February 2013 (a, b: 9.7 mm in notochord length (NL), c: 12.9 mm NL). Lapillus with multiple primordia (pointed by arrows; a) and sagitta (b). Three prominent increments in the sagittal otolith of a wild caught *P. azonus* larva (c). Otolith radii of first, second and third prominent increments (1st, 2nd, and 3rd) were $17 \pm 2.6 \mu\text{m OR}$, $36 \pm 1.9 \mu\text{m OR}$, and $38 \pm 1.5 \mu\text{m OR}$ (mean $\pm$ standard deviation), respectively.

Fig. 3. Relationship between days after fertilization (DAF) and otolith radius (OR) of sagittae. The regression lines for 8 °C: $OR = 0.65DAF - 1.50$ ($r^2 = 0.93$, $p < 0.001$, $N = 21$), 10 °C: $OR = 0.72DAF + 1.86$ ($r^2 = 0.95$, $p < 0.001$, $N = 16$), and 12 °C: $OR = 1.01DAF - 3.78$ ($r^2 = 0.93$, $p < 0.001$, $N = 30$).

Fig. 4. Relationship between water temperature and day of check observed (DCO) after hatching. Numerals show the sample sizes.

Fig. 5. Relationship between days after hatching and notochord length or standard length of *P. azonus* larvae (a), and changes in daily increment widths on sagittae (b).

Fig. 6. Relationship between the number of days after hatching (DAH) and the number of otolith increments after check (NIAC).

Fig. 1. Marannu *et al.*

Fig. 2. Marannu *et al.*

Fig. 3. Marannu *et al.*

Fig. 4. Marannu *et al.*

Fig. 5 Marannu *et al.*

Fig. 6. Marannu *et al.*

Table 1. Temperature and salinity in each rearing tank

Exp.	Rearing condition	Temperature (°C)	Salinity
1	8°C	7.9 ± 0.52	33.3 ± 0.92
	10°C	10.2 ± 0.31	33.1 ± 0.98
	12°C	12.0 ± 0.21	34.4 ± 0.79
2	6°C	6.0 ± 0.17	33.5 ± 1.32
	8°C	7.9 ± 0.24	33.8 ± 0.35
	10°C	9.9 ± 0.10	34.3 ± 1.06

*Values are shown as mean ± SD.

Table 2. The time required for morphological development in the *P. azonus* embryonic stage in days after fertilization (DAF)

Temperature condition	Otolith formation	Eye pigmentation (range)	Mouth opening	Hatching (range)
8°C	23	32–36	49	58–62
10°C	13	21–30	36	46–51
12°C	11	20–27	35	39–48