



Title	Knockdown of DEAD-box helicase 4 (DDX4) decreases the number of germ cells in male and female chicken embryonic gonads
Author(s)	Aduma, Nana; Izumi, Hiroe; Mizushima, Shusei; Kuroiwa, Asato
Citation	Reproduction, Fertility and Development, 31(5), 847-854 https://doi.org/10.1071/RD18266
Issue Date	2019-04
Doc URL	http://hdl.handle.net/2115/75725
Type	article (author version)
File Information	RFD 31 (5) 847.pdf



[Instructions for use](#)

1 **Knockdown of *DDX4* decreases the number of germ cells in male and**
2 **female chicken embryonic gonads**

3

4 Nana Aduma¹, Hiroe Izumi², Shusei Mizushima^{1,2}, Asato Kuroiwa^{1,2}

5

6 ¹ Biosystems Science Course, Graduate School of Life Science, Hokkaido University, Kita 10
7 Nishi 8, Kita-ku, Sapporo, Hokkaido 060-0810, Japan

8

9 ²Division of Reproductive and Developmental Biology, Department of Biological Sciences,
10 Faculty of Science, Hokkaido University, Kita 10 Nishi 8, Kita-ku, Sapporo, Hokkaido
11 060-0810, Japan

12

13

14 Correspondence should be addressed to A. Kuroiwa. E-mail: asatok@sci.hokudai.ac.jp, Kita 10,
15 Nishi 8, Kita-ku, Sapporo, Hokkaido, 060-0810 Japan.

16

17 Short title: Effects of *DDX4* KD on chicken embryonic gonads

18

19 Key words: primordial germ cells, sex differentiation, ovary, testis

20 **Abstract**

21 DEAD-box helicase 4 (*DDX4*, also known as *vasa*) is essential for the proper formation and
22 maintenance of germ cells. Although *DDX4* is conserved in a variety of vertebrates and
23 invertebrates, its roles differ between species. This study investigated the function of *DDX4* in
24 chicken embryos by knocking down its expression using retroviral vectors that encoded
25 *DDX4*-targeting microRNAs. *DDX4* was effectively depleted *in vitro* and *in vivo* via this
26 approach. Male and female gonads of *DDX4*-knockdown embryos contained a decreased
27 number of primordial germ cells, indicating that *DDX4* is essential to maintain a normal level of
28 these cells in chicken embryos of both sexes. *DMRT1* and *SOX9*, which are involved in testis
29 determination and differentiation, were expressed as normal in male gonads of
30 *DDX4*-knockdown embryos. By contrast, expression of *CYP19A1*, which encodes aromatase
31 and is essential for ovary development, was significantly decreased in female gonads of
32 *DDX4*-knockdown embryos. Expression of *FOXL2*, which plays an important role in ovary
33 differentiation, was also slightly reduced, but this was not statistically significant. *FOXL2* was
34 previously hypothesized to regulate aromatase expression based on several pieces of evidence.
35 These results indicate that aromatase expression is also regulated by several additional
36 pathways.

37 **Introduction**

38 DEAD-box helicase 4 (DDX4, also called vasa) is a member of the DEAD-box protein family,
39 which is important for the proper formation of germ cells. The roles of DDX4 are reported to
40 differ between species. Expression of DDX4 has been depleted in *Drosophila melanogaster*
41 (Styhler et al, 1998), *Caenorhabditis elegans* (Roussell and Bennett, 1993; Gruidl et al, 1996;
42 Kawasaki et al, 1998; Kuznicki et al, 2000; Spike et al, 2008), mice (Tanaka et al, 2000), and
43 zebrafish (Braat et al, 2001) via gene knockout (KO), RNA interference (RNAi), and
44 morpholino treatment. DDX4 is essential for oogenesis in most species and is necessary for
45 male germ cells to progress through spermatogenesis in several species, including mice (Tanaka
46 et al, 2000; Kuramochi-Miyagawa et al, 2010; Ewen-Campen et al, 2013). These diverse
47 functions of DDX4 evolved for reproductive adaptation.

48 The chicken *DDX4* homolog (also called chicken vasa homologue, *CVH*) was first cloned
49 by Tsunekawa et al. (2000). This gene encodes 663 amino acids and is located on the Z
50 chromosome (male: ZZ, female: ZW). DDX4 protein localizes to the cytoplasm of germ cells,
51 including presumptive primordial germ cells (PGCs) in uterine-stage embryos and spermatids
52 and oocytes in adult gonads. These findings indicate that the chicken germ lineage is maternally
53 predetermined (Tsunekawa et al 2000). Taylor et al. (2017) efficiently knocked out the *DDX4*

54 locus via homologous recombination mediated by transcription activator-like effector nucleases.
55 Targeted male PGCs were injected into surrogate host chicken embryos. Seventeen targeted G1
56 offspring were generated following mating of a host male cockerel with wild-type (WT) hens.
57 G1 females (hemizygous KO chickens) were sterile and did not exhibit detectable follicles
58 post-hatching. The germ cell lineage was initially formed in early embryos, but PGCs were
59 subsequently lost during meiosis. These findings indicate that *DDX4* plays an important role in
60 the female reproductive system. By contrast, homozygous (null) male KO chickens could not be
61 generated because it was impossible to cross G1 female and male KO chickens (Taylor et al,
62 2017). Therefore, the function of *DDX4* in male PGCs remains unclear in chickens.

63 Herein, we show that *DDX4* is essential for PGCs in chicken embryos of both sexes. We
64 knocked down *DDX4* using retroviral vectors that encoded *DDX4*-targeting microRNAs
65 (miRNAs). There was a decreased number of PGCs in male and female gonads of
66 *DDX4*-knockdown (KD) embryos. *DMRT1* and *SOX9*, which are involved in testis
67 determination and differentiation, were expressed as normal in male gonads of *DDX4*-KD
68 embryos. By contrast, expression of *CYP19A1*, which encodes aromatase, was significantly
69 decreased in female gonads of *DDX4*-KD embryos.

70

71 **Materials and Methods**

72 *Animals and ethics statement*

73 Fertilized chicken eggs (*Gallus gallus domesticus*) were purchased from Takeuchi Hatchery
74 (Nara, Japan). The Hy-Line Maria chicken strain was used in this study. Fertilized eggs were
75 incubated at 37.8°C. The sex of each embryo was determined by PCR genotyping using
76 genomic DNA as the template (Fridolfsson & Ellegren 1999).

77 All animal experiments described in this study were approved by the Institutional Animal
78 Care and Use Committee of National University Corporation Hokkaido University and were
79 performed in accordance with the Guidelines for the Care and Use of Laboratory Animals
80 issued by Hokkaido University. This study did not involve any human participants or
81 specimens.

82

83 *Construction of RNAi vectors*

84 Two miRNAs targeting different regions of chicken *DDX4* mRNA and two non-targeting
85 (scrambled) controls were designed using BLOCK-iT RNAi Designer (Thermo Fisher Scientific,
86 MA) and siRNA Wizard v3.1 (Thermo Fisher Scientific) (Table 1). The miRNA constructs were
87 cloned as described previously (Tanaka et al., 2017). pRFPRNAi.C and RCASRNAi were

88 provided by ARK-Genomics, The Roslin Institute (Das et al, 2006). The vectors encoding
89 *DDX4*-targeting and scrambled miRNAs were named RCASA.miRNA.DDX4.eGFP and
90 RCASA.miRNA.Sc.eGFP, respectively.

91 To test the KD efficiency using DF-1 cells, RCASA.miRNA.DDX4 and
92 RCASA.miRNA.Sc, which lacked the eGFP sequence, were constructed. A vector expressing
93 *DDX4* fused to GFP was also generated for *in vitro* experiments. A forward primer containing a
94 *ClaI* site and a reverse primer containing a *SpeI* site were designed to amplify the coding DNA
95 sequence of *DDX4* (Table 1). Following digestion with *ClaI* and *SpeI*, the PCR product was
96 cloned into the eGFP-harboring RCAS.B vector that had been digested with the same restriction
97 enzymes. This construct was named RCASB.DDX4_eGFP. RCAS.B proviral DNA, RCAS.A,
98 and RCAS.B were kindly provided by Dr. Hughes, National Cancer Institute (Hughes et al,
99 1987).

100

101 ***Preparation and injection of viruses***

102 Endotoxin-free proviral DNA was prepared using a PureYield™ Plasmid Miniprep Kit
103 (Promega, WI). DF-1 cells were transfected with this DNA using Lipofectamine 2000 (Thermo
104 Fisher Scientific) according to the manufacturer's protocol, transferred to 10-mm dishes

105 (surface area of 78.5 cm²), and cultured to sub-confluency in Dulbecco's Modified Eagle's
106 Medium supplemented with 10% (v/v) fetal bovine serum. Active viruses in the pooled medium
107 were concentrated. Briefly, 100 ml of medium was centrifuged overnight at 6,000 × g, and then
108 the supernatant was carefully decanted to leave a small pellet in 200 μl of medium. The
109 resuspended viral solution was aliquoted and stored at -80°C until use. The viral titer was
110 determined as described previously (Nakata et al, 2013).

111 Embryos were injected as described previously (Nakata et al, 2013) with minor
112 modifications. Approximately 3 μl of concentrated viruses containing 0.025% Fast Green
113 tracking dye was injected into the subgerminal cavity of blastoderms at day 0 (Hamburger and
114 Hamilton stage X; Hamburger & Hamilton 1951). Viruses carrying RCASA.miRNA.DDX4
115 were injected into 189 eggs to KD *DDX4*. The eggs were sealed and incubated until embryonic
116 day (E)8.5. The number of embryos developed at ~E8.5 are 18 (ZW) and 15 (ZZ).

117

118 *Reverse-transcription polymerase chain reaction (RT-PCR) and quantitative real-time*
119 *polymerase chain reaction (qRT-PCR)*

120 For RT-PCR, total RNA was extracted from DF-1 cells infected with RCASB.DDX4_eGFP or
121 with RCASA.miRNA.DDX4 followed by RCASB.DDX4_eGFP. For qRT-PCR, total RNA was

122 extracted from gonads of male and female *DDX4*-KD embryos infected with
123 RCASA.miRNA.*DDX4*.eGFP and from gonads of male and female WT embryos at E8.5. Total
124 RNA was extracted using an RNeasy Kit (Invitrogen) according to the manufacturer's
125 instructions. RNA was treated with DNase I and reverse-transcribed using SuperScript III
126 reverse transcriptase (Invitrogen) and an oligo(dT) primer. qRT-PCR was performed using
127 Power SYBR[®] Green PCR Master Mix (Thermo Fisher Scientific) and an ABI 7300 Fast
128 Real-Time PCR System (Thermo Fisher Scientific). To ensure the amplification efficiency was
129 maximal in all reactions, the amplicon size was restricted to 90–101 bp. Reactions were
130 performed in triplicate using 96-well plates in a reaction volume of 10 μ l. Data were analyzed
131 using the $\Delta\Delta$ Ct method, and the mRNA level of each target gene was normalized against that of
132 beta-actin (*ACTB*). The primers used to quantify *DDX4* expression are listed in Table 1, while
133 those used to measure the expression levels of other genes were previously described (Nakata et
134 al, 2013).

135

136 ***Immunohistochemistry***

137 Urogenital tissues of chicken embryos were fixed in formalin for 1 h at room temperature.
138 Paraffin sections (10 μ m thick) were cut along the dorsal-ventral axis and thaw-mounted onto

139 MAS-coated glass slides. The experimental conditions of immunohistochemistry were
140 described previously (Nakata et al, 2013). The anti-DDX4 antibody was kindly provided by Dr.
141 Hattori, Kyushu University (Aramaki et al, 2009).

142

143 *Alkaline phosphatase (AP) staining*

144 Urogenital tissues of chicken embryos were fixed in 4% paraformaldehyde prepared in
145 phosphate-buffered saline (PBS). Serial paraffin sections (10 µm thick) were cut along the
146 anterior-posterior axis and mounted onto MAS-coated glass slides. De-waxed slides were
147 treated with PBS containing 0.5% Triton X-100 for 20 min and incubated with the NBT/BCIP
148 substrate (Promega).

149

150 *In situ hybridization*

151 A fragment of DND microRNA-mediated repression inhibitor 1 (*DNDI*, also known as *CDH*)
152 was amplified by RT-PCR using cDNA obtained from gonads at E8.5 as the template. The
153 primers are listed in Table 1. RNA extraction and cDNA synthesis were performed as described
154 earlier. The PCR product of 568 bp was subcloned using a pGEM T-Easy vector system
155 (Promega). cDNA clones were labeled using Digoxigenin RNA Labeling Mix (Roche, Basel,

156 Switzerland) and T7 or SP6 RNA polymerase (MAXIscript™; Thermo Fisher Scientific).
157 Hybridization to serial frozen sections was performed as described previously (Nakata et al,
158 2013). The images were captured using a cooled CCD camera (DS-Ri1, Nikon corporation,
159 Tokyo, Japan) mounted on a Nikon ECLIPSE E800 microscope, and were analyzed with the
160 NIS ELEMENTS application program of Nikon corporation.

161

162 *Statistical analysis*

163 The data are presented as mean \pm standard deviation. Statistical comparison was made by
164 Student *t* test. $P < 0.005$ was considered statistically significant.

165

166 **Results**

167 *DDX4-targeting miRNAs efficiently deplete DDX4*

168 We first examined the efficiency of *DDX4* KD using *DDX4*-targeting miRNAs in DF-1 cells.
169 These cells were infected with RCASB.DDX4_eGFP (Fig. 1a, b), an RCAS vector expressing
170 *DDX4* fused to eGFP, or with RCASA.miRNA.Sc followed by RCASB.DDX4_eGFP (Fig. 1c,
171 d) as controls. eGFP was observed in both cases. However, eGFP was not detected in DF-1 cells
172 infected with RCASA.miRNA.DDX4, which encoded two *DDX4*-targeting miRNAs, prior to

173 RCASB.DDX4_eGFP (Fig. 1e, f). In addition, *DDX4* was detected by RT-PCR in DF-1 cells
174 infected with RCASB.DDX4_eGFP, but not in those infected with RCASA.miRNA.DDX4
175 prior to RCASB.DDX4_eGFP (Fig. 1g).

176 We performed immunohistochemical staining of *DDX4* in gonads of *DDX4*-KD embryos
177 at E8.5. *DDX4* staining was reduced in female (Fig. 2a–d) and male (Fig. 2e–h) gonads of
178 *DDX4*-KD embryos. *DDX4* expression in gonads of WT and *DDX4*-KD embryos at E8.5 was
179 quantified by qRT-PCR. This demonstrated that *DDX4* expression was significantly reduced in
180 *DDX4*-KD embryos of both sexes (Fig. 2i). These results demonstrate that *DDX4*-targeting
181 miRNAs efficiently deplete *DDX4*.

182

183 ***The number of PGCs is decreased in male and female gonads of DDX4-KD embryos***

184 We performed AP staining and *DND1 in situ* hybridization to label PGCs in gonads of
185 *DDX4*-KD embryos. PGCs exhibit AP staining in many species. *DND1* co-localizes with *DDX4*
186 in germ cell lines generated from chicken embryonic gonads (Kito et al 2010) and is thus a
187 marker of PGCs.

188 PGCs were detected by AP staining in female and male gonads of WT embryos at E8.5
189 (Fig. 3a, b). However, AP staining was not clearly observed in *DDX4*-KD embryos of either sex

190 (Fig. 3c, d). The *DND1 in situ* hybridization experiments yielded similar results (Fig. 3e–h).

191 The total number of PGCs was counted in serial sections subjected to AP staining (Fig.
192 3i) and *DND1 in situ* hybridization (Fig. 3j). Based on AP staining and *DND1 in situ*
193 hybridization, the number of PGCs was decreased by 74% and 79% in *DDX4*-KD female
194 embryos, respectively, and by 61.4% and 62.2% in *DDX4*-KD male embryos, respectively. The
195 detection sensitivity of each method is different: the sensitivity of AP staining is relatively
196 higher than that of *DND1 in situ* hybridization. Although total number of PGCs detected by
197 AP-staining is larger than *DND1 in situ* hybridization, the reduction rate of number of PGCs is
198 almost same between both methods.

199

200 ***KD of DDX4 decreases CYP19A1 expression***

201 We quantified expression of *DMRT1*, *SOX9*, *FOXL2*, and *CYP19A1*, which are involved in
202 gonadal differentiation, in gonads of WT and *DDX4*-KD embryos at E8.5 (Fig. 4). In both WT
203 and *DDX4*-KD embryos, expression of *DMRT1* and *SOX9* was high in male gonads, but low in
204 female gonads (Fig. 4), consistent with previous reports (Smith et al, 1999a; 1999b). Expression
205 of *DMRT1* and *SOX9* did not significantly differ between WT and *DDX4*-KD embryos.

206 Expression of *FOXL2* and *CYP19A1* was reported to be high in female embryonic gonads,

207 but extremely low in male embryonic gonads (Govoroun et al, 2004). We confirmed this finding
208 in WT embryos. However, *CYP19A1* expression in female gonads was significantly lower in
209 *DDX4*-KD embryos than in WT embryos (Fig. 4). Expression of *FOXL2* was slightly lower in
210 female gonads of *DDX4*-KD embryos than in those of WT embryos, but this difference was not
211 statistically significant.

212

213 **Discussion**

214 The *DDX4* gene is conserved in a variety of vertebrates and invertebrates. Although *DDX4* is
215 essential for the development and maintenance of germ cells in a wide range of species, its roles
216 differ between species. A null mutation of *DDX4* in *D. melanogaster*, in which this gene was
217 initially identified, causes female sterility due to severe defects in oogenesis, while males
218 remain fertile (Styhler et al, 1998). However, male *DDX4*-KO mice exhibit a reproductive
219 deficiency; male homozygotes do not produce sperm in the testes, and the proliferation of PGCs
220 is hampered (Tanaka et al, 2000). By contrast, disruption of *DDX4* does not affect the
221 development of female germ cells in mice, despite the fact that this gene is expressed in both
222 male and female germ cells of mice (Tanaka et al, 2000). The current study demonstrated that
223 *DDX4*-targeting miRNAs effectively suppressed *DDX4* expression *in vitro* and *in vivo* (Fig. 1,

224 2). The number of proliferative cells detected by EdU is significantly reduced in female gonads
225 of *DDX4*-KO chicken embryos at E10.5, indicating that germ cells form at an early
226 developmental stage and are subsequently lost (Taylor et al, 2017). Our results confirmed this
227 previous finding and demonstrated that the number of PGCs was also reduced in male gonads of
228 *DDX4*-KD embryos (Fig. 3). Therefore, *DDX4* is also involved in the proliferation of PGCs in
229 males, suggesting it is important for both male and female fertility in chickens.

230 We investigated the expression levels of genes involved in gonadal differentiation and
231 development in *DDX4*-KD embryos (Fig. 4). Expression of *DMRT1* and *SOX9*, which are
232 markers of testis determination and differentiation, were quantified. *DMRT1* is located on the Z
233 chromosome in birds, is a major avian male-determining factor, and begins to be expressed at
234 ~E3.5. Expression of *DMRT1* is detected in the medulla of gonads and is higher in males than in
235 females. Upon high *DMRT1* expression, *SOX9* is continuously expressed for testis development
236 in male chicken embryos after E6.5. Expression of *DMRT1* and *SOX9* was higher in male
237 gonads than in female gonads of both *DDX4*-KD and WT embryos (Fig. 4). This suggests that
238 testis differentiation and development occurred as normal in male gonads with a decreased
239 number of PGCs.

240 By contrast, expression of *CYP19A1* was significantly lower in female gonads of

241 *DDX4*-KD embryos than in those of WT embryos (Fig. 4). Gonadal sex differentiation in female
242 birds is sensitive to the sex steroid hormone estradiol. This hormone is only detected in female
243 embryonic gonads and is required and sufficient for ovarian development (Elbrecht & Smith
244 1992). Aromatase converts androgens into estradiol. Consequently, *CYP19A1* is essential for the
245 female reproductive system. Indeed, treatment with aromatase inhibitors, such as fadrozole, can
246 effectively inhibit to feminization in ZW embryos (Elbrecht & Smith 1992). Aromatase
247 localizes to the cytoplasm of somatic cells in the medulla of female embryonic gonads from
248 E6.5 onwards and its expression increases during ovarian development (Govoroun et al, 2004;
249 Smith et al. 2005). *DDX4* is thought to increase the translation of several proteins required for
250 meiotic progression and the assembly of cytoplasmic granules in germ cells. The
251 ATP-dependent catalytic activity of RNA helicases regulates the translation of multiple mRNAs
252 (Carrera et al, 2000). Considering its expression pattern and function, *DDX4* is not thought to
253 directly regulate expression of *CYP19A1*. Repression of the *DDX4* expression and/or a
254 decreased number of PGCs might indirectly affect aromatase expression.

255 Expression of *FOXL2*, another marker of ovary differentiation, was slightly lower in
256 female gonads of *DDX4*-KD embryos than in those of WT embryos; however, this difference
257 was not statistically significant (Fig. 4). *FOXL2* is an essential factor that is widely conserved in

258 vertebrates, including chickens (Loffler et al, 2003; Wang et al, 2007; Pisarska et al, 2011). The
259 expression patterns of *FOXL2* and *CYP19A1* highly correlate with each other in the developing
260 ovary after 4.7–12.7 days of incubation (Govoroun et al. 2004). The proteins encoded by these
261 genes co-localize at the nuclei of medullar cord cells in female embryonic gonads. *FOXL2* is
262 expressed just prior to *CYP19A1*, suggesting it directly or indirectly regulates transcription of
263 *CYP19A1*, at least in embryos. This hypothesis is supported by the finding that *FOXL2*
264 expression is induced in aromatase-overexpressing male embryos (Lambeth et al, 2013).
265 *FOXL2*-binding sites (5'-CACAACA-3') are located in the promoter region of *CYP19A1* (-668
266 and -3453 bp in the antisense strand and -3642 bp in the sense strand; Govoroun et al, 2004).
267 However, *FOXL2* fails to upregulate *CYP19A1* expression in luciferase assays (Wang and Gong,
268 2017). The expression patterns of *FOXL2* and aromatase differ in the ovaries of adult chickens,
269 with the former protein localizing to the granulosa cell layer and the latter protein localizing to
270 the theca cell layers, indicating that the regulatory mechanism of aromatase differs between
271 embryos and adults (Wang and Gong, 2017). Our data suggest that several pathways regulate
272 expression of *CYP19A1*, in addition to that involving *FOXL2*. Furthermore, treatment with an
273 aromatase inhibitor reduces *FOXL2* expression *in vivo*, suggesting that a regulatory feedback
274 loop exists between *FOXL2* and *CYP19A1* (Hudson et al, 2005). Therefore, the slight reduction

275 in *FOXL2* expression observed in female gonads of *DDX4*-KD embryos might be due to the
276 substantial decrease in *CYP19A1* expression.

277 In summary, this study demonstrates that *DDX4* is important for the female and male
278 reproductive systems of chickens. Furthermore, our data indicate repression of the *DDX4*
279 expression and/or a decreased number of PGCs affect the regulation of *CYP19A1* expression.

280

281 **Declaration of interest**

282 The authors declare that no competing interests exist.

283

284 **Funding**

285 This research was supported by the F3 Project Support Office for Female Researchers at
286 Hokkaido University.

287

288 **Acknowledgments**

289 The authors thank T. Nakata for technical support, C. Nishida for helpful suggestions on cell
290 culture experiments, and M.-A. Hattori for providing the anti-*DDX4* antibody.

291

292 **Authors' contributions**

293 NA performed all the experiments and analyzed all the data. HI constructed the retroviral
294 vectors. AK conceived and designed the study and SM and AK wrote the manuscript. All
295 authors read and approved the final manuscript.

296 **References**

- 297 Aramaki S, Kubota K, Soh T, Yamauchi N & Hattori M-A 2009 Chicken dead end homologue
298 protein is a nucleoprotein of germ cells including primordial germ cells. *Journal of*
299 *Reproduction and Development* **55** 214-218.
- 300 Braat AK, van de Water S, Korving J & Zivkovic DA 2001 zebrafish vasa morphant abolishes
301 vasa protein but does not affect the establishment of the germline. *Genesis* **30** 183-185.
- 302 Carrera P, Johnstone O, Nakamura A, Casanova J, Jäckle H & Lasko P 2000 VASA mediates
303 translation through interaction with a Drosophila yIF2 homolog. *Molecular Cell* **5** 181-187.
- 304 Das RM, Van Hateren NJ, Howell GR, Farrell ER, Bangs FK, Porteous VC, Manning EM,
305 McGrew MJ, Ohyama K, Sacco MA, *et al* 2006 A robust system for RNA interference in
306 the chicken using a modified microRNA operon. *Dev Biol* **294** 554-563.
- 307 Elbrecht A & Smith RG 1992 Aromatase enzyme activity and sex determination in chickens.
308 *Science* **255** 467-470.
- 309 Ewen-Campen B, Donoughe S, Clarke DN & Extavour CG 2013 Germ cell specification
310 requires zygotic mechanisms rather than germ plasm in a basally branching insect. *Current*
311 *Biology* **23** 835-842.

312 Fridolfsson AK & Ellegren H 1999 A simple and universal method for molecular sexing of
313 non-ratite birds. *J Avian Biol* **30** 116-121.

314 Govoroun MS, Pannetier M, Pailhoux E, Cocquet J, Brillard JP, Couty I, Batellier F & Cotinot
315 C 2004 Isolation of chicken homolog of the FOXL2 gene and comparison of its expression
316 patterns with those of aromatase during ovarian development. *Dev Dyn* **231** 859-870.

317 Gruidl ME, Smith PA, Kuznicki KA, McCrone JS, Kirchner J, Roussell DL, Strome S &
318 Bennett KL 1996 Multiple potential germ-line helicases are components of the
319 germ-line-specific P granules of *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* **93**
320 13837-13842.

321 Hamburger V & Hamilton HL 1992 A series of normal stages in the development of the chick
322 embryo. *Developmental Dynamic* **195** 231-272.

323 Hudson QJ, Smith CA & Sinclair AH 2005 Aromatase inhibition reduces expression of FOXL2
324 in the embryonic chicken ovary. *Dev Dyn* **233** 1052-1055.

325 Hughes SH, Greenhouse JJ, Petropoulos CJ & Sutrove P 1987 Adaptor plasmids simplify the
326 insertion of foreign DNA into helper-independent retroviral vectors. *J Virol* **61** 3004-3012.

327 Kato M, Shimada K, Saito N, Noda K & Ohta M 1995 Expression of P450 17
328 alpha-hydroxylase and P450aromatase genes in isolated granulosa, theca interna, and theca

329 externa layers of chicken ovarian follicles during follicular growth. *Biological*
330 *Reproduction* **52** 405-410.

331 Kawasaki I, Shim YH, Kirchner J, Kaminker J, Wood WB & Strome 1998 PGL-1, a predicted
332 RNA-binding component of germ granules, is essential for fertility in *C. elegans*. *Cell* **94**
333 635-645.

334 Kito G, Aramaki S, Tanaka K, Soh T, Yamauchi N & Hattori MA 2010 Temporal and spatial
335 differential expression of chicken germline-specific proteins cDAZL, CDH and CVH
336 during gametogenesis. *J Reprod Dev* **56** 341-346.

337 Kuramochi-Miyagawa S, Watanabe T, Gotoh K, Takamatsu K, Chuma S, Kojima-Kita K,
338 Shiromoto Y, Asada N, Toyoda A, Fujiyama A et al 2010 MVH in piRNA processing and
339 gene silencing of retrotransposons. *Genes & Development* **24** 887-892.

340 Kuznicki KA, Smith PA, Leung-Chiu WM, Estevez AO, Scott HC & Bennett KL 2000
341 Combinatorial RNA interference indicates GLH-4 can compensate for GLH-1; these two P
342 granule components are critical for fertility in *C. elegans*. *Development* **127** 2907-2916.

343 Lambeth LS, Cummins DM, Doran TJ, Sinclair AH & Smith CA 2013 Overexpression of
344 aromatase alone is sufficient for ovarian development in genetically male chicken embryos.
345 *PLoS One* **8** e683622013.

346 Linder P & Lasko P 2006 Bent out of shape: RNA unwinding by the DEAD-box helicase Vasa.
347 *Cell* **125** 219-221.

348 Loffler KA, Zarkower D & Koopman P 2003 Etiology of ovarian failure in blepharophimosis
349 ptosis epicanthus inversus syndrome: FOXL2 is a conserved, early-acting gene in
350 vertebrate ovarian development. *Endocrinology*. **144** 3237-3243.

351 Lorsch JR 2002 RNA chaperones exist and DEAD box proteins get a life. *Cell* **109** 797-800.

352 Nakata T, Ishiguro M, Aduma N, Izumi H & Kuroiwa A 2013 Chicken hemogen homolog is
353 involved in the chicken-specific sex-determining mechanism. *Proc Natl Acad Sci U S A*
354 **110** 3417-3422.

355 Pisarska MD, Barlow G & Kuo FT 2011 Minireview: roles of the forkhead transcription factor
356 FOXL2 in granulosa cell biology and pathology. *Endocrinology* **152** 1199-1208.

357 Roussell DL & Bennett KL 1993 glh-1, a germ-line putative RNA helicase from *Caenorhabditis*,
358 has four zinc fingers. *Proc Natl Acad Sci U S A*. **90** 9300-9304.

359 Sengoku T, Nureki O, Nakamura A, Kobayashi S & Yokoyama S 2006 Structural basis for RNA
360 unwinding by the DEAD-box protein *Drosophila* Vasa. *Cell* **125** 287-300.

361 Smith CA, McClive PJ, Hudson Q & Sinclair AH 2005 Male-specific cell migration into the
362 developing gonad is a conserved process involving PDGF signalling. *Dev Biol* **284**

- 363 337-350.
- 364 Smith CA, McClive PJ, Western PS, Reed KJ & Sinclair AH 1999a Conservation of a
365 sex-determining gene. *Nature* **402** 601-602.
- 366 Smith CA, Smith MJ & Sinclair AH 199b Gene expression during gonadogenesis in the chicken
367 embryo. *Gene* **234** 395-402.
- 368 Spike C, Meyer N, Racen E, Orsborn A, Kirchner J, Kuznicki K, Yee C, Bennett K & Strome S
369 2008 Genetic analysis of the *Caenorhabditis elegans* GLH family of P-granule proteins.
370 *Genetic* **178** 1973-1987.
- 371 Styhler S, Nakamura A, Swan A, Suter B & Lasko P 1998 vasa is required for GURKEN
372 accumulation in the oocyte, and is involved in oocyte differentiation and germline cyst
373 development. *Development* **125** 1569-1578.
- 374 Taylor L, Carlson DF, Nandi S, Sherman A, Fahrenkrug SC & McGrew MJ 2017 Efficient
375 TALEN-mediated gene targeting of chicken primordial germ cells. *Development*. **144**
376 928-934.
- 377 Tanaka R, Izumi H & Kuroiwa A 2017 Androgens and androgen receptor signaling contribute to
378 ovarian development in the chicken embryo. *Mol Cell Endocrinol* **443** 114-120.
- 379 Tanaka SS, Toyooka Y, Akasu R, Katoh-Fukui Y, Nakahara Y, Suzuki R, Yokoyama M & Noce

380 T 2000 The mouse homolog of Drosophila Vasa is required for the development of male
381 germ cells. *Genes & Development* **14** 841-853.

382 Tsunekawa N, Naito M, Sakai Y, Nishida T & Noce T 2000 Isolation of chicken vasa homolog
383 gene and tracing the origin of primordial germ cells. *Development* **127** 2741-2750.

384 Wang J & Gong Y 2017 Transcription of CYP19A1 is directly regulated by SF-1 in the theca
385 cells of ovary follicles in chicken. *Gen Comp Endocrinol* **247** 1-7.

386 Wang DS, Kobayashi T, Zhou LY, Paul-Prasanth B, Ijiri S, Sakai F, Okubo K, Morohashi K &
387 Nagahama Y 2007 Foxl2 up-regulates aromatase gene transcription in a female-specific
388 manner by binding to the promoter as well as interacting with ad4 binding
389 protein/steroidogenic factor 1. *Mol Endocrinol* **21** 712-725.

390 Zhao D, McBride D, Nandi S, McQueen HA, McGrew MJ, Hocking PM, Lewis PD, Sang HM
391 & Clinton M 2010 Somatic sex identity is cell autonomous in the chicken. *Nature* **464**
392 237-242.

393

394 **Figure legends**

395 **Figure 1 *DDX4* expression is efficiently suppressed *in vitro***

396 DF-1 cells were infected with (a, b) RCASB.DDX4_eGFP alone, (c, d) RCASA.miRNA.Sc and
397 RCASB.DDX4_eGFP, or (e, f) RCASA.miRNA.DDX4 and RCASB.DDX4_eGFP. Although
398 eGFP was observed in both cases (a-d), eGFP was not detected in DF-1 cells infected with
399 RCASA.miRNA.DDX4, which encoded two *DDX4*-targeting miRNAs, prior to
400 RCASB.DDX4_eGFP (e, f). Scale bar, 100 μ m. (g) RT-PCR analysis of *DDX4* in DF-1 cells
401 infected with RCASB.DDX4_eGFP alone or together with RCASA.miRNA.DDX4. The *DDX4*
402 expression is not detected in DF-1 cells infected with RCASA.miRNA.DDX4 prior to
403 RCASB.DDX4_eGFP (g). M: molecular marker.

404

405 **Figure 2 *DDX4* expression is efficiently suppressed *in vivo***

406 (a–h) Immunohistochemical staining of *DDX4* in the left gonads of (a–d) female and (e–h) male
407 embryos at E8.5 was performed to examine the KD efficiency of *DDX4*-targeting miRNAs. (a,
408 b) Female and (e, f) male gonads of WT embryos at E8.5 exhibited normal expression of *DDX4*,
409 indicating that PGCs localized to the cortical and interior of the embryonic gonad, respectively.
410 *DDX4* expression was suppressed in (c, d) female and (g, h) male embryos infected with

411 RCASA.miRNA.DDX4. FITC fluorescence is shown in a, c, e, g. Hoechst staining is shown in
412 B, D, F, H. Scale bar, 100 μm . (i) qRT-PCR analysis of *DDX4* in gonads at E8.5. Delivery of
413 *DDX4*-targeting miRNAs effectively suppressed *DDX4* expression. Black, dark gray, white, and
414 light gray bars correspond to WT female, *DDX4*-KD female, WT male, and *DDX4*-KD male
415 embryos, respectively. The data are presented as mean \pm standard deviation. * $P < 0.005$; $n \geq 3$.

416

417 **Figure 3 The number of PGCs is decreased in gonads of *DDX4*-KD embryos**

418 (a–d) AP staining in gonads of WT and KD embryos at E8.5. Gonad sections (dorsal-ventral
419 axis) of (a) WT female, (b) WT male, (c) *DDX4*-KD female, and (d) *DDX4*-KD male embryos.
420 Marked staining was observed in female and male gonads of WT embryos (a, b), but not in
421 those of *DDX4*-KD embryos (c, d). Arrows indicate AP staining. Scale bar, 100 μm . (e–g)

422 *DND1* *in situ* hybridization in frozen gonad sections of WT and KD embryo at E8.5. Gonad
423 sections of (e) WT female, (f) WT male, (g) *DDX4*-KD female, and (h) *DDX4*-KD male
424 embryos. Marked staining was observed in female and male gonads of WT embryos (E, F), but
425 not in those of *DDX4*-KD embryos (g, h). Arrows indicate *DND1* staining. Scale bar, 100 μm .

426 (i) AP staining and (j) *DND1* *in situ* hybridization. Black, dark gray, white, and light gray bars
427 show the numbers of positively stained cells in gonads of WT female, *DDX4*-KD female, WT

428 male, and *DDX4*-KD male embryos at E8.5, respectively.

429

430 **Figure 4 KD of *DDX4* decreases *CYP19A1* expression in female embryonic gonads**

431 qRT-PCR analysis of *CYP19A1*, *FOXL2*, *DMRT1*, and *SOX9* expression in gonads of WT

432 female (black bars), *DDX4*-KD female (dark gray bars), WT male (white bars), and *DDX4*-KD

433 male (light gray bars) embryos at E8.5. Expression of *DMRT1* and *SOX9* did not significantly



434 differ between WT and *DDX4*-KD embryos. *CYP19A1* expression in female gonads was



435 significantly lower in *DDX4*-KD embryos than in WT embryos. Expression of *FOXL2* was

436 slightly lower in female gonads of *DDX4*-KD embryos than in those of WT embryos, but this


437 difference was not statistically significant. The data are presented as mean \pm standard deviation.

438 * $P < 0.005$; $n \geq 3$.







AP staining



DND1 in situ hybridization



■ WT female ■ KD female □ WT male ■ KD male







Table 1 Primer sequences

Experiment	Direction of primer	Sequence
Construction of miRNA vectors DDX4-targeting miRNA #1	F R	5'- GAG AGG TGC TGC TGA GCG AGA TCC TGG TAT GCA AGA TCA TAG TGA AGC CAC AGA TGT A -3' 5'- ATT CAC CAC CAC TAG GCA TGA TCC TGG TAT GCA AGA TCA TAC ATC TGT GGC TTC ACT -3'
DDX4-targeting miRNA #2	F R	5'- CTG GTT CCT CCG TGA GCG AGG TGC TAA TGA AGG ACT TAA TAG TGA AGC CAC AGA TGT A -3' 5'- CCT GAA GAC CAG TAG GCA TGG TGC TAA TGA AGG ACT TAA TAC ATC TGT GGC TTC ACT -3'
Scrambled miRNA #1	F R	5'- GAG AGG TGC TGC TGA GCG AAC CTT CTC AAT TCT CAT CAT TAG TGA AGC CAC AGA TGT A -3' 5'- ATT CAC CAC CAC TAG GCA GAC CTT CTC AAT TCT CAT CAT TAC ATC TGT GGC TTC ACT -3'
Scrambled miRNA #2	F R	5'- CTG GTT CCT CCG TGA GCG ACA TGT TCC CTC GTC ACT TTA TAG TGA AGC CAC AGA TGT A -3' 5'- CCT GAA GAC CAG TAG GCA GCA TGT TCC CTC GTC ACT TTA TAC ATC TGT GGC TTC ACT -3'
Construction of the DDX4_GFP vector	F R	5'- TAA TCG ATG GAG GAG GAC TGG G -3' 5'- GCA CTA GTC TCC CAT GAC TTA AAT GTT G -3'
DND1 in situ hybridization probe	F R	5'- CAA CCG GAC CAA TAA GAT GG -3' 5'- ATT CCC TTC CAC CAG AGC TT -3'
qRT-PCR analysis of DDX4	F R	5'- GTA GCA TCA AGA GGC CTG GA -3' 5'- ACG ACC AGT TCG TCC AAT TC -3'