



Title	Free infinite divisibility for powers of random variables
Author(s)	Hasebe, Takahiro
Citation	ALEA : Latin American Journal of Probability and Mathematical Statistics, 13(1), 309-336
Issue Date	2016
Doc URL	http://hdl.handle.net/2115/75968
Type	article
File Information	ALEA Lat. Am. J. Probab. Math. Stat. 13-1_309-336.pdf



[Instructions for use](#)

Free infinite divisibility for powers of random variables

Takahiro Hasebe

Department of Mathematics, Hokkaido University

Kita 10, Nishi 8, Kitaku, Sapporo 060-0810 Japan

E-mail address: thasebe@math.sci.hokudai.ac.jp

URL: <http://www.math.sci.hokudai.ac.jp/~thasebe>

Abstract. We prove that X^r follows a free regular distribution, i.e. the law of a nonnegative free Lévy process if: (1) X follows a free Poisson distribution without an atom at 0 and $r \in (-\infty, 0] \cup [1, \infty)$; (2) X follows a free Poisson distribution with an atom at 0 and $r \geq 1$; (3) X follows a mixture of some HCM distributions and $|r| \geq 1$; (4) X follows some beta distributions and r is taken from some interval. In particular, if S is a standard semicircular element then $|S|^r$ is freely infinitely divisible for $r \in (-\infty, 0] \cup [2, \infty)$. Also we consider the symmetrization of the above probability measures, and in particular show that $|S|^r \text{sign}(S)$ is freely infinitely divisible for $r \geq 2$. Therefore S^n is freely infinitely divisible for every $n \in \mathbb{N}$. The results on free Poisson and semicircular random variables have a good correspondence with classical ID properties of powers of gamma and normal random variables.

1. Introduction

In classical probability, people have tried to understand if infinite divisibility (ID) can be preserved by powers, products or quotients of (independent) random variables (rvs). Usually the Lévy–Khintchine representation is not useful for this purpose and alternative new ideas are required. One class that behaves well with respect to powers and products is mixtures of exponential distributions (ME), i.e. rvs of the form EX where E, X are independent, E follows an exponential distribution and $X \geq 0$. The Goldie–Steutel theorem says that the class ME is a subset of the class ID (see [Goldie, 1967](#) and [Steutel, 1967](#)). In this class we have the implication

$$X \sim \text{ME} \Rightarrow X^r \sim \text{ME} \text{ for any } r \geq 1, \quad (1.1)$$

and also ME is closed under the product of independent rvs.

Received by the editors October 5, 2015; accepted March 8, 2016.

2010 Mathematics Subject Classification. 46L54, 60E07.

Key words and phrases. free infinite divisibility, free regularity, free Poisson distribution, semicircle distribution, hyperbolic complete monotonicity .

Quite a successful class in the theory of ID distributions is HCM (hyperbolically completely monotone) distributions, see [Bondesson \(1992\)](#). It is known that $\text{HCM} \subset \text{ID}$ and

$$X \sim \text{HCM} \Rightarrow X^r \sim \text{HCM} \text{ for any } |r| \geq 1, \quad (1.2)$$

see [Bondesson \(1992, p. 69\)](#). The class HCM moreover satisfies that if $X, Y \sim \text{HCM}$ and are independent then $XY, X/Y \sim \text{HCM}$, see [Bondesson \(1992, Theorem 5.1.1\)](#). Prior to the appearance of HCM, [Thorin \(1977b,a\)](#) introduced a class GGC (generalized gamma convolutions) which contains HCM (see [Bondesson, 1992, Theorem 5.1.2](#)). [Bondesson \(2015\)](#) recently proved that if $X, Y \sim \text{GGC}$ and X, Y are independent then $XY \sim \text{GGC}$. He also conjectured that $X \sim \text{GGC}$ implies that $X^r \sim \text{GGC}$ for any $r \geq 1$, which is still open. The class GGC is closed with respect to the addition of independent rvs, while ME and HCM are not. It is worth mentioning that [Shanbhag et al. \(1977\)](#) proved a related negative result that the product of two independent positive ID rvs is not always ID.

In free probability, the class of the free regular (FR) distributions, i.e. the laws of nonnegative free Lévy processes, is closed with respect to the product $\sqrt{XY}\sqrt{Y}$ where X, Y are free, see [Arizmendi et al. \(2013, Theorem 1\)](#). However, little is known on powers of rvs except that [Arizmendi et al. \(2013\)](#) showed that if $X \sim \text{FID}$ is even (i.e. having a symmetric distribution) then $X^2 \sim \text{FR}$. The main purpose of this paper is to consider the free infinite divisibility (FID) or more strongly free regularity of powers of rvs.

In this paper we will focus on several examples including free Poisson and semi-circular rvs and prove that: (1) If X follows a free Poisson distribution without an atom at 0, then $X^r \sim \text{FR}$ for any $r \in (-\infty, 0] \cup [1, \infty)$; (2) If X follows a free Poisson distribution with an atom at 0, then $X^r \sim \text{FR}$ for any $r \geq 1$; (3) If X follows some mixtures of (classical) HCM distributions then $X^r \sim \text{FR}$ for any $|r| \geq 1$; (4) If X follows some beta distributions then $X^r \sim \text{FR}$ for r in some interval. Our result has a consequence that $|S|^r \sim \text{FR}$ when S is the standard semicircular element and $r \in (-\infty, 0] \cup [2, \infty)$. We will also consider the symmetrization of powers of beta rvs and mixtures of HCM rvs, and in particular, show that $|S|^r \text{sign}(S) \sim \text{FID}$ for $r \geq 2$. These results imply that S^n are FID for all $n \in \mathbb{N}$. Our result on HCM distributions generalizes [Arizmendi and Hasebe \(2016, Proposition 4.21\(1\)\)](#) where mixtures of positive Boolean stable laws are shown to be FID and part of [Hasebe \(2014, Theorem 1.2\(3\)\)](#) where beta distributions of the 2nd kind are shown to be FID.

The proofs depend on the complex-analytic method which has been developed recently. Until around 2010, one could prove the FID property of a given probability measure only when the R -transform [Voiculescu \(1986\)](#) or Voiculescu transform ([Bercovici and Voiculescu, 1993](#)) is explicit. There had been no way to prove the FID property if the R -transform is not explicit. Actually many probability distributions used in classical probability theory do not have explicit R -transforms, e.g. normal distributions, gamma distributions and beta distributions. By contrast, there are lots of methods in classical probability to show that a probability measure is ID, even if its characteristic function is not explicit.

In 2011, Belinschi et al. changed this situation and they gave the first non-trivial FID probability measure: the normal distribution is FID ([Belinschi et al., 2011](#)). The proof is based on the complex analysis of the Cauchy transform (but there is combinatorial background). Since then, several other people developed the

complex-analytic method. Now many nontrivial distributions are known to be FID: Anshelevich et al. showed that the q -normal distribution is FID for $q \in [0, 1]$ [Anshelevich et al. \(2010\)](#) and the author proved that beta distributions of the 1st kind and 2nd kind, gamma, inverse gamma and Student distributions are FID for many parameters [Hasebe \(2014\)](#). Other results can be found in [Arizmendi and Belinschi \(2013\)](#); [Arizmendi and Hasebe \(2013, 2014, 2016\)](#); [Arizmendi et al. \(2013\)](#); [Bożejko and Hasebe \(2013\)](#). These examples suggest that the intersection of ID and FID is rich. Almost all of these FID distributions belong to a further subclass UI that was introduced in [Arizmendi and Hasebe \(2013\)](#). With this class UI we are able to show the FID property of a given probability measure without knowing the explicit R -transform or free Lévy–Khintchine representation. This class plays an important role in the present paper too.

This paper contains two sections besides this section. In Section 2 we will introduce basic notations and concepts including the classes ID, ME, GGC, HCM, FID, FR, UI and probability measures to be treated. In Section 3 we will state the main results rigorously and then prove them. Section 4 explains similarity between our results on free Poisson and semicircle distributions and the classical results on gamma and normal distributions. Some conjectures are proposed based on this similarity.

2. Preliminaries

Some general notations in this paper are summarized below.

- (1) \mathcal{P}_+ is the set of (Borel) probability measures on $[0, \infty)$.
- (2) For a classical or noncommutative rv X and a probability measure μ on \mathbb{R} , the notation $X \sim \mu$ means that the rv X follows the law μ . A similar notation is used for a subclass of probability measures: for example $X \sim \mathcal{P}_+$ means that $X \sim \mu$ for some $\mu \in \mathcal{P}_+$.
- (3) The function z^p is the principal value defined on $\mathbb{C} \setminus (-\infty, 0]$.
- (4) The function $\arg(z)$ is the argument of z defined in $\mathbb{C} \setminus (-\infty, 0]$, taking values in $(-\pi, \pi)$. We also use another argument $\arg_I(z)$ taking values in an interval I .

2.1. ID distributions and subclasses. Infinitely divisible distributions and subclasses are summarized here. The reader is referred to [Steutel and van Harn \(2004\)](#); [Bondesson \(1992\)](#) for more information on this section.

A probability measure on \mathbb{R} is said to be *infinitely divisible* (ID) if it has an n^{th} convolution root for any $n \in \mathbb{N}$. The class of ID distributions is denoted by ID (and this kind of notations will be adapted to other classes of probability measures too).

Let $\gamma(p, \theta)$ be the gamma distribution

$$\frac{1}{\theta^p \Gamma(p)} x^{p-1} e^{-x/\theta} 1_{(0, \infty)}(x) dx, \quad p, \theta > 0, \quad (2.1)$$

where θ corresponds to the scaling. A probability measure $\mu \in \mathcal{P}_+$ is called a *mixture of exponential distributions* (ME) if there exists $\nu \in \mathcal{P}_+$ such that $\mu = \gamma(1, 1) \otimes \nu$, where \otimes is classical multiplicative convolution. An equivalent definition is that μ is of the form $w\delta_0 + f(x)dx$, where $w \in [0, 1]$ and f is a C^∞ function on $(0, \infty)$ which is completely monotone, i.e. $(-1)^n f^{(n)} \geq 0$ for any $n \in \mathbb{N} \cup \{0\}$. It is known that $\text{ME} \subset \text{ID}$, called the Goldie–Steutel theorem.

A probability measure on $[0, \infty)$ is called a *generalized gamma convolution* (GGC) if it is in the weak closure of the set

$$\{\gamma(p_1, \theta_1) * \cdots * \gamma(p_n, \theta_n) \mid p_k, \theta_k > 0, k = 1, \dots, n, n \in \mathbb{N}\}, \quad (2.2)$$

that is, the class of GGCs is the smallest subclass of ID that contains all the gamma distributions and that is closed under convolution and weak limits.

A pdf (probability density function) $f : (0, \infty) \rightarrow [0, \infty)$ is said to be *hyperbolically completely monotone* (HCM) if for each $u > 0$ the map $w \mapsto f(uv)f(u/v)$ is completely monotone as a function of $w = v + 1/v$. A probability distribution on $(0, \infty)$ is called an HCM distribution if it has an HCM pdf. It turns out that any HCM pdf is the pointwise limit of pdfs of the form

$$C \cdot x^{p-1} \prod_{k=1}^n (t_k + x)^{-\gamma_k}, \quad C, p, t_k, \gamma_k > 0, p < \sum_{k=1}^n \gamma_k, n \in \mathbb{N}. \quad (2.3)$$

This limiting procedure gives us the representation of an HCM pdf

$$f(x) = C \cdot x^{\alpha-1} \exp \left(-b_1 x - \frac{b_2}{x} + \int_{(1, \infty)} \log \left(\frac{t+1}{t+x} \right) \Gamma_1(dt) \right. \\ \left. + \int_{[1, \infty)} \log \left(\frac{t+1}{t+1/x} \right) \Gamma_2(dt) \right) \quad (2.4)$$

where $\alpha \in \mathbb{R}$, $b_1, b_2 \geq 0$ and Γ_1, Γ_2 are measures on $(1, \infty)$ and $[1, \infty)$, respectively, such that $\int (1+t)^{-1} \Gamma_k(dt) < \infty$. However these conditions on the parameters are not sufficient to ensure the integrability of f . The author does not know how to write down necessary and sufficient conditions for $\int_{(0, \infty)} f(x) dx < \infty$ in terms of the parameters $\alpha, b_1, b_2, \Gamma_1, \Gamma_2$. An important fact in the theory of ID distributions is that $\text{HCM} \subset \text{GGC}$.

2.2. FID distributions and subclasses. A probability measure on \mathbb{R} is said to be *freely infinitely divisible* (FID) if it has an n^{th} free convolution root for each $n \in \mathbb{N}$. Let FID be the set of FID distributions on \mathbb{R} . Basic results on the class FID were established in [Bercovici and Voiculescu \(1993\)](#). Connections between the class FID and free Lévy processes were investigated in [Biane \(1998\)](#); [Barndorff-Nielsen and Thorbjørnsen \(2002, 2005\)](#); [Barndorff-Nielsen et al. \(2006\)](#); [Arizmendi et al. \(2013\)](#). [Bercovici and Pata \(1999\)](#) clarified how the FID distributions appear as the limit of the sum of free i.i.d. rvs.

We say that $\mu \in \mathcal{P}_+$ is *free regular* (FR) if μ is FID and $\mu^{\boxplus t} \in \mathcal{P}_+$ for all $t > 0$. This notion was introduced in [Pérez-Abreu and Sakuma \(2012\)](#) in terms of the Bercovici–Pata bijection and then further developed in [Arizmendi et al. \(2013\)](#) in terms nonnegative free Lévy processes. The set of free regular distributions is denoted by FR. The class FR is closed with respect to the weak convergence, see [Arizmendi et al. \(2013, Proposition 25\)](#). A probability measure in $\text{FID} \cap \mathcal{P}_+$ may not be free regular, but we have a criterion.

Lemma 2.1 (Theorem 13 in [Arizmendi et al., 2013](#)). *Suppose $\mu \in \text{FID} \cap \mathcal{P}_+$ and μ satisfies either (i) $\mu(\{0\}) > 0$, or (ii) $\mu(\{0\}) = 0$ and $\int_{(0,1)} x^{-1} \mu(dx) = \infty$. Then $\mu \in \text{FR}$.*

There is a useful subclass of FID, called UI. The idea already appeared implicitly in [Belinschi et al. \(2011\)](#) and the explicit definition was given in [Arizmendi and](#)

Hasebe (2013). The following form of definition is in Bożejko and Hasebe (2013). To define the class UI, let G_μ (or G_X if $X \sim \mu$) denote the Cauchy transform

$$G_\mu(z) = \int_{\mathbb{R}} \frac{1}{z-x} \mu(dx), \quad z \in \mathbb{C}^+, \quad (2.5)$$

where \mathbb{C}^\pm denote the complex upper and lower half-planes respectively. Bercovici and Voiculescu (1993) proved that for any $a > 0$, there exist $\lambda, M, b > 0$ such that G_μ is univalent in the truncated cone

$$\Gamma_{\lambda,M} = \{z \in \mathbb{C}^+ \mid \lambda|\operatorname{Re}(z)| < \operatorname{Im}(z), \operatorname{Im}(z) > M\} \quad (2.6)$$

and $G_\mu(\Gamma_{\lambda,M})$ contains the triangular domain

$$\Delta_{a,b} = \{w \in \mathbb{C}^- \mid a|\operatorname{Re}(w)| < -\operatorname{Im}(w), \operatorname{Im}(w) > -b\}. \quad (2.7)$$

So we may define a right compositional inverse G_μ^{-1} in $\Delta_{a,b}$.

Definition 2.2 (Definition 5.1 in Arizmendi et al., 2013). A probability measure μ on \mathbb{R} is said to be in class UI (standing for univalent inverse Cauchy transform) if the right inverse map G_μ^{-1} , originally defined in a triangular domain $\Delta_{a,b} \subset \mathbb{C}^-$, has univalent analytic continuation to \mathbb{C}^- .

The importance of this class is based on the following lemma proved in Arizmendi and Hasebe (2013, Proposition 5.2 and p. 2763).

Lemma 2.3. UI \subset FID. Moreover, UI is w -closed (i.e. closed with respect to weak convergence).

To consider symmetric distributions, a symmetric version of UI is useful. The following definition is equivalent to Hasebe (2014, Definition 2.5(2)).

Definition 2.4. A symmetric probability measure μ is said to be in class UI_s if (a) the right inverse map G_μ^{-1} , defined in a domain $\Delta_{a,b}$, has univalent analytic continuation to a neighborhood of $i(-\infty, 0)$, and (b) the right inverse G_μ^{-1} , defined in the domain $\Delta_{a,b}$, has univalent analytic continuation to $\mathbb{C}^- \cap i\mathbb{C}^-$ such that $G_\mu^{-1}(\mathbb{C}^- \cap i\mathbb{C}^-) \subset \mathbb{C}^- \cup i\mathbb{C}^-$.

Condition (a) cannot be dropped from the definition: $G_{\frac{1}{2}(\delta_{-1} + \delta_1)}^{-1}(z) = \frac{1 + \sqrt{1 + 4z^2}}{2z}$ satisfies condition (b) but not condition (a).

If a symmetric probability measure μ belongs to UI then $\mu \in \text{UI}_s$ by definition, but the converse is not true. In fact a probability measure $\mu \in \text{UI}_s$ belongs to UI iff $G_\mu^{-1}(\mathbb{C}^- \cap i\mathbb{C}^-) \subset i\mathbb{C}^-$.

The analogue of Lemma 2.3 holds true.

Lemma 2.5. $\text{UI}_s \subset$ FID. Moreover UI_s is w -closed.

Proof: The first claim was proved in Hasebe (2014, Lemma 2.7), but the second claim is new. Suppose that $\mu_n \in \text{UI}_s$ and $\mu_n \rightarrow \mu$ (which implies that $\mu \in \text{FID}$). By Barndorff-Nielsen and Thorbjørnsen (2002, Theorem 3.8) the Voiculescu transform φ_{μ_n} converges to φ_μ uniformly on each compact subset of \mathbb{C}^+ and therefore so does $G_{\mu_n}^{-1}(z) = \varphi_{\mu_n}(z^{-1}) + z^{-1}$ to $G_\mu^{-1}(z) = \varphi_\mu(z^{-1}) + z^{-1}$ on each compact subset of \mathbb{C}^- . Since $G_{\mu_n}^{-1}$ is not a constant, it is univalent in $\mathbb{C}^- \cap i\mathbb{C}^-$ by Hurwitz's theorem. Taking the limit we have $G_\mu^{-1}(\mathbb{C}^- \cap i\mathbb{C}^-) \subset \overline{\mathbb{C}^- \cup i\mathbb{C}^-}$, but since G_μ^{-1} is analytic and not a constant, $G_\mu^{-1}(\mathbb{C}^- \cap i\mathbb{C}^-)$ is an open set by the open mapping theorem, so in fact $G_\mu^{-1}(\mathbb{C}^- \cap i\mathbb{C}^-)$ is contained in $\mathbb{C}^- \cup i\mathbb{C}^-$.

Hurwitz's theorem is not useful for proving the univalence around $i(-\infty, 0)$ since the neighborhoods of $i(-\infty, 0)$ may shrink as $n \rightarrow \infty$, and so an alternative idea is needed. Our assumptions imply that $G_{\mu_n}^{-1}$ are locally univalent, i.e. $(G_{\mu_n}^{-1})'(z) \neq 0$ for all $z \in \mathbb{C}^-$ (note that $G_{\mu_n}^{-1}$ has symmetry with respect to the y -axis). Since a zero of an analytic function changes continuously with respect to the local uniform topology of analytic functions, $(G_{\mu}^{-1})'$ does not have a zero in \mathbb{C}^- ; otherwise $(G_{\mu_n}^{-1})'$ would have a zero for large n . In particular $(G_{\mu}^{-1})'(z) \neq 0$ in $i(-\infty, 0)$. Since μ is symmetric, $G_{\mu}^{-1}(i(-\infty, 0)) \subset i\mathbb{R}$ and hence $(G_{\mu}^{-1})'(i(-\infty, 0)) \subset \mathbb{R}$. Since the derivative $(G_{\mu}^{-1})'(z)$ does not have a zero, the sign does not change (in fact we can show that $(G_{\mu}^{-1})'(z) > 0$ in $i(-\infty, 0)$). Therefore G_{μ}^{-1} is univalent in a neighborhood of $i(-\infty, 0)$ by applying Noshiro–Warschawski's theorem (see [Noshiro, 1934](#), Theorem 12 and [Warschawski, 1935](#), Lemma 1) to a neighborhood of $i(-\infty, 0)$ where $\operatorname{Re}((G_{\mu}^{-1})'(z)) > 0$. \square

2.3. Probability measures to be treated. We introduce several (classes of) probability measures to be treated in this paper.

(1) The semicircle distribution $\mathcal{S}(m, \sigma^2)$ is the probability measure with pdf

$$\frac{1}{2\pi\sigma^2} \sqrt{4\sigma^2 - (x - m)^2} 1_{(m-2\sigma, m+2\sigma)}(x), \quad m \in \mathbb{R}, \sigma > 0. \quad (2.8)$$

By using the R -transform one can compute the inverse Cauchy transform

$$G_{\mathcal{S}(m, \sigma^2)}^{-1}(z) = m + \sigma^2 z + \frac{1}{z}. \quad (2.9)$$

One can prove that $G_{\mathcal{S}(m, \sigma^2)}^{-1}$ is a bijection from \mathbb{C}^- onto $\mathbb{C}^+ \cup (m - 2\sigma, m + 2\sigma) \cup \mathbb{C}^-$, so that $\mathcal{S}(m, \sigma^2) \in \text{UI}$ and in particular $\mathcal{S}(0, \sigma^2) \in \text{UI}_s$.

(2) The free Poisson distribution (or Marchenko–Pastur distribution) $\mathbf{fp}(p, \theta)$ is

$$\max\{1 - p, 0\} \delta_0 + \frac{\sqrt{(\theta(1 + \sqrt{p})^2 - x)(x - \theta(1 - \sqrt{p})^2)}}{2\pi\theta x} 1_{(\theta(1 - \sqrt{p})^2, \theta(1 + \sqrt{p})^2)}(x) dx, \quad (2.10)$$

where $p, \theta > 0$. The parameter θ stands for the scale parameter. Since $\mathbf{fp}(p, \theta)^{\boxplus t} = \mathbf{fp}(pt, \theta) \in \mathcal{P}_+$ for any $t > 0$, $\mathbf{fp}(p, \theta)$ is free regular. The inverse Cauchy transform is given by

$$G_{\mathbf{fp}(p, \theta)}^{-1}(z) = \frac{1}{z} + \frac{p\theta}{1 - \theta z}. \quad (2.11)$$

One can show that $G_{\mathbf{fp}(p, \theta)}^{-1}$ is a bijection from \mathbb{C}^- onto $\mathbb{C}^+ \cup (\theta(\sqrt{p} - 1)^2, \theta(\sqrt{p} + 1)^2) \cup \mathbb{C}^-$ and so $\mathbf{fp}(p, \theta) \in \text{UI}$.

(3) The beta distribution $\beta(p, q)$ is the probability measure with pdf

$$\frac{1}{B(p, q)} \cdot x^{p-1} (1 - x)^{q-1} 1_{(0, 1)}(x), \quad p, q > 0. \quad (2.12)$$

The beta distribution belongs to the class UI if (i) $p, q \geq \frac{3}{2}$, (ii) $0 < p \leq \frac{1}{2}$, $p + q \geq 2$ or (iii) $0 < q \leq \frac{1}{2}$, $p + q \geq 2$ (see [Hasebe, 2014](#)).

(4) The positive Boolean (strictly) stable distribution \mathbf{b}_α , introduced by [Speicher and Woroudi \(1997\)](#), is the distribution with pdf

$$\frac{\sin(\alpha\pi)}{\pi} \cdot \frac{x^{\alpha-1}}{x^{2\alpha} + 2(\cos \pi\alpha)x^\alpha + 1} 1_{(0, \infty)}(x), \quad \alpha \in (0, 1). \quad (2.13)$$

We then consider the classical mixtures of positive Boolean stable distributions

$$\mathfrak{B}_\alpha = \{\mu \circledast \mathbf{b}_\alpha \mid \mu \in \mathcal{P}_+\}, \quad \alpha \in (0, 1), \quad (2.14)$$

where \circledast is classical multiplicative convolution, i.e. $XY \sim \mu \circledast \nu$ when $X \sim \mu, Y \sim \nu$ and X, Y are independent. This class was introduced and investigated in [Arizmendi and Hasebe \(2016\)](#). It is known that $\mathfrak{B}_\alpha \subset \mathfrak{B}_\beta$ when $0 < \alpha \leq \beta < 1$ and $\mathfrak{B}_{1/2} \subset \text{UI} \cap \text{FR} \cap \text{ME}$. The property $\mathfrak{B}_{1/2} \subset \text{FR}$ is not explicitly stated in [Arizmendi and Hasebe \(2016\)](#), but it is a consequence of the fact $\mathfrak{B}_{1/2} \subset \text{FID} \cap \mathcal{P}_+$ proved in [Arizmendi and Hasebe \(2016, Proposition 4.21\)](#), the fact that any measure in $\mathfrak{B}_{1/2} \setminus \{\delta_0\}$ has a pdf which diverges to infinity at 0, and Lemma 2.1 above.

(5) We consider a further subclass of HCM distributions with pdf

$$f(x) = C \cdot x^{p-1} \exp \left(\int_{(0, \infty)} \log \left(\frac{1}{t+x} \right) \Gamma(dt) \right), \quad (2.15)$$

where Γ is a finite measure on $(0, \infty)$, $0 < p < \Gamma((0, \infty))$ and $\int_{(0, \infty)} |\log t| \Gamma(dt) < \infty$ (these conditions ensure the integrability of f and hence we can take the normalizing constant $C > 0$). This subclass is a natural generalization of pdfs of the form (2.3), but does not cover all HCM pdfs. This pdf is of the form of Markov-Krein transform [Kerov \(1998\)](#).

3. Main results

3.1. Statements. Now we are ready to state the main theorems. Independence of random variables means classical independence below.

Theorem 3.1. *Suppose $X \sim \text{HCM}$ whose pdf is of the form (2.15) and satisfies*

$$0 < p \leq \frac{1}{2}, \quad 0 < -p + \Gamma((0, \infty)) \leq \frac{1}{2}. \quad (3.1)$$

If $W \geq 0$ is a rv independent of X and $|r| \geq 1$, then $WX^r \sim \text{FR} \cap \text{UI}$. The case $r \leq -1$ can be considered only when $W > 0$ almost surely. Moreover, X^{-1} also has a pdf of the form (2.15) that satisfies condition (3.1).

Remark 3.2. If one uses [Bondesson \(1992, Theorems 4.1.1, 4.1.4\)](#) and the factorization of a gamma rv as a product of a beta rv and a gamma rv on p. 14 of [Bondesson \(1992\)](#), one can show that $WX^r \sim \text{ME}$ and thus we get $WX^r \sim \text{FR} \cap \text{UI} \cap \text{ME}$.

Since the Boolean stable law $\mathbf{b}_{1/2}$ has the pdf $\pi^{-1}x^{-1/2}(x+1)^{-1}$ which satisfies the assumptions of Theorem 3.1, we have

Corollary 3.3. *If $X \sim \mathfrak{B}_{1/2}$ and $|r| \geq 1$ then $X^r \sim \text{FR} \cap \text{UI}$ ($r \leq -1$ can be considered only when the mixing measure does not have an atom at 0).*

Theorem 3.4. *Suppose that $X \sim \beta(p, q)$. If $r \geq 1$, $q \in [\frac{3}{2}, \frac{5}{2}]$, $0 < 2p \leq r$ and $(q - \frac{3}{2})r \leq p + q - 1 \leq r$, then $X^r \sim \text{FR} \cap \text{UI}$.*

Since $\beta(\frac{1}{2}, \frac{3}{2}) = \mathbf{fp}(1, \frac{1}{4})$, we conclude that if $X \sim \mathbf{fp}(1, 1)$ then $X^r \sim \text{FR} \cap \text{UI}$ for $r \geq 1$. However we will prove a stronger result.

Theorem 3.5. (1) *If $X \sim \mathbf{fp}(p, 1)$, $p \geq 1$ and $r \in (-\infty, 0] \cup [1, \infty)$, then $X^r \sim \text{FR} \cap \text{UI}$.*

(2) *If $X \sim \mathbf{fp}(p, 1)$, $0 < p < 1$ and $r \geq 1$, then $X^r \sim \text{FR} \cap \text{UI}$.*

We avoided the negative powers for $0 < p < 1$ since $\mathbf{fp}(p, 1)$ has an atom at 0. Since $S \sim \mathcal{S}(0, 1)$ implies $S^2 \sim \mathbf{fp}(1, 1)$, the following result is immediate.

Corollary 3.6. *If $S \sim \mathcal{S}(0, 1)$ and $r \in (-\infty, 0] \cup [2, \infty)$ then $|S|^r \sim \text{FR} \cap \text{UI}$.*

Remark 3.7. The result for $r = 4$ was proved in [Arizmendi et al. \(2013\)](#). Recently Chistyakov and Lehner showed the $r = 6$ case (in private communication). A related but negative result is that $(S + a)^2 \not\sim \text{FID}$ for all $a \neq 0$, see [Eisenbaum \(2012\)](#).

We consider the symmetrized versions.

Theorem 3.8. (1) *Suppose $X \sim \text{HCM}$ satisfying condition (3.1) in Theorem 3.1. If W is independent of X and has a symmetric distribution, then $WX^r \sim \text{UI}_s$ for $|r| \geq 1$.*

(2) *Suppose that B, X are independent and $B \sim \frac{1}{2}(\delta_{-1} + \delta_1)$, $X \sim \beta(p, q)$. Under the assumptions on p, q, r in Theorem 3.4, we have $BX^r \sim \text{UI}_s$.*

(3) *Suppose that B, X are independent and $B \sim \frac{1}{2}(\delta_{-1} + \delta_1)$, $X \sim \beta(p, q)$. If either (i) $p > r > 0$ or (ii) $r \leq 0$ then we have $BX^r \not\sim \text{FID}$.*

The case $(p, q) = (1/2, 3/2)$ in (2) and (3) gives us

Corollary 3.9. (1) *If $S \sim \mathcal{S}(0, 1)$ and $r \geq 2$ then $|S|^r \text{sign}(S) \sim \text{UI}_s$.*

(2) *If $S \sim \mathcal{S}(0, 1)$ and $r < 1$ then $|S|^r \text{sign}(S) \not\sim \text{FID}$.*

Remark 3.10. Corollary 3.6 and Corollary 3.9 imply that S^n are FID for all $n \in \mathbb{N}$. In the special case $n = 2$, the distribution of $|S|^2 \text{sign}(S)$ is the symmetrized beta distribution with parameters $1/2, 3/2$ and [Arizmendi et al. \(2010, Proposition 11\)](#) already proved that it is FID.

Discussions and further remarks on main theorems. (1) In Theorem 3.8 only the symmetric Bernoulli rv B is considered for the mixing of beta rvs, while a general W with a symmetric distribution appears in the HCM case. This Bernoulli distribution cannot be generalized to arbitrary symmetric distributions. Actually if W has a symmetric discrete distribution μ_W whose support has cardinality ≥ 4 , then WX is not FID for any beta rv X . The proof is as follows. The pdf of WX is positive at $c := \min\{x \in \text{supp}(\mu_W) \mid x > 0\}$ but is not real analytic at c . This shows that the distribution of WX is not FID by [Belinschi and Bercovici \(2005, Proposition 5.1\)](#). For a similar reason, one cannot take a (positive) scale mixing of beta or free Poisson rvs in Theorem 3.4 or in Theorem 3.5.

(2) Related to Theorem 3.8, a natural question is the symmetrization of powers of free Poisson rvs, i.e. the law of BX^r where B, X are independent, $B \sim \frac{1}{2}(\delta_{-1} + \delta_1)$ and $X \sim \mathbf{fp}(p, 1)$.

- If $p = 1$ and $r \geq 1$ then $BX^r \sim \text{UI}_s$ from Corollary 3.9(1).
- If $p > 1$ and $r \in \mathbb{R}$ then BX^r is not FID from Lemma 3.16 that we will show later.
- If $p < 1$ and $r \geq 1$ then it is not known whether BX^r is FID or not. Actually one can check condition (b) in Definition 2.4 similarly to the proof of Theorem 3.8, but it is not clear how to show condition (a).

(3) In Theorems 3.1, 3.4, 3.8, the assumptions on the parameters of beta and HCM distributions may seem too restrictive. Weakening these assumptions is left to future research.

3.2. *Proofs.* The integral form of the Cauchy transform gives us an analytic function defined outside the support of μ , and we denote it by $\tilde{G}_\mu(z)$:

$$\tilde{G}_\mu(z) = \int_{\mathbb{R}} \frac{1}{z - x} \mu(dx), \quad z \in \mathbb{C} \setminus \text{supp}(\mu). \quad (3.2)$$

In the study of FID distributions, more important is the analytic continuation of the Cauchy transform G_μ from \mathbb{C}^+ into \mathbb{C}^- passing through the support of μ . This analytic continuation is possible when the pdf is real analytic and the explicit formula can be given in terms of $\tilde{G}_\mu(z)$ and the pdf. We state the result in a slightly general form where complex measures are allowed, but the proof is similar to that of [Hasebe \(2014, Proposition 4.1\)](#). Note that $G_\sigma, \tilde{G}_\sigma$ can be defined for complex measures σ by linearity.

Lemma 3.11. *Let $I \subset \mathbb{R}$ be an open interval. Suppose that f is analytic in a neighborhood of $I \cup \mathbb{C}^-$ and that f is integrable on I with respect to the Lebesgue measure. We define the complex measure $\sigma(dx) := f(x)1_I(x)dx$. Then the Cauchy transform G_σ defined on \mathbb{C}^+ has analytic continuation to $\mathbb{C}^+ \cup I \cup \mathbb{C}^-$, which we denote by the same symbol G_σ , and*

$$G_\sigma(z) = \tilde{G}_\sigma(z) - 2\pi i f(z), \quad z \in \mathbb{C}^-. \quad (3.3)$$

The following lemma gives a crucial idea for showing the main theorems. The idea of the proof is to take the curve γ in [Bożejko and Hasebe \(2013, Proposition 2.1\)](#) to be the one that starts from $-\infty + i0$, then goes to 0, turns 180° around 0 and then goes to $-\infty - i0$. This curve is useful since we can easily compute the boundary value of G_μ on $(-\infty, 0) - i0$ thanks to formula (3.3), so that we can check condition (B) in [Bożejko and Hasebe \(2013, Proposition 2.1\)](#).

Lemma 3.12. *Let μ be a probability measure on $[0, \infty)$ which has a pdf of the form*

$$f(x) = x^{p-1}g(x), \quad x > 0, \quad (3.4)$$

where

- (A1) $0 < p \leq \frac{1}{2}$;
- (A2) g is the restriction of an analytic function (also denoted by g) defined in $\{z \in \mathbb{C} \setminus \{0\} \mid \arg(z) \in (-\pi, \theta_0)\}$ for some $\theta_0 \in (0, \pi)$ and the restriction $g|_{\mathbb{C}^-}$ extends to a continuous function on $\mathbb{C}^- \cup (-\infty, 0)$;
- (A3) $\lim_{z \rightarrow 0, \arg(z) \in (-\pi, \theta_0)} g(z) > 0$;
- (A4) $\lim_{|z| \rightarrow \infty, \arg(z) \in (-\pi, 0)} z^\gamma f(z) = 0$ for some $\gamma > 0$;
- (A5) $\text{Re}(f(x - i0)) \leq 0$ for $x < 0$.

Then $\rho \otimes \mu \in \text{FR} \cap \text{UI}$ for all $\rho \in \mathcal{P}_+$.

Proof: We first assume that $\rho = \delta_1$ and $0 < p < 1/2$ and later drop these assumptions. Lemma 3.11 implies that the Cauchy transform G_μ has analytic continuation (denoted by G_μ too) to $\mathbb{C} \setminus (-\infty, 0]$ and $G_\mu(z)$ is given by formula (3.3) in \mathbb{C}^- .

In the following $\eta > 0$ is supposed to be large and $\delta > 0$ is supposed to be small. We consider curves:

- c_1 is the real line segment from $-\eta + i0$ to $-\delta + i0$;
- c_2 is the clockwise circle $\delta e^{i\theta}$ where θ starts from π and ends with $-\pi$;
- c_3 is the line segment from $-\delta - i0$ to $-\eta - i0$;
- c_4 is the counterclockwise circle centered at 0, starting from $-\eta - i0$ and stopping at $-\eta + i0$.

Note that the line segments c_1 and c_3 are meant to be different by considering a Riemannian surface. The left of Fig. 3.1 shows the directed closed curve consisting of $c_k, k = 1, 2, 3, 4$. Let g_k be the image curve $G_\mu(c_k)$ for $k = 1, 2, 3, 4$. More precisely, the curves g_1, g_3 are defined by $G_\mu(c_1 + i0), G_\mu(c_3 - i0)$ respectively, and hence g_1 lies on the negative real line.




FIGURE 3.1. The curves c_k and g_k

Let $\varepsilon > 0$ be supposed to be small. We claim the existence of $\eta > 0$ large enough so that $|G_\mu(z)| < \varepsilon$ for $|z| \geq \eta, z \in \mathbb{C} \setminus (-\infty, 0]$. This can be proved by dividing the region into the two parts $\arg(z) \in (-b, \pi)$ and $\arg(z) \in (-\pi, -b]$ where $b \in (0, \pi/2)$ is arbitrary. First, thanks to assumption (A4) we may change the contour of the integral and obtain

$$G_\mu(z) = \int_{e^{-2ib}(0, \infty)} \frac{1}{z-w} f(w) dw, \quad \arg(z) \in (-b, \pi). \quad (3.5)$$

Using assumption (A4), we can show that $\sup_{\theta \in (-b, \pi)} |G_\mu(re^{i\theta})| \rightarrow 0$ as $r \rightarrow \infty$. For $\arg(z) \in (-\pi, -b]$, we use (3.3) and assumption (A4) to obtain that $\sup_{\theta \in (-\pi, -b]} |G_\mu(re^{i\theta})| \rightarrow 0$ as $r \rightarrow \infty$. This is what we claimed. Therefore, the curve g_4 is contained in the ball centered at 0 with radius ε .

From assumption (A5) it holds that

$$\operatorname{Im}(G_\mu(z)) = \operatorname{Im}(\tilde{G}_\mu(z)) - 2\pi \operatorname{Im}(if(z)) \geq 0, \quad z \in [-\eta, -\delta] - i0 \quad (3.6)$$

since $\operatorname{Im}(\tilde{G}_\mu(z)) = 0$ in $(-\infty, 0) - i0$. Therefore the curve g_3 is contained in $\{w \mid \operatorname{Im}(w) \geq 0\}$.

Assumptions (A2), (A3) allow us to use Hasebe (2014, (5.6)). The assumptions of Hasebe (2014, Theorem 5.1) are a little different from the present case, but the proof is applicable to our case without a change, to conclude that there exists $a > 0$ such that

$$G_\mu(z) = -a(-z)^{p-1} + o(|z|^{p-1}), \quad \text{as } z \rightarrow 0, -\pi < \arg(z) < \pi. \quad (3.7)$$

By asymptotics (3.7), we can take $\delta \in (0, (a\varepsilon/2)^{\frac{1}{1-p}})$ small enough so that

$$|G_\mu(\delta e^{i\theta}) + a(-\delta e^{i\theta})^{p-1}| < \frac{a\delta^{p-1}}{2} \quad (3.8)$$

uniformly on $\theta \in (-\pi, \pi)$. Therefore $|G_\mu(\delta e^{i\theta})| > \frac{1}{2}a\delta^{p-1} > \varepsilon^{-1}$, and so the distance between the curve g_2 and 0 is larger than ε^{-1} . Since we have (3.7), if $\delta, \varepsilon > 0$ are small enough then the final point of g_2 has an argument approximately

equal to $\arg(-(-(-\delta-i0))^{p-1}) = \arg(-(\delta e^{-2\pi i})^{p-1}) = (1-2p)\pi$, which is contained in $(0, \pi)$ because we assumed $p < 1/2$.

From the above arguments, every point of $D_\varepsilon := \{w \in \mathbb{C}^- \mid \varepsilon < |w| < \varepsilon^{-1}\}$ is surrounded by the closed curve $g_1 \cup g_2 \cup g_3 \cup g_4$ exactly once. Hence we can define a univalent inverse function G_μ^{-1} in D_ε . By analytic continuation, we can define a univalent inverse function G_μ^{-1} in \mathbb{C}^- by letting $\varepsilon \downarrow 0$. When $\varepsilon > 0$ is small, the bounded domain surrounded by $c_1 \cup c_2 \cup c_3 \cup c_4$ has nonempty intersection with a truncated cone $\Gamma_{\lambda, M}$ where G_μ is univalent, and $G_\mu(\Gamma_{\lambda, M})$ has intersection with D_ε , and hence our G_μ^{-1} coincides with the original inverse on their common domain. Thus our G_μ^{-1} gives the desired analytic continuation, to conclude that $\mu \in \text{UI}$. The case $p = 1/2$ follows by approximation.

Next we take a discrete measure $\rho = \sum_{k=1}^n \lambda_k \delta_{t_k}$, $\lambda_k, t_k > 0$, $\sum_k \lambda_k = 1$. The multiplicative convolution $\rho \circledast \mu$ has the pdf

$$\begin{aligned} f_\rho(x) &:= \sum_{k=1}^n \lambda_k t_k^{-1} f(x/t_k) = \sum_{k=1}^n \lambda_k t_k^{-1} (x/t_k)^{p-1} g(x/t_k) \\ &= x^{p-1} g_\rho(x), \end{aligned} \quad (3.9)$$

where $g_\rho(x) := \sum_{k=1}^n \lambda_k t_k^{-p} g(x/t_k)$. The pdf f_ρ satisfies all the conditions (A1)–(A5) which follow from the conditions for f . Hence what we proved for f applies to f_ρ without a change, and hence $\rho \circledast \mu \in \text{UI}$. Since $\lim_{x \downarrow 0} g_\rho(x) > 0$ by (A3), the pdf f_ρ satisfies $\lim_{x \downarrow 0} f_\rho(x) = \infty$, so that $\rho \circledast \mu$ satisfies condition (ii) in Lemma 2.1, and so $\rho \circledast \mu \in \text{FR}$. Finally, by using the w-closedness of UI and FR we can approximate a general ρ by discrete measures to get the full result. \square

Remark 3.13. The curve g_1 is a Jordan curve (i.e. a curve with no self-intersection) since $G_\mu(x + i0)$ is decreasing on $(-\infty, 0)$. Moreover, if $\eta > 0$ is large and $\delta > 0$ is small then we can show that g_2, g_4 are also Jordan curves (with e.g. Rouché's theorem). However these properties are not needed to prove the theorem. Also, we do not know if g_3 has a self-intersection or not, but it does not matter for the proof. What we need is only that g_3 is contained in $\mathbb{C}^+ \cup \mathbb{R}$.

The result on powers of HCM rvs is a consequence of the above lemma.

Proof of Theorem 3.1: We further assume that X has a pdf of the form (2.3) and also $r > 1$. Then we define $s := 1/r$. The pdf of X^r is now given by

$$f(x) = sC \cdot x^{ps-1} \prod_{k=1}^n (t_k + x^s)^{-\gamma_k}, \quad (3.10)$$

where

$$0 < p \leq \frac{1}{2}, \quad 0 < -p + \sum_{k=1}^n \gamma_k \leq \frac{1}{2}. \quad (3.11)$$

Then f can be written in the form $x^{ps-1} g(x)$ where

$$g(x) = sC \prod_{k=1}^n (t_k + x^s)^{-\gamma_k}. \quad (3.12)$$

The pdf f is of the form (3.4) in Lemma 3.12 with p now replaced by $ps \in (0, 1/2)$, satisfying assumptions (A1)–(A4). Indeed, assumption (A2) holds since we can extend the function (3.12) analytically by extending x^s to the analytic function z^s

defined in a domain $\{re^{i\theta} : r > 0, \theta \in (-\pi - \delta, \delta)\}$ for some small $\delta > 0$ so that z^s never be a negative real in the domain. Such a $\delta > 0$ exists by our assumption $s \in (0, 1)$. Then (A3) follows immediately, and (A4) can be proved by taking $\gamma > 0$ such that $\gamma < 1 - ps + \sum_{k=1}^n \gamma_k s$. In order to check (A5) we compute

$$f(x - i0) = -sC|x|^{ps-1}e^{-i\pi ps} \prod_{k=1}^n (t_k + |x|^s e^{-i\pi s})^{-\gamma_k}, \quad x < 0. \quad (3.13)$$

Since $\arg(t_k + |x|^s e^{-i\pi s}) \in (-\pi s, 0)$, we have that $\arg(t_k + |x|^s e^{-i\pi s})^{-\gamma_k} \in (0, \pi\gamma_k s)$ and hence $\arg(-f(x - i0)) \in (-\pi ps, -\pi ps + \sum_{k=1}^n \pi\gamma_k s)$. Then (3.11) implies that $\arg(-f(x - i0)) \in (-\frac{\pi}{2}, \frac{\pi}{2})$ and hence assumption (A5) holds true. Thus $X^r \sim \text{FR} \cap \text{UI}$, and we can take the limit $s \uparrow 1$ to get the result for $s \in (0, 1]$.

The law of X^{-1} has the pdf

$$C \cdot x^{-p-1} \prod_{k=1}^n (t_k + x^{-1})^{-\gamma_k} = C' \cdot x^{p'-1} \prod_{k=1}^n (t_k^{-1} + x)^{-\gamma_k}, \quad (3.14)$$

where $C' = C \prod_{k=1}^n t_k^{-\gamma_k}$, $p' = -p + \sum_{k=1}^n \gamma_k$. Our assumption (3.11) guarantees that $p' \in (0, \frac{1}{2}]$ and $-p' + \sum_{k=1}^n \gamma_k = p \in (0, \frac{1}{2}]$. Therefore X^{-1} also satisfies (3.11) with p replaced by p' and so $X^{-r} \sim \text{FR} \cap \text{UI}$ for $r \geq 1$.

Finally we can approximate a pdf of the form (2.15) by pdfs of the form (2.3) in the sense of pointwise convergence. By Scheffé's lemma, we have the weak convergence of probability measures, and hence the full result follows. \square

We then go to the proof of Theorem 3.4 on powers of beta rvs. The idea is similar to the case of HCM rvs, but we need more elaboration since now we have to study the boundary behavior of the Cauchy transform on $(1, \infty) - i0$ in addition to $(-\infty, 0) - i0$.

Proof of Theorem 3.4: We may further assume that $r > 1, p + q - 1 < r, q \in [\frac{3}{2}, 2) \cup (2, \frac{5}{2}]$, $2p < r$ since the general case can be recovered by approximation. We define $s := 1/r \in (0, 1)$. The pdf of X^r is now given by

$$f(x) = \frac{s}{B(p, q)} \cdot x^{ps-1} (1 - x^s)^{q-1} = x^{ps-1} g(x), \quad (3.15)$$

where $g(x) = \frac{s}{B(p, q)} (1 - x^s)^{q-1}$.

Step 1: Analysis of G_{X^r} around $(-\infty, 0)$. We take the same curves c_1, c_2, c_3 depending on $\eta, \delta > 0$ as in Lemma 3.12 and the image curves $g_k = G_{X^r}(c_k)$, $k = 1, 2, 3$. The pdf f is of the form (3.4) with p replaced by $ps \in (0, 1/2)$ and satisfies assumptions (A1), (A3), (A4) in Lemma 3.12. Notice that (A4) holds thanks to $(p + q - 1)s < 1$. We change (A2) to the condition that f analytically continues to $\mathbb{C}^- \cup (0, 1) \cup \mathbb{C}^+$ and $f|_{\mathbb{C}^-}$ extends to a continuous function on $\mathbb{C}^- \cup (-\infty, 0)$. The analytic continuation is given by just replacing x with z in (3.15).

In order to show (A5) in Lemma 3.12, we compute

$$f(x - i0) = -\frac{s}{B(p, q)} \cdot |x|^{ps-1} e^{-i\pi ps} (1 - |x|^s e^{-i\pi s})^{q-1}, \quad x < 0. \quad (3.16)$$

Since $\arg(1 - |x|^s e^{-i\pi s}) \in (0, \pi - \pi s)$, we have that $\arg(-f(x - i0)) \in (-\pi ps, \pi(1 - p - q)s + \pi(q - 1))$. Our assumptions $0 < ps < \frac{1}{2}, q - \frac{3}{2} \leq s(p + q - 1)$ imply that $\arg(-f(x - i0)) \in (-\frac{\pi}{2}, \frac{\pi}{2})$ and hence assumption (A5) holds true. So the curve g_3 lies on $\mathbb{C}^+ \cup \mathbb{R}$.

The asymptotics (3.7) holds in the present case too (the constant a may change and p is replaced by ps), and hence for each $\varepsilon > 0$ there exists $\delta > 0$ small enough so that the distance between g_2 and 0 is larger than ε^{-1} . The maximal argument of the curve g_2 is in $(0, \pi)$ as discussed in the proof of Lemma 3.12.

The above arguments and the proof of Lemma 3.12 show that the curves $g_k, k = 1, 2, 3$ typically behave as in Fig. 3.2.




FIGURE 3.2. The curves c_k and g_k

Step 2: Analysis of G_{X^r} around $[1, \infty)$ and around ∞ . We show the following properties:

- (i) The restriction $f|_{\mathbb{C}^-}$ extends to a continuous function on $\mathbb{C}^- \cup \mathbb{R} \setminus \{0\}$;
- (ii) The analytic function $G := G_{X^r}$ (defined in $\mathbb{C}^+ \cup (0, 1) \cup \mathbb{C}^-$ via Lemma 3.11) extends to a continuous function on $\mathbb{C}^+ \cup (0, 1) \cup \mathbb{C}^- \cup ([1, \infty) + i0) \cup ([1, \infty) - i0)$;
- (iii) $\operatorname{Re}(f(x - i0)) \leq 0$ for $x > 1$.

Note that in (ii) we understand that $[1, \infty) + i0, [1, \infty) - i0$ are different half-lines by cutting the plane along $[1, \infty)$ and going to the Riemannian surface. This property implies that the map $(r, \theta) \mapsto G(re^{i\theta})$ defined in $(0, 1) \times (0, 2\pi)$ extends to a continuous map on $[0, 1) \times [0, 2\pi]$, and in particular that $G(1 + i0) = G(1 - i0)$.

Property (i) is easy to show.

Property (ii) follows from the second asymptotics in Hasebe (2014, (5.6)): there exists $b \in \mathbb{R}$ such that

$$G(z) = b + o(1), \quad z \rightarrow 1, z \in \mathbb{C} \setminus [1, \infty). \quad (3.17)$$

The number b is positive since it equals $\lim_{x \downarrow 1} G(x + i0) > 0$. Note that the proof of this asymptotics required $1 < q < 2$ (in Hasebe (2014) q is denoted by α), but we can give a proof for $2 < q < 3$ too only by using the identity $w^{\alpha-2} = (w-z)w^{\alpha-3} + zw^{\alpha-3}$ in Hasebe (2014, (5.13)). For $q = 2$ a logarithm term appears and so we avoid such a case for simplicity. The continuity on $(1, \infty) - i0$ follows from formula (3.3) and property (i).

Property (iii) follows from the computation

$$f(x - i0) = \frac{s}{B(p, q)} \cdot x^{ps-1} (x^s - 1)^{q-1} e^{i\pi(q-1)}, \quad x > 1 \quad (3.18)$$

and our assumption $q \in [\frac{3}{2}, \frac{5}{2}]$.

We moreover define c_4, \dots, c_7 and the corresponding images $g_k = G(c_k)$ for $k = 4, \dots, 7$:

- c_4 is a counterclockwise curve which lies on $\{z \in \mathbb{C}^- \mid |z| \geq \eta\}$, starting from $-\eta - i0$ (the final point of c_3) and ends at $1 + \eta - i0$;
- c_5 is the line segment from $1 + \eta - i0$ to $1 - i0$;
- c_6 is the line segment from $1 + i0$ to $1 + \eta + i0$;
- c_7 is a counterclockwise curve which lies on $\{z \in \mathbb{C}^+ \mid |z| \geq \eta\}$, starting from $1 + \eta + i0$ and ending at $-\eta + i0$.

Thanks to property (ii), $g_5 \cup g_6$ is a continuous curve, so one need not take a small circle to avoid the point 1.

We can take a large $\eta > 0$ similarly to Lemma 3.12 so that the curves g_4, g_7 lie on the ball $\{z \in \mathbb{C} \mid |z| < \varepsilon\}$ as shown in the right figure. This is easy to prove for g_7 since the measure is compactly supported and so $G(z) = O(1/z)$ ($z \rightarrow \infty, z \in \mathbb{C}^+$). For g_4 we need Lemma 3.11 and our assumption $s(p + q - 1) < 1$. Thanks to property (iii), we have that

$$\operatorname{Im}(G(z)) = \operatorname{Im}(\tilde{G}(z)) - 2\pi \operatorname{Im}(if(z)) \geq 0 \quad (3.19)$$

on c_5 , so the curve g_5 is on $\mathbb{C}^+ \cup \mathbb{R}$.

From the above arguments, every point of $D_\varepsilon = \{w \in \mathbb{C}^- \mid \varepsilon < |w| < \varepsilon^{-1}\}$ is surrounded by the closed curve $g_1 \cup \dots \cup g_7$ exactly once. So we can define a univalent inverse G^{-1} in \mathbb{C}^- as discussed in Lemma 3.12, to conclude that $X^r \sim \text{UI}$. Approximation shows that the result is true for $r = 1$ and $q = 2$ and for the case $p + q - 1 = r$ too. From Lemma 2.1 the law of X^r is free regular. \square

Remark 3.14. The curves g_1 and g_6 are Jordan curves. If $\eta > 0$ is large and $\delta > 0$ is small then we can moreover show that g_2, g_4, g_7 are also Jordan curves (with e.g. Rouché's theorem). However these properties are not needed to prove the theorem.

We will prove the results for powers of free Poisson rvs. The idea is again similar to the previous proofs. A new phenomenon is that the Cauchy transform has a singularity when we take the limit $z \rightarrow 0, z \in \mathbb{C}^-$, and we will see how the previous methods are modified.

Proof of Theorem 3.5: Since approximation is allowed, we only consider parameters (p, r) from an open dense subset of the full set.

Case (a): $p > 1, r > 1$.

(a-UI) We will first show that $X^r \sim \text{UI}$ when $X \sim \mathbf{fp}(p, 1)$. For simplicity we define $G(z) = G_{X^r}(z)$, $s = 1/r$, $a = |\sqrt{p} - 1|^{2/s} > 0$ and $b = (\sqrt{p} + 1)^{2/s} > a$. Then the pdf of X^r is equal to

$$f(x) = \frac{s}{2\pi} \cdot \frac{\sqrt{(b^s - x^s)(x^s - a^s)}}{x} 1_{(a,b)}(x). \quad (3.20)$$

This density extends to an analytic function in $\mathbb{C}^+ \cup (a, b) \cup \mathbb{C}^-$ by replacing x in (3.20) with a complex variable z . The function z^s is the principal value. One can take the square root also as the principal value, but it may not be obvious. To justify this claim, we use the identity

$$(b^s - z^s)(z^s - a^s) = -\left(z^s - \frac{a^s + b^s}{2}\right)^2 + \left(\frac{b^s - a^s}{2}\right)^2. \quad (3.21)$$

If $z \in \mathbb{C}^\pm$ then $z^s - \frac{a^s+b^s}{2} \in \mathbb{C}^\pm$, and hence $-(z^s - \frac{a^s+b^s}{2})^2 \in \mathbb{C} \setminus (-\infty, 0]$, and hence by adding the positive real number $(\frac{b^s-a^s}{2})^2$, we conclude that $(b^s - z^s)(z^s - a^s) \in \mathbb{C} \setminus (-\infty, 0]$. So the principal value is relevant for defining the square root.

The Cauchy transform G extends to $\mathbb{C}^+ \cup (a, b) \cup \mathbb{C}^-$ analytically via Lemma 3.11. The function $f|_{\mathbb{C}^-}$ extends to a continuous function on $\mathbb{C}^- \cup \mathbb{R} \setminus \{0\}$ since $(b^s - z^s)(z^s - a^s)$ continuously extends to $\mathbb{C}^- \cup \mathbb{R}$ without taking the value 0 except at a, b .

Curves $c_k, g_k = G(c_k)$ that we use in the proof are shown in Fig. 3.3.




FIGURE 3.3. The curves c_k and g_k ($p > 1, r > 1$)

We compute the boundary value $f(x - i0)$ for $x < a, x > b$, which is the most crucial part of the proof. For $0 < x < a$, since $(x - i\delta)^s - a^s$ approaches the point $-(a^s - x^s) < 0$ from \mathbb{C}^- as $\delta \downarrow 0$, we should understand that $(x - i0)^s - a^s = (a^s - x^s)e^{-i\pi+i0}$. Hence

$$f(x - i0) = \frac{s}{2\pi} \cdot \frac{e^{-\frac{i\pi}{2}} \sqrt{(b^s - x^s)(a^s - x^s)}}{x}, \quad 0 < x < a, \quad (3.22)$$

and hence $\operatorname{Re}(f(x - i0)) = 0$. So the curve g_2 is on the (negative) real line.

For $x > b$, since $b^s - (x - i\delta)^s$ approaches a point in $(-\infty, 0)$ from \mathbb{C}^+ , we have to understand that $b^s - (x - i0)^s = (x^s - b^s)e^{i\pi-i0}$. Hence

$$f(x - i0) = \frac{s}{2\pi} \cdot \frac{e^{\frac{i\pi}{2}} \sqrt{(x^s - b^s)(x^s - a^s)}}{x}, \quad x > b, \quad (3.23)$$

and hence $\operatorname{Re}(f(x - i0)) = 0$. So the curve g_6 is on the (positive) real line.

As we saw in (3.21), $\arg((b^s - z^s)(z^s - a^s)) \in (-\pi, \pi)$, and hence $\sqrt{(b^s - z^s)(z^s - a^s)} \in \{w \mid \operatorname{Re}(w) > 0\}$. Dividing it by z and taking the limit $z \rightarrow x - i0 = |x|e^{-i\pi}$, we have

$$\operatorname{Re}(f(x - i0)) \leq 0, \quad x < 0 \quad (3.24)$$

and so g_4 lies on $\mathbb{C}^+ \cup \mathbb{R}$. The remaining proof is similar to Theorems 3.1, 3.4, so we only mention important remarks. The proof of Hasebe (2014, Theorem 5.1(5.6)) for $\alpha = \frac{3}{2}, x_0 = a$ and $x_0 = b$ (with reflection) enables us to show that the Cauchy transform G has a continuous extension to

$$\mathbb{C}^+ \cup [a, b] \cup \mathbb{C}^- \cup ((-\infty, a] \cup [b, \infty) + i0) \cup ((-\infty, a] \cup [b, \infty) - i0) \setminus \{0 - i0\}, \quad (3.25)$$

similarly to property (ii) in the proof of Theorem 3.4. This implies that $g_1 \cup g_2$ is a continuous curve and so is $g_6 \cup g_7$. The Cauchy transform has a singularity

at 0 when approaching from \mathbb{C}^- . This singularity is a contribution of x^{-1} of (the analytic continuation of) the pdf f . So we avoid 0 via the curve c_3 . When one draws the picture of g_3 , one should take it into account that $G(z) = -2\pi i f(z) + o(|z|^{-1}) = -\frac{s(p-1)}{z}(1 + o(1))$, $z \rightarrow 0, z \in \mathbb{C}^-$ since $\tilde{G}(z)$ is analytic around 0 and $f(z) = \frac{s}{2\pi} \cdot \frac{-i\sqrt{a^s b^s}}{z}(1 + o(1))$. So the curve g_3 is as shown in Fig. 3.3. Thus one can show that $X^r \sim \text{UI}$.

(a-FR) Since we cannot apply Lemma 2.1, we will directly prove that $X^r \sim \text{FR}$. Let us take the curves $c_1 = (t + i0)_{t \in (-\infty, a]}$ and $c_2 = (a - t - i0)_{t \in [0, a]}$. We can see that for any $t \in (-\infty, a)$ there exists $\delta_1 > 0$ such that $G(t + iy) \in \mathbb{C}^-$ for all $y \in (0, \delta_1)$ (since G maps \mathbb{C}^+ to \mathbb{C}^-) and also for any $t \in (0, a)$ there exists $\delta_2 > 0$ such that $G(a - t - iy) \in \mathbb{C}^-$ for $y \in (0, \delta_2)$ thanks to (3.22). Note also that $G(c_1 + i0) = [-\alpha, 0)$ and $G(c_2 - i0) = (-\infty, -\alpha]$ for $\alpha := \tilde{G}(a - 0) > 0$. These facts imply that our inverse G^{-1} in \mathbb{C}^- satisfies that

$$G^{-1}((-\infty, -\alpha] - i0) = c_2, \quad G^{-1}([-\alpha, 0) - i0) = c_1. \quad (3.26)$$

It also holds that $G^{-1}(-\infty - i0) = 0$ since $G(z)$ tends to ∞ as $z \rightarrow 0, z \in \mathbb{C}^-$. Let $\varphi(z) := G^{-1}(z^{-1}) - z, z \in \mathbb{C}^+$ be the Voiculescu transform. Since $\text{Im}(\varphi(x + i0)) = 0, x < 0$ by (3.26), the free Lévy measure of X^r has no mass on $(-\infty, 0)$ by Stieltjes inversion. Since $\varphi(-0) = G^{-1}(-\infty) = 0$, Arizmendi et al. (2013, Proposition 11) implies that $X^r \sim \text{FR}$.

Case (b): $p < 1, r > 1$. We follow the notations in case (a). Since now the distribution of X^r has an atom at 0 and hence we can apply Lemma 2.1, it suffices to show $X^r \sim \text{UI}$. The probability distribution of X^r is of the form $(1-p)\delta_0 + f(x)dx$, where f is given by the same formula (3.20). The functions f and G have analytic continuation as discussed in case (a).

We will take curves c_k as shown in Fig. 3.4. Now the difference from case (a) is




FIGURE 3.4. The curves c_k and g_k ($p < 1, r > 1$)

that \tilde{G} has a pole at 0, and so we avoid $0 \pm i0$ via the curves c_2, c_5 . Note that $G(z) = \frac{1-p}{z}(1 + o(1))$ as $z \rightarrow 0, z \in \mathbb{C}^+$ and $G(z) = \tilde{G}(z) - 2\pi i f(z) = \frac{(1-p)(1-s)}{z}(1 + o(1))$ as $z \rightarrow 0, z \in \mathbb{C}^-$, so the curves g_2, g_5 look like large semicircles. Thus we can prove the claim $X^r \sim \text{UI}$.

Case (c): $p > 1, r < -1$. For simplicity we use the same notation $G(z) = G_{X^r}(z)$, but now we define $t = -1/r \in (0, 1)$, $A = (\sqrt{p} + 1)^{-2/t} > 0$ and $B = (\sqrt{p} - 1)^{-2/t} >$

A. Then the pdf of X^r is equal to

$$h(x) = \frac{t(p-1)}{2\pi} \cdot \frac{\sqrt{(B^t - x^t)(x^t - A^t)}}{x^{t+1}} 1_{(A,B)}(x). \quad (3.27)$$

The functions h, G continue analytically to $\mathbb{C}^+ \cup (A, B) \cup \mathbb{C}^-$ and $h|_{\mathbb{C}^-}$ extends to a continuous function on $\mathbb{C}^- \cup \mathbb{R} \setminus \{0\}$ as we discussed in case (a).

We will take the same curves c_k as in case (a) except that the points a, b are replaced by A, B respectively (see Fig. 3.5).




FIGURE 3.5. The curves c_k and g_k ($p > 1, r < -1$)

In the present case, the difficulty is to show the inequality corresponding to (3.24), i.e. $\operatorname{Re}(h(x - i0)) \leq 0$ for $x < 0$. The inequality $\operatorname{Re}(h(x - i0)) \leq 0$ (actually equality holds) for $0 < x < a$ and $x > b$ is easy because only the new factor $(p-1)x^{-t} > 0$ is different from (3.22), (3.23).

We show that $\operatorname{Re}(h(x - i0)) \leq 0$ for $x < 0$ by considering two further sub cases.

Case (ci): $0 < t \leq \frac{1}{2}$. We will use the identity (3.21) with (a, b) replaced by (A, B) . First we can easily show that $\arg_{(-2\pi, 0)}((x - i0)^t - (A^t + B^t)/2)^2 \in (-2\pi, -2\pi t)$ for $x < 0$. Then we multiply it by -1 and shift it by $((B^t - A^t)/2)^2$ and so $\arg((B^t - (x - i0)^t)((x - i0)^t - A^t)) \in (-\pi, (1 - 2t)\pi)$. Then we take the square root to get $\arg(\sqrt{(B^t - (x - i0)^t)((x - i0)^t - A^t)}) \in (-\frac{\pi}{2}, (\frac{1}{2} - t)\pi)$. Further we divide it by $(x - i0)^{t+1} = |x|^{t+1}e^{-i\pi(t+1)}$, to get $\operatorname{Re}(h(x - i0)) \leq 0$ for $x < 0$.

Case (cii): $\frac{1}{2} < t < 1$. We can show that the curve $\gamma : ((x - i0)^t - (A^t + B^t)/2)^2, x < 0$ is on the right side of the half-line $L : ((A^t + B^t)/2)^2 + re^{i(2\pi - 2\pi t)}, r > 0$, and then $\arg((B^t - (x - i0)^t)((x - i0)^t - A^t)) \in (-\pi, (1 - 2t)\pi)$ (see Remark 3.15 for more details). The remaining arguments are the same as case (ci).

Thus we have now $\operatorname{Re}(h(x - i0)) \leq 0$ for $x < 0$ and hence g_4 lies on $\mathbb{C}^+ \cup \mathbb{R}$. The remaining proof is almost the same as in case (a), but there are two remarks. One is that as $z \rightarrow 0, z \in \mathbb{C}^-$, the Cauchy transform $G(z) = \tilde{G}(z) - 2\pi i h(z)$ has asymptotics $G(z) = -\frac{t}{z^{t+1}}(1 + o(1))$ which is different from case (a) by the factor z^{-t} and a constant. Therefore the arguments of g_3 change from around $-\pi$ to $t\pi$ as in Fig. 3.5. The other remark is that as $z \rightarrow \infty, z \in \mathbb{C}^-$, the asymptotics of the Cauchy transform is $G(z) = \tilde{G}(z) - 2\pi i h(z) = (1 - t(p-1)) \cdot z^{-1}(1 + o(1))$, so g_5 is

near 0. Thus we are able to show that $X^r \sim \text{UI}$. The proof of $X^r \sim \text{FR}$ is similar to case (a).

Case (d): $p > 1, r \in (-1, 0)$. The pdf is the same as (3.27). We follow the notations in case (c), and so now we have $t > 1$. The function h now analytically continues to $\{z \in \mathbb{C} \setminus \{0\} \mid \arg(z) \in (-\pi/t, 0) \cup (0, \pi/t)\} \cup (A, B)$ and so G extends to $\mathbb{C}^+ \cup (A, B) \cup \{z \in \mathbb{C} \setminus \{0\} \mid \arg(z) \in (-\pi/t, 0)\}$. Moreover h extends to a continuous function on $\{z \in \mathbb{C} \setminus \{0\} \mid \arg(z) \in [-\pi/t, 0]\}$. The curve c_4 in case (c) is not useful now, and instead we take c_4 to be a long line segment with angle $-\pi/t$. So we take curves as shown in Fig. 3.6.




FIGURE 3.6. The curves c_k and g_k ($p > 1, -1 < r < 0$)

We compute the boundary value on c_4 :

$$\begin{aligned} \lim_{\delta \downarrow 0} h(re^{-i\pi/t} + \delta) &= \frac{t(p-1)}{2\pi} \cdot \frac{e^{-i\pi/2} \sqrt{(B^t + r^t)(A^t + r^t)}}{r^{t+1} e^{-i\pi(t+1)/t}} \\ &= \frac{t(p-1) \sqrt{(B^t + r^t)(A^t + r^t)}}{2\pi r^{t+1}} \cdot i e^{i\pi/t}, \quad r > 0, \end{aligned} \quad (3.28)$$

and hence $\text{Re}(h(re^{-i\pi/t} + 0)) < 0$ as desired. This implies that g_4 lies on $\mathbb{C}^+ \cup \mathbb{R}$. The remaining proof is similar to case (c) and the typical behavior of the curves g_k is also similar to case (c). Thus we can show that $X^r \sim \text{UI}$. The proof of $X^r \sim \text{FR}$ is similar to case (a). \square

Remark 3.15 (More details on case (cii)). For simplicity let d denote $(A^t + B^t)/2$. We first show that the curve $\gamma(r) = l(r)^2 = (re^{-it\pi} - d)^2, r > 0$ is on the right side of the half line $L(r) = d^2 + re^{i(2\pi - 2t\pi)}, r > 0$ (see Figs 3.7, 3.8). Our claim is equivalent to $\arg(\gamma(r) - d^2) \in (0, 2\pi - 2t\pi), r > 0$. From Fig. 3.7 and the current assumption $1/2 < t < 1$, we can see that $\arg(\gamma(r) - d^2) \in (0, \pi)$. A simple computation shows that $\gamma(r) - d^2 = (re^{-it\pi} - d)^2 - d^2 = r^2 \cos(2t\pi) - 2dr \cos(t\pi) - 2ir \sin(t\pi)(r \cos(t\pi) - d)$. After some more computations, one has

$$\text{Im}((\gamma(r) - d^2)e^{-i(2\pi - 2t\pi)}) = -2dr \sin(t\pi) < 0. \quad (3.29)$$

Therefore $\arg(\gamma(r) - d^2) < 2\pi - 2t\pi$.

Next, since $d^2 = (\frac{A^t+B^t}{2})^2 > (\frac{B^t-A^t}{2})^2$, the arguments of the curve $\gamma - (\frac{B^t-A^t}{2})^2$ still lie in $(0, 2\pi - 2t\pi)$. Hence the arguments of the curve $-\gamma + (\frac{B^t-A^t}{2})^2$ lie in $(-\pi, (1-2t)\pi)$.




FIGURE 3.7.

$$l : (x-i0)^t - \frac{A^t+B^t}{2}, \\ x < 0$$




FIGURE 3.8.

$$\gamma : ((x-i0)^t - \frac{A^t+B^t}{2})^2, \\ x < 0$$

Finally we prove Theorem 3.8, i.e. the FID properties of symmetric distributions. The negative result in Theorem 3.8(3) on the symmetrized powers of beta rvs entails the following fact.

Lemma 3.16. *Suppose that a probability measure μ on \mathbb{R} is symmetric and $\mu|_{(-\varepsilon, \varepsilon)} = p(x)1_{(-\varepsilon, \varepsilon)}(x) dx$ for some $\varepsilon > 0$ and some continuous function p on $(-\varepsilon, \varepsilon)$. If $p(0) = 0$, then $\mu \notin \text{FID}$.*

Proof: Suppose that $\mu \in \text{FID}$. The Cauchy transform of μ vanishes at 0 since $\text{Re}(G_\mu(iy)) = 0$ and $\lim_{y \downarrow 0} G_\mu(iy) = -i\pi p(0) = 0$. This contradicts the fact that the reciprocal Cauchy transform $1/G_\mu(z)$ extends to a continuous function on $\mathbb{C}^+ \cup \mathbb{R}$ into itself (see Belinschi, 2005, Proposition 2.8). \square

Proof of Theorem 3.8: The negative result (3) holds since if $p > r > 0$ or $r \leq 0$ then the law of BX^r satisfies the assumptions of Lemma 3.16.


We will prove the case (2) and later give some comments on how to prove the HCM case (1). The idea is similar to the proof of Theorem 3.4, so we only mention significant changes of the proof. We follow the notations in the proof of Theorem 3.4. First we will check condition (b) in Definition 2.4. The pdf of BX^r is given by

$$w(x) = \frac{s}{2B(p, q)} \cdot |x|^{ps-1} (1 - |x|^s)^{q-1} 1_{(-1, 1) \setminus \{0\}}(x). \quad (3.30)$$

The restriction $w|_{(0, 1)}$ has the analytic continuation $w(z) = \frac{1}{2}f(z)$ in $\mathbb{C}^+ \cup (0, 1) \cup \mathbb{C}^-$ where $f(x)$ is the pdf of X^r . Note then that $w(x+i0) \neq w(x)$ for $x \in (-1, 0)$. Also $G_{BX^r}(z) = \frac{1}{2}(G_{X^r}(z) - G_{X^r}(-z))$ has analytic continuation to $\mathbb{C}^+ \cup (0, 1) \cup \mathbb{C}^-$ and

$$\begin{aligned} G_{BX^r}(z) &= \frac{1}{2} \left(\tilde{G}_{X^r}(z) - 2\pi i w(z) \right) - \frac{1}{2} \tilde{G}_{X^r}(-z) \\ &= \tilde{G}_{BX^r}(z) - \frac{1}{2} \pi i f(z), \quad z \in \mathbb{C}^-. \end{aligned} \quad (3.31)$$

We take curves c_k in Fig. 3.9 and consider the image curves $g_k = G_{BX^r}(c_k)$.

FIGURE 3.9. The curves c_k, g_k

We have proved $\operatorname{Re}(f(x - i0)) \leq 0$ for $x < 0$ and $x > 1$ in the proof of Theorem 3.4, so the curves g_3, g_5 are contained in $\mathbb{C}^+ \cup \mathbb{R}$.

The function G_{BX^r} extends continuously to

$$\mathbb{C}^+ \cup (0, 1) \cup \mathbb{C}^- \cup ([1, \infty) + i0) \cup ([1, \infty) - i0) \quad (3.32)$$

as we proved in Theorem 3.4, so $g_5 \cup g_6$ is a continuous curve.

We may use the asymptotics (3.7) (we use the same notation a for the constant) and obtain

$$\begin{aligned} G_{BX^r}(z) &= \frac{1}{2}(G_{X^r}(z) - G_{X^r}(-z)) \\ &= \frac{a}{2}(-(-z)^{ps-1} + z^{ps-1}) + o(|z|^{ps-1}) \\ &= a \cos\left(\frac{\pi ps}{2}\right) e^{-i\frac{\pi ps}{2}} z^{ps-1} + o(|z|^{ps-1}) \end{aligned} \quad (3.33)$$

as $z \rightarrow 0, \arg(z) \in (-\pi, \pi)$. So when z goes along a small circle centered at 0 from $\arg(z) = \pi/2$ to $\arg(z) = -\pi$, the Cauchy transform $G_{BX^r}(z)$ goes along a large circle-like curve from the angle $-\pi/2$ to $(1 - \frac{3}{2}ps)\pi$ (> 0 since $ps \leq 1/2$) with small errors. This suggests that the curve g_2 seems as in Fig. 3.9.

The asymptotics (3.33) also implies that $\frac{1}{i}G_{BX^r}(iy) \rightarrow -\infty$ as $y \downarrow 0$. It is elementary to show that $G_{BX^r}(z) \rightarrow 0$ as $|z| \rightarrow \infty, z \in \mathbb{C}^+$. Therefore the curves g_1, g_7 are as shown in Fig. 3.9.

Thus each point of $\mathbb{C}^- \cap i\mathbb{C}^-$ is surrounded by $g_1 \cup \dots \cup g_7$ exactly once when the curve $c_1 \cup \dots \cup c_7$ is sufficiently close to the boundary of $\mathbb{C}^- \cup i\mathbb{C}^- \setminus [1, \infty)$. Therefore $G_{BX^r}^{-1}$ has univalent analytic continuation to $\mathbb{C}^- \cap i\mathbb{C}^-$ such that $G_{BX^r}^{-1}(\mathbb{C}^- \cap i\mathbb{C}^-) \subset \mathbb{C}^- \cup i\mathbb{C}^-$. This is condition (b) in Definition 2.4.

We will check condition (a) in Definition 2.4. By calculus we can show that $w'(x) < 0$ for $x \in (0, 1)$ and hence the distribution of BX^r is unimodal. The map G_{BX^r} is univalent in \mathbb{C}^+ thanks to Kaplan (1952, Theorem 3) (see Avkhadiiev and Aksen'tev, 1975, Theorem 39, for a different proof). Recall that $\frac{1}{i}G_{BX^r}(iy) \rightarrow -\infty$ as $y \downarrow 0$ and that $G_{BX^r}(iy) \rightarrow 0$ as $y \rightarrow \infty$. Therefore, G_{BX^r} maps a neighborhood

of $i(0, \infty)$ onto a neighborhood of $i(-\infty, 0)$ bijectively. Thus we have condition (a) and therefore $BX^r \sim \text{UI}_s$.

For the HCM case, we may assume that W takes only finitely many values. Then the pdf of X is of the form (2.3) and $r \geq 1$. By calculus we can show that the pdf of WX^r is unimodal. Then the proof is similar to the case of beta rvs. \square

According to Mathematica, the Cauchy transform of X^r has an expression in terms of generalized hypergeometric functions when $X \sim \beta(p, q)$ and at least $r = 3/2, 4$. Figs 3.2–3.2 show the numerical computation of the domain $G_{BX^r}^{-1}(\mathbb{C}^- \cap i\mathbb{C}^-)$ which equals a connected component of $\{z \in \mathbb{C}^- \cup i\mathbb{C}^- \setminus [1, \infty) \mid G_{BX^r}(z) \in \mathbb{C}^-\}$. These figures suggest that the law of BX^r belongs to UI since the domain seems to be contained in the right half-plane, but there is no rigorous proof.




FIGURE 3.10. The domain $G_{BX^{3/2}}^{-1}(\mathbb{C}^- \cap i\mathbb{C}^-)$ when $X \sim \beta(1/2, 1.501)$




FIGURE 3.11. The domain $G_{BX^{3/2}}^{-1}(\mathbb{C}^- \cap i\mathbb{C}^-)$ when $X \sim \beta(1/2, 2)$




FIGURE 3.12. The domain $G_{BX^4}^{-1}(\mathbb{C}^- \cap i\mathbb{C}^-)$ when $X \sim \beta(1/3, 1.8)$




FIGURE 3.13. The domain $G_{BX^4}^{-1}(\mathbb{C}^- \cap i\mathbb{C}^-)$ when $X \sim \beta(1/3, 2.4)$

4. Analogy between classical and free probabilities

The free analogue of the normal distribution is the semicircle distribution, and nothing else has been proposed so far. However, three different kinds of “free gamma distributions” have been proposed in the literature: [Anshelevich \(2003, p. 238\)](#) defined a free gamma distribution in terms of orthogonal polynomials; [Pérez-Abreu and Sakuma \(2008\)](#) defined a free gamma distribution in terms of the Bercovici-Pata bijection, whose property was investigated in details by [Haagerup and Thorbjørnsen \(2014\)](#). The third definition of a free gamma distribution is just the free Poisson distribution. This comes from an analogy between a characterization of gamma distributions by Lukacs and that of free Poisson distributions by Szpojankowski. [Lukacs \(1955\)](#) proved that non-degenerate and independent rvs $X, Y > 0$ have gamma distributions with the same scale parameter if and only if $\frac{X}{X+Y}, X+Y$ are independent. The free probabilistic analogue gives us free Poisson distributions.

Namely, in [Szpojankowski \(2015b, Theorem 4.2\)](#) and [Szpojankowski \(2015a, Theorem 3.1\)](#) it was proved that bounded non-degenerate free rvs $X, Y \geq 0$ such that $X \geq \varepsilon I$ for some $\varepsilon > 0$ have free Poisson distributions with the same scale parameter if and only if $(X + Y)^{-1/2} X (X + Y)^{-1/2}, X + Y$ are free. Note that the assumption $X \geq \varepsilon I$ implies that $p > 1$ when $X \sim \mathbf{fp}(p, \theta)$.

From the point of view of the third definition of a free gamma distribution as the free Poisson distribution, our Theorem 3.5 on powers of free Poisson rvs has a good correspondence with the classical case:

Theorem 4.1. *Suppose that Y follows a gamma distribution. Then $Y^r \sim \text{GGC}$ for $r \in (-\infty, 0] \cup [1, \infty)$. If $r \in (0, 1)$ then $Y^r \notin \text{ID}$.*

The result for $|r| \geq 1$ was first proved by [Thorin \(1978\)](#). It also follows from some facts on the class HCM: all gamma distributions are contained in HCM; $\text{HCM} \subset \text{GGC}$; $X \sim \text{HCM}$ implies $X^r \sim \text{HCM}$ for $|r| \geq 1$ [Bondesson \(1992\)](#). The result for $r \in (-1, 0)$ was recently proved by [Bosch and Simon \(2015\)](#). If $r \in (0, 1)$ then Y^r is not ID since the tail decreases in a more rapid way than what is allowed for non-normal ID distributions (see [Ruegg, 1970](#) or [Steutel and van Harn, 2004](#), Chap. 4., Theorem 9.8 for the rigorous statement).

Therefore we have a complete correspondence between the ID property of powers of gamma rvs and the FID property of free Poisson rvs, except $r \in (0, 1)$. We pose a conjecture on this missing interval.

Conjecture 4.2. If $X \sim \mathbf{fp}(p, 1)$, $r \in (0, 1)$ and $p > 0$ then $X^r \not\sim \text{FID}$.

A partial result can be obtained from [Hasebe \(2014, Theorem 5.1\)](#): $X^r \not\sim \text{FID}$ if $X \sim \mathbf{fp}(1, 1)$ and $r \in (\frac{1}{2} - \frac{1}{4n+2}, \frac{1}{2} - \frac{1}{4n+4}) \cup (\frac{1}{2} + \frac{1}{4n}, \frac{1}{2} + \frac{1}{4n-2})$, $n \in \mathbb{N}$. Another way to check the free infinite divisibility is the Hankel determinants of free cumulants. Haagerup and Möller computed the Mellin transform of $X \sim \mathbf{fp}(1, 1)$ in [Haagerup and Möller \(2013, Theorem 3\)](#) and the moments of X^r are given by

$$\int_0^4 x^{rn} \mathbf{fp}(1, 1)(dx) = \frac{\Gamma(1 + 2rn)}{\Gamma(1 + rn)\Gamma(2 + rn)}, \quad n \in \mathbb{N} \cup \{0\}, \quad (4.1)$$

so we can compute free cumulants from these moments. According to Mathematica, the 2nd Hankel determinant $K_4 K_2 - K_3^2$ of the free cumulants $\{K_n\}_{n \geq 2}$ is negative for $0.35 \lesssim r < 1$ and K_6 is negative for $0.335 \lesssim r \lesssim 0.42$ (see Figs. 4.14, 4.15). So it seems that $X^r \not\sim \text{FID}$ for $0.335 \lesssim r < 1$, but this is not a mathematical proof.

Our Corollary 3.6 and Corollary 3.9 on powers of semicircular elements also have a good correspondence with the classical case:

Theorem 4.3. *Suppose $Z \sim N(0, 1)$.*

- (1) *If $r \in (-\infty, 0] \cup [2, \infty)$ then $|Z|^r \sim \text{GGC}$.*
- (2) *If $r \in (0, 2)$ then $|Z|^r \not\sim \text{ID}$.*
- (3) *If $r \geq 2$ then $|Z|^r \text{sign}(Z) \sim \text{ID}$.*
- (4) *If $1 \neq r < 2$ then $|Z|^r \text{sign}(Z) \notin \text{ID}$.*

Proof: (1), (2) are a special case of Theorem 4.1 since $Z \sim N(0, 1)$ implies that Z^2 follows the gamma distribution $(2\pi x)^{-1/2} e^{-x/2} 1_{(0, \infty)}(x) dx$.

(3) follows from [Shanbhag and Sreehari \(1977, Theorem 5\)](#).

(4) When $r < 1$, $r \neq 0$ then the pdf of $|Z|^r \text{sign}(Z)$ is symmetric and vanishes at 0. Such a distribution is not ID by [Rohatgi et al. \(1990, p. 58\)](#). When $r = 0$ the distribution is a symmetric Bernoulli and so not ID.




FIGURE 4.14. $K_4 K_2 - K_3^2$ as a function of r .




FIGURE 4.15. K_6 as a function of r .




FIGURE 4.16. $K_2' K_6' - (K_4')^2$ as a function of r .




FIGURE 4.17. $\det((K'_{2i+2j-2})^4_{i,j=1})$ as a function of r .

When $1 < r < 2$, one may use [Ruegg \(1970, Theorem 2\)](#) or [Steutel and van Harn \(2004, Chap. 4, Theorem 9.8\)](#). \square

The following problems remain to be solved in order to get the complete analogy between classical and free cases.

Conjecture 4.4. Suppose that $S \sim \mathcal{S}(0, 1)$.

- (1) $|S|^r \not\sim \text{FID}$ for $r \in (0, 2)$.
- (2) $|S|^r \text{sign}(S) \not\sim \text{FID}$ for $r \in (1, 2)$.

Note that (1) is a special case of [Conjecture 4.2](#). The odd moments of $|S|^r \text{sign}(S)$ are 0 and the even moments are the same as those of $|S|^r$. So we can compute free cumulants $\{K'_n\}_{n \geq 1}$ of $|S|^r \text{sign}(S)$ from [\(4.1\)](#), and computation in Mathematica suggests that $K'_2 K'_6 - (K'_4)^2 < 0$ for $1 \lesssim r \lesssim 1.8$ and $\det((K'_{2i+2j-2})^4_{i,j=1}) < 0$ for $1.68 \lesssim r \lesssim 1.94$ (in Figs. [4.16](#), [4.17](#)). So the conjecture (2) is likely to hold.

Acknowledgements

This work is supported by JSPS Grant-in-Aid for Young Scientists (B) 15K17549. The author thanks Octavio Arizmendi and Noriyoshi Sakuma for the past collaborations and discussions which motivated this paper. The author thanks Franz Lehner and Marek Bożejko for discussions on powers of semicircular elements and Víctor Pérez-Abreu for information on references. Finally the author thanks the hospitality of organizers of International Workshop “Algebraic and Analytic Aspects of Quantum Lévy Processes” held in Greifswald in March, 2015. This work is based on the author’s talk in the workshop.

References

- M. Anshelevich. Free martingale polynomials. *J. Funct. Anal.* **201** (1), 228–261 (2003). [MR1986160](#).
- M. Anshelevich, S. T. Belinschi, M. Bożejko and F. Lehner. Free infinite divisibility for q -Gaussians. *Math. Res. Lett.* **17** (5), 905–916 (2010). [MR2727617](#).

- O. Arizmendi, O. E. Barndorff-Nielsen and V. Pérez-Abreu. On free and classical type G distributions. *Braz. J. Probab. Stat.* **24** (2), 106–127 (2010). [MR2643561](#).
- O. Arizmendi and S. T. Belinschi. Free infinite divisibility for ultrasphericals. *Inf. Dimens. Anal. Quantum Probab. Relat. Top.* **16** (1), 1350001, 11 (2013). [MR3071453](#).
- O. Arizmendi and T. Hasebe. On a class of explicit Cauchy-Stieltjes transforms related to monotone stable and free Poisson laws. *Bernoulli* **19** (5B), 2750–2767 (2013). [MR3160570](#).
- O. Arizmendi and T. Hasebe. Classical and free infinite divisibility for Boolean stable laws. *Proc. Amer. Math. Soc.* **142** (5), 1621–1632 (2014). [MR3168468](#).
- O. Arizmendi and T. Hasebe. Classical scale mixtures of Boolean stable laws. *Trans. Amer. Math. Soc.* **368** (7), 4873–4905 (2016). [MR3456164](#).
- O. Arizmendi, T. Hasebe and N. Sakuma. On the law of free subordinators. *ALEA Lat. Am. J. Probab. Math. Stat.* **10** (1), 271–291 (2013). [MR3083927](#).
- F. G. Avkhadiyev and L. A. Aksent'ev. The main results on sufficient conditions for an analytic function to be schlicht. *Russian Mathematical Surveys* **30** (4), 1–64 (1975). [DOI: 10.1070/RM1975v030n04ABEH001511](#).
- O. E. Barndorff-Nielsen, U. Franz, R. Gohm, B. Kümmerer and S. Thorbjørnsen. *Quantum independent increment processes. II*, volume 1866 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin (2006). ISBN 978-3-540-24407-3; 3-540-24407-7. Lectures from the School “Quantum Independent Increment Processes: Structure and Applications to Physics” held in Greifswald, March 9–22, 2003. [MR2213451](#).
- O. E. Barndorff-Nielsen and S. Thorbjørnsen. Self-decomposability and Lévy processes in free probability. *Bernoulli* **8** (3), 323–366 (2002). [MR1913111](#).
- O. E. Barndorff-Nielsen and S. Thorbjørnsen. The Lévy-Itô decomposition in free probability. *Probab. Theory Related Fields* **131** (2), 197–228 (2005). [MR2117952](#).
- S. T. Belinschi. *Complex analysis methods in noncommutative probability*. Ph.D. thesis, Indiana University (2005).
- S. T. Belinschi and H. Bercovici. Partially defined semigroups relative to multiplicative free convolution. *Int. Math. Res. Not.* (2), 65–101 (2005). [MR2128863](#).
- S. T. Belinschi, M. Bożejko, F. Lehner and R. Speicher. The normal distribution is \boxplus -infinitely divisible. *Adv. Math.* **226** (4), 3677–3698 (2011). [MR2764902](#).
- H. Bercovici and V. Pata. Stable laws and domains of attraction in free probability theory. *Ann. of Math. (2)* **149** (3), 1023–1060 (1999). With an appendix by Philippe Biane.
- H. Bercovici and D. Voiculescu. Free convolution of measures with unbounded support. *Indiana Univ. Math. J.* **42** (3), 733–773 (1993). [MR1254116](#).
- P. Biane. Processes with free increments. *Math. Z.* **227** (1), 143–174 (1998). [MR1605393](#).
- L. Bondesson. *Generalized gamma convolutions and related classes of distributions and densities*, volume 76 of *Lecture Notes in Statistics*. Springer-Verlag, New York (1992). ISBN 0-387-97866-6. [MR1224674](#).
- L. Bondesson. A class of probability distributions that is closed with respect to addition as well as multiplication of independent random variables. *J. Theoret. Probab.* **28** (3), 1063–1081 (2015). [MR3413969](#).
- P. Bosch and T. Simon. On the infinite divisibility of inverse beta distributions. *Bernoulli* **21** (4), 2552–2568 (2015). [MR3378477](#).

- M. Bożejko and T. Hasebe. On free infinite divisibility for classical Meixner distributions. *Probab. Math. Statist.* **33** (2), 363–375 (2013). [MR3158562](#).
- N. Eisenbaum. Another failure in the analogy between Gaussian and semicircle laws. In *Séminaire de Probabilités XLIV*, volume 2046 of *Lecture Notes in Math.*, pages 207–213. Springer, Heidelberg (2012). [MR2953349](#).
- C. Goldie. A class of infinitely divisible random variables. *Proc. Cambridge Philos. Soc.* **63**, 1141–1143 (1967). [MR0215332](#).
- U. Haagerup and S. Möller. The law of large numbers for the free multiplicative convolution. In *Operator algebra and dynamics*, volume 58 of *Springer Proc. Math. Stat.*, pages 157–186. Springer, Heidelberg (2013). [MR3142036](#).
- U. Haagerup and S. Thorbjørnsen. On the free gamma distributions. *Indiana Univ. Math. J.* **63** (4), 1159–1194 (2014). [MR3263926](#).
- T. Hasebe. Free infinite divisibility for beta distributions and related ones. *Electron. J. Probab.* **19**, no. 81, 33 (2014). [MR3256881](#).
- W. Kaplan. Close-to-convex schlicht functions. *Michigan Math. J.* **1**, 169–185 (1953) (1952). [MR0054711](#).
- S. Kerov. Interlacing measures. In *Kirillov's seminar on representation theory*, volume 181 of *Amer. Math. Soc. Transl. Ser. 2*, pages 35–83. Amer. Math. Soc., Providence, RI (1998). [MR1618739](#).
- E. Lukacs. A characterization of the gamma distribution. *Ann. Math. Statist.* **26**, 319–324 (1955). [MR0069408](#).
- K. Noshiro. On the theory of schlicht functions. *J. Fac. Sci. Hokkaido Imper. Univ.* **02** (3), 129–155 (1934).
- V. Pérez-Abreu and N. Sakuma. Free generalized gamma convolutions. *Electron. Commun. Probab.* **13**, 526–539 (2008). [MR2447839](#).
- V. Pérez-Abreu and N. Sakuma. Free infinite divisibility of free multiplicative mixtures of the Wigner distribution. *J. Theoret. Probab.* **25** (1), 100–121 (2012). [MR2886381](#).
- V. K. Rohatgi, F. W. Steutel and G. J. Székely. Infinite divisibility of products and quotients of i.i.d. random variables. *Math. Sci.* **15** (1), 53–59 (1990). [MR1060698](#).
- A. Ruegg. A characterization of certain infinitely divisible laws. *Ann. Math. Statist.* **41**, 1354–1356 (1970). [MR0267626](#).
- D. N. Shanbhag, D. Pestana and M. Sreehari. Some further results in infinite divisibility. *Math. Proc. Cambridge Philos. Soc.* **82** (2), 289–295 (1977). [MR0448483](#).
- D. N. Shanbhag and M. Sreehari. On certain self-decomposable distributions. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **38** (3), 217–222 (1977). [MR0436267](#).
- R. Speicher and R. Woroudi. Boolean convolution. In *Free probability theory (Waterloo, ON, 1995)*, volume 12 of *Fields Inst. Commun.*, pages 267–279. Amer. Math. Soc., Providence, RI (1997). [MR1426845](#).
- F. W. Steutel. Note on the infinite divisibility of exponential mixtures. *Ann. Math. Statist.* **38**, 1303–1305 (1967). [MR0215339](#).
- F. W. Steutel and K. van Harn. *Infinite divisibility of probability distributions on the real line*, volume 259 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, Inc., New York (2004). ISBN 0-8247-0724-9. [MR2011862](#).
- K. Szpojankowski. A constant regression characterization of a marchenko-pastur. *ArXiv Mathematics e-prints* (2015a). [arXiv: 1503.00917](#).

- K. Szpojankowski. On the Lukacs property for free random variables. *Studia Math.* **228** (1), 55–72 (2015b). [MR3416954](#).
- O. Thorin. On the infinite divisibility of the lognormal distribution. *Scand. Actuar. J.* (3), 121–148 (1977a). [MR552135](#).
- O. Thorin. On the infinite divisibility of the Pareto distribution. *Scand. Actuar. J.* (1), 31–40 (1977b). [MR0431333](#).
- O. Thorin. Proof of a conjecture of L. Bondesson concerning infinite divisibility of powers of a gamma variable. *Scand. Actuar. J.* (3), 151–164 (1978). [MR514311](#).
- D. Voiculescu. Addition of certain non-commuting random variables. *Journal of Functional Analysis* **66** (3), 323–346 (1986). DOI: [10.1016/0022-1236\(86\)90062-5](#).
- S. E. Warschawski. On the higher derivatives at the boundary in conformal mapping. *Trans. Amer. Math. Soc.* **38** (2), 310–340 (1935). [MR1501813](#).