Title	Limit theorems in bi-free probability theory
Author(s)	Hasebe, Takahiro; Huang, Hao-Wei; Wang, Jiun-Chau
Citation	Probability theory and related fields, 172(3-4), 1081-1119 https://doi.org/10.1007/s00440-017-0825-6
Issue Date	2018-12
Doc URL	http://hdl.handle.net/2115/76221
Rights	This is a post-peer-review, pre-copyedit version of an article published in "Probability Theory and Related Fields". The final authenticated version is available online at: http://dx.doi.org/10.1007/s00440-017-0825-6
Type	article (author version)
File Information	Classical prob vs bifree prob_ver3.pdf

[Instructions for use](#)

LIMIT THEOREMS IN BI-FREE PROBABILITY THEORY

TAKAHIRO HASEBE, HAO-WEI HUANG AND JIUN-CHAU WANG

ABSTRACT. In this paper additive bi-free convolution is defined for general Borel probability measures, and the limiting distributions for sums of bi-free pairs of self-adjoint commuting random variables in an infinitesimal triangular array are determined. These distributions are characterized by their bi-freely infinite divisibility, and moreover, a transfer principle is established for limit theorems in classical probability theory and Voiculescu's bi-free probability theory. Complete descriptions of bi-free stability are given and fullness of planar probability distributions are studied. All these results reveal one important feature about the theory of bi-free probability that it parallels the classical theory perfectly well. The emphasis in the whole work is not on the tool of bi-free combinatorics but only on the analytic machinery.

1. INTRODUCTION

The purpose of this paper is to establish an explicit connection between the families of infinitely divisible laws in classical probability theory and bi-free probability theory. As in free probability theory, it is shown that some classical limit theorem has a nice analogue in the bi-free framework.

Denote by $\mathcal{P}_{\mathbb{R}^d}$ the family of Borel probability measures on $\mathbb{R}^d$. The classical convolution $\mu * \nu$ of μ and ν in $\mathcal{P}_{\mathbb{R}^d}$ is the probability distribution of the sum of two independent random vectors whose respective distributions are μ and ν . In the theory of free probability, freeness and free convolution $\boxplus$ are treated as analogues of classical notion of independence and classical convolution for non-commutative random variables, respectively [23]. The latter theory has a *two-faced* extension, which was invented to study pairs of *left* and *right* random variables (also called left and right *faces*) on a free product of complex Hilbert spaces simultaneously [21]. An independence relation put among these pairs is called *bi-freeness*, which gives rise to *bi-free probability theory*. This fascinating theory has grown rapidly and included several interesting findings based on the foundation of free probability theory.

Since the introduction of bi-free probability by Voiculescu in 2013, combinatorial and analytical approaches have been the main research focuses so far [7, 8, 12, 14, 19, 20]. Given a two-faced pair (a, b) in a C^* -probability space $(\mathcal{A}, \varphi)$, its *bi-free partial R -transform* is defined through its *Cauchy transform* $G_{(a,b)}(z, w) = \varphi((z - a)^{-1}(w - b)^{-1})$ as

$$R_{(a,b)}(z, w) = zR_a(z) + wR_b(w) + 1 - \frac{zw}{G_{(a,b)}(R_a(z) + 1/z, R_b(w) + 1/w)}$$

for complex values z and w in a neighborhood of zero, where R_a and R_b are the usual R -transforms of a and b , respectively [22]. This transform plays the same role as the usual R -transform does in the free case. The bi-freeness of (a, b) and (c, d) yields the freeness of the left faces a, c and the freeness of the right ones b, d [21]. The reader is referred to [2, 13] for some recent developments.

Date: December 18, 2017.

¹*2010 Mathematics Subject Classification:* Primary 46L54, Secondary 60E07.

²*Key words and phrases.* bi-free limit theorem, bi-free infinitely divisible distributions, bi-freely stable distributions, full distributions.

In the present paper, we continue the previous work [14] and contribute to the research of bi-free harmonic analysis without any emphasis on Voiculescu's original motivation. To accommodate objects like planar probability distributions or integral representations, it is natural to constrain ourselves to commuting and self-adjoint pairs (a, b) in a certain C^* -probability space, i.e. a, b are self-adjoint elements whose commutator $ab - ba = 0$. This yields an analytic perspective to bi-free probability theory. More precisely, in such a circumstance the Cauchy transform of (a, b) admits an integral form with the joint distribution of (a, b) as its underlying measure. Any measure in $\mathcal{P}_{\mathbb{R}^2}$ with compact support arises in this manner, i.e. it serves as the joint distribution of such a pair. The starting point of relating bi-free probability to the classical situation is that by realizing two given compactly supported $\mu, \nu \in \mathcal{P}_{\mathbb{R}^2}$ as joint distributions of two bi-free and commuting pairs, the bi-free partial R -transform linearizes the bi-free convolution $\boxplus$: for (z, w) near $(0, 0)$,

$$R_{\mu \boxplus \nu}(z, w) = R_{\mu}(z, w) + R_{\nu}(z, w).$$

An important concept in the study of limit theorems in probability theory is the *infinite divisibility*. A probability distribution on $\mathbb{R}^d$ is infinitely divisible with respect to a binary operation $\star$ on $\mathcal{P}_{\mathbb{R}^d}$ if it can be expressed as the $\star$ -convolution of an arbitrary number of copies of identical measures from $\mathcal{P}_{\mathbb{R}^d}$. When $d = 1$, this subject has been thoroughly studied by de Finetti, Kolmogorov, Lévy and Hinčin in classical probability [11, 18]. The free counterpart is also well studied [5]. The theory of infinitely divisible distributions generalizes (free) central limit theorem as they serve as the limit laws for sums of (freely) independent and identically distributed random variables. (Free) Gaussian and (free) Poisson distributions are typical examples of $(\boxplus)$ - $\star$ -infinitely divisible distributions. Distributions of this kind are determined by their characteristic functions or free R -transforms, the so-called Lévy-Hinčin type representations. Measures in $\mathcal{P}_{\mathbb{R}^2}$ which are $\boxplus$ -infinitely divisible were first studied in [12] in the case when they are compactly supported and considered in the general case in [14].

The question under investigation in this paper is to provide the criteria for the weak convergence of the sequence

$$(1.1) \quad \mu_{n1} \boxplus \mu_{n2} \boxplus \cdots \boxplus \mu_{nk_n} \boxplus \delta_{\mathbf{v}_n},$$

where $\{k_n\}_{n=1}^{\infty}$ is a sequence of strictly increasing positive integers, $\{\mu_{nk}\}_{n \geq 1, 1 \leq k \leq k_n}$ is an *infinitesimal triangular array* of measures in $\mathcal{P}_{\mathbb{R}^2}$ and $\delta_{\mathbf{v}_n}$ denotes the dirac measure at the vector $\mathbf{v}_n$ in $\mathbb{R}^2$. Here the infinitesimal condition of a triangular array $\{\mu_{nk}\}$ in $\mathcal{P}_{\mathbb{R}^d}$ means that for any $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \max_{1 \leq k \leq k_n} \mu_{nk}(\{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\| \geq \epsilon\}) = 0,$$

where $\|\cdot\|$ is the Euclidean norm on $\mathbb{R}^d$. When $d = 1$, there is a one-to-one correspondence between the sets of $\star$ -infinitely divisible laws and $\boxplus$ -infinitely divisible laws. Such a correspondence is characterized by the same parameters, a real number and a positive Borel measure on the real line, in the Lévy-Hinčin type formulas [3, 9]. We show in Theorem 5.5 that under the hypotheses mentioned above the weak convergence of the sequence in (1.1) is equivalent to that of the sequence

$$(1.2) \quad \mu_{n1} * \mu_{n2} * \cdots * \mu_{nk_n} * \delta_{\mathbf{v}_n}.$$

Moreover, the limit distribution of (1.1) is shown to be bi-freely infinitely divisible, and the parameters of its bi-free Lévy-Hinčin formula coincide with those in the Lévy-Hinčin formula of the limit

60 distribution of (1.2). These results indicate that a lot of work in the study of limit laws of sums
 61 of independent identically distributed random vectors has its counterpart in bi-free probability
 62 theory.

63 After setting up some basic tools needed for the investigation and proving the generalization of
 64 the bi-free convolution of measures in $\mathcal{P}_{\mathbb{R}^2}$ with compact supports to arbitrary ones in Section 2,
 65 another useful bi-free Lévy-Hinčin integral representation is provided in Section 3. Section 4 and
 66 5 are dedicated to building the parallelism between the families of infinitely divisible laws with
 67 respect to classical and bi-free convolutions. The investigation of $\boxplus\boxplus$ -stable laws on $\mathbb{R}^2$ with their
 68 domains of attraction is carried out in Section 6, while the fullness of $\boxplus\boxplus$ -infinitely divisible laws
 69 is set down in Section 7.

70 2. PRELIMINARIES

71 We begin with reviewing some definitions and results in [21, 22]. The *Cauchy transform* of a
 72 planar Borel probability measure μ , defined as

$$G_\mu(z, w) = \int_{\mathbb{R}^2} \frac{d\mu(s, t)}{(z - s)(w - t)},$$

73 is an analytic function and satisfies the relation

$$G_\mu(\bar{z}, \bar{w}) = \overline{G_\mu(z, w)}$$

74 on $(\mathbb{C} \setminus \mathbb{R})^2$. The underlying measure μ can be recovered from the transform by the *Stieltjes*
 75 *inversion formula*: the family $\{\mu_\epsilon\}_{\epsilon>0}$ of probability measures on $\mathbb{R}^2$ defined by

$$d\mu_\epsilon(s, t) = -\frac{1}{2} \Re[G_\mu(s + i\epsilon, t + i\epsilon) - G_\mu(s + i\epsilon, t - i\epsilon)] \frac{dsdt}{\pi^2}$$

76 converges to μ weakly as $\epsilon \rightarrow 0^+$ [14].

77 A *truncated cone* in the complex plane is a set of the form

$$\Gamma_{\theta, M} := \{x + iy \in \mathbb{C} : |x| \leq \theta|y|, |y| \geq M\}, \quad \theta, M > 0.$$

78 For notational convenience, it will be simply denoted by Γ in the sequel if θ and M are known.
 79 Recall from [5] that free Voiculescu's transform ϕ_ν of a measure $\nu \in \mathcal{P}_{\mathbb{R}}$ is an analytic function and
 80 satisfies the relation $F_\nu(\phi_\nu(z) + z) = z$ on a certain truncated cone Γ , where F_ν is the reciprocal
 81 of the Cauchy transform of ν ,

$$G_\nu(z) = \int_{\mathbb{R}} \frac{d\nu(s)}{z - s}.$$

82 The free Voiculescu transform linearizes the free convolution of probability distributions on $\mathbb{R}$.

83 For a given $\mu \in \mathcal{P}_{\mathbb{R}^2}$, the probability measures defined as $\mu^{(1)}(B) = \mu(B \times \mathbb{R})$ and $\mu^{(2)}(B) =$
 84 $\mu(\mathbb{R} \times B)$ for Borel sets $B \subset \mathbb{R}$ are called the *marginal laws* of μ . The *bi-free ϕ -transform* of μ
 85 defined as

$$(2.3) \quad \phi_\mu(z, w) = \frac{\phi_{\mu^{(1)}}(z)}{z} + \frac{\phi_{\mu^{(2)}}(w)}{w} + 1 - \frac{1}{zwG_\mu\left(F_{\mu^{(1)}}^{-1}(z), F_{\mu^{(2)}}^{-1}(w)\right)}$$

86 is an analytic function and satisfies the relation

$$(2.4) \quad \phi_\mu(\bar{z}, \bar{w}) = \overline{\phi_\mu(z, w)}$$

on Γ^2 (notice that the denominator of the last term in (2.3) never vanishes by shrinking the domain Γ if necessary).

It was shown in [22] that the bi-free ϕ -transform linearizes the bi-free additive convolution of two planar Borel probability measures μ and ν with compact support:

$$(2.5) \quad \phi_{\mu \boxplus \nu}(z, w) = \phi_\mu(z, w) + \phi_\nu(z, w)$$

for (z, w) in the common domain of these transforms. The marginal laws of the bi-free convolution of compactly supported probability measures on $\mathbb{R}^2$ can be expressed in terms of the free convolution of their marginal laws [12]: for $j = 1, 2$,

$$(2.6) \quad (\mu \boxplus \nu)^{(j)} = \mu^{(j)} \boxplus \nu^{(j)}.$$

A sequence $\{\nu_n\}_{n=1}^\infty \subset \mathcal{P}_{\mathbb{R}^d}$ is said to *converge weakly* to $\nu \in \mathcal{P}_{\mathbb{R}^d}$, denoted by $\nu_n \Rightarrow \nu$, if

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} f d\nu_n = \int_{\mathbb{R}^d} f d\nu$$

for any bounded and continuous function f on $\mathbb{R}^d$. For any sequence $\{\mu_n\}_{n=1}^\infty \subset \mathcal{P}_{\mathbb{R}^2}$ converging weakly to $\mu \in \mathcal{P}_{\mathbb{R}^2}$, we have for $j = 1, 2$,

$$(2.7) \quad \mu_n^{(j)} \Rightarrow \mu^{(j)}.$$

A family $\mathcal{F}$ of Borel probability measures on $\mathbb{R}^d$ is called *tight* if

$$\lim_{r \rightarrow \infty} \sup_{\mu \in \mathcal{F}} \mu(\{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\| > r\}) = 0.$$

The correlation of the tightness (or weak convergence) of Borel probability measures on $\mathbb{R}$ and $\mathbb{R}^2$ and the convergence properties of their free and bi-free ϕ -transforms are well known [3, 14]. To make the presentation accessible for readers of different backgrounds, some results with their proofs in this direction are provided below. Recall that points z in $\mathbb{C} \setminus \mathbb{R}$ are said to tend to infinity *non-tangentially*, which is denoted by $z \rightarrow_{\triangleleft} \infty$, if $z \rightarrow \infty$ with $|\Re z / \Im z|$ uniformly bounded.

Proposition 2.1. A family $\mathcal{F} \subset \mathcal{P}_{\mathbb{R}^2}$ is tight if and only if $zwG_\mu(z, w) - 1 = o(1)$ uniformly for $\mu \in \mathcal{F}$ as $z, w \rightarrow_{\triangleleft} \infty$.

Proof. First suppose that $\mathcal{F}$ is tight. If $\xi \in \mathbb{C} \setminus \mathbb{R}$ with $|\Re \xi / \Im \xi|$ bounded by $\theta > 0$, then

$$\left| \frac{c}{\xi - c} \right| \leq \sqrt{1 + \theta^2}, \quad c \in \mathbb{R},$$

which is due to the inequality $(x - c)^2 + x^2/\theta^2 \geq c^2/(1 + \theta^2)$, $x \in \mathbb{R}$. Applying this inequality to the decomposition

$$\frac{zw}{(z - s)(w - t)} - 1 = \frac{s}{z - s} + \frac{t}{w - t} + \frac{st}{(z - s)(w - t)}$$

shows the existence of some constant $c_\theta > 0$ depending on θ only so that for $|\Re z / \Im z|, |\Re w / \Im w| \leq \theta$,

$$\left| \frac{zw}{(z - s)(w - t)} - 1 \right| \leq c_\theta, \quad (s, t) \in \mathbb{R}^2.$$

Hence for any $\mu \in \mathcal{F}$, we have

$$|zwG_\mu(z, w) - 1| \leq \frac{r}{|\Im z|} + \frac{r}{|\Im w|} + \frac{r^2}{|\Im z||\Im w|} + c_\theta \mu(\{\|\mathbf{x}\| > r\}),$$

110 which clearly yields the necessity.

111 Now we prove the sufficiency. Given any $\epsilon > 0$, let $M > 1$ be large enough so that

$$(2.8) \quad \sup_{\mu \in \mathcal{F}} |(iy)(iv)G_\mu(iy, iv) - 1| < \epsilon$$

112 whenever $|y|, |v| \geq M$. Since the inequality

$$\left| \frac{iv}{(iy - s)(iv - t)} \right| \leq 1$$

113 holds for $(s, t) \in \mathbb{R}^2$ and $|y|, |v| \geq M$, it follows that for all $\mu \in \mathcal{F}$,

$$ivG_\mu(iy, iv) = \int_{\mathbb{R}^2} \frac{iv d\mu(s, t)}{(iy - s)(iv - t)} \rightarrow \int_{\mathbb{R}^2} \frac{d\mu(s, t)}{iy - s} = G_{\mu^{(1)}}(iy)$$

114 as $|v| \rightarrow +\infty$ by Dominated Convergence Theorem. Consequently, $iyG_{\mu^{(1)}}(iy) \rightarrow 1$ uniformly for
 115 $\mu \in \mathcal{F}$ as $|y| \rightarrow +\infty$ by letting $|v| \rightarrow +\infty$ in (2.8). This yields the tightness of the marginal
 116 sequence $\{\mu^{(1)} : \mu \in \mathcal{F}\}$ (cf. [5, Proposition 5.1]). The tightness of $\{\mu^{(2)} : \mu \in \mathcal{F}\}$ can be obtained
 117 in a similar way. Now the sufficiency follows since for any $r > 0$,

$$(2.9) \quad \mu(\{\|(s, t)\| \geq r\}) \leq \mu^{(1)}(\{s : |s| \geq r/\sqrt{2}\}) + \mu^{(2)}(\{t : |t| \geq r/\sqrt{2}\}).$$

118

□

119 **Proposition 2.2.** Let $\mathcal{F} \subset \mathcal{P}_{\mathbb{R}^2}$ be a tight family. Then $F_{\mu^{(j)}}$ is univalent on some common
 120 truncated cone Γ in $\mathbb{C}$ with image $F_{\mu^{(j)}}(\Gamma)$ containing some contracted cone $\Gamma_{\alpha, L}$ for every $\mu \in \mathcal{F}$
 121 and $j = 1, 2$. Moreover, $F_{\mu^{(j)}}^{-1}(\xi) = (1 + o(1))\xi$ uniformly for $\mu \in \mathcal{F}$ and $F_{\mu^{(j)}}^{-1}(\xi) \rightarrow_{\triangleleft} \infty$ as $\xi \rightarrow \infty$
 122 with ξ staying in $\Gamma_{\alpha, L}$.

123 *Proof.* The existence of such a truncated cone $\Gamma_{\alpha, L}$ and the asymptotic behaviors can be shown
 124 by the statements and techniques of [5, Proposition 5.4] (see also [3, Proposition 2.6]). □

125 In the following the weak convergence of measures in $\mathcal{P}_{\mathbb{R}^2}$ is translated into the asymptotic
 126 properties of their bi-free ϕ -transforms. Recall that the simplified form Γ^2 denotes the domain of
 127 the bi-free ϕ -transform of a planar probability measure on which (2.4) holds.

128 **Proposition 2.3.** Let $\{\mu_n\}_{n=1}^\infty \subset \mathcal{P}_{\mathbb{R}^2}$. Then the following assertions are equivalent.

- 129 (1) The sequence $\{\mu_n\}_{n=1}^\infty$ converges weakly to a planar Borel probability measure μ .
- 130 (2) Functions in the sequence $\{\phi_{\mu_n}\}_{n=1}^\infty$ are defined on some fixed domain Γ^2 , converge uni-
 131 formly on compact sets of Γ^2 to a function ϕ , and $\phi_{\mu_n}(z, w) = o(1)$ uniformly in n as
 132 $z, w \rightarrow \infty$ with $(z, w) \in \Gamma^2$.
- 133 (3) Functions in the sequence $\{\phi_{\mu_n}\}_{n=1}^\infty$ are defined on some fixed domain Γ^2 , $\lim_{n \rightarrow \infty} \phi_{\mu_n}(iy, iv)$
 134 exists for $(iy, iv) \in \Gamma^2$, and $\phi_{\mu_n}(iy, iv) = o(1)$ uniformly in n as $|y|, |v| \rightarrow \infty$.

135 Moreover, if (1) and (2) are satisfied, then $\phi = \phi_\mu$ in Γ^2 .

136 *Proof.* Throughout the proof, we will use the notations $G_n = G_{\mu_n}$, $G_{jn} = G_{\mu_n^{(j)}}$, $F_{jn} = F_{\mu_n^{(j)}}$,
 137 $F_j = F_{\mu^{(j)}}$, $\phi_{jn} = \phi_{\mu_n^{(j)}}$, and $\phi_j = \phi_{\mu^{(j)}}$ for all n and for $j = 1, 2$.

138 First, suppose $\mu_n \Rightarrow \mu$. According to (2.7) and [5, Proposition 5.7], there exist $\theta, M > 0$ so
 139 that every ϕ_{jn} is defined on $\Gamma := \Gamma_{\theta, M}$, $\phi_{jn} \rightarrow \phi_j$ uniformly on compact sets of Γ as $n \rightarrow \infty$, and
 140 $\phi_{jn}(\xi) = o(|\xi|)$ uniformly in n as $\xi \rightarrow \infty$ with $\xi \in \Gamma$ for $j = 1, 2$. Hence each ϕ_{μ_n} is defined on
 141 Γ^2 . On the other hand, the integrands in G_n are uniformly bounded functions of (s, t) for points

(z, w) lying in compact sets of $(\mathbb{C} \setminus \mathbb{R})^2$. This yields the normality of $\{G_n\}$ by Montel's theorem in complex analysis of several variables. Hence $G_n(F_{1n}^{-1}, F_{2n}^{-1}) \rightarrow G_\mu(F_1^{-1}, F_2^{-1})$ uniformly on compact sets of Γ^2 as $n \rightarrow \infty$ and $\phi = \phi_\mu$. To finish the proof of (1) $\Rightarrow$ (2), it remains to show that

$$\frac{zwG_n(F_{1n}^{-1}(z), F_{2n}^{-1}(w)) - 1}{zwG_n(F_{1n}^{-1}(z), F_{2n}^{-1}(w))} = o(1)$$

uniformly in n as $z, w \rightarrow \infty$ with $(z, w) \in \Gamma^2$, which is equivalent to showing the uniform convergence of $zwG_n(F_{1n}^{-1}(z), F_{2n}^{-1}(w)) - 1 = o(1)$. Note that this can be obtained by applying Proposition 2.1 and Proposition 2.2 to the identity

$$(2.10) \quad zwG_n(F_{1n}^{-1}(z), F_{2n}^{-1}(w)) = F_{1n}^{-1}(z)F_{2n}^{-1}(w)G_n(F_{1n}^{-1}(z), F_{2n}^{-1}(w)) \cdot \frac{z}{F_{1n}^{-1}(z)} \cdot \frac{w}{F_{2n}^{-1}(w)}.$$

Clearly, (2) implies (3). To show (3) $\Rightarrow$ (1), it suffices to verify that the sequence $\{\mu_n\}_{n=1}^\infty$ is tight by the established result (1) $\Rightarrow$ (2). To conclude the proof, observe that

$$\begin{aligned} \phi_{\mu_n}(iy, iv) &= \frac{\phi_{1n}(iy)}{iy} + \frac{\phi_{2n}(iv)}{iv} + 1 - \frac{1}{(iy)\frac{iv}{F_{2n}^{-1}(iv)}F_{2n}^{-1}(iv)G_n(F_{1n}^{-1}(iy), F_{2n}^{-1}(iv))} \\ &\rightarrow \frac{\phi_{1n}(iy)}{iy} + 1 - \frac{1}{(iy)G_{1n}(F_{1n}^{-1}(iy))} \quad \text{as } |v| \rightarrow \infty \\ &= \frac{\phi_{1n}(iy)}{iy}, \end{aligned}$$

where Proposition 2.2 and the fact that $wG_n(z, w) \rightarrow G_{1n}(z)$ for any $z \in \mathbb{C} \setminus \mathbb{R}$ as $w \rightarrow \infty$ non-tangentially are used in the limit. This implies that $\phi_{1n}(iy) = o(|y|)$ uniformly in n as $|y| \rightarrow \infty$. Similarly, $\{\phi_{2n}\}$ has the same asymptotic property. Hence $(iy)(iv)G_n(F_{1n}^{-1}(iy), F_{2n}^{-1}(iv)) - 1 = o(1)$ uniformly in n as $|y|, |v| \rightarrow \infty$ by (2.10), which yields the tightness of $\{\mu_n\}$ by Proposition 2.1. The proof is complete. $\square$

We can now define the bi-free convolution of arbitrary planar Borel probability measures μ and ν . Choose two sequences $\{\mu_n\}_{n=1}^\infty$ and $\{\nu_n\}_{n=1}^\infty$ of planar Borel probability measures with compact support converging to μ and ν weakly, respectively. Then Proposition 2.3 shows that $\mu_n \boxplus \boxplus \nu_n$ weakly converges to a probability measure ρ on $\mathbb{R}^2$ which satisfies

$$(2.11) \quad \phi_\rho(z, w) = \phi_\mu(z, w) + \phi_\nu(z, w)$$

on some Γ^2 . We further deduce from [14, Proposition 2.5] the uniqueness of ρ . These discussions lead into the following definition:

Definition 2.4. For any $\mu, \nu \in \mathcal{P}_{\mathbb{R}^2}$, the unique probability measure ρ satisfying the additive identity in (2.11) is called the *bi-free additive convolution* of μ and ν , and is also denoted by $\mu \boxplus \boxplus \nu$.

Remark 2.5. Our approach to the generalization of bi-free additive convolution is based on analytic tools. An (unbounded) operator model for the bi-free convolution of arbitrary probability measures on $\mathbb{R}^2$ is unknown.

We can also show by Proposition 2.3 that the operation of bi-free convolution is weakly continuous, namely, if $\{\mu_n\}_{n=1}^\infty$ and $\{\nu_n\}_{n=1}^\infty$ are in $\mathcal{P}_{\mathbb{R}^2}$ weakly converging to μ and ν , respectively,

167 then $\mu_n \boxplus \nu_n$ weakly converges to $\mu \boxplus \nu$. Finally, we generalize the result in (2.6) to arbitrary
 168 measures in $\mathcal{P}_{\mathbb{R}^2}$.

169 **Proposition 2.6.** Let $\mu, \nu \in \mathcal{P}_{\mathbb{R}^2}$. Then for $j = 1, 2$,

$$(\mu \boxplus \nu)^{(j)} = \mu^{(j)} \boxplus \nu^{(j)}.$$

170 *Proof.* Choose two sequences $\{\mu_n\}_{n=1}^\infty$ and $\{\nu_n\}_{n=1}^\infty$ of planar Borel probability measures with
 171 compact support converging to μ and ν weakly, respectively. Since the projections onto marginals
 172 and bi-free convolution are weakly continuous, the result (2.6) passes to the conclusion. $\square$

173 3. BI-FREE INFINITE DIVISIBILITY AND LÉVY-HINČIN REPRESENTATION

174 Throughout the remaining part of the paper, points (s, t) in $\mathbb{R}^2$ will be denoted by the bold
 175 letter $\mathbf{x}$ and the origin $(0, 0)$ will be written as $\mathbf{0}$. We will also denote by the real numbers $\mathbf{v}^{(1)}$
 176 and $\mathbf{v}^{(2)}$ the s - and t -coordinate of a given vector $\mathbf{v} \in \mathbb{R}^2$.

177 In classical probability theory, a Borel probability measure μ on $\mathbb{R}^2$ is $*$ -infinitely divisible if and
 178 only if its characteristic function is of the form (called the *Lévy-Hinčin representation*)

$$(3.12) \quad \widehat{\mu}(\mathbf{u}) = \exp \left[i\langle \mathbf{u}, \mathbf{v} \rangle - \frac{1}{2} \langle \mathbf{A} \mathbf{u}, \mathbf{u} \rangle + \int_{\mathbb{R}^2} \left(e^{i\langle \mathbf{u}, \mathbf{x} \rangle} - 1 - \frac{i\langle \mathbf{u}, \mathbf{x} \rangle}{1 + \|\mathbf{x}\|^2} \right) d\tau(\mathbf{x}) \right]$$

179 for some vector $\mathbf{v} \in \mathbb{R}^2$, real positive semi-definite matrix $\mathbf{A}$ and some positive Borel measure τ
 180 on $\mathbb{R}^2$ with the properties that $\tau(\{\mathbf{0}\}) = 0$ and $1 \wedge \|\mathbf{x}\|^2 \in L^1(\tau)$, where $1 \wedge \|\mathbf{x}\|^2 := \min\{1, \|\mathbf{x}\|^2\}$.
 181 Conversely, such a triplet $(\mathbf{v}, \mathbf{A}, \tau)$ generates a probability measure μ for which (3.12) holds. The
 182 triplet $(\mathbf{v}, \mathbf{A}, \tau)$ in the representation is unique and called the (classical) *characteristic triplet* of
 183 μ , while the measure τ is called the (classical) *Lévy measure* of μ . In this case μ is denoted by
 184 $\mu_*^{(\mathbf{v}, \mathbf{A}, \tau)}$. The reader is referred to [16, 18] for more details.

185 Recall that a measure $\mu \in \mathcal{P}_{\mathbb{R}^2}$ is said to be *bi-freely infinitely divisible* if for any $n \in \mathbb{N}$, it can
 186 be expressed as an n -fold bi-free convolution of some $\mu_n \in \mathcal{P}_{\mathbb{R}^2}$:

$$\mu = \underbrace{\mu_n \boxplus \cdots \boxplus \mu_n}_{n \text{ terms}} := \mu_n^{\boxplus n}.$$

187 Such a measure is characterized in terms of the functional properties of its bi-free ϕ -transform [14,
 188 Theorem 4.3]: μ is $\boxplus$ -infinitely divisible if and only if ϕ_μ extends analytically to $(\mathbb{C} \setminus \mathbb{R})^2$ and
 189 admits an integral representation of the form

$$\phi_\mu(z, w) = \frac{1}{z} \left(\gamma_1 + \int_{\mathbb{R}^2} \frac{1 + zs}{z - s} d\sigma_1(\mathbf{x}) \right) + \frac{1}{w} \left(\gamma_2 + \int_{\mathbb{R}^2} \frac{1 + wt}{w - t} d\sigma_2(\mathbf{x}) \right) + \widetilde{D}(z, w),$$

190 where $(\gamma_1, \gamma_2) \in \mathbb{R}^2$, σ_j is a finite, positive Borel measure on $\mathbb{R}^2$ for $j = 1, 2$, and

$$\widetilde{D}(z, w) = \int_{\mathbb{R}^2} \frac{\sqrt{1 + s^2} \sqrt{1 + t^2}}{(z - s)(w - t)} d\widetilde{\sigma}(\mathbf{x})$$

191 for some finite Borel signed measure $\widetilde{\sigma}$ on $\mathbb{R}^2$ satisfying the relations

$$(3.13) \quad \begin{cases} \frac{t}{\sqrt{1+t^2}} \sigma_1 = \frac{s}{\sqrt{1+s^2}} \widetilde{\sigma}, \\ \frac{s}{\sqrt{1+s^2}} \sigma_2 = \frac{t}{\sqrt{1+t^2}} \widetilde{\sigma}, \\ \widetilde{\sigma}(\{\mathbf{0}\})^2 \leq \sigma_1(\{\mathbf{0}\}) \sigma_2(\{\mathbf{0}\}). \end{cases}$$

These parameters $\gamma_1, \gamma_2, \sigma_1, \sigma_2$ and $\tilde{\sigma}$ appearing in the representation are unique. Notice that the first two relations in (3.13) indicate that $a := \sigma_1(\{0\} \times \mathbb{R}) = \sigma_1(\{\mathbf{0}\})$, $b := \sigma_2(\mathbb{R} \times \{0\}) = \sigma_2(\{\mathbf{0}\})$ and $c := \tilde{\sigma}(\{st = 0\}) = \tilde{\sigma}(\{\mathbf{0}\})$. An application of Cauchy-Schwarz inequality gives $c^2 \leq ab$, i.e. the positive semi-definiteness of the matrix

$$(3.14) \quad \mathbf{A} = \begin{pmatrix} a & c \\ c & b \end{pmatrix}.$$

In order to get more insights into $\boxplus$ -infinitely divisible distributions, we will derive another integral representation for them. The representing measure is no longer required to be finite but only positive. First of all, define the positive measure τ on $\mathbb{R}^2$ as

$$(3.15) \quad \tau = \begin{cases} \frac{1+s^2}{s^2} \sigma_1 & \text{on } \{(s, t) \in \mathbb{R}^2 : s \neq 0\}; \\ \frac{1+t^2}{t^2} \sigma_2 & \text{on } \{(s, t) \in \mathbb{R}^2 : t \neq 0\}, \end{cases}$$

and $\tau(\{\mathbf{0}\}) = 0$. The relations among σ_1, σ_2 and $\tilde{\sigma}$ in (3.13) clearly show that τ is well-defined. Moreover, the restriction of τ to the set $\{(s, t) \in \mathbb{R}^2 : st \neq 0\}$ is equal to

$$\frac{\sqrt{1+s^2}\sqrt{1+t^2}}{st} \chi_{\{st \neq 0\}}(s, t) \tilde{\sigma}.$$

It is also easy to verify that the function $1 \wedge \|\mathbf{x}\|^2$ belongs to $L^1(\tau)$. After these setups, we can rewrite ϕ_μ as

$$(3.16) \quad \phi_\mu(z, w) = \frac{\mathbf{v}^{(1)}}{z} + \frac{\mathbf{v}^{(2)}}{w} + \left(\frac{a}{z^2} + \frac{c}{zw} + \frac{b}{w^2} \right) + \mathcal{P}(z, w),$$

where $\mathbf{v} \in \mathbb{R}^2$ and

$$\mathcal{P}(z, w) = \int_{\mathbb{R}^2} \left[\frac{zw}{(z-s)(w-t)} - 1 - \frac{sz^{-1} + tw^{-1}}{1+s^2+t^2} \right] d\tau(s, t).$$

Indeed, by means of the identities

$$(3.17) \quad \frac{zs}{z-s} = \frac{1+zs}{z-s} \frac{s^2}{1+s^2} + \frac{s}{1+s^2}$$

and

$$\frac{s}{1+s^2} = \frac{s}{1+s^2+t^2} + \frac{s}{1+s^2} \frac{t^2}{1+s^2+t^2}$$

which hold for any $z \in \mathbb{C} \setminus \mathbb{R}$ and $s, t \in \mathbb{R}$, we obtain that

$$(3.18) \quad \gamma_1 + \int_{\mathbb{R}^2} \frac{1+zs}{z-s} d\sigma_1(\mathbf{x}) = \phi_{\mu^{(1)}}(z) = \mathbf{v}^{(1)} + \frac{a}{z} + \int_{\mathbb{R}^2} \left[\frac{zs}{z-s} - \frac{s}{1+s^2+t^2} \right] d\tau(\mathbf{x})$$

for some $\mathbf{v}^{(1)} \in \mathbb{R}$. Similarly, one can obtain that

$$(3.19) \quad \gamma_2 + \int_{\mathbb{R}^2} \frac{1+wt}{w-t} d\sigma_2(\mathbf{x}) = \phi_{\mu^{(2)}}(w) = \mathbf{v}^{(2)} + \frac{b}{w} + \int_{\mathbb{R}^2} \left[\frac{wt}{w-t} - \frac{t}{1+s^2+t^2} \right] d\tau(\mathbf{x}).$$

Combining the identities (3.18) and (3.19) with

$$\tilde{D}(z, w) = \frac{c}{zw} + \int_{\mathbb{R}^2} \frac{st}{(z-s)(w-t)} d\tau(s, t)$$

yields the desired expression (3.16). Conversely, a function admitting such an integral form (3.16) with the required properties stated above can be shown to be the bi-free ϕ -transform of some bi-freely infinitely divisible measure. These observations lead to the following result.

Theorem 3.1. (Bi-free Lévy-Hinčin representation) *A probability distribution μ on $\mathbb{R}^2$ is bi-freely infinitely divisible if and only if its bi-free ϕ -transform extends analytically to $(\mathbb{C} \setminus \mathbb{R})^2$ and admits the integral representation (3.16), where $\mathbf{v} \in \mathbb{R}^2$, the matrix $\mathbf{A}$ given as in (3.14) is positive semi-definite, and τ is a positive measure on $\mathbb{R}^2$ with the properties $\tau(\{\mathbf{0}\}) = 0$ and $1 \wedge \|\mathbf{x}\|^2 \in L^1(\tau)$. Moreover, the triplet $(\mathbf{v}, \mathbf{A}, \tau)$ in (3.16) is unique. Conversely, given such a triplet $(\mathbf{v}, \mathbf{A}, \tau)$ there exists a probability measure μ for which (3.16) holds.*

Theorem 3.1 shows that the set $\mathcal{ID}(\boxplus\boxplus)$ of bi-freely infinitely divisible distributions is completely parameterized by the triplet $(\mathbf{v}, \mathbf{A}, \tau)$. In the sequel, a probability measure μ in $\mathcal{ID}(\boxplus\boxplus)$ having the representation (3.16) will be denoted by $\mu_{\boxplus\boxplus}^{(\mathbf{v}, \mathbf{A}, \tau)}$, in which τ is called the *bi-free Lévy measure* and $(\mathbf{v}, \mathbf{A}, \tau)$ is called the *bi-free characteristic triplet* of $\mu_{\boxplus\boxplus}^{(\mathbf{v}, \mathbf{A}, \tau)}$. The classical and bi-free Lévy-Hinčin representations (3.12) and (3.16) have exactly the same characteristic triplets. As a matter of fact, such a one-to-one correspondence also holds true in general limit theorems, see Theorem 5.5.

Example 3.2. Let μ be $\boxplus\boxplus$ -infinitely divisible and let $(\mathbf{v}, \mathbf{A}, \tau)$ be its bi-free characteristic triplet.

(1) Then μ is called a *bi-free Gaussian distribution* if $\tau = 0$.

(2) If τ satisfies

$$\int_{\mathbb{R}^2} \frac{\|\mathbf{x}\|}{1 + \|\mathbf{x}\|^2} d\tau(\mathbf{x}) < \infty,$$

then (3.16) is reduced to the form

$$\phi_\mu(z, w) = \frac{\mathbf{u}^{(1)}}{z} + \frac{\mathbf{u}^{(2)}}{w} + \left(\frac{a}{z^2} + \frac{c}{zw} + \frac{b}{w^2} \right) + \int_{\mathbb{R}^2} \left[\frac{zw}{(z-s)(w-t)} - 1 \right] d\tau(s, t)$$

for some vector $\mathbf{u} \in \mathbb{R}^2$ called the *drift* of μ .

(3) Let ν be in $\mathcal{P}_{\mathbb{R}^2}$ with $\nu(\{\mathbf{0}\}) = 0$. Then μ is called a *bi-free compound Poisson distribution* with rate $\lambda > 0$ and jump distribution ν if

$$(3.20) \quad \mathbf{v} = \int_{\mathbb{R}^2} \frac{\lambda \mathbf{x}}{1 + \|\mathbf{x}\|^2} d\nu(\mathbf{x}),$$

$\mathbf{A} = \mathbf{0}$ and $\tau = \lambda\nu$, and in such a case its bi-free ϕ -transform is given as

$$(3.21) \quad \phi_\mu(z, w) = \lambda \int_{\mathbb{R}^2} \left[\frac{zw}{(z-s)(w-t)} - 1 \right] d\nu(\mathbf{x}).$$

If $\nu = \delta_{\mathbf{p}}$ with $\mathbf{p} \in \mathbb{R}^2 \setminus \{\mathbf{0}\}$, then μ is referred to as a *bi-free Poisson distribution* with rate λ and jump distribution $\delta_{\mathbf{p}}$.

4. ASYMPTOTIC BEHAVIORS OF BI-FREE CONVOLUTIONS

This section treats the asymptotic behavior of the measures

$$(4.22) \quad \mu_n := \mu_{n1} \boxplus \mu_{n2} \boxplus \cdots \boxplus \mu_{nk_n} \boxplus \delta_{\mathbf{v}_n},$$

where $\{k_n\}_{n=1}^\infty$ is a sequence of strictly increasing positive integers, $\{\mu_{nk}\}_{n \geq 1, 1 \leq k \leq k_n}$ is an infinitesimal triangular array in $\mathcal{P}_{\mathbb{R}^2}$, and $\mathbf{v}_n \in \mathbb{R}^2$. In order to cope with the problem, we begin with carrying out the investigation on the asymptotic behavior of the bi-free transforms of μ_{nk} . It turns out that they satisfy certain asymptotic property due to the infinitesimality of $\{\mu_{nk}\}_{n,k}$.

Let S be an unbounded subset of $\mathbb{C}$, and denote by $\mathcal{U}(S^d)$ the collection of triangular arrays of functions $\{\epsilon_{nk}\}_{n \geq 1, 1 \leq k \leq k_n}$ defined on S^d with the following asymptotic properties: the functions

$$\epsilon_n(z_1, \dots, z_d) = \max_{1 \leq k \leq k_n} |\epsilon_{nk}(z_1, \dots, z_d)|, \quad (z_1, \dots, z_d) \in S^d,$$

satisfy that $\lim_{n \rightarrow \infty} \epsilon_n(z_1, \dots, z_d) = 0$ for any $(z_1, \dots, z_d) \in S^d$ and $\epsilon_n(z_1, \dots, z_d) = o(1)$ uniformly in n as $z_1, \dots, z_d \rightarrow \infty$ with $(z_1, \dots, z_d) \in S^d$.

Lemma 4.1. *Let $\{\mu_{nk}\}_{n \geq 1, 1 \leq k \leq k_n} \subset \mathcal{P}_{\mathbb{R}^2}$ be infinitesimal. The following statements hold.*

- (1) *For any $\theta > 0$, there exists a number $M > 1$ so that each $\phi_{\mu_{nk}^{(j)}}$ is defined and satisfies the relation $\phi_{\mu_{nk}^{(j)}}(\bar{\xi}) = \overline{\phi_{\mu_{nk}^{(j)}}(\xi)}$ on $\Gamma := \Gamma_{\theta, M}$.*
- (2) *For $j = 1, 2$,*

$$\left\{ \frac{\phi_{\mu_{nk}^{(j)}}(\xi)}{\xi} \right\}_{n \geq 1, 1 \leq k \leq k_n} \in \mathcal{U}(\Gamma) \quad \text{and} \quad \{\phi_{\mu_{nk}}\}_{n \geq 1, 1 \leq k \leq k_n} \in \mathcal{U}(\Gamma^2).$$

Proof. For notational convenience, write $G_{nk} = G_{\mu_{nk}}$, $F_{jnk} = F_{\mu_{nk}^{(j)}}$, $\phi_{jnk} = \phi_{\mu_{nk}^{(j)}}$ and $\phi_{nk} = \phi_{\mu_{nk}}$ for all j, k, n . Let $\theta > 0$. Then the existence of the number $M > 1$ with the stated properties in (1) is guaranteed by the infinitesimality of $\{\mu_{nk}\}_{n,k}$ [4, Lemma 5]. The relation $\phi_{jnk}(\bar{z}) = \overline{\phi_{jnk}(z)}$ holds for $z \in \Gamma$ because $F_{jnk}(\bar{z}) = \overline{F_{jnk}(z)}$ for $z \in \mathbb{C} \setminus \mathbb{R}$.

Recall from [6, Proposition 2.3] that we can express ϕ_{1nk} as

$$(4.23) \quad \frac{\phi_{1nk}(z)}{z} = [1 + v_{nk}(z)] \int_{\mathbb{R}} \frac{s}{z - s} d\mu_{nk}^{(1)}(s),$$

where $\{v_{nk}\}_{n \geq 1, 1 \leq k \leq k_n} \in \mathcal{U}(\Gamma)$. Applying the techniques used in the proof of Proposition 2.1 to the sequence of functions

$$\epsilon_n(z) := \max_{1 \leq k \leq k_n} \left| \int_{\mathbb{R}} \frac{s}{z - s} d\mu_{nk}^{(1)}(s) \right|,$$

along with the infinitesimality of $\{\mu_{nk}\}_{n,k}$, yields that $\lim_{n \rightarrow \infty} \epsilon_n(z) = 0$ for $z \in \Gamma$ and $\epsilon_n(z) = o(1)$ uniformly in n as $z \rightarrow \infty$ with $z \in \Gamma$. Hence the triangular array $\{\phi_{1nk}(z)/z\}_{n,k}$, as well as $\{\phi_{2nk}(w)/w\}_{n,k}$, is shown to belong to $\mathcal{U}(\Gamma)$.

Finally, in order to show that $\{\phi_{nk}\}_{n \geq 1, 1 \leq k \leq k_n} \in \mathcal{U}(\Gamma^2)$, it suffices to show that $\{(H_{nk} - 1)/H_{nk}\}_{n \geq 1, 1 \leq k \leq k_n} \in \mathcal{U}(\Gamma^2)$, where $H_{nk}(z, w) = zwG_{nk}(F_{1nk}^{-1}(z), F_{2nk}^{-1}(w))$ for $(z, w) \in \Gamma^2$. This is equivalent to showing that $\{H_{nk} - 1\}_{n,k} \in \mathcal{U}(\Gamma^2)$. We further see from Proposition 2.2 that the condition $\{H_{nk} - 1\}_{n,k} \in \mathcal{U}(\Gamma^2)$ is the same as

$$(4.24) \quad \{zwG_{nk}(z, w) - 1\}_{n \geq 1, 1 \leq k \leq k_n} \in \mathcal{U}(\Gamma'^2),$$

where Γ' is some truncated cone in $\mathbb{C}$ on which $F_{\mu_{nk}^{(j)}}$ is univalent. Making use of the techniques in the proof of Proposition 2.1 again one can easily obtain (4.24). The proof is complete. $\square$

Let $L > 0$ be a fixed number. We will use the following functions and measures to study asymptotic properties of μ_n defined in (4.22). Let

$$(4.25) \quad \mathbf{v}_{nk} = \int_{\{\|\mathbf{x}\| < L\}} \mathbf{x} d\mu_{nk}(\mathbf{x})$$

and define a triangular array $\{\dot{\mu}_{nk}\}_{n \geq 1, 1 \leq k \leq k_n} \subset \mathcal{P}_{\mathbb{R}^2}$ as

$$(4.26) \quad \dot{\mu}_{nk}(B) = \mu_{nk}(B + \mathbf{v}_{nk})$$

for any Borel set $B \subset \mathbb{R}^2$. This shifted triangular array $\{\dot{\mu}_{nk}\}$ is also infinitesimal because $\max_{1 \leq k \leq k_n} \|\mathbf{v}_{nk}\| \rightarrow 0$ as $n \rightarrow \infty$. Further define finite positive Borel measures

$$(4.27) \quad \tau_n = \sum_{k=1}^{k_n} \dot{\mu}_{nk},$$

and functions

$$f_{1nk}(z) = \int_{\mathbb{R}^2} \frac{zs}{z-s} d\dot{\mu}_{nk}(s, t) \quad \text{and} \quad f_{2nk}(w) = \int_{\mathbb{R}^2} \frac{wt}{w-t} d\dot{\mu}_{nk}(s, t)$$

for $z, w \in \mathbb{C} \setminus \mathbb{R}$.

The following result, mostly taken from [6, Lemma 2.4, Lemma 3.1, Lemma 3.2], is one of the main ingredients of studying (4.22). For readers' convenience, we provide its proof here.

Lemma 4.2. *With the same notations μ_n , μ_{nk} and $\mathbf{v}_n$ in (4.22), Γ in Lemma 4.1, and τ_n and f_{jnk} defined above, the following statements hold.*

(1) *For $j = 1, 2$ and any fixed $i\ell \in \Gamma$, the sequence*

$$\left\{ \mathbf{v}_n^{(j)} + \sum_{k=1}^{k_n} [\mathbf{v}_{nk}^{(j)} + f_{jnk}(i\ell)] \right\}_{n=1}^{\infty}$$

converges if and only if the sequence $\{\phi_{\mu_n^{(j)}}(i\ell)\}_{n=1}^{\infty}$ converges, in which case they converge to the same value.

(2) *If*

$$V := \sup_{n \geq 1} \int_{\mathbb{R}^2} \frac{\|\mathbf{x}\|^2}{1 + \|\mathbf{x}\|^2} d\tau_n(\mathbf{x}) < \infty,$$

then there exists an $N \in \mathbb{N}$ such that for $|\ell| \geq 1$, $n \geq N$ and $j = 1, 2$, the inequality

$$\sum_{k=1}^{k_n} |f_{jnk}(i\ell)| \leq C_L V |\ell|$$

holds for some constant C_L depending only on L .

Proof. We only prove the assertions for $j = 1$; the proof for $j = 2$ is similar. Applying the formula (4.23) to the triangular array $\{\dot{\mu}_{nk}^{(1)}\}_{n,k}$, we have

$$(4.28) \quad \phi_{\mu_{nk}^{(1)}}(iy) - \mathbf{v}_{nk}^{(1)} = \phi_{\dot{\mu}_{nk}^{(1)}}(iy) = f_{1nk}(iy)[1 + v_{nk}(iy)]$$

for $iy \in \Gamma$. Then the desired result in (1) follows from [6, Lemma 2.4] by choosing $z_{nk} = \phi_{\mu_{nk}^{(1)}}(iy) - \mathbf{v}_{nk}^{(1)}$, $w_{nk} = f_{1nk}(iy)$ and $r_n = \mathbf{v}_n^{(1)} + \sum_{k=1}^{k_n} \mathbf{v}_{nk}^{(1)}$.

For assertion (2), define

$$b_{1nk}(y) = \mathbf{v}_{nk}^{(1)} \int_{\{\|\mathbf{x}\| \geq L\}} d\mu_{nk}(\mathbf{x}) + \int_{\{\|\mathbf{x} + \mathbf{v}_{nk}\| \geq L\}} \frac{y^2 s}{y^2 + s^2} d\dot{\mu}_{nk}(\mathbf{x}).$$

Observe that we have

$$\mathbf{v}_{nk}^{(1)} \int_{\{\|\mathbf{x}\| \geq L\}} d\mu_{nk}(\mathbf{x}) = \int_{\{\|\mathbf{x}\| < L\}} (s - \mathbf{v}_{nk}^{(1)}) d\mu_{nk}(\mathbf{x}) = \int_{\{\|\mathbf{x} + \mathbf{v}_{nk}\| < L\}} s d\dot{\mu}_{nk}(\mathbf{x})$$

and

$$f_{1nk}(iy) = \int_{\mathbb{R}^2} \frac{y^2 s}{y^2 + s^2} d\dot{\mu}_{nk}(\mathbf{x}) - i \int_{\mathbb{R}^2} \frac{ys^2}{y^2 + s^2} d\dot{\mu}_{nk}(\mathbf{x}).$$

288 This shows that for $|y| > 1$,

$$(4.29) \quad |\Re f_{1nk}(iy) - b_{1nk}(y)| = \left| \int_{\{\|\mathbf{x} + \mathbf{v}_{nk}\| < L\}} \frac{s^3}{y^2 + s^2} d\dot{\mu}_{nk}(\mathbf{x}) \right| \leq 2L |\Im f_{1nk}(iy)|,$$

289 where we use the fact that $|\mathbf{v}_{nk}^{(1)}| \leq L$ for any n and k in the last inequality.

290 Now let $N \in \mathbb{N}$ be big enough so that $\sup_{1 \leq k \leq k_n} \|\mathbf{v}_{nk}\| \leq L/2$ for all $n \geq N$. Then for $|y| > 1$
 291 and $n \geq N$ we have

$$(4.30) \quad \begin{aligned} \sum_{k=1}^{k_n} |b_{1nk}(y)| &\leq \sum_{k=1}^{k_n} \frac{L}{2} \int_{\{\|\mathbf{x}\| \geq L/2\}} d\dot{\mu}_{nk}(\mathbf{x}) + |y| \sum_{k=1}^{k_n} \int_{\{\|\mathbf{x}\| \geq L/2\}} \frac{|ys|}{y^2 + s^2} d\dot{\mu}_{nk}(\mathbf{x}) \\ &\leq |y| \sum_{k=1}^{k_n} \int_{\{\|\mathbf{x}\| \geq L/2\}} \frac{1+L}{2} d\dot{\mu}_{nk}(\mathbf{x}) \\ &= |y|(1+L)(4+L^2)(2L^2)^{-1} \int_{\{\|\mathbf{x}\| \geq L/2\}} \frac{L^2/4}{1+L^2/4} d\tau_n(\mathbf{x}) \\ &\leq |y|(1+L)(4+L^2)(2L^2)^{-1} \int_{\{\|\mathbf{x}\| \geq L/2\}} \frac{\|\mathbf{x}\|^2}{1+\|\mathbf{x}\|^2} d\tau_n(\mathbf{x}). \end{aligned}$$

292 Moreover, the estimate

$$(4.31) \quad \sup_{n \geq 1} \sum_{k=1}^{k_n} |\Im f_{1nk}(iy)| = |y| \sup_{n \geq 1} \sum_{k=1}^{k_n} \int_{\mathbb{R}^2} \frac{s^2}{y^2 + s^2} d\dot{\mu}_{nk}(s, t) \leq V|y|$$

293 holds true for $|y| > 1$. Combining (4.29), (4.30) and (4.31) yields assertion (2). $\square$

294 The following result provides an estimation for the bi-free ϕ -transform of measures μ_n in (4.22).

295 **Lemma 4.3.** *Adopt the notations μ_{nk} , $\dot{\mu}_{nk}$ and Γ in (4.22), (4.26) and Lemma 4.1, respectively,*
 296 *and let*

$$(4.32) \quad H_{nk}(z, w) = zw G_{\mu_{nk}} \left(F_{\mu_{nk}^{(1)}}^{-1}(z), F_{\mu_{nk}^{(2)}}^{-1}(w) \right), \quad (z, w) \in \Gamma^2.$$

297 Then $H_{nk} - 1$ can be expressed as

$$\epsilon_{1nk} \int_{\mathbb{R}^2} \frac{s}{z-s} d\dot{\mu}_{nk}(s, t) + \epsilon_{2nk} \int_{\mathbb{R}^2} \frac{t}{w-t} d\dot{\mu}_{nk}(s, t) + (1 + \epsilon_{nk}) \int_{\mathbb{R}^2} \frac{st}{(z-s)(w-t)} d\dot{\mu}_{nk}(s, t),$$

298 where $\{\epsilon_{1nk}(z, w)\}_{n,k}$, $\{\epsilon_{2nk}(z, w)\}_{n,k}$ and $\{\epsilon_{nk}(z, w)\}_{n,k}$ are triangular arrays of functions in $\mathcal{U}(\Gamma^2)$.

299 Consequently, $\{H_{nk} - 1\} \in \mathcal{U}(\Gamma^2)$.

300 *Proof.* For notational convenience, write $\dot{G}_{nk} = G_{\dot{\mu}_{nk}}$, $\dot{F}_{jnk} = F_{\dot{\mu}_{nk}^{(j)}}$ and $\dot{\phi}_{jnk} = \phi_{\dot{\mu}_{nk}^{(j)}}$ for any j , n
 301 and k . Then a simple argument of change of variables shows that

$$H_{nk}(z, w) = zw \dot{G}_{nk}(\dot{F}_{1nk}^{-1}(z), \dot{F}_{2nk}^{-1}(w)), \quad (z, w) \in \Gamma^2.$$

302 To conclude the proof, we first analyze the function

$$H_{nk}(z, w) - 1 = \int_{\mathbb{R}^2} \left[\frac{z}{\dot{\phi}_{1nk}(z) + z - s} \frac{w}{\dot{\phi}_{2nk}(w) + w - t} - 1 \right] d\dot{\mu}_{nk}(s, t).$$

303 Using the identity

$$\frac{\xi}{\dot{\phi}_{jnk}(\xi) + \xi - r} = \frac{\xi}{\xi - r} \left[1 - \frac{\dot{\phi}_{jnk}(\xi)}{\dot{F}_{jnk}^{-1}(\xi) - r} \right],$$

the function $H_{nk} - 1$ can be rewritten as the sum of functions I_{1nk} , I_{2nk} and I_{3nk} , where

$$I_{1nk}(z, w) = \int_{\mathbb{R}^2} \left[\frac{zw}{(z-s)(w-t)} - 1 \right] d\dot{\mu}_{nk}(s, t),$$

$$I_{2nk}(z, w) = - \int_{\mathbb{R}^2} \frac{zw}{(z-s)(w-t)} \left[\frac{\dot{\phi}_{1nk}(z)}{\dot{F}_{1nk}^{-1}(z) - s} + \frac{\dot{\phi}_{2nk}(w)}{\dot{F}_{2nk}^{-1}(w) - t} \right] d\dot{\mu}_{nk}(s, t),$$

and

$$I_{3nk}(z, w) = \int_{\mathbb{R}^2} \frac{zw}{(z-s)(w-t)} \frac{\dot{\phi}_{1nk}(z)}{\dot{F}_{1nk}^{-1}(z) - s} \frac{\dot{\phi}_{2nk}(w)}{\dot{F}_{2nk}^{-1}(w) - t} d\dot{\mu}_{nk}(s, t).$$

For any $z \in \Gamma$ with $|z|$ large enough, $s \in \mathbb{R}$, and n, k , Lemma 4.1 shows that

$$\left| \frac{\dot{F}_{1nk}^{-1}(z) - s}{z} \right| = \left| \frac{z-s}{z} + \frac{\dot{\phi}_{1nk}(z)}{z} \right| \geq \frac{|\Im z|}{|z|} - \left| \frac{\dot{\phi}_{1nk}(z)}{z} \right| \geq \frac{1}{\sqrt{1+\theta^2}} - \left| \frac{\dot{\phi}_{1nk}(z)}{z} \right| \geq c_\theta,$$

where $c_\theta = 1/(2\sqrt{1+\theta^2})$. This implies that

$$(4.33) \quad \frac{1}{\dot{F}_{1nk}^{-1}(z) - s} = \frac{1}{z-s} \left[1 - \frac{\dot{\phi}_{1nk}(z)}{\dot{F}_{1nk}^{-1}(z) - s} \right] = \frac{1}{z-s} [1 + \delta_{1nk}(z, s)],$$

where

$$\max_{1 \leq k \leq k_n} |\delta_{1nk}(z, s)| = \max_{1 \leq k \leq k_n} \left| \frac{\dot{\phi}_{1nk}(z)}{\dot{F}_{1nk}^{-1}(z) - s} \right| \leq c_\theta^{-1} \max_{1 \leq k \leq k_n} \left| \frac{\dot{\phi}_{1nk}(z)}{z} \right| := \delta_{1n}(z).$$

We further obtain from the estimate (4.33) that

$$(4.34) \quad \begin{aligned} & \max_{1 \leq k \leq k_n} \left| \int_{\mathbb{R}^2} \frac{z^2 w}{(z-s)(w-t)} \frac{1}{\dot{F}_{1nk}^{-1}(z) - s} d\dot{\mu}_{nk}(s, t) - 1 \right| \\ &= \max_{1 \leq k \leq k_n} \left| \int_{\mathbb{R}^2} \frac{z^2 w}{(z-s)^2(w-t)} [1 + \delta_{1nk}(z, s)] d\dot{\mu}_{nk}(s, t) - 1 \right| \\ &= \max_{1 \leq k \leq k_n} \left| \int_{\mathbb{R}^2} \left\{ \left[\frac{z^2 w}{(z-s)^2(w-t)} - 1 \right] [1 + \delta_{1nk}(z, s)] + \delta_{1nk}(z, s) \right\} d\dot{\mu}_{nk}(s, t) \right| \\ &\leq \delta_{1n}(z) + [1 + \delta_{1n}(z)] M_n(z, w), \end{aligned}$$

where

$$M_n(z, w) = \max_{1 \leq k \leq k_n} \int_{\mathbb{R}^2} \left| \frac{z^2 w}{(z-s)^2(w-t)} - 1 \right| d\dot{\mu}_{nk}(s, t).$$

Note that the function $\delta_{1n}(z) + [1 + \delta_{1n}(z)] M_n(z, w) = o(1)$ as $n \rightarrow \infty$ for $(z, w) \in \Gamma^2$ and uniformly in n as $z, w \rightarrow \infty$ with $(z, w) \in \Gamma^2$. Indeed, this can be easily obtained by using Lemma 4.1 and applying the techniques employed in the proof of Proposition 2.1 to the identity

$$\frac{z^2 w}{(z-s)^2(w-t)} - 1 = \frac{z}{z-s} \left[\frac{zw}{(z-s)(w-t)} - 1 \right] + \frac{s}{z-s}.$$

Now applying the formula (4.23) to the triangular array $\{\dot{\mu}_{nk}\}_{n \geq 1, 1 \leq k \leq k_n}$ and using (4.34) give

$$\begin{aligned} \int_{\mathbb{R}^2} \frac{zw}{(z-s)(w-t)} \frac{\dot{\phi}_{1nk}(z)}{\dot{F}_{1nk}^{-1}(z) - s} d\dot{\mu}_{nk}(s, t) &= \frac{\dot{\phi}_{1nk}(z)}{z} \int_{\mathbb{R}^2} \frac{z^2 w}{(z-s)(w-t)} \frac{d\dot{\mu}_{nk}(s, t)}{\dot{F}_{1nk}^{-1}(z) - s} \\ &= [1 + v_{1nk}(z, w)] \int_{\mathbb{R}^2} \frac{s}{z-s} d\dot{\mu}_{nk}(s, t), \end{aligned}$$

where $\{v_{1nk}\}_{n,k}$ is a triangular array in $\mathcal{U}(\Gamma^2)$. Similarly, the identity

$$\int_{\mathbb{R}^2} \frac{zw}{(z-s)(w-t)} \frac{\dot{\phi}_{2nk}(w)}{\dot{F}_{2nk}^{-1}(w) - t} d\dot{\mu}_{nk}(s, t) = [1 + v_{2nk}(z, w)] \int_{\mathbb{R}^2} \frac{t}{w-t} d\dot{\mu}_{nk}(s, t)$$

is valid for some triangular array $\{v_{2nk}\}_{n,k} \in \mathcal{U}(\Gamma^2)$. By similar arguments, one can also show that

$$I_{3nk} = v_{3nk} \int_{\mathbb{R}^2} \frac{s}{z-s} d\dot{\mu}_{nk}(s, t) + v_{4nk} \int_{\mathbb{R}^2} \frac{t}{w-t} d\dot{\mu}_{nk}(s, t) + v_{5nk} \int_{\mathbb{R}^2} \frac{st}{(z-s)(w-t)} d\dot{\mu}_{nk}(s, t)$$

for some triangular arrays $\{v_{3nk}\}_{n,k}$, $\{v_{4nk}\}_{n,k}$ and $\{v_{5nk}\}_{n,k}$ in $\mathcal{U}(\Gamma^2)$. Finally, let $\epsilon_{1nk} = v_{3nk} - v_{1nk}$, $\epsilon_{2nk} = v_{4nk} - v_{2nk}$, and $\epsilon_{nk} = v_{5nk}$. Then we conclude the proof of the first assertion by using the integral representations of I_{2nk} and I_{3nk} provided above and the identity

$$I_{1nk}(z, w) = \int_{\mathbb{R}^2} \left[\frac{s}{z-s} + \frac{t}{w-t} + \frac{st}{(z-s)(w-t)} \right] d\dot{\mu}_{nk}(s, t).$$

The fact that $\{H_{nk} - 1\} \in \mathcal{U}(\Gamma^2)$ can be proved by the infinitesimality of $\{\dot{\mu}_{nk}\}$ and the techniques in Proposition 2.1. This finishes the proof. $\square$

Lemma 4.4. *Suppose that the marginal laws of μ_n in (4.22) converge weakly. With the notations $\dot{\mu}_{nk}$, τ_n , Γ as before and H_{nk} as in (4.32), the following statements hold.*

(1) *The positive planar measures $\{\sigma_{1n}\}_{n=1}^\infty$ and $\{\sigma_{2n}\}_{n=1}^\infty$ defined as*

$$\sigma_{1n} = \frac{s^2}{1+s^2} \tau_n \quad \text{and} \quad \sigma_{2n} = \frac{t^2}{1+t^2} \tau_n$$

are uniformly bounded and tight.

(2) *For $(iy, iv) \in \Gamma^2$, the limit*

$$(4.35) \quad \lim_{n \rightarrow \infty} \sum_{k=1}^{k_n} \frac{H_{nk}(iy, iv) - 1}{H_{nk}(iy, iv)}$$

exists if and only if the limit

$$(4.36) \quad \lim_{n \rightarrow \infty} \sum_{k=1}^{k_n} \int_{\mathbb{R}^2} \frac{st}{(iy-s)(iv-t)} d\dot{\mu}_{nk}(s, t)$$

exists, in which case these two limits are equal.

(3) *The function*

$$\sum_{k=1}^{k_n} \frac{H_{nk}(iy, iv) - 1}{H_{nk}(iy, iv)} = o(1)$$

uniformly in n as $|y|, |v| \rightarrow \infty$ if and only if

$$(4.37) \quad \sum_{k=1}^{k_n} \int_{\mathbb{R}^2} \frac{st}{(iy-s)(iv-t)} d\dot{\mu}_{nk}(s, t) = o(1)$$

uniformly in n as $|y|, |v| \rightarrow \infty$.

Proof. For any $\epsilon > 0$, choose a large positive number $y_0 > 1$ so that $|\phi_{\mu_n^{(1)}}(iy_0)| < \epsilon y_0$ for all n by Proposition 2.3. Then we deduce from Lemma 4.2(1) and the identity

$$\int_{\mathbb{R}^2} \frac{s^2}{y_0^2 + s^2} d\tau_n(s, t) = -\frac{1}{y_0} \Im \left(\mathbf{v}_n^{(1)} + \sum_{k=1}^{k_n} [\mathbf{v}_{nk}^{(1)} + f_{1nk}(iy_0)] \right)$$

the existence of a large number $N \in \mathbb{N}$ so that

$$\int_{\mathbb{R}^2} \frac{s^2}{y_0^2 + s^2} d\tau_n(s, t) < 2\epsilon, \quad n \geq N.$$

This, along with the inequalities

$$\frac{s^2}{1 + s^2} \leq \frac{y_0^2 s^2}{y_0^2 + s^2} \quad \text{and} \quad \frac{s^2}{1 + s^2} \leq \frac{2s^2}{y_0^2 + s^2}$$

which hold true for $s \in \mathbb{R}$ and $|s| \geq y_0$, respectively, yields the uniform boundedness and tightness of $\{\sigma_{1n}\}$. Similarly, $\{\sigma_{2n}\}$ is uniformly bounded and tight. This proves (1).

To prove (2), we first argue that the limit in (4.36) exists if and only if so does the limit

$$(4.38) \quad \lim_{n \rightarrow \infty} \sum_{k=1}^{k_n} [H_{nk}(iy, iv) - 1],$$

and show that they are equal. We shall use the integral representation of $H_{nk} - 1$ given in Lemma 4.3 to accomplish this goal. Observe first that the quantity V defined in Lemma 4.2(2) is finite by the established result (1). Hence for $(iy, iv) \in \Gamma^2$ and all large n , we have

$$\begin{aligned} \left| \sum_{k=1}^{k_n} \epsilon_{1nk}(iy, iv) \int_{\mathbb{R}^2} \frac{s}{iy - s} d\mu_{nk}(s, t) \right| &\leq \left[\max_{1 \leq k \leq k_n} |\epsilon_{1nk}(iy, iv)| \right] \frac{1}{|y|} \sum_{k=1}^{k_n} |f_{1nk}(iy)| \\ &\leq C_L V \max_{1 \leq k \leq k_n} |\epsilon_{1nk}(iy, iv)|, \end{aligned}$$

which yields that

$$\lim_{n \rightarrow \infty} \sum_{k=1}^{k_n} \epsilon_{1nk}(iy, iv) \int_{\mathbb{R}^2} \frac{s}{iy - s} d\mu_{nk}(s, t) = 0,$$

as well as

$$\lim_{n \rightarrow \infty} \sum_{k=1}^{k_n} \epsilon_{2nk}(iy, iv) \int_{\mathbb{R}^2} \frac{t}{iv - t} d\mu_{nk}(s, t) = 0$$

by similar arguments. Next notice that the inequality

$$\left| \frac{st}{(iy - s)(iv - t)} \right| \leq \frac{2s^2}{1 + s^2} + \frac{2t^2}{1 + t^2}, \quad |y|, |v| \geq 1, \quad s, t \in \mathbb{R},$$

along with the established result (1), allows us to obtain that

$$\lim_{n \rightarrow \infty} \sum_{k=1}^{k_n} \epsilon_{nk}(iy, iv) \int_{\mathbb{R}^2} \frac{st}{(iy - s)(iv - t)} d\mu_{nk}(s, t) = 0$$

for any point $(iy, iv) \in \Gamma^2$. Hence we have proved that the pointwise convergence of (4.36) is equivalent to that of (4.38), and both limits are the same. These discussions also indicate that $\sum_{k=1}^{k_n} |H_{nk}(iy, iv) - 1|$ is uniformly bounded in n for fixed $(iy, iv) \in \Gamma^2$. Since $\{H_{nk} - 1\}_{nk} \in \mathcal{U}(\Gamma^2)$, one can see that (4.35) converges pointwise if and only if so does (4.38), and they have the same limit. This yields assertion (2). The preceding discussions with a little effort yield assertion (3). $\square$

Before stating the main theorem of this section, let us introduce the following conditions, which play an important role in the asymptotic problem under investigation.

Condition 4.5. Let $\{\tau_n\}_{n=1}^\infty$ be a sequence of finite positive Borel measures on $\mathbb{R}^2$.

(I) The sequences of measures

$$\sigma_{1n} = \frac{s^2}{1+s^2} \tau_n \quad \text{and} \quad \sigma_{2n} = \frac{t^2}{1+t^2} \tau_n$$

converge weakly to finite positive Borel measures σ_1 and σ_2 on $\mathbb{R}^2$, respectively.

(II) The limit

$$\gamma := \lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} \frac{st}{(1+s^2)(1+t^2)} d\tau_n(s, t)$$

exists in $\mathbb{R}$.

Theorem 4.6. *Let $\{\mathbf{v}_n\}_{n=1}^\infty$ be a sequence of vectors in $\mathbb{R}^2$, $\{k_n\}_{n=1}^\infty$ a sequence of strictly increasing positive integers, and let $\{\mu_{nk}\}_{n \geq 1, 1 \leq k \leq k_n}$ be an infinitesimal triangular array in $\mathcal{P}_{\mathbb{R}^2}$. Following the notations in (4.25), (4.26) and (4.27), the following statements are equivalent.*

(1) The sequence

$$\mu_n := \mu_{n1} \boxplus \mu_{n2} \boxplus \cdots \boxplus \mu_{nk_n} \boxplus \delta_{\mathbf{v}_n}$$

converges weakly to some planar probability measure $\mu_{\boxplus}$.

(2) Condition 4.5(I) and (II) are satisfied, and the sequence

$$(4.39) \quad \mathbf{v}_n + \sum_{k=1}^{k_n} \left[\mathbf{v}_{nk} + \int_{\mathbb{R}^2} \frac{\mathbf{x}}{1 + \|\mathbf{x}\|^2} d\mu_{nk}(\mathbf{x}) \right]$$

of vectors in $\mathbb{R}^2$ converges to some vector $\mathbf{v}$.

Proof. We take the set Γ given in Lemma 4.1. Suppose that assertion (1) holds. By Lemma 4.4(1), let $\sigma_{1j_n} \Rightarrow \sigma_1$ and $\sigma_{2j_n} \Rightarrow \sigma_2$ for some subsequences $\{\sigma_{1j_n}\}, \{\sigma_{2j_n}\}$ and for some finite positive Borel measures σ_1, σ_2 . Observe next that the limit in (4.36) exists. Denote this limit by $\tilde{K}(iy, iv)$. Then using the decomposition (3.17) we see that the limit

$$\gamma' := \lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} \frac{st}{(1+s^2)(1+t^2)} d\tau_{j_n}(s, t)$$

must exist and $\tilde{K}$ has an analytic extension (still denoted by $\tilde{K}$) to $(\mathbb{C} \setminus \mathbb{R})^2$. More precisely, the analytic extension $K(z, w) := zw\tilde{K}(z, w)$ can be expressed as the sum of integrals:

$$K(z, w) = \gamma' + \int_{\mathbb{R}^2} \frac{1+zs}{z-s} \left[\frac{t}{1+t^2} + \frac{1+wt}{w-t} \frac{t^2}{1+t^2} \right] d\sigma_1(s, t) + \int_{\mathbb{R}^2} \frac{1+wt}{w-t} \frac{s}{1+s^2} d\sigma_2(s, t).$$

A simple calculation then shows that for any $z = x + iy$ and $w = u + iv$ in $\mathbb{C}^+$,

$$-\frac{1}{2} \Re[K(z, w) - K(\bar{z}, w)] = \int_{\mathbb{R}^2} \frac{yv}{[(s-x)^2 + y^2][(t-u)^2 + v^2]} (1+s^2)t^2 d\sigma_1(s, t).$$

This identity, of course, is also valid for any other weak-limit point σ'_1 of $\{\sigma_{1n}\}$. Combining this result with the Stieltjes inversion formula for two variables shows that

$$(4.40) \quad \frac{t^2}{1+t^2} \sigma_1 = \frac{t^2}{1+t^2} \sigma'_1.$$

On the other hand, according to Lemma 4.2(1) the sequence

$$(4.41) \quad \gamma_{1n} := \mathbf{v}_n^{(1)} + \sum_{k=1}^{k_n} \left[\mathbf{v}_{nk}^{(1)} + \int_{\mathbb{R}^2} \frac{s}{1+s^2} d\mu_{nk}(s, t) \right],$$

converges to some γ_1 and

$$\phi_{\mu_{\boxplus}^{(1)}}(z) = \gamma_1 + \int_{\mathbb{R}^2} \frac{1+zs}{z-s} d\sigma_1(s, t), \quad z \in \mathbb{C} \setminus \mathbb{R}.$$

Hence $\mu_{\boxplus}^{(1)}$ is $\boxplus$ -infinitely divisible and $(\gamma_1, \sigma_1^{(1)})$, as well as $(\gamma_1, \sigma_1'^{(1)})$, is the free generating pair for $\mu_{\boxplus}^{(1)}$ [3]. Therefore we obtain that $\sigma_1^{(1)} = \sigma_1'^{(1)}$. This with (4.40) shows that $\sigma_1 = \sigma_1'$ by [14, Lemma 3.10]. We conclude that σ_1 is the unique weak-limit point of $\{\sigma_{1n}\}_{n=1}^\infty$. Similarly, $\{\sigma_{2n}\}$ has only one weak-limit point. Hence Condition 4.5(I) and 4.5(II) are satisfied. Moreover, the identity

$$\frac{s}{1+s^2} - \frac{s}{1+s^2+t^2} = \frac{st^2}{(1+s^2+t^2)(1+s^2)}$$

shows that the vector defined in (4.39) converges. Hence the proof of (1) $\Rightarrow$ (2) is complete.

Conversely, suppose that assertion (2) holds. Then $\mathbf{v}_n^{(j)} + \sum_{k=1}^{k_n} [\mathbf{v}_{nk}^{(j)} + f_{jnk}(i\ell)]$ converges as $n \rightarrow \infty$ for any $i\ell \in \Gamma$ and $j = 1, 2$, and hence so does $\phi_{\mu_n^{(j)}}(i\ell)$ by Lemma 4.2(1). Employing the identity (3.17) gives that the limit (4.36) must exist, from which we see that ϕ_{μ_n} converges pointwise on Γ^2 by Lemma 4.4(2). To finish the proof of (2) $\Rightarrow$ (1), it remains to show that $\phi_{\mu_n^{(1)}}(iy) = o(|y|)$, $\phi_{\mu_n^{(2)}}(iv) = o(|v|)$ and $\sum_{k=1}^{k_n} [H_{nk}(iy, iv) - 1]/H_{nk}(iy, iv) = o(1)$ uniformly in n as $|y|, |v| \rightarrow \infty$ by Proposition 2.3. First of all, the identities (3.17) and (4.28) show that

$$(4.42) \quad \phi_{\mu_n^{(1)}}(iy) = \gamma_{1n} + \int_{\mathbb{R}^2} \frac{1+iy s}{iy-s} d\sigma_{1n}(s, t) + \sum_{k=1}^{k_n} v_{nk}(iy) f_{1nk}(iy),$$

where γ_{1n} is defined as in (4.41) and $v_{nk} \in \mathcal{U}(\Gamma)$. Since $\{\gamma_{1n}\}$ and $\{y^{-1} \sum_{k=1}^{k_n} |f_{1nk}(iy)| : iy \in \Gamma\}$ are bounded for all large n by the hypotheses in (2) and Lemma 4.2(2), it suffices to show that the second term in (4.42) equals $o(|y|)$ uniformly in n as $|y| \rightarrow \infty$. Notice that for any $r > 0$ and $|y| \geq 1$, we have

$$\begin{aligned} \frac{1}{|y|} \int_{\mathbb{R}^2} \left| \frac{1+iy s}{iy-s} \right| d\sigma_{1n}(s, t) &\leq \frac{1}{|y|} \int_{\{\|\mathbf{x}\| \leq r\}} \frac{1+|sy|}{\sqrt{y^2+s^2}} d\sigma_{1n}(s, t) + \sigma_{1n}(\{\|\mathbf{x}\| > r\}) \\ &\leq \frac{1}{|y|^2} (1+r|y|) \sup_n \sigma_{1n}(\mathbb{R}^2) + \sigma_{1n}(\{\|\mathbf{x}\| > r\}), \end{aligned}$$

which yields the desired result by the uniform boundedness and tightness of $\{\sigma_{1n}\}$. Similarly, $\phi_{\mu_n^{(2)}}(iv) = o(|v|)$ uniformly in n as $|v| \rightarrow \infty$. The last desired result is equivalent to the uniform convergence of (4.37) in n as $|y|, |v| \rightarrow \infty$ by Lemma 4.4(3). The latter uniform convergence can be proved by using Condition 4.5(II) and applying the techniques shown above to the integral in (4.37), which is rewritten as

$$-\frac{1}{yv} \sum_{k=1}^{k_n} \int_{\mathbb{R}^2} \left[\frac{1+iy s}{iy-s} \frac{s^2}{1+s^2} + \frac{s}{1+s^2} \right] \left[\frac{1+iv t}{iv-t} \frac{t^2}{1+t^2} + \frac{t}{1+t^2} \right] d\dot{\mu}_{nk}(s, t).$$

This finishes the proof of (2) $\Rightarrow$ (1). □

Let $\mathbf{u}_n$ be the vector defined in (4.39). From the proof of Theorem 4.6, one can see that for large $|y|$, $\phi_{\mu_n^{(1)}}(iy)$ can also be expressed as

$$\phi_{\mu_n^{(1)}}(iy) = \mathbf{u}_n^{(1)} + \sum_{k=1}^{k_n} \left[(1+v_{nk}(iy)) f_{1nk}(iy) - \int_{\mathbb{R}^2} \frac{s}{1+s^2+t^2} d\dot{\mu}_{nk}(s, t) \right].$$

390 A similar expression also holds for $\phi_{\mu_n^{(2)}}(iv)$ when $|v|$ is large. Since

$$\phi_{\mu_{\boxplus\boxplus}}(iy, iv) = \lim_{n \rightarrow \infty} \left[\frac{\phi_{\mu_n^{(1)}}(iy)}{iy} + \frac{\phi_{\mu_n^{(2)}}(iv)}{iv} + \sum_{k=1}^{k_n} \int_{\mathbb{R}^2} \frac{st}{(iy-s)(iv-t)} d\dot{\mu}_{nk}(s, t) \right],$$

391 a simple calculation shows that

$$(4.43) \quad \phi_{\mu_{\boxplus\boxplus}}(iy, iv) = \frac{\mathbf{v}^{(1)}}{iy} + \frac{\mathbf{v}^{(2)}}{iv} + \mathcal{S}(iy, iv)$$

392 for $(iy, iv) \in \Gamma^2$, where

$$\mathcal{S}(iy, iv) = \lim_{n \rightarrow \infty} \sum_{k=1}^{k_n} \int_{\mathbb{R}^2} \left[\frac{(iy)(iv)}{(iy-s)(iv-t)} - 1 - \frac{(iy)^{-1}s + (iv)^{-1}t}{1+s^2+t^2} \right] d\dot{\mu}_{nk}(s, t).$$

393 In the next section we shall use (4.43) to show that the function $\phi_{\mu_{\boxplus\boxplus}}$ extends analytically to
 394 $(\mathbb{C} \setminus \mathbb{R})^2$ and the analytic extension admits an integral representation of the form (3.16). As a
 395 consequence of Theorem 3.1, $\mu_{\boxplus\boxplus}$ is bi-freely infinitely divisible.

5. TRANSFER PRINCIPLE FOR LIMIT THEOREMS AND BIJECTION BETWEEN $\mathcal{ID}(\ast)$ AND $\mathcal{ID}(\boxplus\boxplus)$

398 This section is mainly devoted to studying the relation between the sets $\mathcal{ID}(\ast)$ and $\mathcal{ID}(\boxplus\boxplus)$ in
 399 terms of classical and bi-free limit theorems. We first introduce another type of convergence on
 400 the set of positive Borel measures on $\mathbb{R}^2$.

401 **Definition 5.1.** Denote by $\mathcal{M}_{\mathbb{R}^2}^0$ the set of positive Borel (not necessarily finite) measures τ on $\mathbb{R}^2$
 402 for which $\tau(B) < \infty$ for any Borel set $B \subset \mathbb{R}^2$ bounded away from zero, i.e. $\inf_{x \in B} \|x\| > 0$. For a
 403 measure τ and a sequence of measures $\{\tau_n\}_{n=1}^\infty$ in $\mathcal{M}_{\mathbb{R}^2}^0$, the convergent situation that $\tau_n(B) \rightarrow \tau(B)$
 404 for any Borel set B which is bounded away from zero and satisfies $\tau(\partial B) = 0$ is denoted by $\tau_n \Rightarrow_{\mathbf{0}} \tau$.

405 We remark here that any finite positive Borel measure on $\mathbb{R}^2$ belongs to $\mathcal{M}_{\mathbb{R}^2}^0$. Also note that
 406 the limiting measure τ in the convergence $\tau_n \Rightarrow_{\mathbf{0}} \tau$ is not necessarily unique since an arbitrary
 407 mass at $\mathbf{0}$ can be added to it. Portmanteau theorem for measures in $\mathcal{M}_{\mathbb{R}^2}^0$ is stated below (see [1]).

408 **Proposition 5.2.** Given $\{\tau_n\}_{n=1}^\infty$ and τ in $\mathcal{M}_{\mathbb{R}^2}^0$, the following are equivalent:

- 409 (1) $\tau_n \Rightarrow_{\mathbf{0}} \tau$;
- 410 (2) for any bounded and continuous function f on $\mathbb{R}^2$ with support bounded away from zero,

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} f d\tau_n = \int_{\mathbb{R}^2} f d\tau;$$

- 411 (3) for any bounded and continuous function f on $\mathbb{R}^2$ and for any Borel set $B \subset \mathbb{R}^2$ which is
 412 bounded away from zero and satisfies $\tau(\partial B) = 0$,

$$\lim_{n \rightarrow \infty} \int_B f d\tau_n = \int_B f d\tau;$$

- 413 (4) for every closed subset C and open subset O of $\mathbb{R}^2$ that are bounded away from zero,

$$\limsup_{n \rightarrow \infty} \tau_n(C) \leq \tau(C) \quad \text{and} \quad \liminf_{n \rightarrow \infty} \tau_n(O) \geq \tau(O).$$

We next introduce two conditions that are used in characterizing the classical limit theorem in multidimensional spaces (see [16]).

Condition 5.3. Let $\{\tau_n\}_{n=1}^\infty$ be a sequence of measures in $\mathcal{M}_{\mathbb{R}^2}^0$.

(III) $\tau_n \Rightarrow_{\mathbf{0}} \tau$ for some measure $\tau \in \mathcal{M}_{\mathbb{R}^2}^0$ with $\tau(\{\mathbf{0}\}) = 0$;

(IV) for every vector $\mathbf{u} \in \mathbb{R}^2$, the limits

$$(5.44) \quad \lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} \langle \mathbf{u}, \mathbf{x} \rangle^2 d\tau_n(\mathbf{x}) \quad \text{and} \quad \lim_{\epsilon \rightarrow 0^+} \liminf_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} \langle \mathbf{u}, \mathbf{x} \rangle^2 d\tau_n(\mathbf{x})$$

exist (as finite numbers), and they are equal.

In the following, we show the equivalence between Condition 4.5 and Condition 5.3, which will play an important role in clarifying the relation between $\mathcal{ID}(\ast)$ and $\mathcal{ID}(\boxplus\boxplus\boxplus)$.

Lemma 5.4. Let $\{\tau_n\}_{n=1}^\infty$ be a sequence of finite positive Borel measures on $\mathbb{R}^2$. Then (I) and (II) in Condition 4.5 hold if and only if (III) and (IV) in Condition 5.3 are satisfied, in which case

$$(5.45) \quad c = \gamma - \int_{\mathbb{R}^2} \frac{st}{(1+s^2)(1+t^2)} d\tau(s, t),$$

is a finite number, the matrix

$$(5.46) \quad \mathbf{A} = \begin{pmatrix} \sigma_1(\{\mathbf{0}\}) & c \\ c & \sigma_2(\{\mathbf{0}\}) \end{pmatrix}$$

is positive semi-definite and the limits in (5.44) define a non-negative quadratic form $\langle \mathbf{A}\mathbf{u}, \mathbf{u} \rangle$.

Proof. Suppose that (I) and (II) in Condition 4.5 are satisfied. With σ_1 and σ_2 at hand, one can define the positive measure τ as in (3.15) with $\tau(\{\mathbf{0}\})$. Then the relation

$$\frac{t^2}{1+t^2} \sigma_1 = \frac{s^2}{1+s^2} \sigma_2,$$

which is obtained from the definition of σ_{1n} and σ_{2n} , ensures that τ is well defined. It is also easy to verify that $\tau(\{\|\mathbf{x}\| \geq \epsilon\}) < \infty$ for any $\epsilon > 0$, whence $\tau \in \mathcal{M}_{\mathbb{R}^2}^0$. Now we claim $\tau_n \Rightarrow_{\mathbf{0}} \tau$. Pick any bounded and continuous function f on $\mathbb{R}^2$ whose support is contained in $\{\|\mathbf{x}\| \geq r\}$ for some $r > 0$. This induces two bounded and continuous functions on $\mathbb{R}^2$ defined as

$$f_1(\mathbf{x}) = \frac{\text{dist}(\mathbf{x}, U_1)}{\text{dist}(\mathbf{x}, U_1) + \text{dist}(\mathbf{x}, U_2)} f(\mathbf{x}) \quad \text{and} \quad f_2(\mathbf{x}) = \frac{\text{dist}(\mathbf{x}, U_2)}{\text{dist}(\mathbf{x}, U_1) + \text{dist}(\mathbf{x}, U_2)} f(\mathbf{x})$$

for $\mathbf{x} \in (U_1 \cap U_2)^c$, and $f_1(\mathbf{x}) = 0 = f_2(\mathbf{x})$ for $\mathbf{x} \in U_1 \cap U_2$, where $U_1 = \{\mathbf{x} : |\mathbf{x}^{(1)}| \leq r/2\}$ and $U_2 = \{\mathbf{x} : |\mathbf{x}^{(2)}| \leq r/2\}$. Clearly, $f = f_1 + f_2$, and the supports of f_1 and f_2 are bounded away from the s - and t -axis, respectively. Then the weak convergence of $\{\sigma_{1n}\}$ and $\{\sigma_{2n}\}$ yields that

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} f(s, t) d\tau_n(s, t) &= \lim_{n \rightarrow \infty} \left(\int_{\mathbb{R}^2} f_1(s, t) d\tau_n(s, t) + \int_{\mathbb{R}^2} f_2(s, t) d\tau_n(s, t) \right) \\ &= \lim_{n \rightarrow \infty} \left(\int_{\mathbb{R}^2} f_1(s, t) \frac{1+s^2}{s^2} d\sigma_{1n}(s, t) + \int_{\mathbb{R}^2} f_2(s, t) \frac{1+t^2}{t^2} d\sigma_{2n}(s, t) \right) \\ &= \int_{\mathbb{R}^2} f_1(s, t) \frac{1+s^2}{s^2} d\sigma_1(s, t) + \int_{\mathbb{R}^2} f_2(s, t) \frac{1+t^2}{t^2} d\sigma_2(s, t) \\ &= \int_{\mathbb{R}^2} f d\tau, \end{aligned}$$

432 which verifies $\tau_n \Rightarrow_0 \tau$.

433 To verify the statement (IV), it suffices to prove the existence of the following limits and the
434 equalities:

$$(5.47) \quad a := \lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} s^2 d\tau_n(s, t) = \lim_{\epsilon \rightarrow 0^+} \liminf_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} s^2 d\tau_n(s, t),$$

435

$$(5.48) \quad b := \lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} t^2 d\tau_n(s, t) = \lim_{\epsilon \rightarrow 0^+} \liminf_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} t^2 d\tau_n(s, t),$$

436 and

$$(5.49) \quad c := \lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} st d\tau_n(s, t) = \lim_{\epsilon \rightarrow 0^+} \liminf_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} st d\tau_n(s, t).$$

437 First of all, the limits

$$\lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} \frac{s^2}{1 + s^2} d\tau_n(s, t) = \lim_{\epsilon \rightarrow 0^+} \liminf_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} \frac{s^2}{1 + s^2} d\tau_n(s, t)$$

exist and equal $\sigma_1(\{\mathbf{0}\})$ by the weak convergence of σ_{1n} to σ_1 . For any $\epsilon > 0$, picking an $\epsilon' \in [\epsilon, 2\epsilon]$ so that $\sigma_1(\{\|\mathbf{x}\| = \epsilon'\}) = 0$ (such an ϵ' exists because σ_1 is a finite measure), we then have

$$\begin{aligned} \limsup_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} \left(s^2 - \frac{s^2}{1 + s^2} \right) d\tau_n(s, t) &= \limsup_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} s^2 d\sigma_{1n}(s, t) \\ &\leq \limsup_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon'} s^2 d\sigma_{1n}(s, t) \\ &= \int_{\|\mathbf{x}\| < \epsilon'} s^2 d\sigma_1(s, t) \leq \epsilon'^2 \sigma_1(\mathbb{R}^2) \rightarrow 0 \text{ as } \epsilon \rightarrow 0^+. \end{aligned}$$

438 Hence (5.47) holds and $a = \sigma_1(\{\mathbf{0}\})$. Similarly, (5.48) holds true and $b = \sigma_2(\{\mathbf{0}\})$.

439 One can also show that the existence of the limits in (5.49) is equivalent to that of the limits

$$(5.50) \quad \lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} \frac{st d\tau_n(s, t)}{(1 + s^2)(1 + t^2)} = \lim_{\epsilon \rightarrow 0^+} \liminf_{n \rightarrow \infty} \int_{\|\mathbf{x}\| < \epsilon} \frac{st d\tau_n(s, t)}{(1 + s^2)(1 + t^2)},$$

440 and all limits are the same if they exist. Next notice that $1 \wedge \|\mathbf{x}\|^2 \in L^1(\tau)$ according to the
441 definition of τ , and therefore

$$(5.51) \quad \int_{\mathbb{R}^2} \frac{|st|}{(1 + s^2)(1 + t^2)} d\tau(s, t) \leq \frac{1}{2} \int_{\mathbb{R}^2} \frac{s^2 + t^2}{(1 + s^2)(1 + t^2)} d\tau(s, t) < \infty.$$

442 We now show that the limits in (5.50) do exist and that the relation of c and γ in (5.45) holds.

443 For any $\epsilon > 0$, choose an $\epsilon' \in (\epsilon, 2\epsilon]$ with the property $\tau(\{\|\mathbf{x}\| = \epsilon'\}) = 0$ (such an ϵ' exists because

444 $\|\mathbf{x}\|^2 \chi_{\{\|\mathbf{x}\| \leq 1\}} \tau$ is a finite positive measure). Consider the difference

$$I_n(\epsilon) := \int_{\mathbb{R}^2} \frac{st d\tau_n(s, t)}{(1 + s^2)(1 + t^2)} - \int_{\mathbb{R}^2} \frac{st d\tau(s, t)}{(1 + s^2)(1 + t^2)} - \int_{\{\|\mathbf{x}\| < \epsilon\}} \frac{st d\tau_n(s, t)}{(1 + s^2)(1 + t^2)}$$

445 and decompose it into the sum of $J_{1n}(\epsilon)$ and $J_2(\epsilon)$, where

$$J_{1n}(\epsilon) = \int_{\{\|\mathbf{x}\| \geq \epsilon\}} \frac{st}{(1 + s^2)(1 + t^2)} d\tau_n(s, t) - \int_{\{\|\mathbf{x}\| \geq \epsilon\}} \frac{st}{(1 + s^2)(1 + t^2)} d\tau(s, t)$$

446 and

$$J_2(\epsilon) = - \int_{\{\|\mathbf{x}\| < \epsilon\}} \frac{st}{(1 + s^2)(1 + t^2)} d\tau(s, t).$$

Denoting by C_ϵ the closed set $\{\epsilon \leq \|\mathbf{x}\| \leq \epsilon'\}$, we have

$$\begin{aligned}
\lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} \left| \int_{C_\epsilon} \frac{st}{(1+s^2)(1+t^2)} d\tau_n \right| &\leq \lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} \int_{C_\epsilon} \frac{|st|}{\sqrt{1+s^2}\sqrt{1+t^2}} d\tau_n \\
&\leq \lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} \left(\int_{C_\epsilon} \frac{s^2}{1+s^2} d\tau_n \right)^{1/2} \left(\int_{C_\epsilon} \frac{t^2}{1+t^2} d\tau_n \right)^{1/2} \\
&= \lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} \sigma_{1n}(C_\epsilon)^{1/2} \sigma_{2n}(C_\epsilon)^{1/2} \\
&\leq \lim_{\epsilon \rightarrow 0^+} \sigma_1(C_\epsilon)^{1/2} \sigma_2(C_\epsilon)^{1/2} = 0,
\end{aligned}$$

where the Cauchy-Schwarz inequality was used in the second inequality and the assumption that $\sigma_{jn} \Rightarrow \sigma_j$ for $j = 1, 2$ was used in the last one. Moreover, by Proposition 5.2 we have

$$\lim_{n \rightarrow \infty} \int_{\{\|\mathbf{x}\| > \epsilon'\}} \frac{st}{(1+s^2)(1+t^2)} d\tau_n(s, t) = \int_{\{\|\mathbf{x}\| > \epsilon'\}} \frac{st}{(1+s^2)(1+t^2)} d\tau(s, t).$$

We now can conclude from the above discussions and (5.51) that $\lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} |J_{1n}(\epsilon)| = 0$, whence $\lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} |I_n(\epsilon)| = 0$ by (5.51) again. Finally, the assumption that the first integral in $I_n(\epsilon)$ converges to γ as $n \rightarrow \infty$ yields that the limits in (5.50) exist and equal, and the relation (5.45) holds. Hence (5.44) is proved.

The positive semi-definiteness of the matrix $\mathbf{A}$ is an easy application of the Cauchy-Schwarz inequality to (5.47), (5.48) and (5.49). It is easy to verify that the limits in (IV) define a non-negative quadratic form $\langle \mathbf{A}\mathbf{u}, \mathbf{u} \rangle$ for any $\mathbf{u} \in \mathbb{R}^2$.

Conversely, suppose that (III) and (IV) in Condition 5.3 hold. Denote by $Q(\mathbf{u})$ the finite quantity in (IV) for any $\mathbf{u} \in \mathbb{R}^2$, and define positive planar measures σ_1 and σ_2 as

$$\sigma_1 = \frac{s^2}{1+s^2} \tau + Q((1, 0))\delta_0 \quad \text{and} \quad \sigma_2 = \frac{t^2}{1+t^2} \tau + Q((0, 1))\delta_0.$$

Note that measures σ_1 and σ_2 are both finite. To see this, it suffices to show that $\chi_{\{\|\mathbf{x}\| \leq 1\}} \|\mathbf{x}\|^2 \in L^1(\tau)$. Take a sequence $\{\epsilon_k\}_{k \geq 1}$ such that $\epsilon_k \downarrow 0$ as $k \rightarrow \infty$ and $\tau(\{\|\mathbf{x}\| = \epsilon_k\}) = 0$ for each k . Then condition (IV) shows that for all $k < j$ large enough, one has

$$\limsup_{n \rightarrow \infty} \int_{\{\epsilon_j < \|\mathbf{x}\| < \epsilon_k\}} s^2 d\tau_n(s, t) \leq Q((1, 0)) + 1,$$

which gives

$$\int_{\{\epsilon_j < \|\mathbf{x}\| < \epsilon_k\}} s^2 d\tau(s, t) \leq Q((1, 0)) + 1$$

by Proposition 5.2. Letting $j \rightarrow \infty$ allows us to obtain that

$$\int_{\{\|\mathbf{x}\| < \epsilon_k\}} s^2 d\tau(s, t) \leq Q((1, 0)) + 1$$

by monotone convergence theorem. This shows that $\chi_{\{\|\mathbf{x}\| \leq 1\}} s^2 \in L^1(\tau)$, as well as $\chi_{\{\|\mathbf{x}\| \leq 1\}} t^2 \in L^1(\tau)$, as desired. Now we are ready to prove that σ_{1n} converges to σ_1 weakly. Let f be a bounded

and continuous function on $\mathbb{R}^2$. Then we have the following estimate

$$\begin{aligned}
& \left| \int_{\mathbb{R}^2} f(\mathbf{x}) d\sigma_{1n}(\mathbf{x}) - \int_{\mathbb{R}^2} f(\mathbf{x}) d\sigma_1(\mathbf{x}) \right| \\
& \leq \int_{\{\|\mathbf{x}\| < \epsilon_k\}} |f(\mathbf{x}) - f(\mathbf{0})| d\sigma_{1n}(\mathbf{x}) + |f(\mathbf{0})| |\sigma_{1n}(\{\|\mathbf{x}\| < \epsilon_k\}) - Q((1, 0))| \\
& + \int_{\{0 < \|\mathbf{x}\| < \epsilon_k\}} |f(\mathbf{x})| d\sigma_1(\mathbf{x}) + \left| \int_{\{\|\mathbf{x}\| \geq \epsilon_k\}} f(\mathbf{x}) d\sigma_{1n}(\mathbf{x}) - \int_{\{\|\mathbf{x}\| \geq \epsilon_k\}} f(\mathbf{x}) d\sigma_1(\mathbf{x}) \right| \\
& =: I_{1n}(k) + I_{2n}(k) + I_3(k) + I_{4n}(k),
\end{aligned}$$

463 First choosing $\mathbf{u} = (1, 0)$ in (5.44) gives that

$$\lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} I_{1n}(k) \leq \lim_{k \rightarrow \infty} \left[\max_{\|\mathbf{x}\| < \epsilon_k} |f(\mathbf{x}) - f(\mathbf{0})| \right] \cdot \left(\limsup_{n \rightarrow \infty} \int_{\{\|\mathbf{x}\| < \epsilon_k\}} s^2 d\tau_n(s, t) \right) = 0.$$

Since

$$\lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} \left| \int_{\{\|\mathbf{x}\| < \epsilon_k\}} \frac{s^2}{1 + s^2} d\tau_n - \int_{\{\|\mathbf{x}\| < \epsilon_k\}} s^2 d\tau_n \right| \leq \lim_{k \rightarrow \infty} \epsilon_k^2 \cdot \left(\limsup_{n \rightarrow \infty} \int_{\{\|\mathbf{x}\| < \epsilon_k\}} s^2 d\tau_n \right) = 0,$$

464 it follows that

$$\lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} |\sigma_{1n}(\{\|\mathbf{x}\| < \epsilon_k\}) - Q((1, 0))| = 0,$$

465 whence $\lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} I_{2n}(k) = 0$. We also have $\limsup_{k \rightarrow \infty} I_3(k) \leq \|f\|_\infty \lim_{k \rightarrow \infty} \sigma_1(\{0 < \|\mathbf{x}\| < \epsilon_k\}) = 0$ and $\limsup_{n \rightarrow \infty} I_{4n}(k) = 0$ because σ_1 is a finite measure and $\tau_n \Rightarrow_{\mathbf{0}} \tau$. Thus we
466 have shown the weak convergence of the sequence $\{\sigma_{1n}\}$ in condition (I). Similarly, $\sigma_{2n} \Rightarrow \sigma_2$.
467

468 Finally, we decompose the desired integral in condition (II) into the sum

$$\int_{\{\|\mathbf{x}\| < \epsilon_k\}} \frac{st}{(1 + s^2)(1 + t^2)} d\tau_n(s, t) + \int_{\{\|\mathbf{x}\| \geq \epsilon_k\}} \frac{st}{(1 + s^2)(1 + t^2)} d\tau_n(s, t).$$

469 As $n \rightarrow \infty$ and $k \rightarrow \infty$, the first integral tends to $[Q((1, 1)) - Q((1, 0)) - Q((0, 1))]/2$, while the
470 second integral tends to

$$\int_{\mathbb{R}^2} \frac{st}{(1 + s^2)(1 + t^2)} d\tau(s, t)$$

471 by Proposition 5.2 and the fact that $1 \wedge \|\mathbf{x}\|^2 \in L^1(\tau)$. Hence condition (II) is verified and the
472 proof is complete. $\square$

473 We are in a position to prove the equivalence between classical and bi-free limit theorems for
474 non-identical distributions.

475 **Theorem 5.5.** *Let $\{k_n\}_{n=1}^\infty \subset \mathbb{N}$ be strictly increasing, $\{\mu_{nk}\}_{1 \leq n, 1 \leq k \leq k_n} \subset \mathcal{P}_{\mathbb{R}^2}$ be an infinitesimal
476 triangular array and $\{\mathbf{v}_n\}_{n=1}^\infty \subset \mathbb{R}^2$. With the notations in (4.25), (4.26), and (4.27), the following
477 are equivalent.*

478 (1) The sequence

$$\mu_{n1} * \mu_{n2} * \cdots * \mu_{nk_n} * \delta_{\mathbf{v}_n}$$

479 converges weakly to some probability measure μ_* on $\mathbb{R}^2$.

480 (2) The sequence

$$\mu_{n1} \boxplus \mu_{n2} \boxplus \cdots \boxplus \mu_{nk_n} \boxplus \delta_{\mathbf{v}_n}$$

481 converges weakly to some probability measure $\mu_{\boxplus}$ on $\mathbb{R}^2$.

482 (3) Condition 4.5(I) and (II) hold, and the vector in (4.39) converges to some vector $\mathbf{v} \in \mathbb{R}^2$.

(4) Condition 5.3(III) and (IV) hold, and the vector in (4.39) converges to some vector $\mathbf{v} \in \mathbb{R}^2$.
 If assertions (1)-(4) hold, then μ_* and $\mu_{\boxplus\boxplus}$ are $*$ -infinitely divisible and $\boxplus\boxplus$ -infinitely divisible
 distributions with $(\mathbf{v}, \mathbf{A}, \tau)$ as the classical and bi-free characteristic triplet, respectively, where $\mathbf{A}$
 is defined as in (5.46).

Proof. The equivalences (2) $\Leftrightarrow$ (3) and (3) $\Leftrightarrow$ (4) were already respectively proved in Theorem
 4.6 and Lemma 5.4, while the equivalence (1) $\Leftrightarrow$ (4) can be obtained by [16, Theorem 3.2.2 and
 (3.52),(3.53),(3.54)]. We remark here that some results cited from [16] contain errors, and the
 reader is referred to the list of errata of the book put on the webpage of one of the authors.

It remains to show that $\phi_{\mu_{\boxplus\boxplus}}$ extends analytically to $(\mathbb{C} \setminus \mathbb{R})^2$ and admits an integral representa-
 tion of the form (3.16). For any $\epsilon > 0$ and $(z, w) \in (\mathbb{C} \setminus \mathbb{R})^2$, let

$$\mathcal{P}_n(z, w, \epsilon) = \int_{\{\|\mathbf{x}\| \geq \epsilon\}} \left[\frac{zw}{(z-s)(w-t)} - 1 - \frac{z^{-1}s + w^{-1}t}{1 + s^2 + t^2} \right] d\tau_n(s, t)$$

and

$$\mathcal{G}_n(z, w, \epsilon) = \int_{\{\|\mathbf{x}\| < \epsilon\}} \left[\frac{zw}{(z-s)(w-t)} - 1 - \frac{z^{-1}s + w^{-1}t}{1 + s^2 + t^2} \right] d\tau_n(s, t).$$

Notice that the integrand in the integral can be rewritten as

$$\frac{1}{z(z-s)} \frac{s^2}{1 + s^2 + t^2} + \frac{1}{w(w-t)} \frac{t^2}{1 + s^2 + t^2} + \frac{st}{(z-s)(w-t)}.$$

Then choosing ϵ so that $\tau(\{\|\mathbf{x}\| = \epsilon\}) = 0$ shows that

$$\mathcal{P}(z, w) := \lim_{\epsilon \rightarrow 0^+} \lim_{n \rightarrow \infty} \mathcal{P}_n(z, w, \epsilon) = \int_{\mathbb{R}^2} \left[\frac{zw}{(z-s)(w-t)} - 1 - \frac{z^{-1}s + w^{-1}t}{1 + s^2 + t^2} \right] d\tau(s, t),$$

where we used Proposition 5.2 and the fact that $1 \wedge \|\mathbf{x}\|^2 \in L^1(\tau)$. On the other hand, (5.47),
 (5.48) and (5.49) yield that

$$\mathcal{G}(z, w) := \lim_{\epsilon \rightarrow 0^+} \lim_{n \rightarrow \infty} \mathcal{G}_n(z, w, \epsilon) = \frac{\sigma_1(\{\mathbf{0}\})}{z^2} + \frac{c}{zw} + \frac{\sigma_2(\{\mathbf{0}\})}{w^2}.$$

Then according to (4.43), $\phi_{\mu_{\boxplus\boxplus}}(z, w)$ agrees with

$$\frac{\mathbf{v}^{(1)}}{z} + \frac{\mathbf{v}^{(2)}}{w} + \mathcal{G}(z, w) + \mathcal{P}(z, w)$$

when $z = iy$ and $w = iv$ with $|y|$ and $|v|$ large, and hence they agree on $(\mathbb{C} \setminus \mathbb{R})^2$ by analytic
 extension. The last assertion regarding $\mu_{\boxplus\boxplus}$ follows from Theorem 3.1. This finishes the proof. $\square$

The classical and bi-free Lévy-Hinčin representations (3.12) and (3.16) establish a bijective
 relation Λ between the sets $\mathcal{ID}(*)$ and $\mathcal{ID}(\boxplus\boxplus)$:

$$(5.52) \quad \Lambda(\mu_*^{(\mathbf{v}, \mathbf{A}, \tau)}) = \mu_{\boxplus\boxplus}^{(\mathbf{v}, \mathbf{A}, \tau)}$$

for any infinitely divisible law $\mu_*^{(\mathbf{v}, \mathbf{A}, \tau)}$ with classical characteristic triplet $(\mathbf{v}, \mathbf{A}, \tau)$. Under this
 bijection, classical Gaussian and (compound) Poisson distributions are respectively mapped to
 bi-free Gaussian and bi-free (compound) Poisson distributions (see Example 3.2). Furthermore,
 Theorem 5.5 and this bijection establish a transfer principle for limit theorems.

The limit theorem for the identically distributed random variables is formulated below.

Theorem 5.6. Let $\{\mu_n\}_{n=1}^\infty$ be a sequence in $\mathcal{P}_{\mathbb{R}^2}$ and let $\{k_n\}_{n=1}^\infty \subset \mathbb{N}$ be strictly increasing. Then the statements (1)–(4) are equivalent.

- (1) The measure $\mu_n^{*k_n}$ converges weakly to some probability measure μ_* on $\mathbb{R}^2$.
- (2) The sequence $\mu_n^{\boxplus k_n}$ converges weakly to some probability measure $\mu_{\boxplus}$ on $\mathbb{R}^2$.
- (3) Condition 4.5(I) and (II) hold with $\tau_n = k_n\mu_n$, and the limit

$$(5.53) \quad \lim_{n \rightarrow \infty} k_n \int_{\mathbb{R}^2} \frac{\mathbf{x}}{1 + \|\mathbf{x}\|^2} d\mu_n(\mathbf{x}) = \mathbf{v}$$

exists.

- (4) Condition 5.3(III) and (IV) hold with $\tau_n = k_n\mu_n$, and the limit in (5.53) exists.

If assertions (1) through (4) hold, then μ_* and $\mu_{\boxplus}$ are $*$ -infinitely divisible and $\boxplus$ -infinitely divisible distributions with $(\mathbf{v}, \mathbf{A}, \tau)$ as the classical and bi-free characteristic triplet, respectively, where $\mathbf{A}$ is defined as in (5.46).

Remark 5.7. Due to the recent work of Gu and Skoufranis [13, Theorem 5.12], the above conditions (1)–(4) are further equivalent to the statement that the sequence $\mu_n^{\boxplus k_n}$ converges weakly to some probability measure $\mu_{\boxplus}$ on $\mathbb{R}^2$, where $\boxplus$ is the bi-boolean convolution. The limit distribution $\mu_{\boxplus}$ may also be characterized by a bi-boolean characteristic triplet.

Proof. The equivalence (1) $\Leftrightarrow$ (2) will follow immediately by choosing $\mu_{nk} = \mu_n$ for $k = 1, \dots, k_n$ and $\mathbf{v}_n = \mathbf{0}$ in Theorem 5.5 once the infinitesimality of $\{\mu_n\}$ is verified in (1) and (2). In (1), we have $\mu_n^{(j)} \Rightarrow \delta_0$ for $j = 1, 2$ by [11] (§14, Theorem 4), whence $\mu_n \Rightarrow \delta_0$ by (2.9). In (2), we see that $\phi_{\mu_n} \rightarrow 0 = \phi_{\delta_0}$ uniformly on compact sets of Γ^2 and $\phi_{\mu_n}(z, w) = o(1)$ uniformly in n as $z, w \rightarrow \infty$ with $(z, w) \in \Gamma^2$ by Proposition 2.3, whence $\mu_n \Rightarrow \delta_0$ by Proposition 2.3 again. Hence the infinitesimality is verified in both situations.

The equivalence (2) $\Leftrightarrow$ (3) was already proved in [14, Theorem 3.2], while the equivalence (3) $\Leftrightarrow$ (4) can be obtained by applying Lemma 5.4 to the positive measures $\tau_n = k_n\mu_n$. That the limiting distribution $\mu_{\boxplus}$ is $\boxplus$ -infinitely divisible distributions with the desired bi-free characteristic triplet follows from the last part of the proof of Theorem 5.5. $\square$

Remark 5.8. In the proof of (2) $\Leftrightarrow$ (3) in Theorem 5.6, a bi-free limit theorem on identical distributions was employed [14]. As one might expect, a more direct proof based on Theorem 5.5 without referring to any other type of limit theorems exists, but it is not a short one. More precisely, what one really needs to show is the equivalence of the following two statements:

- (i) Condition 5.3 holds with $\tau_n = k_n\mu_n$ and there exists some $\mathbf{v} \in \mathbb{R}^2$ so that (5.53) holds;
- (ii) Condition 5.3 holds with $\tau_n = k_n\dot{\mu}_n$ and there exists some $\mathbf{v} \in \mathbb{R}^2$ so that

$$\lim_{n \rightarrow \infty} k_n \left[\mathbf{u}_n + \int_{\mathbb{R}^2} \frac{\mathbf{x}}{1 + \|\mathbf{x}\|^2} d\dot{\mu}_n(\mathbf{x}) \right] = \mathbf{v},$$

where $\dot{\mu}_n$ is the shift of μ_n by the vector $\mathbf{u}_n := \int_{\|\mathbf{x}\| < L} \mathbf{x} d\mu_n(\mathbf{x})$. Some elaboration and techniques are needed to show this equivalence. We leave the proof to the interested reader.

6. STABLE LAWS IN BI-FREE PROBABILITY

In this section we define and study bi-free stable distributions, and show that they arise naturally in limit theorems. The presented result establishes the coincidence of the domains of attraction in classical probability and bi-free probability.

For any $\lambda > 0$, denote by D_λ the dilation operator on measures ρ on $\mathbb{R}^d$, i.e. for any Borel set $B \subset \mathbb{R}^d$,

$$(D_\lambda \rho)(B) = \rho(\{\lambda^{-1} \mathbf{x} : \mathbf{x} \in B\}).$$

Definition 6.1. Let $\star$ be a binary operation on the set $\mathcal{P}_{\mathbb{R}^2}$. A planar probability distribution μ is said to be $\star$ -stable if for any $a, b > 0$, there exist some $c > 0$ and some vector $\mathbf{u} \in \mathbb{R}^2$ so that

$$(D_a \mu) \star (D_b \mu) = (D_c \mu) \star \delta_{\mathbf{u}}.$$

The classification of $\star$ -stable distributions is well known [10, 15]. We wish to thank the referee for bringing the papers to our attention. For the reader's convenience, the statement with a complete proof is provided below. We say that a probability measure is *non-trivial* if it is not a delta measure.

Theorem 6.2. *A non-trivial planar probability measure is $\star$ -stable if and only if either*

- (1) *it is a Gaussian distribution or*
- (2) *it is $\star$ -infinitely divisible and admits the $\star$ -characteristic triplet $(\mathbf{v}, \mathbf{0}, \tau)$ with τ of the form*

$$d\tau(\mathbf{x}) = \frac{1}{r^{1+\alpha}} dr d\Theta(\omega),$$

where $\alpha \in (0, 2)$, Θ is a finite positive measure on the unit circle $\mathbb{T}$ and $\mathbf{x} = r\omega$ with $r > 0$ and $\omega \in \mathbb{T}$.

Proof. Suppose that μ is non-trivial and $\star$ -stable. We may assume that the marginal law $\mu^{(1)}$ is non-trivial. Then for any $a, b > 0$, there exist some $c > 0$ and $\mathbf{u} \in \mathbb{R}^2$ so that

$$(6.54) \quad (D_a \mu^{(1)}) * (D_b \mu^{(1)}) = [(D_a \mu) * (D_b \mu)]^{(1)} = [(D_c \mu) * \delta_{\mathbf{u}}]^{(1)} = (D_c \mu^{(1)}) * \delta_{\mathbf{u}^{(1)}}.$$

This shows that $\mu^{(1)}$ is $\star$ -stable, and hence its Lévy measure is either a zero measure or of the form $d\rho(x) := c_1 x^{-\alpha-1} \chi_{(0, \infty)}(x) dx + c_2 |x|^{-\alpha-1} \chi_{(-\infty, 0)}(x) dx$ for some $\alpha \in (0, 2)$ and $c_1, c_2 \geq 0$ with $c_1 + c_2 > 0$ (see §34 in [11]). If $\rho \neq 0$, one can check from characteristic functions that the constant c is uniquely determined by the relation $c^\alpha = a^\alpha + b^\alpha$. In the first case, $\mu^{(1)}$ is Gaussian, and hence $c^2 = a^2 + b^2$ from characteristic functions again, which is realized as $\alpha = 2$.

To obtain the desired result, let $\alpha \in (0, 2]$ be fixed and consider the family $(\mu_\lambda)_{\lambda > 0}$, where $\mu_\lambda = D_{\lambda^{1/\alpha}} \mu$. Then using the $\star$ -stability of μ and the relation $a^\alpha + b^\alpha = c^\alpha$ shows that

$$(6.55) \quad \mu_{\lambda_1 + \lambda_2} = \mu_{\lambda_1} * \mu_{\lambda_2} * \delta_{\mathbf{u}(\lambda_1, \lambda_2)}$$

for some $\mathbf{u}(\lambda_1, \lambda_2) \in \mathbb{R}^2$, which clearly gives the $\star$ -infinite divisibility of μ and each μ_λ . Let $(\mathbf{v}, \mathbf{A}, \tau)$ be the $\star$ -characteristic triplet of μ . Then μ_λ admits the $\star$ -characteristic triplet $(\mathbf{v}(\lambda), \lambda^{2/\alpha} \mathbf{A}, D_{\lambda^{1/\alpha}} \tau)$ for some $\mathbf{v}(\lambda) \in \mathbb{R}^2$. Moreover, (6.55) yields the following two relations for any $\lambda_1, \lambda_2 > 0$:

$$(6.56) \quad (\lambda_1 + \lambda_2)^{2/\alpha} \mathbf{A} = \lambda_1^{2/\alpha} \mathbf{A} + \lambda_2^{2/\alpha} \mathbf{A}$$

570 and

$$(6.57) \quad D_{(\lambda_1+\lambda_2)^{1/\alpha}}\tau = D_{\lambda_1^{1/\alpha}}\tau + D_{\lambda_2^{1/\alpha}}\tau.$$

571 To continue the proof, let $\Omega \subset \mathbb{T}$ be a fixed Borel set. By restricting the measures appearing in
 572 (6.57) on the set $\{r\Omega : r \geq 1\}$, one can infer that the function $f(\lambda) = \tau(\{r\Omega : r \geq \lambda^{-1/\alpha}\})$ satisfies
 573 Cauchy's functional equation $f(\lambda_1 + \lambda_2) = f(\lambda_1) + f(\lambda_2)$. Since f is increasing on $(0, \infty)$, it is
 574 measurable there, and hence $f(\lambda) = \lambda f(1)$ for any $\lambda > 0$. This allows us to obtain that

$$\tau(\{r\Omega : r \geq \lambda\}) = \lambda^{-\alpha} \tau(\{r\Omega : r \geq 1\})$$

575 for any $\lambda > 0$ and any Borel set $\Omega \subset \mathbb{T}$. Hence the finite positive measure

$$\Theta(\Omega) = \alpha \int_{[1, \infty) \times \Omega} d\tau(r, \omega), \quad \Omega \subset \mathbb{T},$$

576 gives us the desired one. If $\alpha = 2$, then only $\tau = 0$ is allowed in order to fit the condition
 577 $1 \wedge \|\mathbf{x}\|^2 \in L^1(\tau)$, in which case μ is a Gaussian. If $\alpha < 2$, then (6.56) holds for any $\lambda_1, \lambda_2 > 0$ if
 578 and only if $\mathbf{A} = 0$. For the converse, it is clear that μ is $*$ -stable either in the case (1) or (2). $\square$

579 The $\boxplus$ -stable distributions are classified as follows.

580 **Theorem 6.3.** *A non-trivial planar probability measure is $\boxplus$ -stable if and only if either*

- 581 (1) *it is a bi-free Gaussian distribution or*
- 582 (2) *it is $\boxplus$ -infinitely divisible and it has a $\boxplus$ -characteristic triplet $(\mathbf{v}, \mathbf{0}, \tau)$ with τ of the*
 583 *form*

$$d\tau(\mathbf{x}) = \frac{1}{r^{1+\alpha}} dr d\Theta(\omega),$$

584 where $\alpha \in (0, 2)$, Θ is a finite positive measure on the unit circle $\mathbb{T}$ and $\mathbf{x} = r\omega$ with $r > 0$
 585 and $\omega \in \mathbb{T}$.

586 *Proof.* Suppose that μ is non-trivial and $\boxplus$ -stable. Further suppose that $\mu^{(1)}$ is non-trivial.
 587 Then it follows from Proposition 2.6 that

$$(D_a\mu^{(1)}) \boxplus (D_b\mu^{(1)}) = [(D_a\mu) \boxplus (D_b\mu)]^{(1)} = [(D_c\mu) \boxplus \delta_{\mathbf{u}}]^{(1)} = (D_c\mu^{(1)}) \boxplus \delta_{\mathbf{u}^{(1)}}.$$

588 This gives the $\boxplus$ -stability of $\mu^{(1)}$, and hence $\phi'_{\mu^{(1)}}(z) = \beta z^{-\alpha}$ for some $\alpha \in (0, 2]$ and $\beta \in \mathbb{C} \setminus \{0\}$ by
 589 Lemma 7.4 and Theorem 7.5 of [5]. Since $\phi_{D_\lambda\mu^{(1)}}(z) = \lambda\phi_{\mu^{(1)}}(z/\lambda)$ for any $\lambda > 0$, one can conclude
 590 that a, b and c satisfy the relation $c^\alpha = a^\alpha + b^\alpha$.

591 As in the proof of Theorem 6.2, we consider the measures $\mu_\lambda = D_{\lambda^{1/\alpha}}\mu$, $\lambda > 0$. Then the $\boxplus$ -
 592 stability of μ shows that $\mu_{\lambda_1+\lambda_2} = \mu_{\lambda_1} \boxplus \mu_{\lambda_2} \boxplus \delta_{\mathbf{u}(\lambda_1, \lambda_2)}$ for some vector $\mathbf{u}(\lambda_1, \lambda_2) \in \mathbb{R}^2$, which
 593 gives the infinite divisibility of μ and each μ_λ . If $(\mathbf{v}, \mathbf{A}, \tau)$ is the bi-free characteristic triplet of
 594 μ , then the identity $\phi_{D_\lambda\mu}(z, w) = \phi_\mu(z/\lambda, w/\lambda)$, which holds for (z, w) in the common domain of
 595 these transforms, yields that μ_λ admits the bi-free characteristic triplet $(\mathbf{v}(\lambda), \lambda^{2/\alpha}\mathbf{A}, D_{\lambda^{1/\alpha}}\tau)$ for
 596 some $\mathbf{v}(\lambda) \in \mathbb{R}^2$. Then the remaining proof is similar to that of Theorem 6.2. $\square$

597 All $\boxplus$ -stable distributions are $\boxplus$ -infinitely divisible. The number $\alpha \in (0, 2]$ is called *stability*
 598 *index* of μ_* and $\mu_{\boxplus}$. It is shown in the proof that μ_* and $\Lambda(\mu_*)$ have the same stability index,
 599 particularly, the stability index of Gaussian and bi-free Gaussian are both two.

600 In [17], Rvačeva investigated the limiting distribution of random vectors

$$(6.58) \quad \frac{X_1 + \cdots + X_n}{b_n} + \mathbf{u}_n,$$

where $\{X_n\}_{n \geq 1}$ are i.i.d. random vectors, $b_n > 0$ and $\mathbf{u}_n \in \mathbb{R}^2$. It turns out that the set of all possible limiting distributions in (6.58) equals the set of $*$ -stable distributions from the arguments in [11, §33]. The limit theorem of this type in the bi-free setting is considered as follows.

Theorem 6.4. *Let ν be a planar probability distribution, $b_n > 0$ and $\mathbf{u}_n \in \mathbb{R}^2$ for $n = 1, 2, \dots$. Then the following statements are equivalent.*

- (1) *The measures $(D_{1/b_n} \nu^{*n}) * \delta_{\mathbf{u}_n}$ converge weakly to a probability distribution μ_* on $\mathbb{R}^2$.*
 - (2) *The measures $(D_{1/b_n} \nu^{\boxplus n}) * \delta_{\mathbf{u}_n}$ converge weakly to a probability distribution $\mu_{\boxplus}$ on $\mathbb{R}^2$.*
- If (1) and (2) hold, then μ_* and $\mu_{\boxplus}$ are $*$ -stable and $\boxplus$ -stable, respectively, whose respective $*$ -characteristic triplet and $\boxplus$ -characteristic triplet coincide.*

Proof. Applying Theorem 5.6 to the positive integers $k_n = n$ and the measures $\mu_n = (D_{1/b_n} \nu) * \delta_{\mathbf{u}_n/n}$ and $\mu_n = (D_{1/b_n} \nu) \boxplus \delta_{\mathbf{u}_n/n}$ in (1) and (2), respectively, yields the desired equivalence. The last statement is a direct consequence of the mentioned results around (6.58) and the fact $\mu_{\boxplus} = \Lambda(\mu_*)$ established in Theorem 5.6. $\square$

Definition 6.5. A measure $\nu \in \mathcal{P}_{\mathbb{R}^2}$ is said to belong to the $\star$ -domain of attraction of a $\star$ -stable law μ_* if there exist a sequence $\{b_n\}_{n=1}^\infty$ of positive numbers and a sequence $\{\mathbf{u}_n\}_{n=1}^\infty$ of vectors in $\mathbb{R}^2$ so that $(D_{1/b_n} \nu^{*n}) * \delta_{\mathbf{u}_n} \Rightarrow \mu_*$. Denote by $\mathbf{D}_*(\mu_*)$ the $\star$ -domain of attraction of a give $\star$ -stable law μ_* .

The $*$ -domain of attraction was studied in great detail in [17]. One can immediately conclude the following result from Theorem 6.4.

Corollary 6.6. *For any $*$ -stable law μ_* on $\mathbb{R}^2$, $\mathbf{D}_*(\mu_*) = \mathbf{D}_{\boxplus}(\Lambda(\mu_*))$.*

7. FULL DISTRIBUTIONS

We will discuss in this section the concept of fullness which regards the supports of probability distributions introduced below:

Definition 7.1. A Borel measure ρ on $\mathbb{R}^2$ is said to be *full* if it is not supported on a straight line, while ρ is called $\mathcal{M}_{\mathbb{R}^2}^0$ -full if it is in $\mathcal{M}_{\mathbb{R}^2}^0$ and not supported on a line through the origin.

A bivariate normal distribution is full if and only if its symmetric covariance matrix is strictly positive definite, in which case the distribution has a density. If the covariance matrix is not of full rank, then the bivariate normal distribution is non-full and does not have a density.

In the following we relate the fullness of measures in $\mathcal{P}_{\mathbb{R}^2}$ to their Cauchy transforms and bi-free ϕ -transforms.

Lemma 7.2. *A measure $\mu \in \mathcal{P}_{\mathbb{R}^2}$ is non-full if and only if there exist $\alpha, \beta, \gamma \in \mathbb{R}$ so that*

$$(7.59) \quad (\alpha z + \beta w + \gamma) G_\mu(z, w) = \beta G_{\mu^{(1)}}(z) + \alpha G_{\mu^{(2)}}(w)$$

holds for any $(z, w) \in (\mathbb{C} \setminus \mathbb{R})^2$, in which case it is supported on the line $\alpha s + \beta t + \gamma = 0$.

Proof. First notice that for $\alpha, \beta, \gamma \in \mathbb{R}$ and $(z, w) \in (\mathbb{C} \setminus \mathbb{R})^2$, we have

$$\begin{aligned} G(z, w) &:= \int_{\mathbb{R}^2} \frac{\alpha s + \beta t + \gamma}{(z - s)(w - t)} d\mu(s, t) \\ &= (\alpha z + \beta w + \gamma) G_\mu(z, w) - \beta G_{\mu^{(1)}}(z) - \alpha G_{\mu^{(2)}}(w). \end{aligned}$$

This clearly gives (7.59) if μ is supported on $\alpha s + \beta t + \gamma = 0$. Conversely, suppose that $G(z, w) = 0$ holds true for $(z, w) \in (\mathbb{C} \setminus \mathbb{R})^2$. Then a simple computation shows that

$$(7.60) \quad \int_{\mathbb{R}^2} \frac{\alpha s + \beta t + \gamma}{(s^2 + 1)(t^2 + 1)} d\mu(s, t) = -\frac{\Re[G(i, i) - G(-i, i)]}{2} = 0.$$

On the other hand, considering the function $H(z, w) = zG(z, w)$ yields that

$$(7.61) \quad \int_{\mathbb{R}^2} \frac{s(\alpha s + \beta t + \gamma)}{(s^2 + 1)(t^2 + 1)} d\mu(s, t) = -\frac{\Re[H(i, i) - H(-i, i)]}{2} = 0.$$

Similarly, one can obtain that

$$(7.62) \quad \int_{\mathbb{R}^2} \frac{t(\alpha s + \beta t + \gamma)}{(s^2 + 1)(t^2 + 1)} d\mu(s, t) = 0.$$

Multiplying (7.60), (7.61) and (7.62) by γ , α and β , respectively, and then adding them all together shows that

$$\int_{\mathbb{R}^2} \frac{(\alpha s + \beta t + \gamma)^2}{(s^2 + 1)(t^2 + 1)} d\mu(s, t) = 0.$$

Since μ is positive, this clearly shows that it is supported on $\alpha s + \beta t + \gamma = 0$, as desired. $\square$

Proposition 7.3. A measure $\mu \in \mathcal{P}_{\mathbb{R}^2}$ is non-full if and only if there exist $\alpha, \beta, \gamma \in \mathbb{R}$ so that

$$(7.63) \quad zw(\alpha z + \beta w)\phi_\mu(z, w) = \beta w^2\phi_{\mu(1)}(z) + \alpha z^2\phi_{\mu(2)}(w) - \gamma zw$$

holds for $(z, w) \in \Gamma^2$, in which case μ is supported on the line $\alpha s + \beta t + \gamma = 0$.

Proof. With the help of Lemma 7.2, we see that μ is supported on the line $\alpha s + \beta t + \gamma = 0$ if and only if (7.59) holds for $(z, w) \in (\mathbb{C} \setminus \mathbb{R})^2$ or, equivalently, the identity

$$(7.64) \quad (\alpha F_{\mu(1)}^{-1}(z) + \beta F_{\mu(2)}^{-1}(w) + \gamma)G_\mu\left(F_{\mu(1)}^{-1}(z), F_{\mu(2)}^{-1}(w)\right) = \frac{\beta}{z} + \frac{\alpha}{w}$$

is valid for $(z, w) \in \Gamma^2$. Note that the function $G_\mu(F_{\mu(1)}^{-1}, F_{\mu(2)}^{-1})$ never vanishes on Γ^2 by shrinking the domain if necessary. Then we see that μ is supported on $\alpha s + \beta t + \gamma = 0$ if and only if

$$(7.65) \quad (\alpha z + \beta w) \left[1 - \frac{1}{zwG_\mu\left(F_{\mu(1)}^{-1}(z), F_{\mu(2)}^{-1}(w)\right)} \right] = -[\alpha\phi_{\mu(1)}(z) + \beta\phi_{\mu(2)}(w) + \gamma]$$

holds true for $(z, w) \in \Gamma^2$. Apparently, (7.63) and (7.65) are equivalent, concluding the proof. $\square$

Theorem 7.4. Let μ be a $\boxplus$ -infinitely divisible distribution on $\mathbb{R}^2$ with bi-free characteristic triplet $[\mathbf{v}, \mathbf{A}, \tau]$. Then μ is non-full if and only if $\mathbf{A}$ is singular and τ is supported on $\langle \mathbf{u}, (s, t) \rangle = 0$ with some $\mathbf{u} \neq \mathbf{0}$ in the kernel of $\mathbf{A}$, in which case μ is supported on $\langle \mathbf{u}, (s, t) \rangle = \langle \mathbf{u}, \mathbf{v} \rangle$.

Proof. We shall use Proposition 7.3 to conclude the proof. First notice that for any real numbers α, β and γ , the function

$$(7.66) \quad \frac{\alpha z + \beta w}{zw}\phi_\mu(z, w) - \frac{\beta}{z^2}\phi_{\mu(1)}(z) - \frac{\alpha}{w^2}\phi_{\mu(2)}(w) + \frac{\gamma}{zw}$$

can be expressed as

$$(7.67) \quad \frac{\gamma'}{zw} + \frac{\alpha a + \beta c}{z^2 w} + \frac{\alpha c + \beta b}{z w^2} + \int_{\mathbb{R}^2} \left[\frac{\alpha s + \beta t}{(z - s)(w - t)} - \frac{1}{zw} \frac{\alpha s + \beta t}{1 + s^2 + t^2} \right] d\tau(s, t)$$

by (3.16), (3.18) and (3.19), where $\mathbf{A}$ is as in (3.14) and $\gamma' = \alpha \mathbf{v}^{(1)} + \beta \mathbf{v}^{(2)} + \gamma$.

Let $\mathbf{u} = (\alpha, \beta)$ and $\gamma = -\langle \mathbf{u}, \mathbf{v} \rangle$. In this case $\gamma' = 0$. If μ is supported on $\langle \mathbf{u}, (s, t) \rangle = \langle \mathbf{u}, \mathbf{v} \rangle$, then Proposition 7.3 yields that the function in (7.67) vanishes on $(\mathbb{C} \setminus \mathbb{R})^2$. Using the technique employed in Lemma 7.2 we can obtain that

$$\int_{\mathbb{R}^2} \frac{(\alpha s + \beta t)^2}{(1 + s^2)(1 + t^2)} d\tau(s, t) = -(\alpha^2 a + 2\alpha\beta c + \beta^2 b) = -\langle \mathbf{A}\mathbf{u}, \mathbf{u} \rangle.$$

Since $\mathbf{A} \geq 0$, it follows that $\mathbf{A}\mathbf{u} = 0$ and τ is supported on the line $\alpha s + \beta t = 0$. Conversely, if $\mathbf{A}\mathbf{u} = 0$ and τ is supported on the line $\alpha s + \beta t = 0$, then using Proposition 7.3 and (7.67) again shows that μ is supported on $\alpha s + \beta t = \alpha \mathbf{v}^{(1)} + \beta \mathbf{v}^{(2)}$, as desired. $\square$

The following results are both direct consequences of Theorem 7.4.

Corollary 7.5. *A bi-free Gaussian distribution with bi-free characteristic triplet $(\mathbf{v}, \mathbf{A}, 0)$ is non-full if and only if $\mathbf{A}$ is singular, in which case, it is supported on the line $\langle \mathbf{u}, (s, t) \rangle = \langle \mathbf{u}, \mathbf{v} \rangle$, where $\mathbf{u}$ is a nonzero vector in the kernel of $\mathbf{A}$.*

Corollary 7.6. *A bi-free compound Poisson distribution with rate $\lambda > 0$ and jump distribution ν is non-full if and only if ν is $\mathcal{M}_{\mathbb{R}^2}^0$ -nonfull, in which case they are supported on the same line. Consequently, any bi-free Poisson distribution is non-full.*

Proof. Following the notations in (3.20) and (3.21), Theorem 7.4 yields that ν is supported on the line $\langle \mathbf{u}, (s, t) \rangle = 0$ for some $\mathbf{u} \in \mathbb{R}^2$ if and only if μ is supported on the line $\langle \mathbf{u}, (s, t) \rangle = \langle \mathbf{u}, \mathbf{v} \rangle = 0$, as desired. $\square$

Theorem 7.7. *Let $\mu \in \mathcal{P}_{\mathbb{R}^2}$ be $*$ -infinitely divisible. Then μ is full if and only if $\Lambda(\mu)$ is full.*

Proof. Recall that μ is non-full if and only if its characteristic function has the property that $|\hat{\mu}(\lambda \mathbf{u})| = 1$ for all $\lambda \in \mathbb{R}$, where $\mathbf{u}$ is some nonzero vector. If P is the Poisson part of μ , then

$$|\hat{\mu}(\lambda \mathbf{u})| = |\hat{P}(\lambda \mathbf{u})| \exp \left[-\frac{1}{2} \lambda^2 \langle \mathbf{A}\mathbf{u}, \mathbf{u} \rangle \right]$$

yields that μ is non-full if and only if $\mathbf{A}\mathbf{u} = 0$ and $|\hat{P}(\lambda \mathbf{u})| = 1$ for all λ . Then the desired result follows from Theorem 7.4 and (5.52). $\square$

Acknowledgement. The authors would like to thank anonymous reviewers for providing comments and suggestions which improve the quality of this paper. The first-named author is supported by JSPS Grant-in-Aid for Young Scientists (B) 15K17549. The second-named author was supported through a grant from the Ministry of Science and Technology in Taiwan MOST-104-2115-M-110-011-MY2 and a faculty startup grant from the National Sun Yat-sen University. The third-named author was supported by the NSERC Canada Discovery Grant RGPIN-2016-03796.

REFERENCES

- [1] M. Barczy and G. Pap, Portmanteau theorem for unbounded measures, *Statist. Probab. Lett.* **76** (2006), 1831-1835.
- [2] S.T. Belinschi, H. Bercovici, Y. Gu, and P. Skoufranis, Analytic subordination for bi-free convolution, ArXiv: 1702.01673v1
- [3] H. Bercovici and V. Pata, Stable laws and domains of attraction in free probability theory, *Ann. Math.* **149** (1999), 1023-1060.
- [4] H. Bercovici and V. Pata, A free analogue of Hinčin characterization of infinite divisibility, *Proc. Amer. Math. Soc.* **128** (2000), No. 4, 1011-1015.

- [5] H. Bercovici and D. Voiculescu, Free convolutions of measures with unbounded support, *Indiana Univ. Math. J.* **42** (1993), no. 3, 733–773.
- [6] H. Bercovici and J.-C. Wang, The asymptotic behavior of free additive convolution, *Oper. Matrices* **2** (2008), no. 1, 115–124.
- [7] I. Charlesworth, B. Nelson, P. Skoufranis, On two-faced families of non-commutative random variables, *Canad. J. Math.* **67**(6) (2015) 1290–1325.
- [8] I. Charlesworth, B. Nelson, P. Skoufranis, Combinatorics of bi-freeness with amalgamation, *Comm. Math. Phys.* **338**(2) (2015) 801–847.
- [9] G. P. Chistyakov and F. Götze, Limit theorems in free probability theory. I, *Ann. Probab.* **36** (2008), no. 1, 54–90.
- [10] E. Feldheim, *Étude de la stabilité des lois de probabilité*, Ph.D. thesis, Faculté des Sciences de Paris, Paris, France (1937)
- [11] B. V. Gnedenko and A. N. Kolmogorov, *Limit Distributions for Sums of Independent Random Variables*, Addison-Wesley Publishing Company, Inc., 1954.
- [12] Y. Gu, H.-W. Huang, and J. A. Mingo, An analogue of the Lévy-Hinčin formula for bi-free infinitely divisible distributions, *Indiana Univ. Math. J.* **65** (2016), no. 5, 1795–1831.
- [13] Y. Gu and P. Skoufranis, Bi-Boolean independence for pairs of algebras. ArXiv:1703.03072
- [14] H.-W. Huang and J.-C. Wang, Analytic aspects of bi-free partial R -transform, *J. Funct. Anal.* **271** (2016), no. 4, 922–957.
- [15] P. Lévy, *Théorie de l'addition des variables aléatoires*, Monographies des probabilités; 1 (2nd Edition). Paris: Gauthier-Villars (1954).
- [16] M. M. Meerschaert and H.-P. Scheffler, *Limit distributions for sums of independent random vectors: Heavy tails in theory and practice*, Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons, Inc., New York, 2001. xvi+484 pp. The list of errata is available at <http://www.stt.msu.edu/~mcubed/RVbook.html>
- [17] E. L. Rvačeva, On domains of attraction of multi-dimensional distributions, 1962 Select. *Transl. Math. Statist. and Probability* **2** pp. 183–205 American Mathematical Society, Providence, RI.
- [18] K. Sato, *Lévy processes and infinitely divisible distributions*, Cambridge Studies in Advanced Math. **68**, Cambridge, 1999.
- [19] P. Skoufranis, Independences and partial R -transforms in bi-free probability, *Ann. Inst. Henri Poincaré Probab. Stat.* **52** (2016), no.3, 1437–1473.
- [20] P. Skoufranis, *A combinatorial approach to Voiculescu's bi-free partial transforms*, *Pacific J. Math.* **283** (2016), no. 2, 419–447.
- [21] D. V. Voiculescu, Free probability for pairs of faces I, *Comm. Math. Phys.* **332** (2014), 955–980.
- [22] D. V. Voiculescu, Free probability for pairs of faces II: 2-variables bi-free partial R -transform and systems with rank ≤ 1 commutation, *Ann. Inst. Henri Poincaré Probab. Stat.* **52** (2016), no. 1, 1–15.
- [23] D. V. Voiculescu, K. J. Dykema, A. Nica, *Free Random Variables*. CRM Monograph Series, Vol. 1 Am. Math. Soc. Providence, RI, 1992.

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, NORTH 10 WEST 8, KITA-KU, SAPPORO 060-0810, JAPAN

E-mail address: thasebe@math.sci.hokudai.ac.jp

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL SUN YAT-SEN UNIVERSITY, NO. 70, LIENHAI ROAD, KAOHSIUNG 80424, TAIWAN, R.O.C.

E-mail address: hwhuang@math.nsysu.edu.tw

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SASKATCHEWAN, 106 WIGGINS ROAD, SASKATOON, SASKATCHEWAN S7N 5E6, CANADA

E-mail address: jcwang@math.usask.ca