Title	Significance of CCN2 expression in bovine preimplantation development
Author(s)	Akizawa, Hiroki; Yanagawa, Yojiro; Nagano, Masashi; Bai, Hanako; Takahashi, Masashi; Kawahara, Manabu
Citation	Animal science journal, 90(1), 49-54 https://doi.org/10.1111/asj.13126
Issue Date	2019-01
Doc URL	http://hdl.handle.net/2115/76453
Rights	This is the peer reviewed version of the following article: Hiroki Akizawa, Yojiro Yanagawa, Masashi Nagano, Hanako Bai, Masashi Takahashi, Manabu Kawahara (2019). Significance of CCN2 expression in bovine preimplantation development. Animal science journal, 90(1), 49-54, which has been published in final form at https://doi.org/10.1111/asj.13126 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
Type	article (author version)
File Information	ASJ akizawa.pdf

[Instructions for use](#)

Significance of *CCN2* expression in bovine preimplantation development

Hiroki Akizawa¹, Yojiro Yanagawa², Masashi Nagano², Hanako Bai¹, Masashi Takahashi¹, Manabu Kawahara^{1*}

¹Laboratory of Animal Breeding and Reproduction, Research Faculty of Agriculture, Hokkaido University, Kita-ku Kita 9 Nishi 9, Sapporo 060-8589, Japan

²Laboratory of Theriogenology, Department of Veterinary Clinical Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

***Corresponding author:** Manabu Kawahara, Ph.D.

Laboratory of Animal Genetics and Reproduction, Research Faculty of Agriculture

Hokkaido University, Kita-ku Kita 9 Nishi 9, Sapporo 060-8589, Japan

E-mail: k-hara@anim.agr.hokudai.ac.jp

Tel. & Fax: +81-11-706-2541

Running title: CCN2 expression dynamics in bovine blastocysts

ABSTRACT

In mammalian preimplantation development, the first cell lineage segregation occurs during the blastocyst stage, when the Inner Cell Mass and Trophectoderm (TE) differentiate. Species-specific analyses are essential to elucidate the molecular mechanisms that underlie this process, since they differ between various species. We previously showed that the reciprocal regulation of *CCN2* and *TEAD4* is required for proper TE differentiation in bovine blastocysts; however, the function of *CCN2* during early embryogenesis has remained otherwise elusive. The present study assessed the spatiotemporal expression dynamics of *CCN2* in bovine embryos, and evaluated how changes to *CCN2* expression (using a *CCN2* knockdown (KD) blastocyst model) regulate the expression of pluripotency-related genes such as *OCT4* and *NANOG*. The conducted quantitative PCR analysis revealed that *CCN2* mRNA was expressed in bovine oocytes (at the metaphase stage of their second meiosis), and embryos. Similarly, immunostaining detected both cytoplasmic and nuclear *CCN2* at all analyzed oocyte and embryonic stages. Finally, both *OCT4* and *NANOG* expression levels were shown to be significantly reduced in *CCN2* KD blastocysts. Together, these results demonstrate that bovine *CCN2* exhibits unique expression patterns during preimplantation development, and is required for the proper expression of key regulatory genes in bovine blastocysts.

- 1 **Key words:** *CCN2*, pluripotency-related genes, bovine, blastocyst, cell-lineage specification.

INTRODUCTION

During early embryonic cleavage, blastomeres are morphologically indistinguishable, and have the potential to differentiate into all cell lineages (Kelly 1977). As development progresses however, the blastomeres gradually lose their developmental plasticity, and cell lineage segregation begins, and is thereafter maintained via a fine-tuned transcriptional network (Arnold & Robertson 2009). By the blastocyst stage, embryos are comprised of two distinct cell lineages; namely, the Inner Cell Mass (ICM), and the Trophectoderm (TE), which subsequently give rise to embryo-proper and extraembryonic tissues, respectively. Hence, defective segregation of these two cell lineages leads to diverse developmental defects.

The transcription factor *TEAD4* plays critical roles in TE differentiation by activating the expression of core TE regulators, such as *Cdx2* in mice (Yagi *et al.* 2007; Nishioka *et al.* 2009). Murine embryos lacking *TEAD4* expression exhibit failed blastocyst development, and resultantly, a loss of trophoblast stem cells (Yagi *et al.* 2007). To evaluate whether *TEAD4* is similarly required for bovine blastocyst formation, we previously knocked down *TEAD4* transcripts using short hairpin RNA (shRNA)-mediated RNA interference in preimplantation embryos (Akizawa *et al.* 2018). *TEAD4* knockdown (KD) significantly reduced the expression of important TE-expressed genes such as *CDX2*, *GATA2*, and *CCN2*. Of these, *CCN2* was most markedly suppressed; however, while it has been previously shown to be essential for fetal morphogenesis by the fact that *CCN2*-mutant

1 mice die soon after birth due to severe skeletal dysmorphisms (Ivkovic *et al.* 2003), its role in early
2 embryogenesis, including bovine preimplantation development, is not yet clear. Structurally, CCN2
3 is a signaling protein composed of four cysteine-rich domains, comprising an insulin-like growth
4 factor binding protein (IGFBP) domain, a von Willebrand type C (VWC) repeat, a thrombospondin
5 (TSP) repeat, and a C-terminal (CT) motif that contains a cysteine knot (Hall-Glenn & Lyons 2011).
6 In most mammals, CCN2 is uniformly expressed in the blastocyst ICM and TE (Gulnaar *et al.* 1998;
7 Moussad *et al.* 2002; Munoz *et al.* 2017), but its expression patterns vary between species after this
8 stage of development (Moussad *et al.* 2002; Oh *et al.* 2009). For example, CCN2 expression in pig
9 trophoblast cells diminishes soon after the blastocyst stage, while in humans, it is maintained until
10 implantation. CCN2 expression dynamics during bovine preimplantation development have not yet
11 been analyzed.

12 Similarly, little is known about the function of CCN2 in bovine preimplantation
13 development. Previously, we demonstrated that *CCN2* KD blastocysts exhibit significantly reduced
14 *TEAD4* mRNA expression, suggesting that *TEAD4* and CCN2 may function together (i.e.
15 non-redundantly) to direct correct TE differentiation in bovine embryos. Interestingly,
16 immunofluorescent staining of CCN2 production at the blastocyst stage showed that CCN2 localizes
17 to both TE and ICM cells; however, *TEAD4* KD-mediated CCN2 suppression occurs in the TE, but
18 not the ICM. Thus, while the underlying molecular mechanisms are not yet clear, these data suggest

that *CCN2* likely plays important roles not only in bovine TE formation, but also in the overall development of the bovine blastocyst, including the ICM.

In the present study, we first investigated *CCN2* expression dynamics during bovine preimplantation development via both a quantitative PCR analysis, and an immunostaining assay for *CCN2*. We then examined the effects of *CCN2* KD on the expression of pluripotency-related genes, and showed that in *CCN2* KD blastocysts, the expression levels of the core pluripotency related transcription factors *OCT4* and *NANOG* are significantly reduced. Taken together, these findings support that *CCN2* is required for proper lineage segregation in bovine blastocysts, to regulate the expression of both TE-expressed, and core pluripotency-related genes.

1 MATERIALS and METHODS

2 *Preparation of bovine embryos*

3 Bovine oocyte retrieval, and *in vitro* oocyte maturation, fertilization, and subsequent *in vitro*
 4 embryo culture were performed as previously described (Nagatomo *et al.* 2013). Briefly,
 5 cumulus-oocyte complexes (COCs) collected from slaughterhouse-derived ovaries were matured via
 6 culture (20–22 h, at 38.5°C, in a humidified atmosphere with 5% CO₂ in air) in TCM-199 medium
 7 (Thermo Fisher Scientific, Inc., Waltham, MA, USA) containing 5% (v/v) FBS. *In vitro*-matured
 8 oocytes were transferred to Brackett and Oliphant (BO) medium (Brackett & Oliphant 1975)
 9 containing 2.5 mM theophylline (Wako Pure Chemical Industries, Ltd., Osaka, Japan), and 7.5
 10 µg/mL Heparin Sodium Salt (Nacalai Tesque, Inc., Kyoto, Japan). Subsequently, frozen-thawed
 11 semen was centrifuged at 600 × g for 7 min in BO medium, and the isolated spermatozoa were
 12 added to the COCs at a final concentration of 5 × 10⁶ cells/mL. After 12 h of incubation, the
 13 presumptive zygotes were denuded by pipetting, and cultured (8 days, at 38.5°C, in a humidified
 14 atmosphere with 5% CO₂ and 5% O₂ in air) in mSOFai medium (Aono *et al.* 2013). A Holstein cow
 15 on campus of Hokkaido University was administered 25 mg of prostaglandin F2α (Pronalgon F;
 16 Schering-Plough K.K., Tokyo, Japan) 20 days prior to artificial insemination (AI) followed by
 17 intravaginal indwelling of the controlled intravaginal drug release (CIDR; 1.9 g of progesterone;
 18 Eazi-Breed CIDR Cattle Insert; Pfizer Animal Health, New York, NY, USA) device from 9 to 3 days

prior to AI. Estradiol benzoate (2 mg; Ovahormon, Injection; Aska Animal Health, Tokyo) was simultaneously injected with CIDR insertion. From 5 to 3 days prior to AI, follicle stimulating hormone (Antrin R-10; Kyoritsu Seiyaku, Tokyo, Japan) consisted of twice-daily (morning and afternoon) with a decreasing dose (6, 6, 5, 5, 4, and 4 AU per injection) were administered. At the fifth and sixth FSH treatments, 30 and 20 mg of prostaglandin F2 α were injected, respectively. On the day of AI, 200 μ g of gonadotropin releasing hormone (Conceral; ASKA Pharmaceutical Co.,Ltd., Tokyo, Japan) was administrated. The superstimulated cow was inseminated (day 0), and day-14 conceptuses were subsequently collected via non-surgical flushing in warm lactated Ringer's solution (Nippon Zenyaku Kogyo Co.,Ltd, Fukushima, Japan).

Quantitative Reverse-Transcription PCR

The total RNA from five oocytes or embryos per biological replicate was isolated using the ReliaPrep RNA Cell Miniprep System (Promega, Fitchburg, WI, USA), according to the manufacturer's instructions. cDNA was then synthesized from the oocyte/embryo RNA using ReverTra Ace qPCR RT Master Mix (Toyobo, Osaka, Japan). Quantitative PCR (qPCR) was performed using THUNDERBIRD SYBR qPCR Mix (Toyobo), and the specified primers (Supplemental Table S1). The utilized thermal cycling conditions consisted of one cycle of 95°C for 30 s (denaturation), followed by 50 cycles of 95°C for 10 s (denaturation), the appropriate annealing

temperature for each primer set for 15 s (primer annealing), and 72°C for 30 s (extension). Relative mRNA abundance was calculated using the $\Delta\Delta C_t$ method, where *H2A histone family member Z* (*H2AFZ*) was used as a reference gene for each sample. All experiments were performed in triplicate.

Immunofluorescence and Confocal Microscopy

CCN2 immunofluorescence was analyzed in both bovine oocytes and embryos as previously described (Akizawa *et al.* 2018). Briefly, the zona pellucida was removed from bovine oocytes and embryos using 0.05% (w/v) Pronase (Sigma-Aldrich, St. Louis, MO, USA). Both oocytes and embryos were fixed in 4% (w/v) paraformaldehyde (Wako Pure Chemical Industries, Ltd.) in phosphate-buffered saline (PBS) for 60 min, and then permeabilized for 60 min using 0.2% (v/v) Triton X-100 in PBS. They were then blocked for 45 min in Blocking One (1:5; Nacalai Tesque, Inc.) solution diluted in 0.05% (v/v) Tween 20 in PBS (blocking buffer), and incubated in rabbit anti-CCN2 antibody (1:300, ab6992; Abcam, Cambridge, UK) overnight at 4°C. After five 10-min washes in 0.1% (v/v) Triton X-100 and 0.3% (w/v) bovine serum albumin (Sigma-Aldrich), the oocytes and embryos were incubated with Alexa Fluor 555 goat anti-Rabbit IgG polyclonal antibodies (diluted to 1:400 in blocking buffer; ab150082; Thermo Fisher Scientific, Inc.) for 30 min at room temperature. Nuclei were counterstained with 25 mg/mL Hoechst 33342 stain

(Sigma-Aldrich) prepared in 0.2% (w/v) polyvinyl alcohol in PBS. Fluorescence signals were observed using a TCS SP5II confocal laser-scanning microscope (Leica, Wetzlar, Germany). In each experiment, at least five oocytes/ embryos were observed.

Microinjection of CCN2-shRNA expression vectors into bovine embryos

shRNA-mediated RNA interference (RNAi) for *CCN2* was induced as previously described (Akizawa *et al.* 2018). Briefly, an shRNA containing antisense/sense regions, a 19-bp loop (5'-CTGTGAAGCCACAGATGGG-3'), and a 6-bp terminator element (5'-TTTTTT-3'), was designed to target nucleotides 665–683 of the *CCN2* mRNA sequence (NCBI Reference Sequence NM_174030), and ligated downstream of the U6 promoter in the pBasi/mU6 Neo vector (Stratagene, CA, USA). Embryos injected with pBasi/mU6 Neo plasmids lacking the shRNA insert (empty vector) were used as controls, as previously described (Nagatomo *et al.* 2013; Akizawa *et al.* 2016). Twelve hours after insemination, the synthesized shRNA-expression constructs (diluted to a final concentration of 10 ng/mL with mSOFai medium) were injected into denuded presumptive zygotes using a FemtoJet injection device (Eppendorf, Hamburg, Germany). These presumptive zygotes were then cultured to the blastocyst stage to examine the effect of *CCN2* KD on the expression of pluripotency-related genes.

1 *Statistical analysis*

2 In the result of *CCN2* mRNA expression dynamics (Fig. 1A), data were statistically analyzed
3 using a one-way analysis of variance followed by Tukey's multiple comparison test. Excluding Fig.
4 1A, significant differences were compared using Student's *t* test. Statistical analyses were conducted
5 using StatView statistical-analysis software (Abacus Concepts, Inc., Berkeley, CA, USA). A *P* value
6 < 0.05 was considered to indicate statistical significance.

RESULTS

CCN2 mRNA expression levels during preimplantation development

The conducted qPCR analysis of *CCN2* mRNA expression dynamics in bovine oocytes and embryos detected *CCN2* mRNA at all of the examined stages, (i.e. metaphase MII oocytes, and 2-, 4-, 8-, 16-, morula, and blastocyst-stage embryos) (Fig. 1A). Notably, *CCN2* expression significantly decreased between the 16-cell and the blastocyst stages of development ($P < 0.05$) (Fig. 1A). Next, elongating embryos were collected from cows at day 14 after artificial insemination via uterine flushing, to investigate *CCN2* mRNA expression after the blastocyst stage. The results of this analysis revealed that *CCN2* expression levels in the elongating conceptuses was significantly increased compared to that observed in blastocyst-stage embryos ($P < 0.05$) (Fig. 1B).

Subcellular CCN2 protein localization patterns during preimplantation development

Next, immunostaining was performed to visualize and evaluate the subcellular localization of *CCN2* during early bovine embryogenesis. As for *CCN2* mRNA expression, *CCN2* was detected at all of the examined stages (Fig. 2), and shown to consistently localize to both the cytoplasm and nucleus of each blastomere from the 2-cell to the blastocyst stage (Fig. 2). Notably, at the blastocyst stage, the *CCN2* fluorescent signals appeared strongest in the nucleus.

1 *Effects of CCN2 mRNA suppression on the expression of pluripotency-related genes in blastocysts*

2 We previously constructed a *CCN2* KD system in bovine embryos (Akizawa *et al.* 2018),
3 which demonstrated that *CCN2* KD is sufficient to disrupt the expression patterns of primary
4 TE-expressed genes, such as *CDX2*, *GATA2*, and *TEAD4*. Since *CCN2* was found in the present
5 study to be ubiquitously (i.e. in both the ICM and TE) expressed throughout the analyzed embryos
6 (Fig. 2), we evaluated the effects of *CCN2* KD on the blastocyst expression of pluripotency-related
7 genes. The results of this analysis showed that the expression of both *OCT4*, (which is normally
8 detected uniformly in both ICM and TE blastomeres), and *NANOG*, (which is predominantly
9 expressed in ICM blastomeres), were significantly decreased in *CCN2* KD blastocysts ($P < 0.05$)
10 (Fig. 3). These factors are known to be required for ICM-cell self-renewal and pluripotency in both
11 mice and cattle (Wu *et al.* 2016; Bogliotti *et al.* 2018); thus, our results indicate that *CCN2*
12 expression mediates *OCT4* and *NANOG* expression in bovine blastocysts.

DISCUSSION

CCN2 has been shown to be strongly involved in several major biological disorders, such as fibrosis, and various malignancies; thus, it has been extensively studied as a possible therapeutic target for the last few decades (Kubota & Takigawa 2015). However, because homozygous mutant-*CCN2* neonates are recovered at the expected Mendelian ratio (Ivkovic *et al.* 2003), the function(s) of *CCN2* in preimplantation development have thus far been overlooked. Thus, the present study is the first to evaluate both *CCN2* mRNA and protein expression dynamics, and the contribution of *CCN2* to the proper expression of pluripotency-related genes in bovine preimplantation embryos.

Firstly, we revealed that *CCN2* mRNA is detectable throughout early bovine embryogenesis. *CCN2* expression levels were higher in oocytes and embryos prior to major zygotic genome activation (ZGA) (which occurs in cattle after the 8- to 16-cell stages) (Khan *et al.* 2012) than in the subsequent-staged embryos. These results suggest that maternal *CCN2* mRNA may persist, and direct bovine embryogenesis until ZGA, in a similar manner to maternal SMAD2/3 (which are known to be upstream regulators of *CCN2* in bovine embryos) (Zhang *et al.* 2015). The fluorescent signals for *CCN2* proteins detected after the 16-cell stage in Fig. 2 might mainly be derived from maternally expressed *CCN2* mRNA transcribed before the 8-cell stage. Conversely, we observed a significant increase in *CCN2* expression in elongating conceptuses compared to blastocysts. The

1 elongation of bovine conceptus is accompanied by a remarkable proliferation of trophoblast cells
2 (Betteridge and Fléchon 1988). In fact, *CCN2* has been shown to be associated with cell proliferation
3 in both cattle and other mammals (Zhou *et al.* 2008); thus, it may mediate the elongation of bovine
4 embryos after the blastocyst stage.

5 Immunostaining revealed that *CCN2* localized within both the cytoplasm and nucleus in
6 bovine embryos. *CCN2* is secreted, and generally detected in the Golgi apparatus (Zhou *et al.* 2008),
7 supporting that it performs important roles in the cytoplasm. However, *CCN2* is detected in both
8 cytoplasmic and nuclear fractions in some somatic cells and nuclear-localized *CCN2* has been shown
9 to modulate RNA transcription (Wahab *et al.* 2001). Although the mechanisms by which nuclear
10 translocation of *CCN2* is enhanced in bovine embryos are not yet clear, the results of the conducted
11 knockdown experiments in the present study strongly suggest that nuclear *CCN2* regulates
12 transcriptional activity in blastomeres during bovine preimplantation development.

13 We previously demonstrated that *CCN2* KD affects the expression of TE-expressed genes,
14 and decreases the ratio of TE to ICM cells in bovine embryos, indicating that *CCN2* mediates TE
15 development (Akizawa *et al.* 2018). However, to date, the role(s) played by *CCN2* in the
16 establishment of the complex transcriptional networks required for blastocyst formation have not yet
17 been elucidated. In the present study, *CCN2* KD suppressed both *OCT4* and *NANOG* expression in
18 bovine blastocysts. A recent *OCT4* knockout study in bovine embryos revealed that *OCT4* is

essential for maintaining NANOG expression at the blastocyst stage (Simmet *et al.* 2018). Consistent with this, the remarkable reduction of *NANOG* observed in the *CCN2* KD embryos in the present study may be a result of downregulated *OCT4* expression. *CCN2* has also been shown to interact with various cell signaling pathways, including the FGFR and TGFB pathways, during murine blastocyst-stage cell lineage segregation (Abreu *et al.* 2002; Crean *et al.* 2004). Further studies are needed to elucidate the pathways by which *CCN2* contributes to cell lineage segregation in bovine embryos.

In conclusion, the present study demonstrates that *CCN2* is expressed throughout development until implantation, and contributes to the transcriptional regulation of pluripotency-related genes in both TE and ICM cell lineages in bovine blastocysts. In both cattle and other mammals, the correct establishment and function of complex transcriptional networks enables preimplantation embryos to develop to the blastocyst stage. Continued research is needed to decipher the role(s) of *CCN2* in bovine-specific preimplantation after the blastocyst stage of development.

Acknowledgements

This study was supported by a Grant-in-Aid for Scientific Research (B) (No. 18H02321) to M.K., and a Grant-in-Aid for JSPS Fellows (No. 17J01122) to H.A. from the Japan Society for the Promotion of Science. We are grateful to all members and technical staff at the Field Science Center

- 1 for the Northern Biosphere, Hokkaido University, Sapporo, Japan, for their support during the
- 2 collection of bovine embryos.

References

- Abreu J.G., Ketpura N.I., Reversade B. & Robertis E.M. De. 2002. Connective-tissue growth factor (CTGF) modulates cell signalling by BMP and TGF-beta. *Nature Cell Biology* 4, 599–604.
- Akizawa H., Kobayashi K., Bai H., Takahashi M., Kagawa S., Nagatomo H. & Kawahara M. 2018. Reciprocal regulation of TEAD4 and CCN2 for the trophectoderm development of the bovine blastocyst. *Reproduction* 155, 563–571.
- Akizawa H., Nagatomo H., Odagiri H., Kohri N., Yamauchi N., Yanagawa Y., Nagano M., Takahashi M. & Kawahara M. 2016. Conserved roles of fibroblast growth factor receptor 2 signaling in the regulation of inner cell mass development in bovine blastocysts. *Molecular Reproduction and Development* 83, 516–525.
- Aono A., Nagatomo H., Takuma T., Nonaka R., Ono Y. & Kawahara M. 2013. Dynamics of intracellular phospholipid membrane organization during oocyte maturation and successful vitrification of immature oocytes retrieved by ovum pick-up in cattle. *Theriogenology* 79, 1146–1152.e1.
- Arnold S.J. & Robertson E.J. 2009. Making a commitment: cell lineage allocation and axis patterning in the early mouse embryo. *Nature Reviews Molecular Cell Biology* 10, 91–103.
- Betteridge K.J. & Fléchon J.E. 1988. The anatomy and physiology of pre- attachment bovine embryos. *Theriogenology* 29, 155–187.
- Bogliotti Y.S., Wu J., Vilarino M., Okamura D., Soto D.A., Zhong C., Sakurai M., Sampaio R.V., Suzuki

- 1 K., Izpisua Belmonte J.C. & Ross P.J. 2018. Efficient derivation of stable primed pluripotent
2 embryonic stem cells from bovine blastocysts. *Proceedings of the National Academy of Sciences*
3 115, 2090-2095.
- 4 Brackett B.G. & Oliphant G. 1975. Capacitation of rabbit spermatozoa in vitro. *Biology of Reproduction*
5 274, 260–274.
- 6 Crean J.K., Furlong F., Finlay D., Mitchell D., Murphy M., Conway B., Brady H.R., Godson C. & Martin
7 F. 2004. Connective tissue growth factor [CTGF]/CCN2 stimulates mesangial cell migration
8 through integrated dissolution of focal adhesion complexes and activation of cell polarization.
9 *FASEB Journal* 18, 1541–1543.
- 10 Gulnaar A.S., Amy K.W. & David R.B. 1998. Localization of connective tissue growth factor during the
11 period of embryo implantation in the mouse. *Biology of Reproduction* 59, 1207–1213.
- 12 Hall-Glenn F. & Lyons K.M. 2011. Roles for CCN2 in normal physiological processes. *Cellular and*
13 *Molecular Life Sciences* 68, 3209–3217.
- 14 Ivkovic S., Yoon B.S., Popoff S.N., Safadi F.F., Libuda D.E., Stephenson R.C., Daluiski A. & Lyons
15 K.M. 2003. Connective tissue growth factor coordinates chondrogenesis and angiogenesis during
16 skeletal development. *Development* 130, 2779–2791.
- 17 Kelly S.J. 1977. Studies of the developmental potential of 4- and 8-cell stage mouse blastomeres. *Journal*
18 *of Experimental Zoology* 200, 365–376.

- Khan D.R., Dubé D., Gall L., Peynot N., Ruffini S., Laffont L., Bourhis D. Le, Degrelle S., Jouneau A. & Duranthon V. 2012. Expression of pluripotency master regulators during two key developmental transitions: EGA and early lineage specification in the Bovine Embryo. *PLoS ONE* 7, 1–12.
- Kubota S. & Takigawa M. 2015. Cellular and molecular actions of CCN2/CTGF and its role under physiological and pathological conditions. *Clinical Science* 128, 181–196.
- Moussad E.E.-D.A., Rageh M.A.E., Wilson A.K., Geisert R.D. & Brigstock D.R. 2002. Temporal and spatial expression of connective tissue growth factor (CCN2; CTGF) and transforming growth factor β type 1 (TGF- β 1) at the utero– placental interface during early pregnancy in the pig. *Molecular Pathology* 55, 186–193.
- Munoz M., Martin D., Carrocera S., Alonso-Guervos M., Mora M.I., Corrales F.J., Peynot N., Giraud-Delville C., Duranthon V., Sandra O. & Gomez E. 2017. Localisation of stem cell factor, stanniocalcin-1, connective tissue growth factor and heparin-binding epidermal growth factor in the bovine uterus at the time of blastocyst formation. *Reproduction, Fertility and Development* 29, 2127–2139.
- Nagatomo H., Kagawa S., Kishi Y., Takuma T., Sada A., Abe Y., Wada Y., Takahashi M. & Kono T. 2013. Transcriptional wiring for establishing cell lineage specification at the blastocyst stage in cattle. *Biology of Reproduction* 88, 10.1095/biolreprod.113.108993.
- Nishioka N., Inoue K. ichi, Adachi K., Kiyonari H., Ota M., Ralston A., Yabuta N., Hirahara S.,

- Stephenson R.O., Ogonuki N., Makita R., Kurihara H., Morin-Kensicki E.M., Nojima H., Rossant J., Nakao K., Niwa H. & Sasaki H. 2009. The hippo signaling pathway components Lats and Yap pattern Tead4 activity to distinguish mouse trophectoderm from inner cell mass. *Developmental Cell* 16, 398–410.
- Oh S.Y., Song S.E., Seo E.S., Kim K.H., Choi S.J., Suh Y.L., Sadovsky Y. & Roh C.R. 2009. The Expression of Connective Tissue Growth Factor in Pregnancies Complicated by Severe Preeclampsia or Fetal Growth Restriction. *Placenta* 30, 981–987.
- Simmet K., Zakhartchenko V., Philippou-Massier J., Blum H., Klymiuk N. & Wolf E. 2018. OCT4/POU5F1 is required for NANOG expression in bovine blastocysts. *Proceedings of the National Academy of Sciences* 115, 2770-2775.
- Wahab N.A., Brinkman H. & Mason R.M. 2001. Uptake and intracellular transport of the connective tissue growth factor: a potential mode of action. *The Biochemical Journal* 359, 89–97.
- Wu X., Song M., Yang X., Liu X., Liu K., Jiao C., Wang J., Bai C., Su G., Liu X. & Li G. 2016. Establishment of bovine embryonic stem cells after knockdown of CDX2. *Scientific Reports* 6, 1–12.
- Yagi R., Kohn M.J., Karavanova I., Kaneko K.J., Vullhorst D., DePamphilis M.L. & Buonanno A. 2007. Transcription factor TEAD4 specifies the trophectoderm lineage at the beginning of mammalian development. *Development* 134, 3827–3836.

- 1 Zhang K., Rajput S.K., Lee K.-B., Wang D., Huang J., Folger J.K., Knott J.G., Zhang J. & Smith G.W.
2 2015. Evidence supporting a role for SMAD2/3 in bovine early embryonic development: potential
3 implications for embryotropic actions of follistatin. *Biology of Reproduction* 93, 86.
- 4 Zhou Y., Capuco A. V. & Jiang H. 2008. Involvement of connective tissue growth factor (CTGF) in
5 insulin-like growth factor-I (IGF1) stimulation of proliferation of a bovine mammary epithelial cell
6 line. *Domestic Animal Endocrinology* 35, 180–189.
- 7
- 8

Figure legends

Figure 1. *CCN2* mRNA expression dynamics in bovine oocytes and embryos. (A) *CCN2*

expression levels were assessed in oocytes undergoing metaphase during their second meiosis (MII), and in embryos at the 2- (2C), 4- (4C), 8- (8C), 16-cell (16C), morula (MO), and blastocyst (BL) stages, via quantitative RT-PCR (qPCR). Three independent experiments were performed using five oocyte/embryo samples per assay. (B) *CCN2* expression levels were compared between blastocysts and day-14 conceptuses (D14), (where day 0 = artificial insemination). Data are presented as the mean $\pm$ SEM (error bars) of three replicate experiments. Superscript letters indicate statistically significant differences between groups ($P < 0.05$).

Figure 2. *CCN2* protein localization in bovine oocytes and preimplantation embryos.

Representative confocal images showing bovine oocytes (in metaphase of their second meiosis (MII)), and embryos (at the 2- (2C), 4- (4C), 8- (8C), 16-cell (16C), morula (MO), and blastocyst (BL) stages) in which *CCN2* was visualized via immunofluorescent staining (middle column). Nuclei (left column) were visualized via staining with Hoechst 33342. Bar = 75 μ m.

Figure 3. Effects of *CCN2* knockdown (KD) on the expression of pluripotency-related genes.

The relative expression levels of the pluripotency-related genes *NANOG* and *OCT4* in empty

1 vector-injected control, and *CCN2* KD blastocysts (*NANOG*, 7.5%; and *OCT4*, 39.1% compared to
2 the empty control vector, respectively) are shown. Three independent experiments were performed
3 using five embryo samples per assay. Data are presented as the mean $\pm$ SEM (error bars) of three
4 replicate experiments. Superscript letters indicate statistically significant differences between groups
5 ($P < 0.05$).

A **B**

□ Empty ■ *CCN2KD*

Supplemental Table S1 Primer sets for the quantitative RT-PCR analysis

Gene	Accession number	Primer sequence (5'-3')	Annealing temperature	Product length (bp)
<i>OCT4</i>	NM_174580	F: ACATGTGTAAGCTGCGGCCC R: CTTTCGGGCCTGCACAAGGG	58	107
<i>NANOG</i>	NM_001025344	F: CTCGCAGACCCAGCTGTGTG R: CCCTGAGGCATGCCATTGCT	58	198
<i>H2AFZ</i>	NM_174809	F: AGAGCCGTTTGCAGTTCCCG R: TACTCCAGGATGGCTGCGCTGT	58	116