Title	Microsatellite markers developed using a next-generation sequencing technique for <i>Neotrogla</i> spp. (Psocodea: Prionoglarididae), cave dwelling insects with sex-reversed genitalia
Author(s)	Kamimura, Yoshitaka; Abe, Jun; Ferreira, Rodrigo L.; Yoshizawa, Kazunori
Citation	Entomological Science, 22(1), 48-55 https://doi.org/10.1111/ens.12339
Issue Date	2019-03
Doc URL	http://hdl.handle.net/2115/76855
Rights	This is the peer reviewed version of the following article: "Microsatellite markers developed using a next generation sequencing technique for <i>Neotrogla</i> spp. (Psocodea: Prionoglarididae), cave dwelling insects with sex reversed genitalia" Entomological Science 22(1) pp48-55; March 2019, which has been published in final form at https://doi.org/10.1111/ens.12339 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
Type	article (author version)
File Information	2019microsat.pdf

[Instructions for use](#)

**Microsatellite markers developed using a next
generation sequencing technique for *Neotrogla* spp.
(Psocodea: Prionoglarididae), cave dwelling insects
with sex-reversed genitalia**

**Yoshitaka KAMIMURA^{1*}, Jun ABE^{2,3}, Rodrigo L. FERREIRA⁴, and Kazunori
YOSHIZAWA⁵**

*¹ Department of Biology, Keio University, Yokohama, Japan, ² Faculty of Liberal Arts,
Meijigakuin University, Yokohama, Japan, ³ Research Institute for Integrated Science,
Kanagawa University, Hiratsuka, Japan, ⁴ Biology Department, Federal University of
Lavras, Lavras (MG), Brazil, ⁵ Systematic Entomology, School of Agriculture, Hokkaido
University, Sapporo, Japan*

Correspondence: Yoshitaka Kamimura, Department of Biology, Keio University,
Yokohama 223-8521, Japan.

Email: kamimura@fbc.keio.ac.jp

Abstract

The genus *Neotroglia* (Psocodea: Prinoglarididae) comprises four named species from Brazil. Females of this cave-dwelling insect are characterized by a conspicuous penis-like intromittent organ, termed a gynosome, which is inserted into the vagina-like male genitalia during copulation. Another evolutionarily novel structure, the spermathecal plate, enables a female to simultaneously store two freshly deposited spermatophores (consisting of sperm and possibly nutritious substances) in her sperm storage organ (spermatheca). It is unknown whether the two spermatophores are derived from two different males. To investigate the mating ecology and population genetic structures of these insects with sex-reversed genitalia, 16 novel highly polymorphic microsatellite loci were isolated and characterized based on ~2,275 Mbp genomic sequences from an undescribed *Neotroglia* species. Our first screening detected 99,888 candidate loci. Similar to other hemipteroid insects studied thus far, AAT motif microsatellites were conspicuously dominant. We further screened 99 sequences, for which 50 pairs of PCR primers were successfully designed. Sixteen of these primers successfully amplified products of the expected size in the 11 *Neotroglia* sp. individuals collected from two caves. The number of alleles per loci varied from two to nine, with no significant deviation from Hardy–Weinberg equilibrium in either population. Although the caves sampled were only approximately 1 km apart, significant genetic differentiation was detected between the two populations. In total, 13, 12, 13, and 11 loci were cross-amplified in *N. aurora*, *N. brasiliensis*, *N. curvata*, and *N. truncata*, respectively, indicating the applicability of these microsatellite loci for metapopulation genetic studies in multiple *Neotroglia* species.

42 **Key words:** cave populations, genetic differentiation, nuptial gift, sex role reversal,

43 simple sequence repeat (SSR)

44

INTRODUCTION

The cave insect genus *Neotrogla* (Psocodea: Prinoglarididae) comprises four named species from Brazilian caves (Lienhard et al. 2010; Lienhard & Ferreira 2013). Female insects of this genus are characterized by a conspicuous penis-like intromittent organ, termed a gynosome, which is inserted into the male genitalia during copulation (Lienhard et al. 2010; Lienhard & Ferreira 2013; Yoshizawa et al. 2014). The male genitalia or “vaginas,” are relatively simple structures, but possess multiple species-specific pouch-like invaginations, which accommodate the elaborate spines of the gynosome (Yoshizawa et al. 2014). The correlated male and female genital morphologies result in a firm coupling of mating pairs, and copulation lasts ~40–70 h (at least for *N. curvata* Lienhard & Ferreira), during which voluminous materials are transferred to the female sperm storage organ, the spermatheca, to form gigantic spermatophores (Yoshizawa et al. 2014). The spermatophore is formed as a hardened capsule, and females usually possess multiple (up to 11 in *N. brasiliensis* Lienhard) empty capsules in the spermatheca (Yoshizawa et al. 2014). Thus, female *Neotrogla* likely engage in polyandrous mating to obtain multiple spermatophores, not only for fertilization but also as nutrition. Interestingly, *Neotrogla* spp. females have evolved another specialized structure, the spermathecal plate, on their spermatheca. This organ enables females to store two freshly deposited spermatophores simultaneously (Yoshizawa et al. 2014), whereas the females of other psocids (e.g., *Lepinotus patruelis* Pearman) can store only one undigested spermatophore at a time (Wearing-Wilde 1995).

As *Neotrogla* inhabit nutritionally poor cave environments, the evolutionarily novel female structures, the gynosome and spermathecal plate, likely represent

adaptations to obtain as many nuptial gifts from males as possible in an efficient manner. To investigate the mating system of *Neotrogla* underlying the evolution of their sex-reversed genitalia, parentage analysis by means of highly polymorphic genetic markers is indispensable. There are two possible functions of the spermathecal plate. A female may coerce her mating partner to transfer a large volume of ejaculate corresponding to two fresh spermatophores by holding him with the gynosome for a long period of time. A second possibility is that males may provide only one spermatophore at a single bout of copulation, and females may mate with multiple males in succession to obtain more spermatophores. To discriminate between these two possibilities, analyzing the paternal DNA of spermatophores offers a promising approach.

To date, both nymphs and adult *Neotrogla* have been found exclusively in caves. The adults possess fully developed wings and are capable of flight (Lienhard et al. 2010; Yoshizawa & Kamimura, personal observation). However, their wing structures (and observation of living specimens) suggest that they are poor fliers (Ogawa & Yoshizawa 2018), and that migrations may occur only among nearby cave populations. The distributions of the four named species of this genus do not overlap (Lienhard et al. 2010; Lienhard & Ferreira 2013), nor do those of populations that possibly represent undescribed species. Highly polymorphic genetic markers can also be used to reveal their (meta)population genetic structures, providing the basis for their conservation.

Next-generation sequencing techniques are increasingly used to develop large numbers of genetic markers, such as single nucleotide polymorphisms (SNPs) and simple sequence repeats (SSRs: microsatellites), in non-model organisms (reviewed in Ekblom & Galindo 2011; Guichoux et al. 2011; McCormack et al. 2013; Lemmon & Lemmon 2013; Hodel et al. 2016; Wachi et al. 2017). Here, we report the development of 16

microsatellite markers for an undescribed species of *Neotrogla*. Cross-amplification tests were also conducted in all four named species of the genus.

MATERIALS AND METHODS

Genomic DNA extraction and sequencing

Total genomic DNA was extracted from the head and thorax of 11 ethanol-preserved samples (six males and five females) of *Neotrogla* sp. using a DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. The samples were collected on 19 January 2015 from two caves in Varzelândia, Minas Gerais, Brazil: Gruta da Madeira (GM; eight individuals; sample nos. 150119GM1–8) and Lapa do Índio (LI; three individuals; 150119LI1–3). The horizontal projections of these caves are approximately 400 m and 30 m, respectively. The sampled species is apparently very close to *N. aurora* Lienhard, but, owing to their considerable difference in body size, we tentatively treated it as an undescribed species. Some of the DNA extracted from the 11 individuals (20 to 40 µL each) was pooled to make a required volume (350 µL) of DNA solution in AE buffer (10.5 ng/µL) for sequencing with an Illumina HiSeq 2500 platform (Illumina, Inc., San Diego, CA, USA). Library construction and sequencing were provided as a custom service of Eurofins Genomics, Inc. (Tokyo, Japan). A paired-end library of ca. 200-bp inserts was sequenced for 125 bp from both ends. After sequencing, low-quality sequences were trimmed using bcl2fastq Conversion Software v.1.8.4 (Illumina, Inc.).

Microsatellite mining and primer design

Forward and reverse reads of each insert were assembled using BioConductor (Gentleman

et al. 2004) implemented in R v. 3.4.1 (R Core Team 2017), and the text-processing language awk (Aho et al. 1987). Assembled reads shorter than 150 bp were excluded. Our mining strategy of microsatellite markers consisted of two steps. In the first step, all reads were scanned for the presence of microsatellite regions using MSATCOMMANDER 1.08 (Faircloth 2008). Following the methods of several previous studies on other insect groups (Pannebakker et al. 2010; Abe and Pannebakker 2017), we searched for microsatellites with motif lengths of 2–6 bp with minimum repeat numbers of eight, five, five, five, and five for di-, tri-, tetra-, penta-, and hexanucleotides, respectively. Repeats with unit patterns that were circular permutations and/or reverse complements were categorized as one motif type (e.g., an AGC pattern includes AGC, GCA, CAG, GCT, TGC, and CTG; Jurka & Pethiyagoda 1995). In the second step, reads containing microsatellite regions with 13 or more repeats and 70-bp or longer flanking sequences were screened to facilitate subsequent primer designs.

Primer pairs for the amplification of microsatellites were designed using BatchPrimer3 ver. 1.0 (You et al., 2008), using the following search parameters: primer size, 18–22 bp (opt., 20 bp); primer T_m, 58–62°C (opt., 60°C); max T_m difference, 2°C; primer GC%, 40–60%; and other parameters of the default setting.

PCR amplification and genotyping

PCR amplification of microsatellite loci was performed in the 11 *Neotroglia* sp. individuals using an ABI 2720 thermal cycler (Life Technologies, Carlsbad, CA, USA). We adopted Schuelke's (2000) procedure of nested PCR with the M13 universal primer. For this purpose, the 5' ends of forward primers were attached to M13-tails (5'-TGTAACGACGGCCAGT-3') for annealing with fluorescence-labeled M13

universal primers. Nested PCR reactions were conducted in a 10 μ L volume containing 0.1 μ L forward and 0.4 μ L reverse primers (10 μ M each), 0.4 μ L M13 primer (10 μ M) labeled with a fluorescent dye (6-FAM, VIC, NED, or PET), 1 μ L 10 $\times$ PCR buffer, 0.8 μ L 25 mM $MgCl_2$, 0.8 μ L dNTPs (2.5 mM each), 0.05 μ L AmpliTaq Gold 360 DNA polymerase (Applied Biosystems, Carlsbad, CA, USA), and 1 μ L of genomic DNA. The PCR temperature profile consisted of 5 min at 94°C, then 30 cycles of 30 s at 94°C, 45 s at 60°C, and 45 s at 72°C, followed by 8 cycles of 30 s at 94°C, 45 s at 53°C and 45 s at 72°C, and a final extension for 10 min at 72°C. Following electrophoresis with an ABI 3130 capillary sequencer (Life Technologies), the amplified fragments were analyzed using the Peak Scanner software v. 1.0 (Life Technologies).

For cross-amplification tests, DNA was also extracted from the head and thorax of an ethanol-preserved individual of each of the four named species of *Neotroglia*: *N. aurora*, *N. brasiliensis*, *N. truncata* Lienhard, and *N. curvata*. The *N. truncata* sample was collected for observing the genital coupling during copulation (Yoshizawa et al. 2014), and thus initially fixed in hot water (~80°C) and then transferred to 100% ethanol.

Genotypic analysis

We calculated the observed and expected heterozygosity (H_O and H_E) and tested deviations from Hardy–Weinberg equilibrium (HWE) for each cave population (GM and LI) and for the combined data set using GENEPOP 4.6.9 (Rousset 2008). Although the sample sizes were small (eight and three individuals), genetic differentiation between the cave populations was evaluated as a preliminary analysis. For this purpose, F_{ST} and R_{hoST} , which take the allelic identity or the difference between microsatellite allelic sizes, respectively, into account (Weir & Cockerham 1984; Michalakis & Excoffier 1996), were

calculated using GENEPOP 4.6.9 software, and the former was tested using the method of Raymond and Rousset (1995). Linkage disequilibrium was also calculated and tested for all possible combinations of the examined loci. For the GM cave population, non-exclusion probability (NEP), the probability that two unrelated males cannot be discriminated as sperm donors, was calculated for each locus using CERVUS version 3.0.7 (Kalinowski et al. 2007). Significance thresholds in multiple comparisons were corrected using the false discovery rate (FDR; Benjamini & Hochberg 1995).

RESULTS AND DISCUSSION

By sequencing with an Illumina platform, we obtained 32,317,868 reads of 125 bp, corresponding to a 4,040-Mb genomic sequence of the *Neotroglia* sp. The raw sequence data are deposited in the NCBI Sequence Read Archive (accession no.: SPR6871534). After connecting of the corresponding forward and reverse reads (with more than 25 bp overlap), we obtained 13,376,080 reads (with an average length of 170.1 bp; a range of 150–225 bp; and a total of 2,274,799,820 bp) for mining microsatellites.

For this data set, our first screening detected 29,132 (29.2%), 66,086 (66.2%), 3,151 (3.2%), 1,049 (1.1%), and 470 (0.5%) reads (99,888 total) containing di-, tri-, tetra-, penta-, or hexanucleotide repeats, respectively, as candidate sequences for developing microsatellite markers. Pannebakker et al. (2010) mined microsatellites in the genomes of 10 insect species using essentially the same strategy, and found that the compositions of microsatellites varied extensively both between and within insect orders. For example, dinucleotide motifs are more than twice as abundant as trinucleotide motifs in Hymenoptera, whereas trinucleotide motifs are usually more abundant than dinucleotide

motifs in other groups (Pannebakker et al. 2010; Abe & Pannebakker 2017). The microsatellite composition of *Neotrogla* sp. was characterized by the almost two-fold overrepresentation of trinucleotide repeats (66.2%) compared with dinucleotide repeats (29.2%), and the low representation of longer motifs (less than 5% total). Of the trinucleotide repeats, the AAT motif was notably dominant over the other motifs (Fig. 1B). Among the insects studied thus far, these characteristics are shared by the aphid, *Acyrtosiphon pisum* Harris (Homoptera: Aphidoidea: Aphididae) (Pannebakker et al. 2010). For the brown planthopper (*Nilaparvata lugens* (Stål): Homoptera: Delphacidae), Jing et al. (2012) also reported that AAT repeats were the most abundant, but represented only 15.1% of the trinucleotide motifs. Hemipteroid insects, or Paraneoptera (Psocodea + Hemiptera + Thysanoptera) showed an almost consistent A-T bias, especially in the mitochondrial genes of many Psocodea and some Hemiptera species (Yoshizawa & Johnson 2003, 2013). In accordance with this tendency, the CCG motif was the rarest among the trinucleotide repeats (Fig. 1B). The GC-motif is especially rare among dinucleotide motifs in *Neotrogla* sp. (Fig. 1A) as well as in *A. pisum* (Pannebakker et al. 2010).

We further screened 99 reads containing 13 or more repeats and 70-bp or longer flanking sequences, for 50 of which a pair of primers was successfully designed (47 and 3 primer pairs for di- and trinucleotide repeats, respectively; expected PCR product size: 138–198 bp; Table S1).

As we required markers potentially amplifiable with only a small amount of low quality DNA (e.g., genotyping sperm in female storage organs), we selected 16 primer pairs among 50 primer pairs tested that successfully amplified products of expected sizes in all 11 *Neotrogla* sp. individuals for further characterization. All 16 loci were

polymorphic, with two to nine alleles (Table 1). The observed heterozygosity was significantly lower than those expected from the HWE at two loci (Neosp18 and Neosp23) when the data from the two cave populations were merged, but not when they were analyzed separately (Table 1). Although the two caves are only approximately 1 km apart and only a small number of individuals (eight and three) were sampled, these two populations showed significant genetic differentiation ($F_{ST} = 0.043$, Chi2, 34 = 75.3, $P = 0.000059$; $Rho_{ST} = 0.048$). Thus, the observed deviations from HWE, in the direction of overabundance of homozygotes, in the combined data set are likely due to the presence of population-specific alleles: in the LI population, for which only three individuals were sampled, we detected a total of 22 microsatellite alleles at 14 loci specific to this cave population (Table 1).

Based on the analysis of multiple microsatellite loci, a comparative value of F_{ST} (0.042) was reported for the fruitfly *Drosophila americana* Spencer (Diptera: Drosophilidae) between Howell Island (Missouri, USA) and Lake Ashbaugh (Arkansas, USA) populations, which are more than 200 km apart (Schäfer et al. 2006). Similarly, in the ant *Formica exsecta* Nylander (Hymenoptera: Formicidae) Switzerland and Ural populations, which are more than 3,000 km apart, show genetic differentiation measured as $F_{ST} = 0.043$ (Goropashnaya et al. 2007). Thus, although results obtained by different sets of molecular markers cannot be simply compared, the observed genetic differentiation between the *Neotroglia* populations supports the hypothesis that the adults are poor fliers (Ogawa & Yoshizawa 2018) and migration between caves is limited. However, both caves (LI and GM) are located in a continuous limestone outcrop, and several shelters with suitable microclimate conditions certainly exist in between those caves. Accordingly, the moderate genetic differentiation observed may alternatively

indicate that populations present high allegiance in relation to the caves, and dispersion may occur only (or especially) if the environmental conditions of their habitat become unsuitable or hazardous (e.g. by extreme oligotrophic conditions).

After correcting for multiple comparisons using the FDR, no significant linkage disequilibrium was detected among the 16 loci in both the combined data set and in the GM cave population. Due to the small sample size, linkage disequilibrium could not be tested in the LI population.

SNPs are widely used as molecular markers in contemporary genetic studies, including parentage analyses, of non-model organisms. Microsatellites and SNPs both have advantages and disadvantages (Hodel et al. 2016). Each locus of a SNP has up to four alleles (A, T, C, or G). Thus, microsatellites, which usually show much higher allelic diversity, can provide a much more efficient and economical method for analyzing mixed DNA samples (Clayton et al. 1998; Gill 2001). Genetic analyses of sperm stored in a female body, which are likely subject to contamination with female DNA, is such an example. Among the 16 microsatellite loci developed in this study, we observed 6.4 ± 1.9 alleles (mean $\pm$ standard deviation; Table 1). This higher allelic diversity compared to SNPs (usually only two alleles per site; e.g., Lai 2001) resulted in notably low NEPs: 0.1513 on average for the 15 loci (excluding Neosp41, which is monomorphic for the GM population; Table 1). When conjointly used, these markers yield NEP of 4.33×10^{-14} , enabling efficient estimation of the number of males that inseminated the focal females.

Of the 16 primer pairs, five successfully amplified fragments within the expected size range in all four related species examined. The other eight, two, and one primer pairs exhibited successful amplification in three, two, or only one species, respectively (Table 2). The applicability of microsatellite markers usually correlates with the relatedness of

the species (e.g., Jarne & Lagoda 1996). Our preliminary molecular analysis of their phylogeny suggests that the *Neotrogla* sp. sampled in this study is closer to *N. aurora* than to *N. brasiliensis*, *N. curvata*, and *N. truncata* (K. Yoshizawa, unpublished data). However, the number of applicable primers did not vary much among species (from 11 to 13), suggesting their rapid divergence. The lowest amplification success in *N. truncata* (Table 2) may have been due to DNA deterioration owing to fixation and short-term preservation in hot water (see Materials and Methods). The microsatellite markers developed in this study show promise as powerful tools for analyzing the inter- and intraspecific genetic structures of *Neotrogla*.

ACKNOWLEDGMENTS

We are grateful to M. Shinya and I. Yao for advice during the course of this study. This research was supported by a JSPS KAKENHI grant (number 15H04409) to K.Y. and Y. K. R. L. F. is thankful to CNPq (National Council of Technological and Scientific Development) for the research grant (process No. 304682/2014-4).

REFERENCES

- Abe J, Pannebakker BA (2017). Development of microsatellite markers and estimation of inbreeding frequency in the parasitoid wasp *Melittobia*. *Scientific Reports* **7**, Article ID 39879. <https://doi.org/10.1038/srep39879>
- Aho AV, Kernighan BW, Weinberger PJ (1987). *The AWK programming language*. Addison-Wesley Longman Publishing, Boston.
- Benjamini Y, Hochberg Y. (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society* **B57**, 289–300.
- Clayton TM, Whitaker JP, Sparkes R, Gill P (1998) Analysis and interpretation of mixed forensic stains using DNA STR profiling. *Forensic Science International* **91**, 55–70.
- Eklom R, Galindo J (2011) Applications of next generation sequencing in molecular ecology of non-model organisms. *Heredity* **107**,1–15.
- Faircloth BC (2008) MSATCOMMANDER: detection of microsatellite repeat arrays and automated, locus-specific primer design. *Molecular Ecology Resources* **8**, 92–94.
- Gentleman RC, Carey VJ, Bates DM *et al.* (2004) Bioconductor: open software development for computational biology and bioinformatics. *Genome Biology* **5**, Article ID R80. <https://doi.org/10.1186/gb-2004-5-10-r80>
- Gill P (2001) An assessment of the utility of single nucleotide polymorphisms (SNPs) for forensic purposes. *International Journal of Legal Medicine* **114**, 204–210.
- Goropashnaya AV, Fedorov VB, Seifert B, Pamilo P (2007). Phylogeography and population structure in the ant *Formica exsecta* (Hymenoptera, Formicidae) across Eurasia as reflected by mitochondrial DNA variation and microsatellites. *Annales*

301 *Zoologici Fennici* **44**, 462–474.

302 Guichoux E, Lagache L, Wagner S *et al.* (2011) Current trends in microsatellite
303 genotyping. *Molecular Ecology Resources* **11**, 591–611.

304 Hodel RG, Segovia-Salcedo MC, Landis JB *et al.* (2016) The report of my death was an
305 exaggeration: A review for researchers using microsatellites in the 21st century.
306 *Applications in Plant Sciences* **4**, Article ID 1600025.
307 <https://doi.org/10.3732/apps.1600025>

308 Jarne P, Lagoda PJ (1996) Microsatellites, from molecules to populations and back.
309 *Trends in Ecology & Evolution* **11**, 424–429.

310 Jurka J, Pethiyagoda C (1995) Simple repetitive DNA sequences from primates:
311 compilation and analysis. *Journal of Molecular Evolution* **40**, 120–126.

312 Jing S, Liu B, Peng L *et al.* (2012) Development and use of EST-SSR markers for
313 assessing genetic diversity in the brown planthopper (*Nilaparvata lugens* Stål).
314 *Bulletin of Entomological Research* **102**, 113–122.

315 Kalinowski ST, Taper ML, Marshall TC (2007) Revising how the computer program
316 CERVUS accommodates genotyping error increases success in paternity
317 assignment. *Molecular Ecology* **16**, 1099–1106.

318 Lai E (2001) Application of SNP technologies in medicine: lessons learned and future
319 challenges. *Genome Research* **11**, 927–929.

320 Lemmon EM, Lemmon AR (2013) High-throughput genomic data in systematics and
321 phylogenetics. *Annual Review of Ecology, Evolution, and Systematics* **44**, 99–121.

322 Lienhard C, Do Carmo TO, Ferreira RL (2010). A new genus of Sensitibillini from
323 Brazilian caves (Psocodea: 'Psocoptera': Prinoglarididae). *Revue suisse de*
324 *Zoologie* **117**, 611–635.

325 Lienhard C, Ferreira RL (2013). A new species of *Neotrogla* from Brazilian caves
 326 (Psocodea: 'Psocoptera': Prinoglarididae). *Revue suisse de Zoologie* **120**, 3–12.
 327 McCormack JE, Hird SM, Zellmer AJ, Carstens BC, Brumfield RT (2013) Applications
 328 of next-generation sequencing to phylogeography and phylogenetics. *Molecular*
 329 *Phylogenetics and Evolution* **66**, 526–538.
 330 Michalakis Y, Excoffier L (1996) A generic estimation of population subdivision using
 331 distances between alleles with special reference for microsatellite loci. *Genetics*
 332 **142**, 1061–1064.
 333 Ogawa N, Yoshizawa K (2018) Origin and transformation of the in-flight wing-coupling
 334 structure in Psocodea (Insecta: Paraneoptera). *Journal of Morphology* **279**, 517–
 335 530.
 336 Pannebakker BA, Niehuis O, Hedley A, Gadau J, Shuker DM (2010) The distribution of
 337 microsatellites in the *Nasonia* parasitoid wasp genome. *Insect Molecular Biology*
 338 **19**, 91–98.
 339 R Core Team (2017). *R: A language and environment for statistical computing*. R
 340 Foundation for Statistical Computing, Vienna, Austria. Available from URL:
 341 <https://www.R-project.org/>.
 342 Raymond M, Rousset F (1995) An exact test for population differentiation. *Evolution* **49**,
 343 1283–1286.
 344 Rousset F (2008) Genepop'007: a complete re-implementation of the Genepop software
 345 for Windows and Linux. *Molecular Ecology Resources* **8**, 103–106.
 346 Schäfer MA, Orsini L, McAllister BF, Schlötterer C (2006) Patterns of microsatellite
 347 variation through a transition zone of a chromosomal cline in *Drosophila*
 348 *americana*. *Heredity* **97**, 291–295.

349 Schuelk M (2000) An economic method for the fluorescent labeling of PCR fragments.
 350 *Nature Biotechnology* **18**, 233–234.
 351 Yoshizawa K, Johnson KP (2003) Phylogenetic position of Phthiraptera (Insecta:
 352 Paraneoptera) and elevated rate of evolution in mitochondrial 12S and 16S rDNA.
 353 *Molecular Phylogenetics and Evolution* **29**, 102–114.
 354 Yoshizawa K, Johnson KP (2013) Changes in base composition bias of nuclear and
 355 mitochondrial genes in lice (Insecta: Psocodea). *Genetica* **141**, 491–499.
 356 Yoshizawa K, Ferreira RL, Kamimura Y, Lienhard C (2014) Female penis, male vagina,
 357 and their correlated evolution in a cave insect. *Current Biology* **24**, 1006–1010.
 358 You FM, Huo N, Gu YQ *et al.* (2008) BatchPrimer3: a high throughput web application
 359 for PCR and sequencing primer design. *BMC Bioinformatics* **9**, 253.
 360 Wachi N, Matsubayashi KW, Maeto K (2017) Application of next-generation sequencing
 361 to the study of non-model insects. *Entomological Science* Published online: 17
 362 Sep 2017; <https://doi.org/10.1111/ens.12281>
 363 Wearing-Wilde J (1995) The sclerotized spermatophore of the barklouse *Lepinotus*
 364 *patruelis*. *Tissue & Cell* **27**, 1006–1010.
 365 Weir BS, Cockerham CC (1984) Estimating F–statistics for the analysis of population
 366 structure. *Evolution* **38**, 1358–1370.
 367

368 **SUPPORTING INFORMATION**

369

370 Additional Supporting Information may be found online in the supporting information
371 tab for this article.

372 **Table S1** The primer pairs designed and tested in *Neotroglia* sp.

373

374 **Figure Legends**

375

376 **Figure 1.** Relative abundance of (A) di- and (B) trinucleotide repeat motifs found in ca.
377 2,275 Mbp genome sequences of *Neotroglia* sp.