Title	Accumulation patterns and risk assessment of metals and metalloid in muscle and offal of free-range chickens, cattle and goat in Benin City, Nigeria
Author(s)	Ogbomida, Emmanuel Temiotan; Nakayama, Shouta M. M.; Bortey-Sam, Nesta; Oroszlany, Balazs; Tongo, Isioma; Enuneku, Alex Ajeh; Ozekeke, Ogbeide; Ainerua, Martins Oshioriamhe; Fasipe, Iriagbonse Priscillia; Ezemonye, Lawrence Ikechukwu; Mizukawa, Hazuki; Ikenaka, Yoshinori; Ishizuka, Mayumi
Citation	Ecotoxicology and environmental safety, 151, 92-102 https://doi.org/10.1016/j.ecoenv.2017.12.069
Issue Date	2018-04-30
Doc URL	http://hdl.handle.net/2115/77809
Rights	© 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	171228_revised_heavy_metal_manuscript.pdf

[Instructions for use](#)

Accumulation Patterns and Risk Assessment of Metals and Metalloid in Muscle and Offal of
free-Range Chickens, Cattle and Goat in Benin City, Nigeria

Emmanuel Temiotan Ogbomida¹, Shouta M.M. Nakayama², Nesta Bortey-Sam², Balazs
Oroszlany², Isioma Tongo³, Alex Ajeh Enuneku³, Ogbeide Ozekeke³, Martins Oshioriamhe
Ainerua³, Iriagbonse Priscillia Fasipe¹, Lawrence Ikechukwu Ezemonye³, Hazuki Mizukawa²,
Yoshinori Ikenaka^{2,4}, *Mayumi Ishizuka²

¹Ecotoxicology and Environmental Forensic Unit, National Centre for Energy and
Environment, Energy Commission of Nigeria, University of Benin, P.M.B 1154, Benin City,
Nigeria

²Laboratory of Toxicology, Department of Environmental Veterinary Sciences, Graduate
School of Veterinary Medicine, Hokkaido University, Kita 18, Nishi 9, Kita ku, Sapporo
0600818, Japan

³Department of Animal and Environmental Biology (AEB), University of Benin, P.M.B 1154,
Benin City Nigeria

⁴Water Research Group, Unit for Environmental Sciences and Management, North-West
University, Potchefstroom, South Africa

* Corresponding author: Mayumi Ishizuka, e-mail: ishizum@vetmed.hokudai.ac.jp

Laboratory of Toxicology, Department of Environmental Veterinary Sciences, Graduate
School of Veterinary Medicine, Hokkaido University, N18, W9, Kita-ku, Sapporo 060-0818,
Japan

Tel: +81-11-706-6949; Fax: +81-11-706-5105

Abstract

The use of free range animals for monitoring environmental health offers opportunities to
detect exposure and assess the toxicological effects of pollutants in terrestrial ecosystems.
Potential human health risk of dietary intake of metals and metalloid via consumption of offal
and muscle of free range chicken, cattle and goats by the urban population in Benin City was
evaluated. Muscle, gizzard, liver and kidney samples were analyzed for Cr, Mn, Fe, Co, Ni,
Cu, Zn, As, Cd, and Pb concentrations using inductively coupled plasma mass spectrometer
(ICP-MS) while Hg was determined using Hg analyzer. Mean concentrations of metals
(mg/kg ww) varied significantly depending upon the tissues and animal species. Human
health risk estimations for children and adults showed estimated daily intake (EDI) values of
tissues below oral reference dose (RfD) threshold for non essential metals Cd, As, Pb and Hg
thus strongly indicating no possible health risk via consumption of animal based food.
Calculated Hazard quotient (THQ) was less than 1 (<1) for all the metals analysed for both
adult and children. However, Cd and As had the highest value of THQ suggestive of possible
health risk associated with continuous consumption of Cd and As contaminated animal based
foods. Hazard Index (HI) for additive effect of metals was higher in chicken liver and gizzard
for children and chicken liver for adults. Thus, HI indicated that chicken liver and gizzard
may contribute significantly to adult and children dietary exposure to heavy metals. Principal
component analysis (PCA) showed a clear species difference in metal accumulation between

chickens and the ruminants. This study provides baseline data for future studies and also valuable evidence of anthropogenic impacts necessary to initiate national and international policies for control of heavy metal and metalloid content in food items.

Keywords: Heavy metals, Offal, Muscles, Hazard Quotient, Hazard Index

1.0 Introduction

Heavy metals and metalloids are pollutants that pose a great potential threat to the environment and human health on a global scale. They are intrinsic, natural constituents of the environment (Simone *et al.*, 2012) and humans may promote their pollution through anthropogenic activities (Gall *et al.*, 2015) such as extensive application of fertilizers or sewage sludge and pesticides in agriculture, waste disposal, atmospheric deposition, burning of fossil fuels, smelting, mining operations, electroplating, and discharge of wastewater from manufacturing industries including textile industry (Gall *et al.*, 2015).

Heavy metals are widespread and highly persistent in the ecosystems due to their stability and non-degradable properties (Ali *et al.*, 2013). They can enter the food chain via anthropogenic or natural contaminations of air, water or soil and accumulate in animals at the top of the food chain through bio-magnification effects to pose chronic toxicity and serious health risk to man. Heavy metal toxicity in man can diminish mental and central nervous system function, elicit damage to blood composition, as well as the kidneys, lungs, and liver, and reduces energy levels (Amirah *et al.*, 2013). Long-term exposure may result in slow progressive physical, muscular and neurological degeneration that mimics Alzheimer's disease, Parkinson's disease, muscular dystrophy and multiple sclerosis (Amirah *et al.*, 2013).

In recent years, human exposures to heavy metals in Africa have risen as a result of an exponential increase in industrialization, urbanization, municipal solid waste generation and agricultural processes. Also uncontrolled illegal mining activities have added to the enormous amount of heavy metal contamination. The impact of these heavy metals on the environment can be a serious threats to the stability of the ecosystem and human health especially the African children as vulnerable group. In Sub-Sahara Africa environmental pollution by heavy metals has become a serious problem due to non-compliance and enforcement of existing environmental laws and regulations, inadequate monitoring capabilities, weak institutional structures and poor legal framework. In March 2010 large scale of lead poisoning due to artisanal/illegal gold mining activities occurred in Anka and Bukkuyum local government area of Zamfara state North-Western Nigeria leading to 163 deaths of which 111 were children under 5 years of age (MSF, 2010). Similarly, in May 2015, 28 children also died in Angwan Maijero and Angwan Karo, Madaka district of Rafi local government area of Niger State, Nigeria as a result of lead poisoning (MMSD, 2015).

Contamination of the environment and dietary intake of contaminated meat and meat products from food chain has been the most common and principal pathways of human exposure to heavy metals. Animals at the top of food chain may generally accumulate a large amount of heavy metals in their tissue, according to their age, size and feeding habits (Mahmood *et al.*, 2012). In Nigeria, traditional rural animal production systems are mainly based on free-range (FMAWR, 2008). These animals could potentially pick heavy metals through grazing pesticide treated vegetation land, scavenging in open waste dumps for fodder, drinking polluted water from drains and streams and exposure to atmospheric depositions especially from automobile fumes and open burning of solid waste. These heavy metals can accumulate in organs and other fatty tissues, thus providing a major route for human

exposure upon consumption. A recent study conducted by Bortey-Sam et al. (2015) evaluated human health risks from metals and metalloid via consumption of food animals in Tarkwa municipality in the western region of Ghana, an area predominantly known for artisanal gold mining activities. They indicated accumulation and distribution of heavy metals and metalloid in offal and muscles of chicken, goat and sheep with sufficient emphasis on species sensitivity and public health risk through consumption of these animal based foods. The study shows evidence of metal transfer from artisanal gold mining sites to free range animals thus, leading to accumulation of different metals in offal and muscles (Cd, Cr, Cu, Ni, Pb and Zn). They observed that metal accumulation was more pronounced in the liver and kidney of chicken than in muscles. This signifies sensitivity of chicken to heavy metals accumulation and the unique role of animal based food play in metals and metalloid transfer to man with possible health risks. While Bortey-Sam *et al.* (2015) succinctly compare health risks of metals and metalloid via intake of offal and muscles of free range animals from Tarkwa artisanal gold mining sites, recent studies in Nigeria focused mainly on human health risk assessment of some selected heavy metals via drinking water (Maigari et al., 2016), consumption of fishes (Orosun et al., 2016), cow meat (Ihedioha and Okoye, 2013), and illegal gold mining site (Olujimi et al., 2015). However, potential health impacts of municipal solid waste (MSW) in urban areas with specific reference to free range animals reared in proximity to the dumpsites have not been fully evaluated. Causal linkages between exposure to waste and health outcomes for some particular types waste are well established, but the impacts of MSW on free range animals still remain unclear or not prioritized as public health issues. MSW is a growing major challenge to many rapidly urbanizing Africa countries. Furthermore, the full extent of the burden of ill health attributable to exposure of free range animals and the consumption of their muscles and offal has not also be elucidated.

The present study address health risks of metals and metalloid of offal and muscles of free range animals from municipal solid waste sites in Benin City Nigeria, a major environmental problem of urbanisation, population growth and economic development. Free range animals are important in heavy metal studies as bio-indicators of the general environmental pollution status (Roggeman *et al.*, 2013). Therefore, comprehensive study of heavy metal levels in tissues and offal of free range animals are needed to documented the safety of meat and meat products in Nigeria. Although heavy metal contamination have been documented in literatures, information concerning human health risk via consumption of meat and meat products contaminated with heavy metal in Benin City, Edo State still remain very scarce. Therefore, this study describes and compares accumulation, distribution and species sensitivity to heavy metals under different land use pattern. The accumulation of metals and metalloid in free range animals from different locations and ecological regions present a clear picture of environmental pollution status. The distribution pattern of metals in different organs using principal component analysis (PCA) showed species sensitivity to metal and possible human exposure through dietary intake. The outcome of this study will aid in the development of sustainable environmental management options of municipal solid wastes and health intervention policies for affected areas and many other developing countries.

2.0 MATERIALS AND METHODS

2.1 Study area

The study was conducted in Benin City, Edo State around markets, abattoirs and dumpsite areas (Fig 1). Benin City is located 6.3176°N, 5.6145°E and is the capital and largest City of

138 Edo State in Southern Nigeria. The city has a total area of 1,225 km² with an estimated
139 human population of 1,147,188 (NPC, 2006). The city is a commercial centre strategically
140 positioned as the gateway to the northern, eastern and western states of Nigeria. The notable
141 economic activities include breweries, wood carving, traditional brass and bronze casting,
142 wood and timber processing, printing and publishing. Major markets located within the city
143 are Oba, New Benin, Oliha, Uselu, Agbado and Edaiken markets. Domestic and industrial
144 wastes are often generated and discharged into the environment causing public health
145 nuisances. Heaps of solid wastes are often seen littering the streets and market places. The
146 disposal sites are capable of releasing large amounts of harmful chemicals such as heavy
147 metals into the soil which easily found their ways into the food chain.

Fig 1: Map of Benin City showing sampling locations

2.2 Sample Collection

This study was carried out between August 2013 to March 2014 in Benin City, Edo State of Nigeria. Samples were collected from six locations in the three Local Government Areas

which includes Ovia North-East Local Government Area (Oluku abattoir), Egor Local Government Area (Benin Technical College Road abattoir, Uselu market, University of Benin residential quarters, Ekiuwa market) and Oredo Local Government Area (Oliha market) of Edo State (Fig 1). Fresh samples of kidney, liver and muscle of free-range goats (*Capra hircus*) and cattle (*Bos Taurus*) were collected from the abattoirs while kidney, liver and muscle of free-range chickens (*Gallus gallus domesticus*) were collected from live adult chickens after exsanguination and dissection in the laboratory. Samples were kept frozen in labelled plastic bags in the National Centre for Energy and Environment laboratory, University of Benin and were later transported to the Laboratory of Toxicology, Graduate School of Veterinary Medicine, Hokkaido University, Japan and stored in -20°C until analysis. All procedures used in this experiment were performed according to the guidelines of the Committee of Animal Care and Use, University of Benin.

2.3 Sample Preparation and Metal Extraction

All materials and instruments used in metal extraction were washed in 2 % nitric acid (HNO₃) and rinsed at least twice with distilled water. Metals were extracted from liver, kidney and muscle by acid digestion using a closed microwave digestion system (Speed wave MWS-2; Berghof, Germany) according to the method of Nakayama *et al.* (2011). Briefly, 0.5 g of thawed liver, kidney and muscle samples each were placed in a prewashed DAP-60K digestion vessel. 5 ml of 65 % HNO₃ (Kanto Chemical Corp., Tokyo, Japan) and 1ml of (30 %) hydrogen peroxide, H₂O₂ (Kanto Chemical Corp., Tokyo, Japan) spectrometry grade were added. The digestion vessels were capped and placed into a 10-position turntable conditions followed by a ramped temperature program: ramp to 160°C (5 min hold); and increase to 190°C (15 min hold). After digestion in the microwave for 52 min, the samples

were cooled and transferred into plastic tubes and diluted to final volume of 10 ml with distilled deionized water (Milli-Q water). A reagent blank was also prepared using the same procedure.

2.4 Metal Analysis

The quantification of metals in the digested samples were determined using an Inductive Coupled Plasma-Mass Spectrometer (ICP-MS) model 7700 series, Agilent technologies, Tokyo, Japan). The instrument was calibrated using standard solutions of respective metals to establish standard curves before metal analysis. All chemicals and standard stock solutions were analytical reagent-grade (Wako Pure Chemicals, Osaka, Japan), while the water was bi distilled and deionized (Milli-Q; Merck Millipore). The detection limits ($\mu\text{g/L}$) of chromium (Cr), manganese (Mn), Iron (Fe), cobalt (Co), nickel (Ni), copper (Cu), zinc (Zn), arsenic (As), cadmium (Cd), and lead (Pb) were 0.0024, 0.0118, 0.0187, 0.0008, 0.0081, 0.0042, 0.0161, 0.0022, 0.0005 and 0.0022 respectively. Concentrations of metals were expressed in mg/kg wet weight (mg/kg ww).

2.4.1 Total Mercury Analysis

The concentration of total mercury (Hg) was measured by thermal decomposition, gold amalgamation and atomic absorption spectrophotometry (Mercury Analyzer, MA-3000; Nippon Instruments Corporation, Tokyo, Japan), after preparation of calibration standards. Recovery rates of Hg for the certified reference material, DOLT-4 (Dogfish liver, the National Research Council, Canada) ranged from 92 % to 103 % (94.3 ± 4.2 %). The detection limit of total mercury (Hg) was 2.0 pg.

200

201 2.5 Quality Assurance and Quality Control

202 All analytes were subjected to stringent quality control methods. Before sample analysis,
203 appropriate quality assurance procedures and precautions were carried out to ensure
204 reliability of the results. Double distilled deionized water was used throughout the study.
205 Glassware were properly cleaned, and reagents were of analytical grade. Reagents blank
206 determinations were used to correct the instrument readings. Replicate blanks and standard
207 reference materials (SRM), DORM-3 (Fish protein, the National Research Council, Canada)
208 and DOLT-4 (Dogfish liver, the National Research Council, Canada) were used for method
209 validation and quality control. Replicate analysis of these reference materials showed good
210 accuracy (relative standard deviation, RSD, $\leq 3\%$) and recovery rates ranged from 80 % to
211 115 %.

212

213 2.6. Human health risk assessment

214 Risk assessment was done based on the mean concentrations of carcinogenic and non-
215 carcinogenic metals determined in the meat samples using United States Environment
216 Protection Agency (USEPA, 2015) human health risk assessment models. It was assumed
217 that the ingested dose is equal to the adsorbed contaminant dose and that cooking has no
218 effect on the toxicity of heavy metal (Copat *et al.*, 2013). Health risk estimates for heavy
219 metals concentrations in cattle, goat and chicken tissues were computed using three basic
220 standard indices: Estimated Daily Intake (EDI), Hazard Quotient (THQ) and Hazards index
221 (HI). Two hypothetical age/weight categories were used. 30 kg was assumed for children and
222 60 kg for adults (Ihedioha and Okoye, 2013).

223

224

225

226
227
228
229

230

231
232
233
234
235
236
237
238
239

240

241

242
243
244

2.6 Data Analysis

2.6.1 Estimated daily intake (EDI) of toxic metals

The estimated daily intakes (EDI) of heavy metals (Cr, Mn, Fe, Co, Ni, Cu, Zn, As, Cd, Hg and Pb) was expressed as mg/kg/day and was dependent on both the concentration of heavy metals in offal or muscle and the amount of consumption. The EDI of metals for children and adults were determined by the following equation according to Bortey-Sam *et al.* (2015).

$$EDI = \frac{MC \times FDC}{BW} \dots\dots\dots \text{Eqn. 1}$$

where MC is average concentration of metal in food (µg/g, on fresh weight basis); FDC represents the average offal/muscle daily consumption in this region (g/person/d); BW is the average body weight. It was assumed that local inhabitants consumed an average liver, muscle and gizzard of 150 and 100 g/day for adult (60kg in BW) and children (30 kg in BW) respectively (Bortey-Sam *et al.*, 2015). For kidney, it was assumed that inhabitants consumed 10 and 8 g/day for adults and children respectively (Ihedioha and Okoye, 2013), because the size of kidneys are relatively smaller and is not the favorite compared with liver, muscle, or gizzard. The metal intakes were compared with the tolerable daily intakes for metals recommended by FAO/WHO (2010).

2.6.2 Target Hazard Quotient (THQ)

The Target Hazard Quotient (THQ) was used to assess risk associated with non-carcinogenic and carcinogenic health effect. The health risks from consumption of offal and muscle of chicken, goat and cattle by the local inhabitants were assessed based on the hazard quotient

(THQ). The THQ is a ratio of determined dose of a pollutant to a reference dose level. If the ratio is less than 1, the exposed population is unlikely to experience obvious adverse effects. The method of estimating risk using THQ was provided in the USEPA Region III risk-based concentration table (USEPA, 2007), and is based on the equation below:

$$THQ = \frac{Efr \times ED \times FIR \times MC}{RfD \times BW \times AT} \times 10^{-3} \quad \text{Eqn. 2}$$

where THQ is target hazard quotient; EFr is exposure frequency (365 days/year); ED is exposure duration (70 years); FIR is food ingestion rate (g/person/d); MC is average concentration of metal in food (µg/g, on fresh weight basis); RfD is the oral reference dose (mg/kg/d); BW is the average body weight, adult (60 kg); children (30 kg); AT is the average exposure time (365 days/year number of exposure years, assuming 70 years in this study). Oral reference doses (RfD) were based on 3E-04, 1E-03, 5E-04, 4E-03, 3E-01, 4E-02, 1.4E-01, 2E-02, 7E-01, 15E-1 mg/kg/d for As, Cd, Hg, Pb, Zn, Cu, Mn, Ni, Fe, Cr respectively (USEPA IRIS, 2007). There is no consensus about the RfD for Co, However, the RfD for Co was estimated as 4.3E-02 (Food and Nutrition Board, 2004).

2.6.3. Hazardous Index (HI)

To estimate the human health risk for more than one heavy metal, the hazard index was calculated by the summation of the hazard quotients for all heavy metals in the Equation below.

$$HI = \sum THQ_n \quad \text{Eqn 3}$$

It assumes that the magnitude of the adverse effect will be proportional to the sum of multiple metal exposures. It also assumes similar working mechanisms that linearly affect the target

organ. When HI is equal to or less than one ($HI \leq 1$), it indicates no appreciable health risk, while if $HI > 1$, then it indicates a reason for health concern (Cao *et al.*, 2015). Hence, $HI < 1$ means no hazard; $1 > HI < 10$ means moderate hazard while greater than 10 means high hazard or risk (Ukoha *et al.*, 2014).

2.6.4 Statistical analysis

Statistical analyses were performed using SPSS version 16.0 statistical software (SPSS Inc., Illinois, USA) after data were normalized by log transformation. To analyze differences in the distribution of metals in tissues of free-range chickens, cattle and goats, analysis of variance (ANOVA) and Tukey was done and differences were considered statistically significant with p value < 0.05 . Principal component analysis (PCA) based on log transformed data was done to determine the distribution pattern of metals in organs, using JMP statistical software v.10 (SAS Institute). The principal components were extracted with eigen values > 1 . Statistical differences between the different organs (liver, muscle, gizzard and kidney) were determined by one-way analysis of variance (ANOVA). The limit of significant level was accepted at $p < 0.05$.

3.0. Results and Discussion

3.1 Concentration of Metals and Metalloid in Offal and Muscle

The mean concentrations of these metals in offal and muscle of free-range chickens, cattle and goats from Benin city are presented in Table1. The composition profile of the metals in the three animal food species are Cr (0.050 - 0.29), Mn (0.13 - 3.6), Fe (10 - 310), Co (0.0040

289 - 0.054), Ni (0.013 - 0.049), Cu (0.50 - 88), Zn (4.4 - 47), As (0.0040 - 0.081), Cd (0.0010 -
290 0.89), Pb (0.0090 - 0.59) and Hg (nd - 0.019) mg/kg ww. The results showed that chicken
291 liver accumulated the highest levels of Fe, Mn and Hg, chicken kidney (Pb, Cd and As) and
292 chicken gizzard (Ni and Cr) while cattle liver accumulated Co, Cu and Zn. This findings
293 conform with Demirezen and Uruc (2006) who reported that the primary site of heavy metals
294 are the liver and kidney due to exposure and physiological responses of animals to detoxify
295 the system. The reason for high accumulation of metals in the gizzard could be attributed to
296 the function of the gizzard which is the primary reservoir of food materials and organ for
297 grinding up food partilces. Free range local chickens according to Nesheim *et al.* (1979) have
298 been found to harbour the greatest pollutants in their body system as they have no oral
299 discrimination for food hence pick whatever they find edible in the environment. In this study
300 free-range chicken accumulated more metals than the cattle and goat which can be attributed
301 to their foraging dietary habits. The high levels of Pb, Cd, Hg, As, Ni, Cr, Mn and Fe in the
302 offal of the chickens could have come from anthropogenic activities in the environment and
303 feeds fed on by the chickens. Several studies have shown that diet is the primary pathway for
304 metal accumulation in animals and man. For example, toxicological investigations on bovines,
305 ducks and poultry have shown a direct correlation between metal concentrations in animal
306 feed and animal tissues accumulation (Sedki et al., 2003; Kim and Koo, 2007). Also, it has
307 been reported that rice and vegetables grown in mining areas contaminated by heavy metals
308 have posed a great potential health risk for consumers, who suffered from serious cancers
309 (Zhuang et al., 2009). Plants uptake from soil contribute to the circulation of heavy metal in
310 the food chain through their active and passive absorption, accumulation in tissues as well as
311 subsequent grazing by animals that depends on them for nutrients or consumption by man.
312 This reflect that metal-rich diets lead to high tissue metal accumulation. Metal accumulation

in liver and kidney also reflect short-term exposure and are most appropriate target organ to test metals.

3.1.1 Human Essential Metals

The 11 elements of highest concern within the European Community are: As, Cd, Co, Cr, Cu, Hg, Mn, Fe, Zn, Ni, and Pb (Stankovic and Stankovic, 2013). Some of these elements are actually necessary for humans in trace amounts (Co, Cu, Cr, Mn, Ni), while more than necessary amounts of these trace elements and some others are carcinogenic or toxic, affecting among others, the central nervous system (Hg, Pb, As, Mn), the kidneys or liver (Hg, Pb, Cd, Cu) or skin, bones, or teeth (Ni, Cd, Cu, Cr) (Rai and Pal, 2002; Chen et al., 2008; Lavery et al., 2009; Jovic et al., 2012; Stankovic and Jovic 2012; Markovic et al., 2012). Essential metals are significant to humans because they are important in various metabolic enzymes and constituents of cells (Maret, 2016). The frequent consumption of animal based food contaminated with heavy metals could increase the level in humans. Large doses of essential metals can damage living organisms or in some cases results in toxic effects (Nagajyoti *et al.*, 2010).

Cu is a component of various enzymes (Amaral et al., 2008; Osredkar and Sustar, 2011). It is involved in collagen synthesis and in the normal development of connective tissues, nerves and immune system (Amaral et al., 2008). High intake has been recognized to cause adverse health problems such as liver and kidney damage (Amaral *et al.*, 2008). Concentration of Cu was highest in cattle and chicken liver (Table 1). This is because the liver act as the primary storage organ for Cu and maintains Cu homeostasis for animal. In the present study the results shows that the average concentration of Cu in cattle liver (88.016 ± 36.349 mg/kg

ww) is higher than the permissible limits of 20 mg/kg set by USDA (2006) and ECR (2006) and also 30 mg/day maximum limit set by WHO (1996).

Mn is an essential element for both animals and plants; its deficiency results in severe skeletal and reproductive abnormalities in mammals (Sivaperumal *et al.*, 2007). Mn concentration was found to be high in livers and kidneys of chicken and cattle liver. The mean concentrations of Mn ranged from 0.130 ± 0.037 (chicken muscle) to 3.607 ± 1.198 mg/kg ww (chicken liver). Mn levels in chicken liver, kidney and cattle liver and were above the WHO (1996) reference standard of 0.5 mg/kg. Mn has been found to be toxic in excess; in brain it can cause a Parkinson-type syndrome (Aschner, 2000).

Fe is one of the most abundant transition element, and probably the most well known metal in biologic systems participating in a wide variety of metabolic processes, including oxygen transport, DNA synthesis, and electron transport. High levels of iron may lead to tissue damage, as a result of the formation of free radicals. The concentration of Fe in this study was highest in chicken liver and kidney. The mean concentration of Fe ranged from 10.349 mg/kg ww in cattle muscle to 305.484 mg/kg ww in chicken liver (wet weight). This result reflect the dominance of Fe concentration observed in other animal based foods by Medeiro *et al.* (2012). The most serious forms of Fe overload is acute poisoning while chronic Fe intoxication is associated with genetic and metabolic diseases leading to repeated blood transfusions (Fraga and Oteiza, 2002).

Zn is the most abundant intracellular component which is needed to maintain some important biological functions such as genetic stability, gene expression, DNA repair and programmed cell death (Hussein and Khaled, 2014). High levels of zinc can cause pancreatitis, anemia, muscle pain, and acute renal failure (Pais and Benton Jones, 1997). In this study the mean concentration of Zn ranged from 4.430 mg/kg ww in chicken muscle to

46.704 mg/kg ww in cattle liver. This result conform with Okoye and Ugwu (2010) who reported elevated zinc levels in goats bred in Nigeria. The high concentrations of Zn in animal based food can be attributed to food chain transfer. The values of zinc reported in this study were below the Codex Alimentarium Commission maximum permissible limit of 50 mg/kg for muscle and 80 mg/kg for edible offals (Codex, 1995).

Cr is an important element known to enhance the action of insulin in man (Mertz, 1998), a hormone critical to the metabolism and storage of carbohydrate, fat, and protein in the body (Porte *et al.*, 2003). The mean concentrations of Cr in this study ranged from 0.051 (chicken kidney) to 0.286 mg/kg ww (chicken gizzard). Our data is similar to Iwegbue, *et al.* (2008) who reported 0.33 ± 0.14 mg/kg in chicken meat consumed in southern Nigeria. Cr detected in the gizzard of chicken could be attributed to chicken diet.

Ni plays an important role in the biology of plants and animals. Overexposure to Ni can cause decreased body weight, heart and liver damage and skin irritation (Homady *et al.*, 2002). The average concentration of Ni observed in this study ranged from $(0.017 \pm 0.011 - 0.049 \pm 0.037$ mg/kg ww) in chicken $(0.019 \pm 0.003 - 0.030 \pm 0.020$ mg/kg ww) in cattle and $(0.013 \pm 0.005 - 0.028 \pm 0.020$ mg/kg ww) in goat. The levels of Ni in the kidneys of chicken, cattle and goat showed no significant difference ($P > 0.05$) while the livers and muscles of cattle showed significant difference when compared with chicken and goat. Chicken gizzard and muscle had the highest value of Ni. The observed levels in this study were below the 0.5 mg/kg maximum permissible limit of Ni in food according to FAO/WHO (2000).

3.1.2 Human Non-Essential Metals

Human non essential metals Cd, As, Pb and Hg have no biological role in living organisms but are tolerated at very low concentration and also inhibit metabolic activity at higher concentrations (Pandey and Madhuri, 2014) . They are potentially toxic to living organisms. Animals are constantly exposed to these toxicant through dietary intake. As shown in Table 1 the concentration of Cd ranged from 0.890 ± 1.035 mg/kg ww in chicken kidney to 0.001 ± 0.00 mg/kg ww in muscle of cattle and goat.

The range of Cd level in this study is lower than result obtained by Ihedioha and Okoye (2013) in swine and broiler chicken ($ND - 40.14$ mg/kg ww). The levels of Cd observed in free range chicken kidney were below the maximum permissible limit of 1.0 mg/kg (FAO/WHO, 2000). However, Doganoc (1996) observed higher levels of Cd in the liver and kidney of chicken which exceeded the official tolerance levels. The ability of liver and kidney to accumulate high concentration of Cd is a common feature of chickens and many other animals and may be due to the detoxification function of the organs (Stoyke *et al.*, 1995). Similar levels of Cd has also been reported by Okoye and Ugwu (2010) in liver (0.35 mg/kg) and kidney (0.83 mg/kg) of goats from Nigeria. Cd is presently listed as number 7 of 275 of the most hazardous substances in the environment, behind As, Pb and Hg (ATSDR, 2013). Chronic Cd poisoning impair calcium metabolism leading to softening of bones, fractures, and skeletal deformations (Stankovic *et al.*, 2011). Liver and kidney tissues are the two main sites of Cd storage and these organs accumulate considerable amounts of Cd, about 40 - 80 % of the body burden (WHO, 2007).

Pb is the second element (after arsenic) on the ASTDR's top 20 list of the most poisoning heavy metals (ASTDR, 2013). Its target organs are bones, brain, blood, kidneys, reproductive and cardiovascular systems, and thyroid gland (Homady *et al.*, 2002). The levels of Pb in the offal and muscle of free range cattle, goat and chicken ranged from 0.58826 mg/kg ww in

chicken kidney to 0.009 mg/kg ww in goat muscle (Table 1). Pb is the most frequently reported causes of accidental poisoning in domestic animals, with cattle as the most commonly affected species because they are very inquisitive and commonly ‘taste test’ new finds (Khalafalla *et al.*, 2015). Absorbed Pb is stored mainly in the liver and kidney, and like Cd, it accumulates in tissues of animals (Khalafalla *et al.*, 2015). However, there was significant differences when the concentration of Pb in chicken kidney and liver were compared with goat kidney and liver (Tukey test: $p < 0.05$) Table 1. Similarly, the level of Pb in the muscles were statistically different (Tukey test: $p < 0.05$). Apart from the chicken kidney, the gizzard was second highest in the accumulation of Pb. This could be attributed to the feeding pattern of chicken. In this study, concentration of Pb in the kidney (0.588 ± 0.703 mg/kg ww), and gizzard (0.239 ± 0.327 mg/kg ww) of free range chicken were slightly above the recommended limits of 0.5 and 0.1 mg/kg ww for offal and muscle by ECR (2006) and USDA (2006) respectively.

The result of this study indicated that the highest concentration of Hg was found in chicken liver (0.034 ± 0.036 mg/kg ww) and kidney (0.030 ± 0.031 mg/kg ww) while the lowest concentrations was found in cattle liver (0.001 ± 0.000) and goat muscle (0.0010 ± 0.002) mg/kg ww respectively. This is in conformity with Landis *et al.* (2010) that birds accumulate Hg in the kidney and liver with less in the muscle. The concentrations of Hg in the kidney of free-range chicken, cattle and goat were not statistically different (Tukey test; $p > 0.05$) However, levels of Hg in chicken liver were significantly different ($P < 0.05$) when compared with the levels in the livers of cattle and goat. In the muscle, the levels were significantly different ($p < 0.05$) with high value in chicken muscle. In this study the values of Hg in offal and muscles were below the permissible limit of 0.05 mg/kg established by ECR (2006) and USDA (2006).

The metalloid As was detected in all the tissues and offal samples of the three animal foods. Concentrations of As in muscle and offal of free-range chicken, goat and cattle ranged from (0.040 ± 0.034-0.081 ± 0.073 mg/kg ww), (0.007 ± 0.011-0.008 ± 0.011 mg/kg ww) and (0.007 ± 0.011 -0.008 ± 0.004 mg/kg ww) respectively (Table1). The distribution of As in liver, kidney, gizzard and muscle of chicken was found to be higher than that of goat and cattle ($p < 0.05$). Mean concentration of 0.08 mg/kg ww in the kidney of free-range chicken in Benin City was found to be higher than the USDA (2006), ECR (2006) and SAC/MOHC (2005) standard of 0.05 mg/kg.

3.1.3 Principal Component Analysis (PCA) of Metals Distribution

Principal component analysis (PCA) of heavy metals and metalloid distribution in offal and muscle samples of animal based food was performed and the results are presented in Fig. 2. PCA was used to determine the multivariate structure of the data, to highlight the possible trend and to further identify the potential source of heavy metals contamination. In the interpretation of PCA patterns, factor loadings greater than 0.71 are typically considered excellent while those less than 0.32 are regarded as very poor (Nowak, 1998). Factor loading determines the relationships between the variables and each factor. The first principal component (PC1) of metals and metalloid in liver of cattle (Fig. 2) showed 38.7 % of the total variance with the loading of the variables Mn, Fe, Ni, Zn and Pb. Fe and Zn in PC1 were highly associated, displaying high factor loading values (0.827 and 0.843) suggesting metals may come from the same source. Also, zinc and iron share common dietary sources. The second component (PC2) accounts for 14.7 % of the total variance of heavy metals in the chicken liver and is dominated by Co, As and Cd while the third component PC3 accounts for 12.5 % of the total variance in goat liver with a positive loading for Cu. The first principal

component which explains 38.7 % may be controlled by a long term anthropic activity such as municipal solid waste found to be seriously contaminated by toxic metals (Naveedullah *et al.*, 2013) such as electronic waste in the area for a long period of time in agreement with the clustering of variables in Group I. Also metallic zinc and iron are used in a wide variety of applications but when exposed to the atmosphere may be subject to corrosion that may result in the slow release of small amounts of zinc and iron into the environment. The PCA results also showed a clear separation between chicken (F) grouped on one side and the ruminants; goat (G) and cattle (C) clustered on the other side. The results obtained from PCA of kidney of the three animal based foods Fig 2 indicates that Mn, Co, As, Cd and Pb are associated with high factor loadings of 0.857, 0.823 and 0.776 (Co, Cd and As) in cattle kidney. This association strongly suggests that these variables have a similar source and may be attributed to cattle grazing in polluted environment or drinking wastewater with heavy metals. The first PC which explains 33.5 % of the total variance and loads in the cattle kidney confirm the interpretation of their anthropogenic origin. The occurrence of Cd and Hg likely reflects the increased use of nickel-cadmium batteries in electronic items and a variety of Hg-containing devices and lamps such as compact florescence lamps (CFLs) which are disposed in municipal solid waste. The second component (PC2) showed 15.6 % total variance of heavy metals in chicken kidney which contributes Cu and Zn with factor loading of 0.840 and 0.784. Cu and Zn may be controlled by a long-term anthropic activity such as pesticides usage (Yang *et al.*, 2014). Copper is usually considered as a marker element of environmental waste dumps, domestic waste water, combustion of fossil fuels and wastes, wood production, phosphate fertilizer production, and natural sources. The third component (PC3) alone explains 11.6 % of the variance of Cr and Ni in the goat kidney while Hg showed negative loadings. In the cattle muscle the first component (PC1) with 33 % of variance comprises positive factor loading of Mn, Fe, Cu and Zn which could be linked to

anthropogenic and natural sources with some contributions from the municipal waste system. The second component (PC2) explains 19.3 % of the total variation, which exhibited a high positive factor loading of 0.755, 0.815 and 0.955 on Ni, Cd and Pb. These are significant environmental pollutants with anthropogenic origin especially from industry, production of lead-acid batteries, metal products (solder and pipes) and poor urban waste management which are common in developing countries. The third component (PC3) showed a high positive factor loading with 14 % variance of As and Hg. This suggest that levels of these toxic metals can be primarily attributed to anthropogenic influences. Animals are constantly exposed to As and Hg through contaminated drinking water, feedstuff, grasses, vegetables etc. However, arsenic has been one of the most common causes of inorganic chemical poisoning in farm animals and also a main threat to human health. Africa today is becoming a dumping ground for electronic waste especially used computers, mobile phones etc from developed countries which contains hazardous metals such as lead, cadmium and mercury. All these electronic wastes subsequently end up in municipal waste dump sites which consists of a mix of combustible and non-combustible materials and subsequently over time contribute to increased environmental problem.

3.2 Human Health Risk Assessment

3.2.1 Estimated Dietary Intake of Metals

The Estimated Daily Intakes (EDI) of metals and metalloid taken by an adult or children per day through the consumption of free range chicken, cattle and goat offal and tissue are presented in tables 2. Meat and edible offal are considered a significant part of most diet in Nigeria. They serve as major sources of animal derived protein, vitamins, essential trace

elements and fatty acids. However, meat and edible offal consumption had been identified as an important pathway of human exposure to pollutants. The degree of heavy metal toxicity to human population depends upon their daily intake. The EDI values for human non essential metals Cd, As, Pb and Hg for adults person of 60 kg body weight show that chicken liver had the highest EDI value. The EDI values obtained were generally lower than the respective oral reference dose (RfD) threshold for human non essential metals Cd, As and Pb except in Hg. Our results strongly indicate no possible health risk associated with animal based food consumed by the local population of Benin city. The RfD threshold dose or intake is an estimate of a daily exposure to the human population that is likely to be without an appreciable adverse risk of deleterious noncancer effects when exposure is long-term (lifetime) intake. If the predicted intake is less than the oral reference dose ($THQ < 0.1$), it indicates almost no possibility of an adverse health effects. However if the intake exceeds the reference dose ($THQ > 0.1$) does not necessarily imply that adverse health effects are expected only that a conservative reference oral dose (RfDo) is exceeded. The results of this study are similar to those obtained in an earlier study on offal and muscles of sheep, goat and chicken (Bortey-Sam *et al.*, 2015) where they reported low daily intakes of As, Cd, Hg, Pb and Mn in food animals compared to the provisional tolerable daily intake guidelines. Chronic exposure to low doses of Cd, As, Pb and Hg above their safe threshold in humans and animals could therefore result to many non- carcinogenic hazards such as neurologic involvement, headache and liver malfunctions. For human essential heavy metals Co, Cr, Cu, Mn, Fe, Zn and Ni, the estimated daily intake of Co, Cr, Cu, Mn, Fe, Zn and Ni in this study ranged from 0.005 - 0.135, 0.008 - 0.716, 0.698 - 220.041, 0.185 - 9.018, 13.808 - 229.838, 4.117 - 116.760 and 0.005 - 0.123 $\mu\text{g/kg bw/day}$ respectively. Based on this values, the EDI is lower than the RfD guidelines which strongly showed that there is no serious health risk via consumption of these food based animals. The calculated EDIs of As, Cd, Hg, Pb and Mn

in offal and muscle samples of the three food animals from Benin city were below the WHO (2000), FSA (2006) and FAO/WHO (2010) tolerable daily intakes (Table 2). However, the EDI values in children were higher than the adult values. Therefore, children are especially more vulnerable to acute, sub-acute and chronic effects of ingestion of chemical pollutants, since they (children) consume more (twice of the amount) of food per unit of body weight as adults (ENHIS, 2007). As a result, intakes of these toxic metals through food could be higher for children in Benin city. The EDI values from other studies agreed with our findings. Specifically, the findings of Tongo and Ezemonye (2015) on predicted EDI values for children associated with residual pesticide levels in edible tissues of slaughtered cattle in Benin City conform with our current findings. Also, Onwukeme et al. (2014) and Oforka et al. (2102) predicted the risk of heavy metals via consumption of cow meat and offal in Enugu and chicken meat in Port-Harcourt, Nigeria. However, our findings in this study regarding metals contents in meat and offal suggest that their consumption are not free of risks as they may present detrimental health concerns through a lifetime. Due to interspecific differences however, the EDI levels of heavy metals were higher in chicken than other species.

3.2.2 Target Hazard Quotient (THQ)

Estimated health risk from consumption of offal and muscle from chicken, cattle and goat were assessed based on the THQ. A THQ value greater than 1 would indicate that a potential health risk may exist. THQ value in this study was less than 1 which means the exposed population is unlikely to experience any obvious adverse effects (USEPA, 2000) from the intake of individual metals through the consumption of offal and muscle (Fig. 3). Generally, consumption of muscle tissue poses less health risk to man compared to kidney and liver because it accumulates low level of metal. However, muscle tissue seldom come in direct

contact with metals as it is totally covered externally by the skin that in many ways prevent the penetration of metals. It is also not an active site for detoxification, and therefore, transport of metals from other tissues to muscle does not seem to arise (Elnabris *et al.*, 2013). Our findings also indicates that the severity of metals and metalloids are more in liver and kidney. THQ for Cd and As were highest in chicken liver and gizzard but less than 1 in both adults and children. This may indicate potential significant health risk associated with continuous consumption of chicken liver and gizzard. Therefore, some attention should be paid for the Cd and As content in chicken because their THQ values for adults and children are not far below the threshold value of 1. The higher levels of the cadmium shown in the chicken liver relative to other tissues may be attributed to the high coordination of metallothionein protein with the cadmium. In addition, the liver act as the principal organ responsible for the detoxification, transportation, and storage of toxic substances. Also, feed habits and composition can influence the retention of heavy metals in the liver. Therefore continuous consumption of chicken liver may lead to biomagnification suggesting possible significant risk to man. The of the three animal based showed lowest values an indication that there might be no public health risk from the consumption of muscle of free-range chicken, cattle and goats. Our findings agrees with Tyokumbur (2016) and Ihedioha and Okoye (2013) who reported THQ value less than 1 for chicken and cow meat. The essential metals Cu, Zn, Fe and Mn were also clearly below all the permissible limits for human consumption.

3.2.3 Estimated Hazard Index (HI)

The hazard index (HI) of the mixture of different non-essential metals Cd, As, Pb and Hg was calculated by adding together all the component hazard quotients (THQ). The potential health effects of the mixture were further analyzed and investigated if the HI value is equal to or

greater than 1. Hence, $HI < 1$ means no hazard; $1 > HI < 10$ means moderate hazard while greater than 10 means high hazard or risk (Ukoha *et al.*, 2014). This approach assumes that all components have similar joint action for residents ingesting metals via consumption of tissues and offal of free range chicken, cattle and goat. The calculated HI values for both adults and children are given in Figure. 4. The HI for human non essential metals Cd, As, Pb and Hg revealed that chicken liver (2.19) and chicken gizzard (1.17) for children were greater than 1. This suggests that there is overtly adverse health effects from the consumption of chicken liver and gizzard. In adult, HI value showed that chicken liver (1.64) and gizzard (0.88) suggests higher health risk from the consumption of chicken liver than chicken gizzard. This implies that when metals combined they pose a greater health risk than any lone metal. All heavy metals analyzed had a hazard quotient of less than one ($THQ < 1$). The greater the value of (THQ) and (HI) (if > 1), the greater the level of risk associated with offal and muscle consumption. Our results agreed with MahMoud *et al.* (2014) that reported HI greater than 1 for Cd indicating high potential risk to the Egyptian consumers.

4.0 Conclusion

This study was developed to provide baseline information on heavy metals and metalloid concentration in free range animals; chicken, cattle and goats sampled in Benin City, Edo State, Nigeria. As expected, metal concentrations showed a great variability. The levels of metals in the offal and muscle suggested that significant differences existed in the concentrations of elements across offal and muscles of animal species. It was observed that free-range chickens reared in Benin City accumulated large quantities of Cd above maximum permissible limits compared to cattle and goats. This may be largely related to the organism mobility, food preferences, or other factors with respect to environment pollution most likely

from dumpsites. The estimation of risk showed that adverse health effect may occur from prolonged consumption of chicken liver with high concentrations of Cd and may lead to accumulation in the human body thereby causing metal toxicity. Overall, our findings show that consumption of chicken liver and gizzard present more health than muscle tissue especially to children. Contamination is due mainly to environmental pollution as the animals are free ranging. Therefore, it is strongly recommended that appropriate steps should be taken to identify ways to reduce the health risk via consumption of contaminated chicken offal especially the gizzard and liver as well as restrict chickens from roaming and scavenging for food in dumpsites in order to safeguard the food chain. Further research is required to investigate the dietary intake of heavy metals and possible health risks from other communities where animal free ranging is predominant. This will help to provide consumption advisories to the public about the risks of eating contaminated leaner meat products in an effort to avoid the dangers of high cholesterol.

5.0 Acknowledgments

This work was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan awarded to M. Ishizuka (No. 24405004 and No. 24248056) and Y. Ikenaka (No. 26304043, 15H0282505, 15K1221305), S.M.M. Nakayama (No. 16K16197) and the foundation of JSPS Core to Core Program (AA Science Platforms) and Bilateral Joint Research Project (PG36150002 and PG36150003). We also acknowledge financial support from The Soroptimist Japan Foundation, The Nakajima Foundation, The Sumitomo foundation and The Nihon Seimei Foundation. This research was supported by JST/JICA, SATREPS (Science and Technology Research Partnership for Sustainable Development). We are grateful to Mr. Takahiro Ichise (Laboratory of Toxicology,

Graduate School of Veterinary Medicine, Hokkaido University) for technical support. We acknowledged the National Centre for Energy and Environment (Energy Commission of Nigeria), University of Benin, Benin City, Edo State Nigeria for support and provision of its ancillary facilities used in the preparation and preservation of samples. We gratefully appreciate the assistance of Prof Lawrence N I Ezemonye and staff of the National Centre for Energy and Environment during field sampling.

REFERENCES

Agency for Toxic Substances and Diseases Registry (ATSDR), 2013. The Priority List of Hazardous Substances that will be the subject of toxicological profiles. <http://www.atsdr.cdc.gov/SPL/index.html>

Ali, H., Khan, E., Sajad, M.A., 2013. Phytoremediation of Heavy Metals-Concepts and Applications. *Chemosphere* 91:869-881

Amaral, A.F.S., Arruda, M., Cabral, S., Rodrigues, A.S., 2008. Essential and non-essential trace metals in scalp hair of men chronically exposed to volcanogenic metals in the Azores, Portugal. *Environment International* 34: 1104–1108

Amirah, M.N., Afiza, A.S., Faizal, W.I.W., Nurliyana, M.H., Laili, S., 2013. Human Health Risk Assessment of Metal Contamination through Consumption of Fish. *Journal of Environment Pollution and Human Health* 1(1): 1-5.

Aschner, M., 2000. Manganese: Brain Transport and Emerging Research Needs *Environ. Health Perspect* 108 (3):429-432.

644 Bortey-Sam, N., Nakayama, S.M.M., Ikenaka, Y., Akoto, O., Baidoo, E., Yohannes, Y.B.,
645 Mizukawa, H., Ishizuka, M., 2015. Human health risks from metals and metalloid via
646 consumption of food animals near gold mines in Tarkwa, Ghana: Estimation of the daily
647 intakes and target hazard quotients (THQTHQs). *Ecotoxicology and Environmental*
648 *Safety* 111:160–167.

649 Cao, L.L., Yang, J., Wang, P., Wang, L., Tian, H. T., Zou, M. K., Xie, J., 2015. Potentially
650 Toxic Metal and Metalloid Fractionation Contamination in Sediments of Daya Bay, South
651 China Sea. *Kemija u industriji*. 64(5-6):255–262

652 Chen, Y.C., Serrell, N., Evers, C.D., Fleishman, J.B., Lambert, F.K., Weiss, J., Mason, R.P.,
653 Bank, S.M., 2008. Methylmercury in marine ecosystems - from sources to seafood consumers.
654 *Environ Health Perspect* 116(12):1706–1712.

655 Codex, 1995. Alimentarius Commission (CAC) Doc. no. CX/FAC 96/17. Joint FAO/WHO
656 food standards programme. Codex general standard for contaminants and toxins in foods;

657 Copat, C., Conti, G. O., Signorelli, C., Marmioli, S., Sciacca, S., Vinceti, M., Ferrante, M.,
658 2013. “Risk Assessment for Metals and PAHs by Mediterranean Seafood,” *Food and*
659 *Nutrition Sciences*, 4. 10-13.

660 Demirezen, D., Uruc, K., 2006. Comparative Study of Trace Elements in Certain Fish, Meat
661 and Meat Products. *Food Chemistry*, 32:215-222.
662 <http://dx.doi.org/10.1016/j.meatsci.2006.03.012>

663 Doganoc, D.Z., 1996. Lead and cadmium concentrations in meat, liver and kidney of
664 Slovenian cattle and pigs from 1989-1993. *Food Additives and Contaminants*. 13:237-241.

665 Elnabris, K.J., Muzyed, S.K., El-Ashgar, N.M., 2013. Heavy metal concentrations in some
666 commercially important fishes and their contribution to heavy metals exposure in Palestinian
667 people of Gaza Strip (Palestine) *Journal of the Association of Arab Universities for Basic and*
668 *Applied Sciences*, 13:44–51

669 European Commission Regulation(ECR)., 2006. No 1881/2006, Setting maximum levels for
670 certain contaminants in foodstuff. *Official Journal. of the European Union*, L 364: 5-24

671 European Environment and Health Information System (ENHIS)., 2007. Exposure of
672 children to chemical hazards in food. Fact Sheet No.4.4, CODE: RPG4-Food-EX1. World
673 Health Organization.

674 FAO/WHO, 2000. Report of the 32nd Session of the codex committee of the food additives
675 Contaminants. Beijing People's Republic of China, 20-24 March.

676 FAO/WHO, 2010. Summary Report of the Seventy-third Meeting of JECFA. Joint
677 FAO/WHO Expert Committee on Food Additives, Geneva.

678 FMAWR, 2008. National Programme for Food Security. Federal Ministry of Agriculture and
679 Water Resources (FMA&WR), Abuja, p.107.

680 Food and Nutrition Board., 2004. Dietary Reference Intakes [DRIs]: Recommended Intakes
681 for Individuals. National Academy of Sciences, Washington, DC, USA.

682 Fraga, C.G., Oteiza, P.I., 2002. Iron toxicity and antioxidant nutrients. *Toxicology* 180: 23–32.

683 Gall, J. E., Boyd, R.S., Rajakaruna, N., 2015. Transfer of heavy metals through terrestrial
684 food webs: a review. *Environmental Monitoring and Assessment*. 187(4):201-222

685 Homady, M., Hussein, H., Jiries, A., Mahasneh, A., Al-Nasir, F., Khleifat, K., 2002. Survey
686 of some heavy metals in sediments from vehicular service stations in Jordan and their effects
687 on social aggression in prepubertal male mice. *Environmental Research* 89: 43–49.

688 Hussein, A., Khaled, A., 2014. Determination of Metals in Tuna Species and Bivalves from
689 Alexandria, Egypt. *Egyptian Journal of Aquatic Research*, 40:9-17.
690 <http://dx.doi.org/10.1016/j.ejar.2014.02.003>

691 Ihedioha, J.N., Okoye, C.O.B., 2013. Dietary intake and health risk assessment of lead and
692 cadmium via consumption of cow meat for an urban population in Enugu State, Nigeria,
693 *Ecotoxicology and Environmental Safety*. 93:101–106.

694 Iwegbue, C.M.A., Nwajei, G.E., Iyoha, E. H., 2008. Heavy Metal Residues of Chicken Meat
695 and Gizzard and Turkey Meat Consumed In Southern Nigeria. *Bulgarian Journal of*
696 *Veterinary Medicine*. 11(4):275-280.

697 Jovic, M., Onjia, A., Stankovic, S., 2012. Toxic metal health risk by mussel consumption.
698 *Environ Chem Lett* 1:69–77.

699 Khalafalla, F. A., Abdel-Atty, N. S., Abd-El- Wahab, M. A., Ali, O. I., Rofaida, B. A., 2015.
700 Assessment of heavy metal residues in retail meat and offals. *J Am Sci*. 11(5):50-54.

701 Kim, J., Koo, T.H., 2007. Heavy metal concentrations in diet and livers of black-crowned
702 night heron *Nycticorax nycticorax* and grey heron *Ardea cinerea* chicks from Pyeongtaek,
703 Korea. *Ecotoxicology* 16:411–416.

704 Landis, W., Sofield, R., Yu, M.H., Landis, W.G., Sofield, R.M., 2010. Introduction to
705 Environmental Toxicology: Molecular Substructures to Ecological Landscapes, Fourth
706 Edition 542 pages

707 Lavery, J.T., Kemper, M.C., Sanderson, K., Schultz, G.C., Coyle, P., James, G., Mitchell,
 708 G.J., Seuront, L., 2009. Heavy metal toxicity of kidney and bone tissues in South Australian
 709 adult bottlenose dolphins (*Tursiops aduncus*). *Mar Environ Res* 67:1–7.

710 Mahmood, Q., Rashid, A., Ahmad S.S., Azim, M.R., Bilal, M., 2012. Current status of toxic
 711 metals addition to environment and its consequences. In: Anjum NA, *et al.*, editors. The Plant
 712 Family Brassicaceae: Contribution Towards Phytoremediation, *Environmental Pollution*.

713 Maigari, A.U., Ekanem, E.O., Garba, I.H., Harami, A., Akan, J.C., 2016. Health Risk
 714 Assessment for Exposure to Some Selected Heavy Metals via Drinking Water from
 715 Dadinkowa Dam and River Gombe Abba in Gombe State, Northeast Nigeria. *World Journal*
 716 *of Analytical Chemistry* 4(1): 1-5.

717 Maret, W., 2016. The Metals in the Biological Periodic System of the Elements: Concepts
 718 and Conjectures. *Int. J. Mol. Sci.* 17(1): 66. doi:10.3390/ijms17010066.

719 Markovic, J., Joksimovic, D., Stankovic, S., 2012 Trace elements concentrations determined
 720 in collected wild mussels in the coastal area of southeastern Adriatic, Montenegro. *Arch Biol*
 721 *Sci Belgrade* 64(1):265–275.

722 Médecins Sans Frontières (MSF)., 2010. Field News; Lead Poisoning In Zamfara State,
 723 Nigeria; Nigeria 16.07.10. Available from corresponding website;
 724 [http://www.msf.org.au/from-the-field/fieldnews/](http://www.msf.org.au/from-the-field/fieldnews/field-news/article/lead-poisoning-inzamfara-state-igeria.html) field- news/article/lead-poisoning-
 725 inzamfara-state-igeria.html.

726 Medeiros, R.J., dos Santos, L.M.G., Freire, A.S., Santelli, R.E., Braga, A.M.C.B., Krauss,
 727 T.M., Jacob, S.C., 2012. Determination of inorganic trace elements in edible marine fish from
 728 Rio de Janeiro State, Brazil. *Food Control* 23: 535-541.

729 Mertz, W., 1998. Interaction of chromium with insulin: a progress report. *Nutrition Reviews*;
730 56:174-7.

731 Ministry of Mines and Steel Development (MMSD)., 2015. Lead poisoning in Angwan
732 Maijero and Angwan Karo, Madaka districts of Rafi LGA, Niger State, Nigeria. A report by
733 the Federal Ministry of Mines and Steel Development

734 Nagajyoti, C.P., Lee, D.K., Sreekanth, M.V.T., 2010. Heavy metals, occurrence and toxicity
735 for plants: a review. *Environmental Chemistry Letters* 8:199–216

736 Nakayama, S.M.M., Ikenaka, Y., Hamada, K., Muzandu, K., Choongo, K., Teraoka, H.,
737 Mizuno, N., Ishizuka, M., 2011. Metal and metalloid contamination in roadside soil and wild
738 rats around a Pb–Zn mine in Kabwe, Zambia. *Environ. Pollut.* 159:175–181.

739 National Population Commission (NPC)., 2006. The report of National population
740 commission on the census conducted in Nigeria.

741 Naveedullah, M.Z.H., Yu, C., Shen, H., Duan, D., Shen, C., Lou, L., Chen, Y., 2013. Risk
742 Assessment of Heavy Metals Pollution in Agricultural Soils of Siling Reservoir Watershed in
743 Zhejiang Province, *China BioMed Research International* Volume 2013, Article ID 590306,
744 10 pages <http://dx.doi.org/10.1155/2013/590306>

745 Nesheim, M.C., Austic, R.E., Cand, L., 1979. Poultry Production. 12th Edition, Lea and
746 Febiger, Philadelphia, 123-125.

747 Nowak, B., 1998. Contents and relationship of elements in human hair for a non-
748 industrialised population in Poland. *Science of the Total Environment* 209(1):59–68. Oforka,
749 N.C., Osuji, L.C., Onwuachu, U.I., 2102. Estimation of Dietary intake of Cadmium, Lead,

750 Manganese, Zinc and Nickel due to consumption of chicken meat by inhabitants of Port-
 751 Harcourt Metropolis, Nigeria. *Archives of Applied Science Research*, 4 (1):675-684.

752 Okoye, C.O.B., Ugwu, J.N., 2010. Impact of environmental cadmium, lead, copper and zinc
 753 on quality of goat meat in Nigeria. *Bull. Chem. Soc. Ethiop.* 24(1):133-138.

754 Olujimi, O.O., Oputu, O., Fatoki, O., Opatoyinbo, O.E., Aroyewun, O.A., Baruani, J., 2015.
 755 Heavy Metals Speciation and Human Health Risk Assessment at an Illegal Gold Mining Site
 756 in Igun, Osun State, Nigeria. *J Health Pollution* 8: 19-32

757 Onwukeme, V.I., Nwako, P.M., Mgbemena, M.N., Onuoha, M.U., 2014. Dietary Exposure to
 758 Heavy Metals via Consumption of Some Vegetables and Parts of Cow Meat by Adults in
 759 Awka, Nigeria. *IOSR Journal of Environmental Science, Toxicology and Food Technology*
 760 *(IOSR-JESTFT)*. 8 (5): 01-06

761 Osredkar, J., Sustar, N., 2011. Copper and Zinc, Biological Role and Significance of
 762 Copper/Zinc Imbalance. *J Clinic Toxicol* S3:001. doi: 10.4172/2161-0495.S3-001. Pais, I.,
 763 Benton Jones Jr., J., 1997. The Handbook of Trace Elements. Saint Lucie Press, Boca Raton,
 764 Florida, 223p.

765 Orosun, M.M., Tchokossa, P., Orosun, R.O., Akinyose, F.C., Ige, S.O., Oduh, V.O., 2016.
 766 Determination of Selected Heavy Metals and Human Health Risk Assessment in Fishes from
 767 Kiri Dam and River Gongola, North eastern Nigeria. *J Phys Chem Biophys* 6: 229.
 768 doi:10.4172/2161-0398.1000229

769 Pandey, G., Madhuri, S., 2014. Heavy Metals Causing Toxicity in Animals and Fishes. *Res. J.*
 770 *Animal, Veterinary and Fishery Science*. 2(2): 17-23,

771 Porte Jr., D., Sherwin, R.S., Baron, A., 2003. Ellengerg and Rifkin's Diabetes Mellitus, 6th
 772 Edition. McGraw-Hill, New York,

773 Rai, N., Pal, A., 2002. Health hazards of heavy metals. EnviroNews ISEB India vol 8 No. 1.
 774 In: International conference on plants and environmental pollution (ICPEP-2). Proceedings of
 775 the 2nd international conference on plants and environmental pollution, Lucknow, India, 4–9
 776 Feb 2002. <http://isebindia.com/issuewise.html>

777 Roggeman S., Van Den Brink, N., Van Praet, N., Blust, R., Bervoets, L., 2013. Metal
 778 exposure and accumulation patterns in free-range cows (*Bos taurus*) in a contaminated natural
 779 area: Influence of spatial and social behaviour. *Environmental Pollution* 172:186-199

780 SAC/MOHC (The Standardization Administration China/Ministry of Health, China), 2005.
 781 Maximum levels of contaminants in foods (GB2762-2005). Beijing.

782 Sedki, A., Lekouch, N., Gamon, S., Pineau, A., 2003. Toxic and essential trace metals in
 783 muscle, liver and kidney of bovines from a polluted area of Morocco. *Sci. Total Environ.*
 784 317: 201–205

785 Simone, M., Fernando, G.C., Maria, L.P., 2012. Heavy Metals and Human Health,
 786 Environmental Health - Emerging Issues and Practice, Prof. Jacques Oosthuizen (Ed.), ISBN:
 787 978-953-307-854-0, In Tech, p227-246

788 Sivaperumal, P., Sankar, T.V., Viswanathan Nair, P.G., 2007. Heavy metal concentrations in
 789 fish, shellfish and fish products from internal markets of India vis-a-vis international
 790 standards. *Food Chemistry*. 102:612–620

791 Stankovic, S., Jovic, M., 2012. Health risks of heavy metals in the Mediterranean mussels as
 792 seafood. *Environ Chem Lett* 2: 119–130.

793 Stankovic, S., Jovic, M., Stankovic, R.A., Katsikas, L., 2011. Heavy metals in seafood
 794 mussels. Risks for human health. In: Lichtfouse E et al (eds) *Environmental Chemistry for a*
 795 *Sustainable World*, 1(64):311–375.

796 Stankovic, S., Stankovic, A.R., 2013. Bioindicators of Toxic Metals. In: Lichtfouse E *et al*
 797 (eds), Green Materials for Energy, Products and Depollution. *Environmental Chemistry for a*
 798 *Sustainable World*, 3: 151-228

799 Stoyke, M., Doberschütz, K.D., Lusky, K., 1995. Heavy metal contents (cadmium, lead and
 800 mercury) in selected feedstuff, organs and tissue in cattle found different sites of
 801 Brandenburg. *Mengen. Spurenelemente. Arbeitstag* 15: 269-276.

802 Tongo, I., Ezemonye L., 2015. Human health risks associated with residual pesticide levels in
 803 edible tissues of slaughtered cattle in Benin City, Southern Nigeria. *Toxicology Reports* 2:
 804 1117–1135

805 Tyokumbur, E.T., 2016. Evaluation of Cadmium (Cd) in Domestic Chicken Meat and Offal
 806 and Associated Health Risk Assessment in Ibadan. International Journal of Pure and Applied
 807 Zoology. 4 (2): 155-160

808 Ukoha, P.O., Ekere, N.R., Udeogu, U.V., Agbazue, V.E., 2014. Potential health risk
 809 assessment of heavy metals concentrations in some imported frozen fish species consumed in
 810 Nigeria. *Int. J. Chem. Sci.* 12 (2): 366–374.

811 USDA, 2006. Foreign Agricultural Service GAIN Report Global Agriculture Information
 812 Network Voluntary Report-public distribution GAIN Report Number CH 6064, China
 813 Peoples Republic of FAIRS Product. Specific maximum levels of contaminants in foods, Jim
 814 Butterworth and Wu Bugang.

815 USEPA, 2000. Risk-based Concentration Table. Philadelphia PA: United States
 816 Environmental Protection Agency, Washington DC.

817 USEPA, 2007. Integrated Risk Information System-Database (IRIS), Philadelphia PA;
 818 Washington, DC.

819 USEPA, 2015. Integrated Risk Information System. Available online:
 820 <http://www.epa.gov/iris/rfd.htm>

821 Uzairu, A., Harrison, G.F.S., Balarabe, M.L., Nnaji, J.C., 2009 “Concentration levels of trace
 822 metals in fish and sediment from Kubanni River, Northern Nigeria,” *Bulletin of the Chemical*
 823 *Society of Ethiopia*, 23 (1): 9–17.

824 WHO, 1996. Trace Elements in Human Nutrition and Health. Geneva.

825 WHO., 2007. Health risks of heavy metals from long-range transboundary air pollution.
 826 World Health Organization 2007. WHO Regional Office for Europe Copenhagen, Denmark
 827 ISBN 978 92 890 7179 6.

828 Yang, P., Yang, M., Mao, R., Shao, H., 2014. Multivariate-Statistical Assessment of Heavy
 829 Metals for Agricultural Soils in Northern China. *The Scientific World Journal* Volume 2014
 830 Article ID 517020, 7 pages <http://dx.doi.org/10.1155/2014/517020>.

831 Zhuang, P., Zou, B., Li, N.Y., Li, Z.A., 2009. Heavy metal contamination in soils and food
 832 crops around Dabaoshan mine in Guangdong, China: implication for human health. *Environ.*
 833 *Geochem. Health* 31:707–715.

Figures and Tables

Table 1: Mean concentrations (mg/kg ww) of heavy metal in different tissues of free range chicken (*Gallus gallus domesticus*), Cattle (*Bos taurus*) and Goat (*Capra hircus*)

		Cr	Mn	Fe	Co	Ni	Cu	Zn	As	Cd	Pb	Hg
CL	Mean $\pm$ SD	0.064 $\pm$ 0.014	2.785 $\pm$ 0.440*	91.935 $\pm$ 20.228	0.054 $\pm$ 0.023	0.019 $\pm$ 0.003	88.016 $\pm$ 36.349	46.704 $\pm$ 5.400	0.006 $\pm$ 0.003	0.044 $\pm$ 0.013	0.036 $\pm$ 0.018	0.001 $\pm$ 0.000*
CHL	Mean $\pm$ SD	0.077 $\pm$ 0.053	3.607 $\pm$ 1.198*	305.484 $\pm$ 187.966	0.049 $\pm$ 0.026	0.018 $\pm$ 0.009	4.362 $\pm$ 1.017	43.363 $\pm$ 13.084	0.069 $\pm$ 0.091*	0.291 $\pm$ 0.338*	0.171 $\pm$ 0.193*	0.034 $\pm$ 0.036*
GL	Mean $\pm$ SD	0.051 $\pm$ 0.030	1.656 $\pm$ 0.526*	65.328 $\pm$ 27.470	0.033 $\pm$ 0.030	0.013 $\pm$ 0.005	11.883 $\pm$ 8.118	22.731 $\pm$ 3.197	0.005 $\pm$ 0.005*	0.023 $\pm$ 0.009*	0.044 $\pm$ 0.032*	0.003 $\pm$ 0.002*
CK	Mean $\pm$ SD	0.051 $\pm$ 0.017	1.109 $\pm$ 0.137*	82.850 $\pm$ 16.001	0.030 $\pm$ 0.008	0.030 $\pm$ 0.020	4.187 $\pm$ 0.400	30.463 $\pm$ 11.204	0.008 $\pm$ 0.004*	0.238 $\pm$ 0.065	0.052 $\pm$ 0.002	0.006 $\pm$ 0.003
CHK	Mean $\pm$ SD	0.089 $\pm$ 0.076	2.468 $\pm$ 0.565*	101.521 $\pm$ 28.623*	0.049 $\pm$ 0.028	0.020 $\pm$ 0.004	2.904 $\pm$ 0.326	24.700 $\pm$ 3.368	0.081 $\pm$ 0.073*	0.890 $\pm$ 1.035*	0.588 $\pm$ 0.703*	0.030 $\pm$ 0.031
GK	Mean $\pm$ SD	0.060 $\pm$ 0.097	0.821 $\pm$ 0.302	80.520 $\pm$ 22.191*	0.025 $\pm$ 0.012	0.028 $\pm$ 0.020	2.765 $\pm$ 0.298	15.232 $\pm$ 2.026	0.008 $\pm$ 0.011*	0.123 $\pm$ 0.158*	0.072 $\pm$ 0.066*	0.019 $\pm$ 0.012
CM	Mean $\pm$ SD	0.261 $\pm$ 0.216	0.163 $\pm$ 0.067	28.918 $\pm$ 8.863	0.004 $\pm$ 0.002	0.023 $\pm$ 0.008	0.978 $\pm$ 0.312	40.658 $\pm$ 20.102	0.007 $\pm$ 0.011*	0.001 $\pm$ 0.000	0.013 $\pm$ 0.010	ND*
CHM	Mean $\pm$ SD	0.148 $\pm$ 0.166	0.130 $\pm$ 0.037*	10.349 $\pm$ 5.706	0.004 $\pm$ 0.002	0.035 $\pm$ 0.032	0.499 $\pm$ 0.156	4.430 $\pm$ 0.673	0.023 $\pm$ 0.018*	0.003 $\pm$ 0.003*	0.024 $\pm$ 0.029*	0.004 $\pm$ 0.003*
GM	Mean $\pm$ SD	0.106 $\pm$ 0.159	0.187 $\pm$ 0.050*	24.792 $\pm$ 6.287	0.006 $\pm$ 0.006	0.017 $\pm$ 0.011	0.825 $\pm$ 0.137	24.541 $\pm$ 6.737	0.007 $\pm$ 0.011*	0.001 $\pm$ 0.001*	0.009 $\pm$ 0.006*	0.001 $\pm$ 0.002*
CHG	Mean $\pm$ SD	0.286 $\pm$ 0.131	0.547 $\pm$ 0.204	80.856 $\pm$ 36.112	0.010 $\pm$ 0.006	0.049 $\pm$ 0.037	3.069 $\pm$ 1.412	24.690 $\pm$ 4.774	0.047 $\pm$ 0.026	0.126 $\pm$ 0.111	0.239 $\pm$ 0.327	0.003 $\pm$ 0.003

CL: Cattle liver; CHL: Chicken Liver; GL: Goat Liver; CK: Cattle kidney; CHK: Chicken kidney; GK: Goat kidney; CM: Cattle muscle; CHM: Chicken muscle; GM: Goat muscle. Letter (*) between columns in the same organ indicates statistical difference (Tukey's test; $p < 0.05$).

ND = Not detectable

Table 2: Estimated Daily Intake ($\mu\text{g/kg bw/day}$) of Heavy Metals.

Sample Name		Cr	Mn	Fe	Co	Ni	Cu	Zn	As	Cd	Pb	Hg
Cattle Liver	Adult	0.161	6.962	229.838	0.135	0.048	220.041	116.760	0.014	0.109	0.089	0.002
	Child	0.214	9.282	306.451	0.180	0.065	293.388	155.681	0.018	0.145	0.119	0.002
Cattle Muscle	Adult	0.653	0.408	72.296	0.011	0.057	2.445	101.644	0.009	0.002	0.034	0
	Child	0.871	0.545	96.394	0.015	0.076	3.260	135.526	0.012	0.003	0.045	0
Cattle Kidney	Adult	0.008	0.185	13.808	0.005	0.005	0.698	5.077	0.001	0.040	0.009	0.001
	Child	0.013	0.296	22.093	0.008	0.008	1.117	8.123	0.002	0.063	0.014	0.002
Chicken Liver	Adult	0.192	9.018	763.710	0.122	0.045	10.905	108.409	0.172	0.728	0.427	0.085
	Child	0.257	12.024	1018.281	0.163	0.060	14.540	144.545	0.229	0.970	0.569	0.113
Chicken Muscle	Adult	0.369	0.326	25.874	0.010	0.088	1.248	11.076	0.057	0.007	0.061	0.009
	Child	0.492	0.434	34.498	0.014	0.117	1.664	14.768	0.076	0.009	0.081	0.013
Chicken Kidney	Adult	0.015	0.411	16.920	0.008	0.003	0.484	4.117	0.014	0.148	0.098	0.005
	Child	0.024	0.658	27.072	0.013	0.005	0.774	6.587	0.022	0.237	0.157	0.008
Chicken Gizzard	Adult	0.716	1.367	202.140	0.026	0.123	7.673	61.725	0.117	0.314	0.596	0.008
	Child	0.954	1.823	269.520	0.034	0.164	10.231	82.300	0.156	0.419	0.795	0.010
Goat Liver	Adult	0.127	4.140	163.320	0.083	0.032	29.707	56.828	0.012	0.058	0.111	0.007
	Child	0.169	5.521	217.760	0.111	0.043	39.610	75.771	0.016	0.077	0.147	0.009
Goat Muscle	Adult	0.264	0.468	61.980	0.014	0.044	2.063	61.353	0.018	0.002	0.022	0.003
	Child	0.352	0.624	82.640	0.018	0.058	2.750	81.804	0.024	0.003	0.030	0.003
Goat Kidney	Adult	0.025	0.342	33.550	0.010	0.012	1.152	6.347	0.003	0.051	0.030	0.008
	Child	0.040	0.547	53.680	0.017	0.019	1.843	10.155	0.005	0.082	0.048	0.012

Liver

Kidney

Muscle

Figure 2: Distribution patterns of heavy metals in tissues of free range chicken (*Gallus gallus domesticus*), Cattle (*Bos taurus*) and Goat (*Capra hircus*) characterized by PCA (C: Cattle; F: Chicken; G: Goat).

Table 3: Rotated component matrix of heavy metal distribution in tissues of Cattle, Chicken and Goat

Metals	LIVER			KIDNEY			MUSCLE		
	Factor 1	Factor 2	Factor 3	Factor 1	Factor 2	Factor 3	Factor 1	Factor 2	Factor 3
Cr	0.57295	0.442054	0.3478	0.125716	0.171982	0.586742	0.322429	0.076322	0.560515
Mn	0.746037	0.006849	-0.16617	0.634222	0.451342	-0.13242	0.788645	-0.16787	0.019669
Fe	0.826638	0.164758	-0.30036	0.597652	0.21665	-0.02082	0.860973	-0.23712	-0.11556
Co	0.152608	0.787065	0.167181	0.857003	-0.07441	0.03258	0.563319	0.349646	0.119911
Ni	0.601517	0.471661	0.41585	-0.08658	-0.52952	0.525105	-0.07011	0.754954	0.005269
Cu	0.014404	-0.0496	0.761673	-0.15836	0.839743	0.129631	0.647074	0.065516	-0.49891
Zn	0.843317	0.115863	0.118478	0.385707	0.784202	0.149639	0.702072	-0.23161	-0.39122
As	0.063011	0.784748	-0.27553	0.775946	0.027437	-0.2946	-0.19852	0.069948	0.832405
Cd	0.220286	0.782195	-0.38159	0.823136	-0.0745	-0.10271	-0.1393	0.81524	0.170016
Pb	0.683165	0.19457	-0.12312	0.618564	0.168359	0.287615	-0.03435	0.955143	0.012434
Hg	0.324645	0.200308	-0.54916	0.223473	0.005372	-0.69775	-0.22616	0.055939	0.769395

Figure 3: Hazard quotient (THQ) of heavy metals (Cr, Mn, As, Cd, Pb and Hg) in children and adults via consumption of cattle liver CL, chicken liver CHL, goat liver GL, cattle kidney CK, chicken kidney CHK, goat kidney GK, cattle muscle CM, chicken muscle CHM, goat muscle GM, chicken gizzard CHG

Fig 4: Hazard Index (HI) for children and adults through the consumption of offal and muscle