



Title	Synthesis of Chiral Sphingolipids and their Stereochemical Effects on Induction of Neurite Outgrowth and Sphingomyelin Synthase Activity [an abstract of dissertation and a summary of dissertation review]
Author(s)	SAJEER, KOOLATH
Citation	北海道大学. 博士(生命科学) 甲第13943号
Issue Date	2020-03-25
Doc URL	<a href="http://hdl.handle.net/2115/78171">http://hdl.handle.net/2115/78171</a>
Rights(URL)	<a href="https://creativecommons.org/licenses/by/4.0/">https://creativecommons.org/licenses/by/4.0/</a>
Type	theses (doctoral - abstract and summary of review)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	SAJEER_KOOLATH_abstract.pdf (論文内容の要旨)



[Instructions for use](#)

## Abstract of Doctoral Dissertation

Degree requested: Doctor of Life Science

Applicant's name: Koolath Sajeer

### Title of Doctoral Dissertation

Synthesis of Chiral Sphingolipids and their Stereochemical Effects on Induction of Neurite Outgrowth and Sphingomyelin Synthase Activity

(キラルスフィンゴ脂質の合成とその立体化学による神経突起伸長及びスフィンゴミエリン合成酵素活性への影響について)

Sphingolipids are essential components of cell membrane having significant structural and functional roles in the cells. They are first discovered from the brain extract in 1870s. Sphingolipids are named after the mythological Sphinx because of their enigmatic nature. They are known to protect cell surface by forming plasma membrane lipid bilayer. They contain backbone of sphingoid bases, derivative of amino alcohol sphingosine, sphinganine or phytosphingosine. Sphingolipids help to define the structural properties of membranes and lipoproteins and also have important roles in cell signaling, cell-cell interaction, cell recognition, and help to regulate cell growth and differentiation. Sphingolipids are important biomolecules and they are chiral. Chirality of sphingolipids are very important because they play very crucial role in the biological system. Most of the drugs are chiral. Naturally the stereochemistry of most common sphingolipid ceramide has *D-erythro* stereochemistry. So, the Chirality of sphingolipids play very crucial role in the drug discovery because one enantiomer of drug may be a useful medicine for a disease, but another isomer may be inactive or may be toxic to that disease.

Sphingolipids, ceramide and ganglioside GM3 have attracted intense research interest in drug discovery due to their role in metabolic disorders. Recent study revealed that, sphingolipid metabolizing enzymes such as sphingomyelin synthase (SMS) and GM3 synthase are potential drug target to cure metabolic disorders. The deficiency of SMS has been reported to control the development of obesity, fatty liver and type 2 diabetes, since then researchers are trying to establish selective and potent SMS inhibitors for the therapeutic application of metabolic disorders. On the other hand, gangliosides play an important role in the case induction of neurite out growth. This phenomenon is very important for the recovery of nervous system after injury.

Initial goal of our study was to discover the stereochemical effects of chiral ganglioside GM3 on induction of neurite outgrowth. To synthesize the stereoisomers of GM3, initially, we succeeded in synthesizing and purifying all stereoisomers of sphingosine. Further by using these isomers and sialyllactose we succeeded the synthesis of four stereoisomers of GM3 and performed the cell viability assay. Two different concentrations (10  $\mu$ M and 40 $\mu$ M) of GM3 isomers are used to perform the neurite outgrowth assay with or without NGF in PC12 cells. Neurite outgrowth activity was measured by staining with Coomassie brilliant blue (CBB) and fluorescence imaging with an inverted microscope and fluorescence microscopy. All four isomers are enhancing the neurite outgrowth in the presence of NGF and without NGF. Surprisingly, *L-erythro* GM3 inducing more neurite outgrowth as compared to other three isomers.




Fig.1. (a) Chiral GM3 induce neurite outgrowth (b) Neurite outgrowth CBB staining assay of chiral GM3s 10  $\mu$ M

Further study was continued to understand the effect of sphingolipid chirality on sphingomyelin synthase activity. Successfully synthesized 128 unique stereoisomers of ceramides by efficient solid-phase synthesis. Further experiment was performed to discover the stereochemical effects of these chiral ceramide on sphingomyelin synthase activity. The cell-based assay of sphingomyelin synthase inhibition in the presence of chiral ceramide suggested that these ceramides have good inhibitory activities ( $IC_{50} = 0.2 \sim 1 \mu M$ ) for sphingomyelin synthases 1 and 2. Furthermore, according to heatmap analysis of  $IC_{50}$  values, we confirmed that most of the unnatural (L)-*threo* stereoisomer ceramide derivatives showed strong inhibitory activities towards SMS1 and SMS2 respectively compared to the inhibitory activities of other stereoisomers.




Fig.2. (c) Unnatural stereoisomer (L)-*threo* ceramides possesses a highest potential for inhibition of lipid metabolizing enzyme (SMS) comparing to other stereoisomers including natural one. (d) Heatmap analysis of  $IC_{50}$  values of each of the four unique chiral ceramides for SMS1 and SMS2

In conclusion, synthesis the four stereoisomers of chiral GM3 and chiral ceramide library was successfully completed. Stereochemical study of four GM3 isomers described that they are inducing the neurite outgrowth and *L-erythro* induces more as compared to the other three isomers. On the other hand, stereochemical effects of chiral ceramide on sphingomyelin synthase activity suggested that, these ceramides show excellent inhibitory activity on SMS1 and 2, especially the unnatural *L-threo* ceramide derivatives showed strong inhibitory activities towards SMS1 and SMS2 respectively compared to the inhibitory activities of other stereoisomers. In the future, these chiral GM3 and chiral ceramides are better therapeutic target for metabolic disorder and nervous system injuries.