



Title	First molecular identification of Trypanosoma evansi from cattle in Syria
Author(s)	Megasari, Marsela; Kyoko, Hayashida; Alaa, Terkawi; Xuenan, Xuan; Chihiro, Sugimoto; Junya, Yamagishi
Citation	Japanese Journal of Veterinary Research, 68(2), 117-127
Issue Date	2020-05
DOI	10.14943/jjvr.68.2.117
Doc URL	<a href="http://hdl.handle.net/2115/78622">http://hdl.handle.net/2115/78622</a>
Type	bulletin (article)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	JJVR68-2_117-127_MegasariMarsela.pdf



[Instructions for use](#)

# First molecular identification of *Trypanosoma evansi* from cattle in Syria

Megasari Marsela<sup>1)</sup>, Kyoko Hayashida<sup>1)</sup>, Alaa Terkawi<sup>2,3)</sup>,  
Xuenan Xuan<sup>3)</sup>, Chihiro Sugimoto<sup>1,4)</sup> and Junya Yamagishi<sup>1,4,\*)</sup>

<sup>1)</sup> Division of Collaboration and Education, Research Center for Zoonosis Control, Hokkaido University, Kita-20, Nishi-10, Kita-ku, Sapporo, Hokkaido 001-0020, Japan.

<sup>2)</sup> Department of Orthopedic Surgery, Hokkaido University, Faculty of Medicine and Graduate School of Medicine, Kita-15, Nishi-7, Kita-ku, Sapporo, Hokkaido 060-8638, Japan.

<sup>3)</sup> National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine, Inada-cho, Obihiro, Hokkaido 080-8555, Japan.

<sup>4)</sup> Global Station for Zoonosis Control, Global Institution for Collaborative Research and Education (GI-CoRE), Hokkaido University, Kita-20, Nishi-10, Kita-ku, Sapporo, Hokkaido 001-0020, Japan.

Received for publication, November 8, 2019; accepted, March 17, 2020

## Abstract

*Trypanosoma evansi*, the “surra” disease-causing agent, is a blood protozoan parasite that infects a wide range of mammalian species within an unlimited geographical region. It causes anemia, weight loss, and even death of the infected livestock that heavily affect animal husbandry. However, the full epidemiological information of *T. evansi* is lacking, especially in developing countries, and the risk of the disease is largely underestimated. In this study, 207 samples of blood DNA collected from Holstein Friesian crossbred cattle in the central region of Syria in May 2010 were screened for *T. evansi*, aiming to determine the prevalence of the parasite. *T. evansi* was screened by PCR targeting the internal transcribed spacer (ITS) 1 region, and 27 samples were found positive out of 207 (13%), which is relatively high considering that no clinical symptoms were observed. The ITS1 amplicons were later subjected to *RoTat1.2*-PCR for detection of *T. evansi* type A. This is the first report of molecular detection of *T. evansi* in Syria. Our study suggests that advanced investigations in cattle and other domestic animals are necessary in Syria.

Key Words: Epidemiology, ITS1-PCR, Syria, *Trypanosoma evansi*

## Introduction

*Trypanosoma evansi*, the causative agent of “surra” disease classified in the subgenus *Trypanozoon*, is a flagellated protozoan parasite<sup>14)</sup>. It shares some characteristics with other *Trypanozoon* species, including *Trypanosoma brucei* and *Trypanosoma equiperdum*, in terms of morphology and genome sequences<sup>8,30)</sup>. However,

while *T. brucei* undergoes a complex cycle of differentiation and multiplication in tsetse flies, *T. evansi* does not have a vector stage and is only transmitted mechanically. Unlike *T. brucei*, *T. evansi* has lost the maxicircles of kinetoplast mitochondrial DNA, which are required to undergo the procyclic form in tsetse flies. This makes *T. evansi* unable to reproduce in tsetse flies<sup>27)</sup>. *T. evansi* is transmitted mechanically by

\*Corresponding author: Junya YAMAGISHI Division of Collaboration and Education Research Center for Zoonosis Control, Hokkaido University, Kita-20, Nishi-10, Kita-ku, Sapporo, Hokkaido 001-0020, Japan  
Tel: +81-11-706-9516 Fax: +81-11-706-7307 email: junya@czc.hokudai.ac.jp  
doi: 10.14943/jjvr.68.2.117

a wide range of biting flies including tabanids and stomoxes, vampire bats, and ingestion of raw meat<sup>12,13</sup>. Since *T. evansi* is no longer restricted to tsetse flies, it has the largest geographical distribution and host range among salivarian trypanosomes<sup>11,29</sup>. *T. evansi* infects a broad range of domestic animals and wildlife, which are more pathogenic to camelids and equids<sup>14</sup>. It has been described that the susceptibility of *T. evansi* is highly variable depending on the host species and presumably according to the region<sup>12</sup>.

To differentiate *Trypanozoon* from other trypanosomes, polymerase chain reaction (PCR) targeting the internal transcribed spacer (ITS) 1 region of ribosomal RNA (rRNA) has been commonly used<sup>16,33,41</sup>. Identification of each species in *Trypanozoon*, including *T. evansi* and *T. brucei*, is complicated because of their twinning characteristics<sup>9,30</sup>. Although *T. evansi* has no maxicircles, they still possess short minicircle DNA that encodes guide RNAs for RNA editing<sup>6,7</sup>. Based on its minicircle restriction digestion profile, *T. evansi* is divided into type A and type B<sup>6,34</sup>. *T. evansi* type A is the most common species found in Africa, South America, and Asia<sup>4</sup>. In contrast, *T. evansi* type B is found in Eastern Africa. To date, the prevalence of *T. evansi* type B remains largely unknown, especially in Asia<sup>3,6,18,32,41</sup>. Further, it has not been studied whether the presence of the *Rode Trypanozoon antigen type 1.2 (RoTat1.2)* gene of *T. evansi* is related to pathogenesis or other biological phenotypes.

The variant surface glycoprotein (VSG) of *RoTat1.2* is specifically present in *T. evansi*, particularly in type A, but not in *T. brucei* strains, and has been utilized to differentiate *T. evansi* from other *Trypanozoon* members<sup>9,49,50</sup>. Because some of the diagnostic strategies for *T. evansi* rely on the presence of the *RoTat1.2*, such as *RoTat1.2* loop-mediated isothermal amplification (LAMP) and the serological card agglutination test for *T. evansi* (CATT)/*RoTat1.2*, only *T. evansi* type A, not type B, can be detected using these methods<sup>2,15,422</sup>.

For specific diagnosis of *T. evansi* type B,

several tests have been developed, including EVAB-PCR, targeting a type B-specific minicircle DNA sequence; a PCR; and LAMP targeting type B-specific *VSG JN 2118Hu*<sup>32,35,36</sup>. Once the presence of *T. evansi* type B infection is confirmed, CATT or *RoTat1.2*-PCR test should be avoided as the current CATT/*RoTat1.2*-PCR tests cannot detect *RoTat1.2*-negative *T. evansi* type B infection. In such cases, nucleic acid amplification methods targeting other genes, such as ITS1-PCR used in this study, are recommended to detect all *T. evansi* infections. *T. brucei* and *T. evansi* can switch their diverse transferrin receptors, and the two genes encoding the transferrin receptors, expression-site-associated gene (*ESAG*) 6 and *ESAG7*, display high genetic diversity<sup>20-22,31,44,52</sup>. This diversity has been speculated to contribute to sufficient iron uptake from different mammalian species and escape from anti-transferrin receptor antibodies<sup>5,17,46</sup>. However, differences in transferrin receptors of animals are not always responsible for differences in trypanosome growth in the sera<sup>43</sup>. In addition, the antibody against the transferrin receptor failed to prevent iron uptake of trypanosome<sup>45</sup>. Hence, it is not clear whether the genetic diversity of *ESAG6/7* within *T. evansi* is related to the biology or infectivity of the trypanosome in mammalian host and whether it is in any way related to countries/regions or host species<sup>44,52</sup>.

It is widely believed that *Trypanozoon*, including *T. evansi*, originated in Africa and spread across the world<sup>12</sup>. It has been suggested that *T. evansi* evolved from a *T. brucei* infection in camels that had temporarily entered the sub-Saharan tsetse belt and adapted to mechanical transmission by biting flies<sup>19</sup>. However, a recent study explained the independent origins of *T. evansi* from *T. brucei* strains, where the ability of *T. evansi* to be transmitted mechanically occurred repeatedly<sup>23</sup>. The parasite spread from North Africa toward the Middle East, Turkey, India, up to Russia, across all of Southeast Asia, down to Indonesia and the Philippines, and was also introduced by the conquistadores into Latin

America<sup>19,28,40</sup>). However, there is no molecular study of *T. evansi* in Syria). Syria, as one of the Middle Eastern countries situated between Africa and Asia and Europe might be a gate to learn about *T. evansi* evolution, where the parasite moved out of its place of origin and started to spread worldwide. Collectively, the aim of this study was to determine the prevalence and characterize the genetic diversity of *T. evansi* in Syria. This epidemiology study and genetic diversity information of the parasites will help in improving the control measures in this region and understanding host tropism and adaptation of *T. evansi* in different animals and regions worldwide.

## Materials and Methods

**DNA Samples:** A total of 207 blood DNA samples were used, which were collected from clinically healthy Holstein Friesian crossbred cattle in the central region of Syria in May 2010. The sampling sites were Huleh ( $n = 51$ ), Hama ( $n = 73$ ), Qyser ( $n = 28$ ), Ghab ( $n = 32$ ), and Salmia ( $n = 23$ ). The collection of field samples was approved by the Syrian government through the Ministry of Agriculture and supported by veterinarians and staff working at the Society for Protection of Animals Abroad in Syria. Sample collection methods were followed as described previously<sup>47</sup>. Readily prepared DNA samples were previously provided and used by Terkawi *et al.* (2012)<sup>47</sup>. In this study, microscopic parasite examination and serological tests for *T. evansi* diagnosis were not conducted.

**PCR:** PCR amplification of the ITS1 region of the rRNA was conducted to screen trypanosomes, including *Trypanozoon*. For amplification of the ITS1 region, the primer set CF 5'-CCGGAAGTTCACCGATATTG-3' and BR 5'-TGCTGCGTTCTTCAACGAA-3' was used<sup>33</sup>. Each reaction included 5  $\mu$ l Ampdirect plus (Shimadzu, Japan), 0.05  $\mu$ l BIOTAQ HS DNA

Polymerase (5 U/ $\mu$ l) (Bioline, UK), 0.5  $\mu$ l of each 10 mM primer, 2.95  $\mu$ l RNase-free water, and 1  $\mu$ l extracted DNA. The thermocycling profile started with an initial hold for 10 min at 95 °C, followed by 40 cycles at 94 °C for 30 sec, 55 °C for 1 min, 72 °C for 1 min, and a final extension at 72 °C for 7 min. PCR products were electrophoresed on 1.2% agarose S (Nippongene, Japan) in Tris-acetate EDTA buffer and stained using GelRed (Biotium, USA) dye before being visualized under UV light.

The ITS1-positive samples were subjected to PCR specific for *T. evansi* (type A), which amplifies 151 bp of the *RoTat1.2 VSG* fragment by using the primer set TeRoTat920F 5'-CTGAAG AGGTTGGAAATGGAGAAG-3' and TeRoTat1070R 5'-GTTTCGGTGGTTCT GTTGTGTTA-3'<sup>25</sup>). The *RoTat1.2* amplicons were electrophoresed on a 2.2% agarose gel. PCR amplification of the 740 bp fragment of *ESAG6* was performed using *ESAG7* F455 5'-CATTCCAGCAGGAGTTGGAGG-3' and *ESAG6* R1045 5'-TTGTTCACTCACTCTCTTTGACAG-3' primers as described by Isobe *et al.*<sup>21</sup>). *ESAG6* amplicons were electrophoresed on a 1.2% agarose gel. The reaction was performed for 35 cycles at 58 °C annealing temperature in *RoTat1.2*-PCR and 40 cycles at 60 °C in *ESAG6*-PCR; the other thermocycling conditions were as mentioned above. The master mix conditions were the same as those for ITS1-PCR. Increased sensitivity of *ESAG6* amplification was achieved using the newly designed nested PCR. The second round of PCR was performed using the inner primer set, forward 5'-GCAGGAGTTGGAGGAAATGA-3' and reverse 5'-TGAGCTCAGCCTCTTTCTGC-3. The reaction mixture was the same as that used for the initial *ESAG6*-PCR. The thermal cycling conditions used were the same as that for the initial PCR except for the modification to 35 cycles.

**Sequence and phylogenetic analysis of ESAG6:** The *ESAG6*-PCR products were purified by ExoSAP-IT (GE healthcare/USB, USA) following the manufacturer's instructions. Purified PCR

products were sequenced using the Big-Dye Terminator v3.1 (Applied Biosystems, USA) on an automated capillary sequencer (Applied Biosystems 3130 Genetic Analyzer; Applied Biosystems Japan Ltd., Tokyo, Japan). DNA sequences were edited using ApE<sup>10</sup>. The DNA sequence data were aligned against 47 sequences of *ESAG6* deposited in GenBank using the ClustalW program in the MEGA7 software<sup>26</sup>. A phylogenetic tree was constructed using neighbor-joining (NJ) algorithms. For the trees provided by NJ methods, bootstrap branch supports were calculated from 1000 pseudo-replicates following the rule of branch consistency.

*Statistical analysis:* Chi-square test was used to evaluate significant differences ( $P < 0.05$ ) in the infection rate in animals of different ages, locations, anemia, and co-infection with *Babesia* spp. Statistical analysis was conducted to compare the infection rate of *T. evansi* in animals and co-infection with *Babesia* spp. using the data from Terkawi et al. (2012)<sup>47</sup>.

*PCR sensitivity test:* Sensitivity of the PCR systems for ITS1, *RoTat1.2*, and *ESAG6* was validated using primer sets of CF and BR, TeRoTat920F and TeRoTat1070R, and *ESAG7* F455 and *ESAG6* R1045, respectively. The detection limit was examined in four replicates using 10-fold serial dilutions of DNA extracted from *T. evansi* IL3354 isolate cultured *in vitro*.

## Results

*Molecular detection of T. evansi by PCR:* DNA samples were subjected to PCR amplification of the ITS1 region and *RoTat1.2* for molecular identification and *ESAG6* for genetic diversity characterization of *T. evansi*. Sanger sequencing was carried out to validate the results of ITS1-PCR, and it was confirmed that all 27 ITS1-positive samples contained *T. evansi*. The present study showed a 13.0% prevalence of *T. evansi*

infection in cattle by ITS1-PCR screening (Table 1). *RoTat1.2*-PCR amplified in 17 samples that were ITS1-positive, but it failed to amplify in the other 10 samples.


*Prevalence of T. evansi according to location, packed cell volume (PCV) value, age, and co-infection with Babesia spp.:* The highest prevalence of *T. evansi* was observed in Ghab (34.4%,  $n = 11$ ), followed by Qyser (25%,  $n = 7$ ) and was significantly higher than that in the other regions ( $P < 0.0001$ , Table 2). In contrast, *T. evansi* infection in cattle had no correlation ( $P > 0.05$ ) with anemia (Table 3), age (Table 4), and *Babesia* spp. co-infection (Table 5).

*Sequencing and phylogenetic analysis based on ESAG6:* Sixteen samples were positive in *ESAG6*-PCR, and these were successfully sequenced. Phylogenetic analysis based on the *ESAG6* revealed that Syrian sequences clustered together, with the majority in clade 10 (9 sequences) and clade 9 (4 sequences), while the remaining sequences were distributed in clade 1 (1 sequence) and clade 5 (2 sequences). This indicated that some of the major genotypes were shared within the three sampling locations in Syria, and genetic diversity was observed in our samples (Figure 1).

*Analytical sensitivity test of ITS1, RoTat1.2, and ESAG6-PCR:* The minimum amount of DNA detectable by PCR was  $1 \times 10^{-4}$  ng for ITS1-PCR (two positives out of four experiments),  $1 \times 10^{-2}$  ng for *RoTat1.2*-PCR (one positive out of four experiments), and  $1 \times 10^{-4}$  ng for *ESAG6* (one positive out of four experiments; Supp Fig 1 and Supp Table 1). ITS1-PCR and *ESAG6*-PCR showed higher sensitivity than *RoTat1.2*-PCR.

## Discussion

In the present study, we described for the first time the prevalence of *T. evansi* infection among cattle in Syria. ITS1-PCR was used to determine



**Figure 1**

Phylogenetic tree of the *ESAG6* was constructed based on the nucleotide sequences of *Trypanosoma brucei* and *Trypanosoma evansi*. The entry data were represented according to their countries of origin—Peru, Philippine, Egypt, India, Thailand, Bolivia, and Syria; the hosts—cattle, camel, horse, donkey, buffalo, deer, and pig; and the GenBank accession numbers. The dotted entry names are the nucleotide sequences obtained from this study. The bootstrap confidence values at the nodes illustrate the percentage of times the group occurred out of 1000 trees, and the bar depicts the genetic distance.

**Table 1. PCR screening result for *T. evansi***

ITS1	<i>RoTat1.2</i>	<i>ESAG6</i>	Number of samples	
		+	9	10
	+	-	1	
+		+	7	17
	-	-	10	
-	-	-	180	

(+) = PCR positive, (-) = PCR negative.

**Table 2. Comparisons of the prevalence of *T. evansi* infections on the basis of sampling region.**

City	Total number of samples	ITS1 positives with 480bp (putative <i>T. evansi</i> )	Positive rate
Huleh	51	1	2.0%
Hama	73	6	8.2%
Qyser	28	7	25.0%
Ghab	32	11	34.4%
Salmia	23	2	8.7%
Total	207	27	13.0%

**Table 3. Comparisons of the prevalence of *T. evansi* infections on the basis of PCV value.**

PCV	Status	Total number of samples	ITS1 positives with 480bp (putative <i>T. evansi</i> )	Positive rate
<24%	Anemic	9	2	22.2%
24-46%	Normal	198	25	12.6%
Total		207	27	13.0%

**Table 4. Comparisons of the prevalence of *T. evansi* infections on the basis of age.**

Age (years)	Category	Total number of samples	ITS1 positives with 480bp (putative <i>T. evansi</i> )	Positive rate
1-2	Young	55	6	10.9%
3-5	Old	99	14	14.1%
Above 5	Older	53	7	13.2%
Total		207	27	13.0%

**Table 5. Comparisons of the prevalence of *T. evansi* infections on the basis of coinfection with *Babesia* spp.**

Category	Total number of samples	ITS1 positives with 480bp (putative <i>T. evansi</i> )	Positive rate
Absence of <i>Babesia</i> spp.	160	19	11.9%
Presence of <i>B. bovis</i>	16	1	6.2%
Presence of <i>B. bigemina</i>	31	7	22.6%
Total	207	27	13.0%

the presence of *T. evansi* in the studied samples because Syria is not a habitat of the tsetse fly, with the *T. brucei* vector found only in sub-Saharan Africa<sup>51</sup>). The positive rate determined by ITS1-PCR screening was 13.0% and relatively high, considering that no clinical symptoms were observed. In our analysis, 37% of the ITS1-PCR positive samples could not be detected by *RoTat1.2*-PCR possibly owing to low parasitemia, presence of *T. evansi* type B, or the performance gap between ITS1-PCR and *RoTat1.2*-PCR in our samples. To confirm the *RoTat1.2* negativity in ITS1-PCR positive samples, further analyses are required, including microscopic parasite examination, serology assay to detect *RoTat1.2* antigen, and highly sensitive nucleic acid detection system.

Geographic factors played a role in the infection rate distribution in this study. Ghab, followed by Qyser, were the most prevalent areas of *T. evansi* infection in this study. Both these regions were green areas dedicated to the agriculture sector in the country, where the population of livestock was dense and animal movements were frequent. Poor farm hygiene and weather conditions in the areas provided the best environment for horseflies (tabanus flies) to reproduce extensively. Farmers in Ghab and Qyser practiced a free-range management system where animals were released during the day to graze freely and returned home at dusk to sleep, which increased the exposure to horseflies. These factors elevated the possibility of disease transmission in Ghab and Qyser. Given the significant differences in occurrence of *T. evansi* among the studied regions, it is necessary to investigate the geographic distribution of horseflies as parasite vectors in Syria. The higher altitude of Hama has possibly become a geographical barrier for wildlife trespasses, which might explain the difference in prevalence<sup>47</sup>). In addition, the indoor management farm in Salmia contributed to lesser animal contact with fly vectors, which was one of the transmission factors.

In Asia, Holstein Friesian cattle are

susceptible to *T. evansi* infection, and infected cattle frequently exhibit a significant decrease in PCV profiles and body weight as well as a negative effect on milk yield and fertility, including abortion<sup>24,37-39</sup>). In this study, the cattle samples were a mixed breed of Syrian local and Holstein Friesian. Syrian local cattle are known to be resistant to trypanosome infection. This may explain the lack of correlation between infection in cattle and anemia. Considering the observed high prevalence and mild symptoms, these cattle might be one of the potential reservoirs of *T. evansi* in Syria.

We observed that all age groups of cattle were equally exposed and affected by surra. The studied regions were endemic to other blood parasites of *Babesia* spp.; therefore, we also analyzed the correlation between *Babesia* spp. and *T. evansi* using the molecular analysis data of Terkawi *et al.*<sup>47</sup>) in the same samples collected. We did not find any significant correlation in co-infection of *Babesia* spp. and *T. evansi*. This is possibly because of limited interaction in terms of vector, lifecycle (one is an intracellular parasite and the other is extracellular), and immunogenicity (presumably acquired immunity does not cross-react with each other).

Sequence analysis of the *ESAG6* of *T. evansi* in Syria showed genetic diversity. The major genotypes were clades 10 and 5, and the genotypes were found in three sampling locations. An association of genotypes with countries, regions, or host species was not observed in our study. The diversity of transferrin receptors has been shown to possibly relate with the need for antigenic variation to escape from host immune responses<sup>48</sup>). This is the first report of molecular detection of *T. evansi* in Syria. Therefore, additional epidemiological study of the parasites is necessary. Further investigations in cattle and other livestock animals is also required to improve the control measures against *T. evansi* in Syria.



## Acknowledgements

This study was funded by the AMED program of the Japan Initiative for Global Research Network on Infectious Diseases (J-GRID; JP18fm0108008) and the International Collaborative Research Program for Tackling Neglected Tropical Disease (NTD) Challenges in African countries (JP18jm0510001). The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. The authors would like to thank the staff at the Division of Collaboration and Education, Research Center for Zoonosis Control, Hokkaido University, Japan, for their support during the experiments.

## Compliance with ethical standards

The authors declare that they have no conflicts of interest.

## Availability of data and materials

The obtained sequences are available at GenBank under the following accession numbers: LC440031.1, LC519566.1, LC519567.1, LC519568.1, LC519569.1, LC519571.1, LC519572.1, LC519573.1, LC519574.1, LC519575.1, LC519576.1, LC519577.1, LC519578.1, LC519579.1, and LC519580.1.

## Supplemental data

Supplemental data associated with this article can be found, in the online version, at <http://dx.doi.org/10.14943/jjvr.68.2.117>

## References

- 1) Aregawi WG, Agga GE, Abdi RD, Büscher P. Systematic review and meta-analysis on the global distribution, host range, and prevalence of *Trypanosoma evansi*. *Parasit Vectors* 12, 1–25, 2019.
- 2) Bajyana Songa E, Hamers R. A card agglutination test (CAT) for veterinary use based on an early VAT RoTat 1/2 of *Trypanosoma evansi*. *Ann Soc Belg Med Trop* 68, 233–240, 1988.
- 3) Birhanu H, Fikru R, Said M, Kidane W, Gebrehiwot T, Hagos A, Alemu T, Dawit T, Berkvens D, Goddeeris BM, Büscher P. Epidemiology of *Trypanosoma evansi* and *Trypanosoma vivax* in domestic animals from selected districts of Tigray and Afar regions, Northern Ethiopia. *Parasit Vectors* 8, 1–11, 2015.
- 4) Birhanu H, Gebrehiwot T, Goddeeris BM, Büscher P, Van Reet N. New *Trypanosoma evansi* Type B Isolates from Ethiopian Dromedary Camels. *PLoS Negl Trop Dis* 10, e0004556, 2016.
- 5) Bitter W, Gerrits H, Kieft R, Borst P. The role of transferrin-receptor variation in the host range of *Trypanosoma brucei*. *Nature* 391, 499–502, 1998.
- 6) Borst P, Fase-Fowler F, Gibson WC. Kinetoplast DNA of *Trypanosoma evansi*. *Mol Biochem Parasitol* 23, 31–38, 1987.
- 7) Borst P, Hoeijmakers J H. Kinetoplast DNA. *Plasmid* 2, 20–40, 1979.
- 8) Carnes J, Anupama A, Balmer O, Jackson A, Lewis M, Brown R, Cestari I, Desquesnes M, Gendrin C, Hertz-Fowler C, Imamura H, Ivens A, Kořený L, Lai DH, MacLeod A, McDermott SM, Merritt C, Monnerat S, Moon W, Myler P, Phan I, Ramasamy G, Sivam D, Lun ZR, Lukeš J, Stuart K, Schnauffer A. Genome and Phylogenetic Analyses of *Trypanosoma evansi* Reveal Extensive Similarity to *T. brucei* and Multiple Independent Origins for Dyskinetoplasty. *PLoS Negl Trop Dis* 9, e3404, 2015.
- 9) Claes F, Radwanska M, Urakawa T, Majiwa PAO, Goddeeris B, Büscher P. Variable Surface Glycoprotein RoTat1.2 PCR as a

- specific diagnostic tool for the detection of *Trypanosoma evansi* infections. *Kinetoplastid Biol Dis* 3, 1–6, 2004.
- 10) Davis MW. ApE: a plasmid editor. <http://www.biology.utah.edu/jorgensen/wayned/ape/>. 2012.
  - 11) Desquesnes M, Dargantes A, Lai DH, Lun Z, Holzmuller P, Jittaplapong S. *Trypanosoma evansi* and Surra: A Review and Perspectives on Transmission, Epidemiology and Control, Impact, and Zoonotic Aspects. *Biomed Res Int* 1-20, 2013a.
  - 12) Desquesnes M, Holzmuller P, Lai DH, Dargantes A, Lun ZR, Jittapalapong S. *Trypanosoma evansi* and Surra: A Review and Perspectives on Origin, History, Distribution, Taxonomy, Morphology, Hosts, and Pathogenic Effects. *Biomed Res Int* 1–22, 2013b.
  - 13) Desquesnes M. Livestock Trypanosomoses and their Vectors in Latin America. CIRAD-EMVT publication, OIE, Paris, France, 2004.
  - 14) Donelson JE, Artama WT. Diagnosis of *Trypanosoma evansi* by the polymerase chain reaction (PCR). *J Protozool Res* 8, 204-213, 1998.
  - 15) Elsaid HM, Nantulya VM, Hilali M. Diagnosis of *Trypanosoma evansi* Infection Among Sudanese Camels Imported to Egypt Using Card Agglutination Test (CATT) and Antigen Detection Latex Agglutination Test (Suratex). *J Protozool Res* 8, 194-200, 1998.
  - 16) Gaithuma AK, Yamagishi J, Martinelli A, Hayashida K, Kawai N, Marsela M, Sugimoto C. A single test approach for accurate and sensitive detection and taxonomic characterization of Trypanosomes by comprehensive analysis of internal transcribed spacer 1 amplicons. *PLoS Negl Trop Dis* 13, e0006842, 2019.
  - 17) Gerrits H, Mußmann R, Bitter W, Kieft R, Borst P. The physiological significance of transferrin receptor variations in *Trypanosoma brucei*. *Mol Biochem Parasitol* 119, 237–247, 2002.
  - 18) Hagos A, Yilkal A, Esayass T, Alemu T, Fikru R, Ab Feseha G, Goddeeris BM, Claes F. Parasitological and serological survey on trypanosomiasis (surra) in camels in dry and wet areas of Bale Zone, Oromyia Region, Ethiopia. *Rev Med Vet (Toulouse)* 160, 569–573, 2009.
  - 19) Hoare CA. The trypanosomes of mammals. A zoological monograph. Blackwell Scientific Publications, Oxford. 1972.
  - 20) Holland WG, Claes F, My LN, Thanh NG, Tam PT, Verloo D, Büscher P, Goddeeris B, Vercruyssen J. A comparative evaluation of parasitological tests and a PCR for *Trypanosoma evansi* diagnosis in experimentally infected water buffaloes. *Vet Parasitol* 97, 23–33, 2001.
  - 21) Isobe T, Holmes EC, Rudenko G. The Transferrin Receptor Genes of *Trypanosoma equiperdum* Are Less Diverse in Their Transferrin Binding Site than Those of the Broad-Host Range *Trypanosoma brucei*. *J Mol Evol* 56, 377–386, 2003.
  - 22) Kabiri M, Steverding D. *Trypanosoma evansi*: Demonstration of a Transferrin Receptor Derived from Expression Site-Associated Genes 6 and 7. *J Parasitol* 87, 1189–1191, 2001.
  - 23) Kamidi CM, Saarman NP, Dion K, Mireji PO, Ouma C, Murilla G, Aksoy S, Schnauffer A, Caccone A. Multiple evolutionary origins of *Trypanosoma evansi* in Kenya. *PLoS Negl Trop Dis* 11, e0005895, 2017.
  - 24) Kashiwazaki Y, Pholpark M, Polsar C, Pholpark S. Haemoparasite infections in newly introduced dairy cattle in Loei Province, Thailand: *Trypanosoma evansi* antigen levels by ELISA referring to abortion. *Vet Parasitol* 80, 99–109, 1998.
  - 25) Konnai S, Mekata H, Mingala CN, Abes NS, Gutierrez CA, Herrera JR, Dargantes AP, Witola WH, Cruz LC, Inoue N, Onuma M, Ohashi K. Development and application of a quantitative real-time PCR for the diagnosis of Surra in water buffaloes. *Infect Genet Evol* 9, 449–452, 2009.

- 26) Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 33, 1870-4, 2016.
- 27) Lai DH, Hashimi H, Lun ZR, Ayala FJ, Lukes J. Adaptations of *Trypanosoma brucei* to gradual loss of kinetoplast DNA: *Trypanosoma equiperdum* and *Trypanosoma evansi* are petite mutants of *T. brucei*. *Proc Natl Acad Sci USA* 105, 1999–2004, 2008.
- 28) Luckins AG. *Trypanosoma evansi* in Asia. *Parasitol Today* 4, 137–142, 1988.
- 29) Lun ZR, Desser SS. Is the Broad Range of Hosts and Geographical Distribution of *Trypanosoma evansi* Attributable to the Loss of Maxicircle Kinetoplast DNA? *Parasitol Today* 11, 131-133, 1995.
- 30) Masiga DK, Gibson WC. Specific probes for *Trypanosoma (Trypanozoon) evansi* based on kinetoplast DNA minicircles. *Mol Biochem Parasitol* 40, 279-284, 1990.
- 31) Mekata H, Konnai S, Witola WH, Inoue N, Onuma M, Ohashi K. Molecular detection of trypanosomes in cattle in South America and genetic diversity of *Trypanosoma evansi* based on expression-site-associated gene 6. *Infect Genet Evol* 9, 1301–1305, 2009.
- 32) Ngaira JM, Olembo NK, Njagi ENM, Ngeranwa JJN. The detection of non-RoTat 1.2 *Trypanosoma evansi*. *Exp Parasitol* 110, 30–38, 2005.
- 33) Njiru ZK, Constantine CC, Guya S, Crowther J, Kiragu JM, Thompson RCA, Dávila AMR. The use of ITS1 rDNA PCR in detecting pathogenic African trypanosomes. *Parasitol Res* 95, 186–192, 2005.
- 34) Njiru ZK, Constantine CC, Masiga DK, Reid SA, Thompson RC, Gibson W. Characterization of *Trypanosoma evansi* type B. *Infect Genet Evol* 6, 292–300, 2006.
- 35) Njiru ZK, Constantine CC, Masiga DK, Reid SA, Thompson RC, Gibson WC. Characterization of *Trypanosoma evansi* type B. *Infect Genet Evol* 6: 292–300, 2006.
- 36) Njiru ZK, Ouma JO, Enyaru JC, Dargantes AP. Loop-mediated Isothermal Amplification (LAMP) test for detection of *Trypanosoma evansi* strain B. *Exp Parasitol* 125: 196–201, 2010.
- 37) Payne RC, Sukanto IP, Bazeley K, Jones TW. The effect of *Trypanosoma evansi* infection on the oestrous cycle of Friesian Holstein heifers. *Vet Parasitol* 51, 1–11, 1993.
- 38) Payne RC, Sukanto IP, Partoutomo S, Polytedi F. Experimental infection of Friesian Holstein calves with an Indonesian isolate of *Trypanosoma evansi*. *Trop Med Parasitol* 43, 115-117, 1992.
- 39) Pholpark S, Pholpark M, Polsar C, Charoenchai A, Paengpassa Y, Kashiwazaki Y. Influence of *Trypanosoma evansi* infection on milk yield of dairy cattle in northeast Thailand. *Prev Vet Med* 42, 39-44, 1999.
- 40) Reid SA. *Trypanosoma evansi* control and containment in Australasia. *Trends Parasitol* 18, 219–224, 2002.
- 41) Salim B, Bakheit MA, Kamau J, Nakamura I, Sugimoto C. Molecular epidemiology of camel trypanosomiasis based on ITS1 rDNA and RoTat 1.2 VSG gene in the Sudan. *Parasit Vectors* 4, 2–6, 2011.
- 42) Salim B, Hayashida K, Mossaad E, Nakao R, Yamagishi J, Sugimoto C. Development and validation of direct dry loop mediated isothermal amplification for diagnosis of *Trypanosoma evansi*. *Vet Parasitol* 260, 53–57, 2018.
- 43) Salmon D, Paturiaux-Hanocq F, Poelvoorde P, Vanhamme L, Pays E. *Trypanosoma brucei*: growth differences in different mammalian sera are not due to the species-specificity of transferrin. *Exp Parasitol* 109, 188-194, 2005.
- 44) Sarkhel SP, Gupta SK, Kaushik J, Singh J, Saini VK, Kumar S, Kumar R. Intra and inter species genetic variability of transferrin receptor gene regions in *Trypanosoma evansi* isolates of different livestock and geographical regions of India. *Acta Parasitol* 62, 133–140, 2017.
- 45) Steverding D. On the significance of host

- antibody response to the *Trypanosoma brucei* transferrin receptor during chronic infection. *Microbes Infect* 8, 2777-2782, 2006.
- 46) Steverding D. The significance of transferrin receptor variation in *Trypanosoma brucei*. *Trends Parasitol* 19, 125–127, 2003.
  - 47) Terkawi MA, Alhasan H, Huyen NX, Sabagh A, Awier K, Cao S, Goo YK, Aboge G, Yokoyama N, Nishikawa Y, Kalb-Allouz AK, Tabbaa D, Igarashi I, Xuan X. Molecular and serological prevalence of *Babesia bovis* and *Babesia bigemina* in cattle from central region of Syria. *Vet Parasitol* 187, 307–311, 2012.
  - 48) Trevor CE, Gonzalez-Munoz AL, Macleod OJ, Woodcock PG, Rust S, Vaughan TJ, Garman EF, Minter R, Carrington M, Higgins MK. Structure of the trypanosome transferrin receptor reveals mechanisms of ligand recognition and immune evasion. *Nature microbiology* 4, 2074-2081, 2019.
  - 49) Urakawa T, Verloo D, Moens L, Büscher P, Majiwa PA. *Trypanosoma evansi*: Cloning and Expression in *Spodoptera fugiperda* Insect Cells of the Diagnostic Antigen RoTat1.2. *Exp Parasitol* 99, 181–189, 2001.
  - 50) Verloo D, Magnus E, Büscher P. General expression of RoTat 1.2 variable antigen type in *Trypanosoma evansi* isolates from different origin. *Vet Parasitol* 97, 183-189, 2001.
  - 51) Vreysen MJ, Seck MT, Sall B, Bouyer J. Tsetse flies: Their biology and control using area-wide integrated pest management approaches. *J Invertebr Pathol* 112, S15–S25, 2013.
  - 52) Witola WH, Sarataphan N, Inoue N, Ohashi K, Onuma M. Genetic variability in ESAG6 genes among *Trypanosoma evansi* isolates and in comparisons to other *Trypanozoon* members. *Acta Trop* 93, 63-73, 2005.