Title	Effects of ozone (O ₃) and ethylenediurea (EDU) on the ecological stoichiometry of a willow grown in a free-air exposure system
Author(s)	Agathokleous, Evgenios; Kitao, Mitsutoshi; Qingnan, Chu; Saitanis, Costas J.; Paoletti, Elena; Manning, William J.; Watanabe, Toshihiro; Koike, Takayoshi
Citation	Environmental pollution, 238, 663-676 https://doi.org/10.1016/j.envpol.2018.03.061
Issue Date	2018-07
Doc URL	http://hdl.handle.net/2115/78852
Rights	© 2018 Elsevier Ltd. All rights reserved., © 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	Manuscript.pdf

[Instructions for use](#)

Effects of ozone (O₃) and ethylenediurea (EDU) on the ecological stoichiometry of a willow grown in a free-air exposure system

Evgenios Agathokleous^{1,2*}, Mitsutoshi Kitao¹, Chu Qingnan^{2,3}, Costas J. Saitanis⁴, Elena Paoletti⁵, William J. Manning⁶, Toshihiro Watanabe², Takayoshi Koike²

¹Hokkaido Research Center, Forestry and Forest Products Research Institute (FFPRI), Forest Research and Management Organization, 7 Hitsujigaoka, Sapporo, Hokkaido, 062-8516, Japan.

²Research Faculty of Agriculture, School of Agriculture, Hokkaido University, Kita ku Kita 9 Nishi 9, Sapporo, Hokkaido, 060-8589, Japan.

³Institute of Agricultural Resources and Environment, Jiangsu Academy of Agricultural Science, Nanjing, 210014, China

⁴Lab of Ecology and Environmental Science, Agricultural University of Athens, Iera Odos 75, Athens, 11855, Greece

⁵Institute of Sustainable Plant Protection, National Council of Research, Via Madonna del Piano 10, Sesto Fiorentino, Florence, 50019, Italy.

⁶Department of Plant, Soil and Insect Sciences, University of Massachusetts, Amherst, MA, USA.

*Corresponding author: Tel: +81-11-851-4131, Fax: +81-11-851-4167, globalscience@frontier.hokudai.ac.jp; evgenios@ffpri.affrc.go.jp

ABSTRACT

Ground-level ozone (O₃) concentrations have been elevating in the last century. While there has been a notable progress in understanding O₃ effects on vegetation, O₃ effects on ecological stoichiometry remain unclear, especially early in the oxidative stress. Ethylenediurea (EDU) is a chemical compound widely applied in research projects as protectant of plants against O₃ injury, however its mode of action remains unclear. To investigate O₃ and EDU effects early in the stress, we sprayed willow (*Salix sachalinensis*) plants with 0, 200 or 400 mg EDU L⁻¹, and exposed them to either low ambient O₃ (AOZ) or elevated O₃ (EOZ) levels during the daytime, for about one month, in a free air O₃ controlled exposure (FACE); EDU treatment was repeated every nine days. We collected samples for analyses from basal, top, and shed leaves, before leaves develop visible O₃ symptoms. We found that O₃ altered the ecological stoichiometry, including impacts in nutrient resorption efficiency, early in the stress. The relation between P content and Fe content seemed to have a critical role in maintaining homeostasis in an effort to prevent O₃-induced damage. Photosynthetic pigments and P content appeared to play an important role in EDU mode of action. This study provides novel insights on the stress biology which are of ecological and toxicological importance.

Keywords: iron homeostasis, nutrient cycling, nutrient resorption, ozone, re-translocation

Key message: Elevated O₃ alters ecological stoichiometry, and Fe homeostasis plays an important role early in O₃ stress. Photosynthetic pigments and phosphorus are important components in EDU mode of action.

1. INTRODUCTION

The levels of tropospheric ozone (O_3) are nowadays elevated at wide areas of the Northern Hemisphere (Feng et al., 2015; Hendriks et al., 2016; Kalabokas et al., 2017; Sicard et al., 2017; Tian et al., 2016; Trieu et al., 2017), and are potentially phytotoxic at least to O_3 -sensitive vegetation (Feng et al., 2008; Hayes et al., 2007; Saitanis et al., 2015). When elevated O_3 doses are taken up by plants, plants undergo a responsive process which involves a series of physiological and biological adjustments ranging from single cell to whole plant level (Ashmore, 2005; Bussotti et al., 2007; Fiscus et al., 2005; Jolivet et al., 2016; Matyssek and Innes, 1999; Munne-Bosch et al., 2013). Exposure of vegetation to elevated O_3 doses poses a risk for food supplies, ecosystem health and biosphere sustainability (Andersen, 2003; Broberg et al., 2015; Felzer et al., 2004; Fuhrer, 2009; Lindroth, 2010; Lu et al., 2015; Peñuelas and Staudt, 2010; Tian et al., 2016; Wang et al., 2016; Wilkinson et al., 2012).

The protection of vegetation against O_3 adverse effects is thus an important matter. Plenty of potential agrochemicals have been tested as to their efficacy to protect plants against O_3 phytotoxicity (Agathokleous et al., 2016c; Manning et al., 1973a, 1973b; Runeckles and Resh, 1975; Saitanis et al., 2015; Taylor, 1974). The most effective phytoprotectant currently known is the synthetic chemical ethylenediurea (EDU, $C_4H_{10}N_4O_2$) (Agathokleous, 2017). EDU has been a matter of study in dozens of investigations over the past four decades (Carnahan et al., 1978; Oksanen et al., 2013; Paoletti et al., 2009; Salvatori et al., 2017; Singh et al., 2015), and it is now recognized that EDU applied repeatedly at low concentrations of up to 400 mg L^{-1} sufficiently protects against O_3 phytotoxicity across plant species (Agathokleous et al., 2015; Manning et al., 2011a; Oksanen et al., 2013; Paoletti et al., 2009; Singh et al., 2015). EDU can protect numerous biological or physiological endpoints; nonetheless, literature review studies suggest that the EDU

72 mode of action in protecting against O₃ phytotoxicity remains unclear (Agathokleous et al.,
73 2015; Manning et al., 2011a; Oksanen et al., 2013; Paoletti et al., 2009; Singh et al., 2015). It
74 was recently proposed that EDU may protect plants through hormesis, by activating plant
75 defense at low doses via homeostatic disruptions, and the necessity for studying plant
76 metabolism was highlighted (Agathokleous, 2017). Homeostatic re-adjustments after disruption
77 may imply alterations in leaf metabolism and in quantitative and qualitative traits of leaf
78 nutrients. However, there is hitherto no evidence on leaf stoichiometry under EDU treatments.
79 Ecological stoichiometry is critically important for predicting stress consequences in biochemical
80 cycles and other ecological processes. Leaf stoichiometry and nutrient resorption efficiency
81 (NRE) have been mainly studied in plants experiencing prolonged O₃ stress (Cao et al., 2016;
82 Nakaji et al., 2004; Oksanen et al., 2005; Shang et al., 2018; Shi et al., 2016, 2017; Zhuang et al.,
83 2017) but not at an early stage of the stress so as to be utilized as an early diagnostic tool. Leaf
84 age is an important factor determining O₃ effects on stoichiometry. Younger or late leaves may
85 be more resistant to O₃ than older, mature leaves (Li et al., 2017; Moreno Pina et al., 2017) as a
86 result of adaptive responses which may take place at plant level. For example, relative to mature
87 leaves, younger leaves may: emit greater concentrations of volatile organic compounds (Bison et
88 al., 2018); take up less amount of O₃ (Craker and Starbuck, 1973); have greater carbon (C)
89 assimilation rate and water use efficiency (Moreno Pina et al., 2017); and overall have greater
90 efficiency in apoplastic O₃ detoxification as indicated by avoidance of accumulation of reactive
91 oxygen species in the chloroplasts (Oksanen et al., 2005). Besides, shed leaves may be more
92 affected than mature leaves as they experience O₃ pressure for longer time and the nutrient re-
93 translocation may also be affected (Shang et al., 2018; Shi et al., 2017; Uddling et al., 2006).
94 Despite NRE was studied at later stages of oxidative stress (Shang et al., 2018; Shi et al., 2017;

Uddling et al., 2006), leaf nutrient cycling across the crown has not been explicitly investigated under O₃ pressure; none of the previous studies included all leaf classes (i.e. mature, young, shed). Effects of EDU on ecological stoichiometry, including NRE and leaf nutrient cycling across the crown, remain elusive too.

The principal aim of this study is to investigate the ecological stoichiometry in willow (*Salix sachalinensis* F. Schmidt) plants sprayed with 0, 200 or 400 mg EDU L⁻¹ and exposed to ambient O₃ or an O₃-enriched atmosphere. The main research question is whether EDU and O₃ affect the leaf stoichiometry and leaf nutrient cycling across the crown (basal, top and shed leaves) early in oxidative stress, before leaves develop visible O₃ symptoms. Based on previous studies where O₃ affected leaf stoichiometry and NRE at later stages of oxidative stress (Shang et al., 2018; Shi et al., 2017; Uddling et al., 2006), it is hypothesized that leaf stoichiometry and nutrient cycling across the crown may be altered at early stages of oxidative stress when plants are under homeostatic disruption. An additional aim is to assess photosynthetic pigments and leaf mass per area (LMA) in mature leaves, as a reference of the stress status, and to assess essential and non-essential elements in an analysis of patterns in the dataset of the elements. Photosynthetic pigments and LMA are known as simple and effective indices for assessing health status in mature leaves (Agathokleous et al., 2017; Onoda et al., 2017; Ronen and Galun, 1984). Mature leaves experience O₃ exposure for longer time and are less affected by ontogenic adjustments than young leaves, and are widely assessed as to gas exchange and other traits under stress.

2. MATERIALS AND METHODS

2.1. Experimental site

The experiment was conducted at Sapporo Experimental Forest of Hokkaido University, Sapporo, Japan (43°04' N, 141°20' E, 15 m a.s.l.), in the year 2015. For the period June-September, mean monthly average of air temperature was 17.83 (± 1.83 s.d.) °C; daily maximum temperature was 22.52 (± 1.75 s.d.) °C; daily minimum temperature was 14.07 (± 2.02 s.d.) °C; wind speed was 3.52 (± 0.17 s.d.) m s⁻¹; relative humidity was 68.83 (± 1.58 s.d.) %; mean monthly total sunshine duration was 186.25 (± 9.98 s.d.) h; and mean monthly precipitation was 120.50 (± 23.19 s.d.) mm, respectively (43°03.6'N 141°19.7'E; Japan Meteorological Agency, 2016; <http://www.jma.go.jp/jma/indexe.html>).

2.2. Plant material & Design of the Experiment

The plant materials were current-year cuttings of *S. sachalinensis* (= *S. udensis* Trautv. et C.A. Mey.), originated from the river basin of Ebetsu city, grown in 15-L pots filled with Akadama (well-weathered volcanic ash) and Kanuma (well-weathered pumice) soils at a ratio of 1:1 (DCM Homac CO., LTD., Sapporo, JP). This soil substrate is free from organic matter and poor in phosphorus (P) and nitrogen (N) (for further information please see Electronic Supplementary Material, Table 2S). This willow was selected as biological model due to prior knowledge with regards to its exposure to EDU and O₃ (Agathokleous et al., 2018, 2016a, 2016b). Eighty uniform plants were transplanted in the 15-L pots on June 9th, when they were well rooted. On August 14th, 72 pots were randomly allocated to six plots (12 pots per plot), of which three served as background O₃ (ambient) and three as elevated O₃ treatment. The pots in each plot were further randomly assigned three EDU treatments (four pots per EDU treatment). These plants were used for a different study and thus the full, detailed procedure has been previously

described (Agathokleous et al., 2016a). The eight remaining plants were harvested on August 6th, for an approximate image of the initial leaf traits. The length and width of each leaf was non-destructively measured, and the leaf size was calculated using a simple linear model as described earlier (Agathokleous et al., 2016b). Leaves ($n = 549$) were air-dried at 65 °C until a constant mass, and the dry matter of each leaf was measured using a digital balance. The specific leaf area (SLA, or its inverse, LMA) was calculated as the one-side area of a fresh leaf to the dry matter of the leaf. The average leaf size was $7.20 \pm 0.26 \text{ cm}^2$, average leaf dry matter $0.09 \pm 0.01 \text{ g}$, and SLA $134.27 \pm 11.78 \text{ cm}^2 \text{ g}^{-1}$.

2.3. EDU treatments

The EDU treatments were 0 (EDU0), 200 (EDU200), and 400 (EDU400) mg L⁻¹ (Agathokleous et al., 2016a). EDU (100% a.i., source W.J. Manning, University of Massachusetts, USA) was prepared using an electric hotplate 30 min before each application. The required EDU amount was dissolved in partitions of 500 mL, so as the target concentration to be achieved in the final desired volume by adding cool pure water. The initial 500 mL stock for each EDU concentration was prepared using gently-warmed water (Manning et al., 2011) with continuous stirring. The initial stocks were prepared with pure water, and surfactant was not added. EDU was applied using an electric sprayer using Venturi effect, with two nozzles spraying simultaneously. Both abaxial and adaxial leaf surfaces were sprayed until run-off.

The first EDU application was carried out 50 days after transplanting (July 29), when the plants had 63 ± 2 leaves. EDU application was applied on a 9-day interval (Agathokleous, 2017). The last (5th) EDU treatment was applied on September 3rd. All the applications were conducted during the hours 10:00-11:00 am, JST.

2.4. Ozone treatments

O₃ treatments were carried out in the free air O₃ controlled exposure (FACE) system of Hokkaido University, Japan, located in the Sapporo Experimental Forest, Sapporo, Japan. This FACE system has been previously described (Agathokleous et al., 2017). The O₃ treatments were ambient O₃ (AOZ) and elevated O₃ (EOZ). O₃-enrichment lasted from August 15th to September 11th, 2015, during daylight hours, when the photosynthetic photon flux density (PPFD) was >70 $\mu\text{mol m}^{-2} \text{s}^{-1}$. This PPFD threshold is considered the light compensation point of photosynthesis of targeted plants (Koike, 1988). Background O₃ mixing ratios at the experimental site were continuously monitored (1-min interval) by an ultraviolet absorption O₃ analyzer (TUV-1100; Tokyo Industries Inc., Tokyo, Japan), from August 15th to September 11th, 2015. Data were averaged per hour, and the 10-hour (07:00 am – 17:00 pm, JST) means were calculated. The mean background O₃ mixing ratio for the period August 15th to September 11th was $14.52 \pm 0.34 \text{ nmol mol}^{-1}$, whereas the index of Accumulated exposure to O₃ Over the Threshold of 40 nmol mol^{-1} (AOT40) (Mills et al., 2007) was practically 0 $\mu\text{mol mol}^{-1} \text{h}$. The mean O₃ mixing ratio in the three EOZ plots for the monitored period was $68.84 \pm 1.12 \text{ nmol mol}^{-1}$, which can be translated to AOT40 of $7.79 \pm 0.30 \mu\text{mol mol}^{-1} \text{h}$ until September 10th and $6.34 \pm 0.25 \mu\text{mol mol}^{-1} \text{h}$ until September 5th.

2.5. Samplings, sample analysis and data collection

2.5.1 Leaf mass per area (LMA)

On September 4th, 2 mature (basal), 2 young (top) and 2-4 fallen leaves (after gently shaking of shoots) were collected from each plant, and air-dried at 65 °C until a constant weight, under dark conditions. Prior to drying, randomly-selected mature leaves ($n=89$, 4-25 leaves per experimental

condition) were scanned (Canon LIDE 40, Tokyo), and the leaf area was measured using the LAI32 v.0.377 software (© Kazukiyo Yamamoto). LMA was calculated for each leaf.

2.5.2 Non-photosynthetic elements

The leaves collected on September 4th were pooled per maturity level per plant so as to give one more robust sample per plant to be used for elemental analyses. Leaf samples were manually grounded into powder. An amount of grounded powder was used for digestion, as previously described (Chu et al., 2015): Leaf powder was digested with 2 mL of 61% nitric acid (HNO₃) (EL grade; Kanto Chemical, Tokyo) at 110 °C, in a DigiPREP apparatus (SCP Science, Quebec, Canada) for 3 h. Then, 0.5 mL of hydrogen peroxide (H₂O₂) (semiconductor grade; Santoku Chemical, Tokyo, Japan) was added two times, at a 30-min interval, and heated at 110 °C until the solution became clear. Tubes were left to cool to room temperature and afterwards filled to 15 mL with 2% HNO₃. The analyses were conducted with an Inductively Coupled Plasma Mass Spectrometer (ICP-MS) (Elan, DRC-e; PerkinElmer, Waltham, MA, USA), with ultratrace-level detection limits. The ICP multielement standard solution IV (Merck, Tokyo, Japan) was used as standard for ionic determination. The essential elements magnesium (Mg), manganese (Mn), iron (Fe), calcium (Ca), nickel (Ni), P, potassium (K), and the non-essential elements chromium (Cr) and aluminum (Al) were quantified, and the ratios K/Mg, K/Ca, Fe/Mn, Ca/Mg, P/Fe, and P/Mn were calculated. The quantified elements were selected in accordance with a previous study under similar environmental conditions and with a similar soil (Shi et al., 2017). NRE was calculated for each element as $NRE = ((C_{\text{basal}} + C_{\text{top}}) - (C_{\text{shed}})) / (C_{\text{basal}} + C_{\text{top}})$, where C is the content in the leaf. This was considered a more robust calculation of NRE than the classical one (Aerts, 1996) because it reflects the nutrient content throughout the crown, i.e. from base to top. Nitrogen and C content was analyzed in leaf powder using a third generation Vario EL III

204 Element Analyzer (Elementar Analysensysteme GmbH, Hanau-Germany). In total, 204 samples
205 were analyzed for all the chemical analyses, except K for which only 91 samples were analyzed.

206 **2.5.3 Photosynthetic pigments**

207 On September 9th, two mature, basal leaves were randomly selected from each plant, and one
208 leaf disk (area=2.26 cm²) was sampled from each leaf and stored in deep freeze. Five days later,
209 photosynthetic pigments were measured. The two leaf disks consisted a sample per plant, whose
210 pigments were extracted in dimethyl sulfoxide (DMSO) and stored in dark condition at 65 °C
211 until the appearance of a “ghost-like” thallus (Barnes et al., 1992). Measurements were taken for
212 the optical densities (OD) 415, 435, 470, 648, 665 nm (GeneSpec III; Hitachi Genetic Systems;
213 MiraiBio, Alameda, CA). The OD 470 was used to estimate total carotenoid content (TCar)
214 following the classical Lichtenthaler formula (Lichtenthaler, 1987). This formula is given for
215 acetone 100% extracts and thus the actual TCar values may differ. The ODs 648 and 665 were
216 used to estimate the chlorophyll *a* content (Chl_a), chlorophyll *b* content (Chl_b) and total
217 chlorophylls, *a+b*, content (TChl) (Barnes et al., 1992). The ratios of OD₄₃₅/OD₄₁₅, *Chl_a/Chl_b*
218 and *TChl/TCar* were calculated as potential indices of stress. OD₄₃₅/OD₄₁₅, has been proposed as
219 an effective index of chlorophyll degradation to phaeopigments (Ronen and Galun, 1984). Since
220 OD₄₃₅/OD₄₁₅ is underexplored in higher plants, phaeophytinization was also assessed by the
221 classical method of acidification (Ronen and Galun, 1984). Overall, after extracts were measured
222 (before acidification), 60 µl of 1 M HCl were added per 10 ml of extract, stored in 4 °C for 12
223 min, and measured again (after acidification). The OD value at 665 nm before the acidification
224 (OD₆₆₅) was divided by the OD value at 665 nm after the acidification (OD_{665a}) (OD₆₆₅/OD_{665a}).

225 Leaf sampling was conducted at the onset of O₃-induced visible foliar injury, when only very
226 few leaves showed a small number of tiny spots of discoloration visible to naked eye.

2.5. Data handling & Statistics

The threshold for statistical significance was predefined at a level of significant $\alpha=0.05$. The data of all the response variables were averaged per EDU treatment per experimental unit, i.e. O₃ plot ($n = 3$).

Data of pigments, ICP elements, and elements ratios were transformed by a Box-Cox power transformation (Box and Cox, 1964), as described in an earlier report (Agathokleous et al., 2016c). Data transformation was unnecessary for N, C and N/C data.

Data of pigments and LMA were analyzed by simple contrasts, based on prior theoretical knowledge, to answer only the most biologically meaningful questions (Agathokleous et al., 2016a; Ruxton and Beauchamp, 2008). A contrast represents a linear combination of experimental conditions whose sum of coefficients equals to zero. The contrasts were conducted among the following six combinational groups: AOZ×EDU0 (AOZ0), AOZ×EDU200 (AOZ200), AOZ×EDU400 (AOZ400), EOZ×EDU0 (EOZ0), EOZ×EDU200 (EOZ200), EOZ×EDU400 (EOZ400). The five degrees of freedom were partitioned to the following contrasts:

C₁: (AOZ0) *vs.* (AOZ200+AOZ400), which tests the null hypothesis that EDU does not affect plants under the ambient O₃ condition

C₂: AOZ200 *vs.* AOZ400, which tests the hypothesis EDU400 plants do not perform better than EDU200 plants under the ambient O₃ condition

C₃: (EOZ0) *vs.* (EOZ200+EOZ400), which tests the hypothesis EDU-treated plants do not perform better than plants not treated with EDU under the elevated O₃ condition

248 C₄: EOZ200 vs. EOZ400, which tests the hypothesis that EDU400 plants do not perform better
 249 than EDU200 plants under the elevated O₃ condition

250 C₅: AOZ0 vs. EOZ0, which tests the hypothesis that elevated O₃ does not affect plants.

251 To test the question of whether EDU and O₃ affect the cycling of leaf chemical elements across
 252 the crown, data were analyzed by a General Linear Model (GLM) where leaf position (three
 253 levels: basal (mature), top (young), shed) was set as within effects dependent variable and EDU
 254 (three levels: EDU0, EDU200, EDU400) and O₃ (two levels: AOZ, EOZ) categorical predictors.
 255 Analysis was based on Type 6 Sums of Squares with σ -restricted coding of effects that provides
 256 unique effect estimates even for lower-order effects (straightforward computing method). For
 257 aposteriori comparisons, the post-hoc Bonferroni test was utilized (at $\alpha=0.05$).

258 To emphasize variation and observe patterns in the dataset of the response variables assessed in
 259 mature leaves, a Principal Component Analysis (PCA) was utilized. Data of the main variables
 260 (Mn, Fe, Mg, Ca, Ni, P, Al, K, Cr, N, C, Chla, Chlb, CarT, OD₆₆₅/OD_{665a}) were submitted to
 261 PCA. The derivative variables (i.e. the sums or ratios of the main variables: K/Mg, K/Ca, Fe/Mn,
 262 Ca/Mg, P/Fe, P/Mn, Chla/Chlb, ChlT, ChlT/CarT, N/C, and OD₄₃₅/OD₄₁₅) were considered
 263 redundant and thus, along with EDU and Ozone, were passively only included as supplementary
 264 variables (not considered in the calculations). The Centroid of AOZ and EOZ ozone groups are
 265 also additionally illustrated. Prior to PCA analysis, data were standardized so all the variables to
 266 have equal units with s.d. = 1 (Saitanis et al., 2015).

267 MS EXCEL 2010 (Microsoft ©), XLSTAT 2017 (© Addinsoft), and STATISTICA v.10
 268 (StatSoft Inc. ©) software were used for data processing and statistics.

3. RESULTS

3.1. Photosynthetic pigments and LMA in mature leaves

In AOZ, EDU200 and EDU400 together increased TChl compared to EDU0 (C1). This increase was mainly upon EDU400 which led to greater TChl than EDU200 (C2) (Table 1, Fig. 1). EDU400 also led to higher Chla, Chlb and TCar than EDU200 in AOZ (C1 and C2), indicating superiority of EDU400 over EDU0 and no difference between EDU0 and EDU200. EDU200 and EDU400 had no effect on Chla/Chlb, OD435/OD415, OD665/OD665a, and TChl/TCar, compared to EDU0, in AOZ (Table 1).

In EOZ, EDU200 and EDU400 together increased Chla, Chlb and TChl compared to EDU0 (C3). EDU200 and EDU400, when pooled, had no effect on TCar, Chla/Chlb, OD435/OD415, OD665/OD665a and TChl/TCar in EOZ (C3). However, EDU400 plants had greater Chla, Chla/Chlb, and TChl/TCar than EDU200 and EDU0 plants in EOZ (Table 1).

EOZ decreased Chla, Chlb, TChl and Chla/Chlb but did not significantly affect TCar, OD435/OD415, OD665/OD665a and TChl/TCar (C5).

The LMA means were 8.0 ± 0.23 , 9.9 ± 0.54 , 8.6 ± 0.63 , 8.8 ± 0.39 , 9.3 ± 0.1 and $8.1 \pm 0.12 \mu\text{g mm}^{-2}$ for AOZ0, AOZ200, AOZ400, EOZ0, EOZ200 and EOZ400, respectively. Only EDU in AOZ significantly affected LMA, and this was only upon an increase induced by EDU200 (Table 1).

3.2. Chemical elements across leaf types

N and C

Nitrogen differed significantly (Bonferroni) among the leaves in the order basal>shed>top (Table 2, Fig. 2). C/N also differed significantly among the leaves, but in the opposite order (basal<shed<top). There were no other significant effects in N, C and C/N (Table 2, Fig 2).

Two-factorial GLM revealed that NRE of C and N was not affected by Ozone ($F < 0.01$, $P = 0.966$),
EDU ($F = 0.17$, $P = 0.845$) or Ozone $\times$ EDU ($F = 0.27$, $P = 0.765$) (data not shown).

Mn, Fe, Mg, Ca, Ni, P, Al, K, and Cr

EOZ led to 119.3% greater P compared to AOZ (Fig 3A, Table 2).

EDU was a significant factor for only Cr which was increased by EDU200 (115%) compared to
EDU0 and EDU400; means of EDU0 and EDU400 were same (Fig 4D).

Leaf position (i.e. maturity) was significant for all the elements. At an $\alpha = 0.05$, in Mn (Fig 3B),
Mg (Fig 3D) and Ca (Fig 3E), leaf position ranked Top < Basal = Shed. In Fe (Fig 3C) and Al
(Fig 4B), the rank was Top < Basal < Shed, and, in Ni (Fig 4A) and K (Fig 4C), the rank was
Shed < Basal = Top. In P (Fig 3A), ranking was in the order Shed < Top < Basal, whereas, in Cr
(Fig 4D), Top was lower than Shed but Mature was non-different from Top and Shed. In many
of the elements, the means of Shed had much higher variance than Basal and Top.

Ozone $\times$ EDU was significant only for P (Table 2), with the only difference being a 140.7%
greater P content in EOZ*EDU0 than in AOZ*EDU0 (Fig 3A).

Ozone $\times$ Leaf position was significant only for Fe, Ni, K and Cr (Table 2). In AOZ, Fe (Fig 3C),
Ni (Fig 4A) and Cr (Fig 4D) were non-different among leaf types, and K (Fig 4C) was lower in
Shed than Basal leaves. In EOZ, Fe (Fig 3C) varied among leaf types with the order
Shed>Basal>Top. Ni was greater in Basal than in Shed and Top (Fig 4A), K was greater in Basal
than Shed (Fig 4C), and Cr was lower in Basal than Shed in EOZ (Fig 4D). When comparing
each leaf type between AOZ and EOZ, Fe (Fig 3C) was lower in Top leaves of EOZ than Top
leaves of AOZ and Cr (Fig 4D) much lower in Shed leaves of AOZ than in Shed leaves in EOZ;
Cr was also much greater in Basal and Top leaves of AOZ than the pairwise leaves in EOZ.

EDU $\times$ Leaf position was significant only for Ni and Cr (Table 2). The only significant differences in Ni were lower means of Top than Basal leaves of EDU200 and EDU400 in both AOZ and EOZ; however the variance was high, especially for EDU0 (Fig 4A).

Ozone $\times$ EDU $\times$ Leaf position interaction was significant for all the elements except Fe, Al and Cr (Table 2). Regarding Mn (Fig 3B), EDU treatment of 0, 200 and 400 mg L⁻¹ in basal leaves was not significantly different from the pairwise EDU treatments in top leaves in AOZ; there was a high variance in Shed leaves. However, Mn in EDU400 was 206.7 and 218.0% greater in Shed and Basal than in Top leaves in EOZ. Mn differences within EDU400 were similar as within EDU200 in AOZ and as within EDU0 and EDU200 in EOZ, however the variance was higher (Fig 3B). In Ca, EDU0 was not different among leaf positions in both AOZ and EOZ (Fig 3E). EDU200 was 256.7 and 299.1% higher in Shed than in Top in AOZ and EOZ but not different from Basal in both AOZ and EOZ. EDU400 was not different among leaf positions in AOZ, however it was lower in Top than in Shed (2.6 times) and Basal (2.8 times) leaves and non-different between Shed and Basal leaves in EOZ (Fig 3E). Ni displayed very large variability in Shed in AOZ (Fig 4A). The only significant differences in Ni content worth mentioning are a lower mean in Top than Basal of EDU400 in AOZ and lower means in Shed than in Basal of EDU0 and EDU400 in EOZ (Fig 4A). Regarding P, the only significant difference of interest in AOZ was a lower content in Shed than in Basal and Top leaves treated with EDU400; these comparisons were non-different in EDU0 and EDU200 which had higher variance (Fig 3A). In EOZ, P means of EDU0, EDU200 and EDU400 were lower in Shed than in Basal and Top, and non-different between Basal and Top.

NRE was greater in AOZ than in EOZ for Fe ($F=5.92$, $P<0.05$) and Cr ($F=9.63$, $P<0.01$). However, NRE was greater in EOZ than in AOZ for Ni ($F=4.12$, $P<0.05$). There were no other

Ozone, EDU or Ozone×EDU significant differences in NRE (two-factorial GLM, $P>0.05$, data not shown).

Ratios of chemical elements across leaf types

EOZ increased P/Fe by 115.0% (Fig 4E, Table 2).

EDU was non-significant factor for all the ratios (Table 2).

Leaf position was significant for all the ratios. At an $\alpha=0.05$, leaf position ranked Top > Basal > Shed in K/Ca (Fig 5B), K/Mg (Fig 5A), P/Fe (Fig 4E), and P/Mn (Fig 5E), and Shed > Basal > Top in Ca/Mg (Fig 5D). In Fe/Mn, mean of Shed was greater than means of Basal and Top, and mean of Basal was non-different from mean of Top (Fig 5C).

Ozone × EDU was significant only for P/Fe (Table 2), with the only difference being a 103.7% greater ratio in EOZ*EDU0 than in AOZ*EDU0 (Fig 4E).

EDU × Leaf position was significant only for Ca/Mg and P/Fe (Table 2). Ca/Mg was lower in Top than Basal and Shed leaves for all the three EDU treatments, however, in EDU400, it was also lower in Basal than in Shed leaves (Fig 5D). P/Fe was greater in Top and Basal than Shed leaves for all the three EDU treatments, and greater in Top than Basal leaves only for EDU200 (Fig 4E).

Ozone × EDU × Leaf position interaction was significant for K/Ca and P/Fe (Table 2).

Variation & patterns in chemical composition of mature leaves

Pairwise Pearson correlation analysis revealed several significant ($P<0.05$) correlations (Electronic Supplementary Material, Table 2S). Some of the significant positive correlations are those between the pairs Mn_Ca, P_K, Mg_P, Mn_Mg, Ca_C, Ca_Mn and Ca_Mg and between the pairs of pigments Chla_Chlb, Chla_TCar.

The dataset submitted to PCA was highly structured (Table 3, Fig. 6). The variability of the entire data set can be explained, with a high degree of satisfaction (68%), by six factors (F1-F6) with eigenvalues >1 , each of which explain 21, 15, 10, 8.5, 6.6 and 7.2% respectively (Electronic Supplementary Material, Fig 1S). However, only the first three (F1-F3) explain more than 10% of the total variability. Fe, C, Ni, Pha, Al and LMA were totally separated, having high loadings on the factors F3, F4, F5, F6, F7 and F8 respectively (Table 3A). Only one response variable had a significant contribution to the variability explained by the extracted factors F4-F8; these autonomous response variables were C, Ni, OD665/OD665a, Al and LMA (Table 3B). Taken into account this observation as well as the fact that their eigenvalues were <1 , the factors F4-F8 were considered non-important. The only significant contribution to F3 was upon Fe. The elements Mn, Mg, Ca, K, and P had high positive loadings whereas Cr had a high negative loading on the first PC. The pigments Chla, Chlb and TCar had high positive loadings with the second factor. Biplot analysis shows high correlation of Ca, Mg, P, K, Mn and Ni with the first factor and of N, Chla, Chlb, TCar with the second factor (Fig. 6). This analysis visually separates the samples treated with EOZ and those under AOZ, and confirms the high inter-correlation between pigments. EDU seems to be highly correlated with N, Chla, Chlb, Tchl and TCar and anticorrelated with OD665/OD665a. The chlorophyll pigments are also anticorrelated with ozone.

4. DISCUSSION

In this study, a willow was used as a biological model with prior knowledge about its responses to EDU and ozone. EDU and O₃ effects were studied early in the stress, before O₃ symptoms widely appeared on the leaves. We demonstrated the single and interactive effects of O₃ and EDU on leaf stoichiometry and leaf nutrient cycling across the crown by analyzing for the first

time the chemical composition of leaves of all maturity stages (mature, young, shed) treated with EDU and under O₃ stress.

4.1. Status of EDU-induced stress

The fact that EDU did not significantly affect the ratios Chla/Chlb, OD435/OD415, OD665/OD665a and TChl/TCar suggests that EDU did not induce severe stress in plants in AOZ (Agathokleous et al., 2017). EDU-induced stimulation of the main and accessory pigments content may imply increase in energy absorption and photosynthesis efficiency and is in agreement with previous studies showing EDU-induced stimulation in photosynthetic pigments or other endpoints in O₃-clean air such as in closed chambers (Agathokleous, 2017; Agathokleous et al., 2016c, 2014; Eckardt and Pell, 1996; Whitaker et al., 1990). The present study, indeed, suggests a hormetic effect of EDU in LMA which was increased by EDU200 but not significantly affected by EDU400. Hormesis is a phenomenon where low levels of stress induce stimulatory effects whereas high levels of stress induce adverse effects (Calabrese, 2013, 2011; Calabrese and Baldwin, 2001).

4.2. Status of O₃-induced stress

EOZ decreased TChl but decreased more Chla than Chlb, thus resulting into a lower Chla/Chlb, in line with other studies dealing with O₃ stress (Agathokleous et al., 2017b; Caregnato et al., 2013; Feng et al., 2008; Knudson et al., 1977; Neufeld et al., 2006; Saitanis et al., 2001). In a stress-induced need for photoprotection and chlorophyll breakdown, photooxidation occurs with an imbalance in the relation between carotenoids and chlorophylls which is found also under O₃ stress (Agathokleous et al., 2016c, 2014; Pellegrini, 2014; Peñuelas et al., 1995). In the present study, EOZ did not affect TChl/TCar and phaeophytinization (neither Mg content) as O₃ stress

was in the early stages. Direct stimulation by EDU200 and EDU400 might have helped plants maintain greater content of photosynthetic pigments in EOZ.

4.3. EDU and O₃ effects on stoichiometry and cycling of N and C

In earlier research with willow, it was found that 800 and 1600 mg EDU L⁻¹ given as soil drench increased the N content in mature leaves when soil was infertile (Agathokleous et al., 2016b) but did not increase it when soil was fertile (Agathokleous et al., 2018). Based on findings from such high EDU doses, it was suggested that EDU at low concentrations applied for phytoprotection against O₃ damage (up to 400 mg L⁻¹) is not expected to increase foliar N content of willow plants, even in infertile soil (Agathokleous et al., 2016b). The present study verifies this assumption by showing that despite EDU was associated with N, it had no direct effect on C and N contents or C/N ratio. In the present study, where EDU was applied as foliar spray, plant leaf area was not a factor influencing the EDU amount available to leaves (Agathokleous, 2017). Hence, the findings verify the assumption that the EDU mode of action in protecting plants against O₃ is not upon N contribution (Agathokleous, 2017; Manning et al., 2011b, see also charcoal-filter treatment in Ali and Abdel-Fattah (2006)). Ozone did not affect C and N content and cycling or C/N balance across leaves at the early stages of plant stress in this study with relatively low O₃ exposure. In a study where European aspen (*Populus tremula* L.) was exposed to O₃-free air, 50 nmol O₃ mol⁻¹, or 100 nmol O₃ mol⁻¹ for two growing seasons and leaf N content was analyzed in mid- or late summer of the second growing season, the only significant effect was an increase in N content of O₃ symptomatic leaves by 100 nmol O₃ mol⁻¹; leaves in 50 nmol O₃ mol⁻¹ had no visible injury like in the present study (Matyssek et al., 1993). No changes in N content were also found when European silver birch (*Betula pendula* Roth) clones were exposed to twice-ambient O₃ concentration for three growing seasons (Oksanen et al., 2005) and when Japanese red pine (*Pinus densiflora*) was exposed to 60 nmol O₃ mol⁻¹ for two

growing seasons (Nakaji et al., 2004). Some other studies showed O₃-induced alterations in C, N or C/N in mature leaves of different species (Cao et al., 2016; Shang et al., 2018), however these observations represent later stages of O₃-induced stress. In the present study, NRE of C and N was not affected by EOZ. NRE of N remained also unaffected in three deciduous tree species exposed to elevated O₃ for two growing seasons (Shi et al., 2017). It was also affected only in one from two poplar clones exposed to elevated O₃ for about three months (Shang et al., 2018); however it was impaired in birch, *Betula pendula* Roth, exposed to elevated O₃ for two growing seasons (Uddling et al., 2006). The present study verified that N is not involved in plant response mechanisms early in O₃-induced stress and that the mode of action of EDU in protecting plants against O₃ injury is not upon its contribution with N. Overall, EDU and O₃ did not affect C and N stoichiometry and cycling throughout the crown early in the stress, suggesting that C and N are not primarily involved at the early stages of the detoxification process.

4.4. EDU and O₃ effects on stoichiometry and cycling of other elements

EOZ significantly increased P content and EDU increased Cr content. The fact that P was increased by EOZ when plants were treated with EDU0 but were not affected by EOZ when plants were treated with EDU200 and EDU400 indicates that EDU200 and EDU400 contributed in increasing P content in AOZ in a similar way with EOZ. In previous studies with prolonged O₃ stress, elevated O₃ had no effect on P content in evergreen or deciduous trees (Cao et al., 2016; Karlsson et al., 2002; Nakaji et al., 2004; Shi et al., 2016, 2017), decreased it (Shi et al., 2017) or increased it (Matyssek et al., 1993; Shang et al., 2018). More P in leaves relates to chlorosis (Dekock et al., 1960), which occurs under O₃ stress. For example, increased P content was found in yellowish leaves of European aspen exposed to 100 nmol O₃ mol⁻¹ for two growing seasons; no increase was observed in leaves with no visible injury or in bronzing leaves (Matyssek et al., 1993). In the present study, chlorosis was not observed by naked eye; however

EOZ-induced decreases in the photosynthetic pigments were observed. This study suggests a role of P in homeostatic adjustments early in the stress, perhaps to prevent oxidative damage (Ganz, 2013; Hernández and Munné-Bosch, 2015). It remains unknown at present if EDU can increase P content by improving uptake or utilization efficiency.

Along with P, Fe and Cr seem to have an important role in homeostasis and metabolism early in stress as NRE of Fe and Cr was lower in EOZ than in AOZ, suggesting a higher need for reducing Fe and Cr retranslocation under EOZ; nonetheless, it should be mentioned that Cr dynamics in plant have hitherto uncertain physiological roles. The EOZ-induced need for reducing Fe was reflected into the increased P/Fe which shows stoichiometric modifications to adjust to the stress. It seems that Fe homeostasis was involved in the stress response of young, top, leaves which had lower Fe in EOZ than in AOZ, perhaps to adapt to the stress. While Fe was not different among leaf types in AOZ, it presented re-adjustments in EOZ where more Fe was maintained as leaves were aging. Decreased P content was also found in yellowish leaves, but not in non-injured or bronzing leaves, of European aspen exposed to 100 nmol O₃ mol⁻¹ for two growing seasons (Matyssek et al., 1993). EOZ led also to alterations in the stoichiometry of Ni, Cr and K among leaf types, indicating effects of EOZ on nutrient cycling and suggesting implications of EOZ in the aging process. Considering that EOZ also led to lower Fe in top leaves, lower Cr in basal and top leaves, and greater Cr in shed leaves compared to the pairwise leaf types in AOZ, it is suggested that assessing elevated O₃ impacts in mature leaves only is not realistic and may lead to erroneous conclusions regarding O₃ impacts. This assumption is in line with a series of studies providing evidence for different O₃ effects on different leaf ages (Bison et al., 2018; Craker and Starbuck, 1973; Moreno Pina et al., 2017; Shang et al., 2018; Shi et al., 2017; Uddling et al., 2006). The regulation of Cr may be a mechanism for maintaining carbon

474 metabolism (Samantaray et al., 1998). Fe homeostasis holds an important role in plant nutrition
475 as an element necessary for biochemical processes (Pich et al., 2001; Vigani et al., 2013). In the
476 tobacco cultivars Bel-W3 (O_3 -sensitive) and Virginia (O_3 -insensitive), basal leaves of both
477 cultivars grown in ambient air had greater Fe and indifferent Mn and Zn than control plants
478 grown in growth chambers, even if Virginia plants had no O_3 -induced visible symptoms
479 (Alcántara et al., 2006). In further assessments in Bel-W3 and Virginia, Fe accumulation was
480 correlated with ethylene production and associated with O_3 tolerance (Alcántara et al., 2006).
481 This indicates that the mechanisms regulating Fe homeostasis interfere with stress response
482 mechanisms, in agreement with the present findings which suggest a need for reducing stress-
483 induced Fe accumulation in the attached leaves. However, greater NRE of Ni in EOZ than in
484 AOZ suggests a higher need of attached leaves for more Ni, a phenomenon whose physiological
485 explanations need further investigations.

486 EDU affected the stoichiometry and nutrient cycling across the crown as indicated by the
487 significant interaction $EDU \times Leaf\ position$ for Ni, Ca/Mg and P/Fe. The significantly lower
488 Ca/Mg in Basal than in Shed leaves of EDU400 and the greater P/Fe in Top than Basal leaves of
489 EDU200 were not observed in the rest of EDU treatments for each case. These results suggest
490 that the EDU mode of action differed between EDU200 and EDU400. EDU400 might have
491 altered Ca and Mg homeostasis across leaves whereas EDU200 seem to have contributed to
492 maintain less Fe, relatively to P, in Top than Basal leaves. The greater P/Fe in Top than Basal
493 leaves only for EDU200 may be attributed to the hormetic stimulation of health by EDU200.
494 Since only EDU200 affected Cr and P/Fe (and LMA), it can be postulated that the EDU-
495 conferred protection against O_3 was not only upon stimulation in Cr (and LMA) or changes in

P/Fe because EDU400 had an equally or even more effective protection than EDU200 at the end of the experiment (Agathokleous et al., 2016a).

4.5. Potential connections to activation of defence enzymes and/or signalling pathways under O₃ stress

Many nutrients have important roles as co-factors for several physiological functions. Absence of significant EOZ-induced decreases in concentrations of elements such as K may indicate that plants did not undergo an O₃ avoidance mechanism by decreasing stomatal conductance and transpiration rate (Oksanen et al., 2005; Riikonen et al., 2005). Furthermore, no significant effects of EOZ on LMA may suggest absence of dilution effect (Oksanen et al., 2005). The fact that EOZ did not affect N may also suggest that the quantity and activity of RuBisCO enzyme was not involved in the early stress mechanisms (Riikonen et al., 2003). EOZ decreased chlorophyll contents, and, as O₃ primarily reduces the maximum rate of RuBisCO carboxylation ($V_{c_{max}}$) (Cotrozzi et al., 2016), maintaining the capacity of electron transport with adequate Fe would not be of much relevance to prevent photoinhibition.

Different essential elements have different roles in plants. Phosphorus, which was higher in EOZ than in AOZ, has numerous critical roles in plants (Vance et al., 2003). Here, it may be speculated that plants in EOZ maintained a greater content of P in an effort to prevent O₃-induced photo-oxidative stress. As we have discussed, young, top, leaves had lower Fe in EOZ than in AOZ. This might relate to a mechanism for activating jasmonate signaling (Kobayashi et al., 2016) and ethylene production, which are known as regulators of plant response to stress (Hindt and Guerinot, 2012; Iqbal et al., 2013) such as for modulating O₃-induced hypersensitive cell death (Craker, 1971; Rao et al., 2002, 2000). However, Fe deficiency may be more apparent in young leaves than in old leaves because Fe is not easily transferred within a plant (immobile). Therefore, if O₃ causes a reduction in photosynthesis and an impairment of phloem loading, a

reduction in photosynthates transferred into roots may decrease the activity of Fe absorption by roots as well as the activity of siderophore production by rhizosphere microbes (Qiu et al., 1993; Rasheed et al., 2017; Wang et al., 2011). Reduced release of photosynthates (sugar, organic acid, etc.) into rhizosphere would associate with reduced production of siderophore (Amin et al., 2009). Such effects can lead to Fe deficiency especially in young leaves. At the same time, these malfunctions may result in lower NRE of Fe, a relatively immobile element with upward movement (from root to leaves) through xylem (Marschner, 2012; Shen et al., 2011). On the other hand, Ni and K, whose cycling was altered by EOZ, are water soluble elements and mobile in phloem translocated from old to younger leaves (Marschner, 2012; Shi et al., 2017).

These associations between nutrients, enzymes and physiological functions provide a preliminary line of suggestions that should be further experimentally tested.

5. CONCLUSIONS

EDU did not contribute significantly to plants with N at concentrations within the range applied to plants against O₃ damage. Therefore, it is concluded that EDU-conferred protection of plants against O₃ injury was not upon N contained in EDU. While EDU affected Cr, it had more profound effects on P which seems to be affected by EDU similarly as it is affected by EOZ. EDU200 and EDU400 displayed different effects on Ca/Mg and P/Fe across leaf types. It is concluded that EDU may affect stoichiometry and element cycling. However, the EDU mode of action may differ upon EDU concentration, even within the low-dose zone, thus, a need for more dose-response studies is highlighted.

EOZ altered the cycling of Fe, Ni, Cr and K throughout crown leaves, decreased NRE of Fe and Cr, and increased P/Fe and NRE of Ni. Interestingly, Fe homeostasis and regulations in Cr

seemed to play an important role early in O₃ stress, and this should be further investigated in future studies. It is concluded that EOZ may alter leaf stoichiometry and element cycling throughout the crown early in stress. This may affect decomposition process and nutrient input in the soil via alterations in the chemical composition of shed leaves, thus with potential ecological implications in the long-term.

ACKNOWLEDGEMENTS

The authors are grateful to Mr. Tatsushiro Ueda of Dalton Co. (Hokkaido Branch) for managing the O₃ FACE system. Evgenios Agathokleous is an International Research Fellow (ID No: P17102) of the Japan Society for the Promotion of Science (JSPS). This study was supported by JSPS KAKENHI Grants (JP17F17102, to E.A. and M.K, and Type B: 26292075, to T.K). JSPS is a non-profit organization. The authors declare that there are no conflicts of interest.

CITED LITERATURE

- Aerts, R., 1996. Nutrient resorption from senescing leaves of perennials: are there general patterns? *The Journal of Ecology* 84, 597–608. doi:10.2307/2261481
- Agathokleous, E., 2017. Perspectives for elucidating the ethylenediurea (EDU) mode of action for protection against O₃ phytotoxicity. *Ecotoxicology and Environmental Safety* 142, 530–537. doi:10.1016/j.ecoenv.2017.04.057
- Agathokleous, E., Koike, T., Watanabe, M., Hoshika, Y., Saitanis, C.J., 2015. Ethylene-di-urea (EDU), an effective phytoproctectant against O₃ deleterious effects and a valuable research tool. *Journal of Agricultural Meteorology* 71, 185–195. doi:10.2480/agrmet.D-14-00017
- Agathokleous, E., Paoletti, E., Manning, W.J., Kitao, M., Saitanis, C.J., Koike, T., 2018. High

563 doses of ethylenediurea (EDU) as soil drenches did not increase leaf N content or cause
 564 phytotoxicity in willow grown in fertile soil. *Ecotoxicology and Environmental Safety* 147.
 565 doi:10.1016/j.ecoenv.2017.09.017

566 Agathokleous, E., Paoletti, E., Saitanis, C.J., Manning, W., Sugai, T., Koike, T., 2016a. Impacts
 567 of ethylenediurea (EDU) soil drench and foliar spray in *Salix sachalinensis* protection
 568 against O₃-induced injury. *Science of The Total Environment* 573, 1053–1062.
 569 doi:10.1016/j.scitotenv.2016.08.183

570 Agathokleous, E., Paoletti, E., Saitanis, C.J., Manning, W.J., Shi, C., Koike, T., 2016b. High
 571 doses of ethylene diurea (EDU) are not toxic to willow and act as nitrogen fertilizer.
 572 *Science of the Total Environment* 566–567, 841–850. doi:10.1016/j.scitotenv.2016.05.122

573 Agathokleous, E., Saitanis, C.J., Papatheohari, Y., 2014. Evaluation of di-1-*p*-menthene as
 574 antiozonant on Bel-W3 tobacco plants, as compared with ethylenediurea. *Water, Air, & Soil*
 575 *Pollution* 225, 2139. doi:10.1007/s11270-014-2139-y

576 Agathokleous, E., Saitanis, C.J., Stamatelopoulos, D., Mouzaki-Paxinou, A.-C., Paoletti, E.,
 577 Manning, W., 2016c. Olive oil for dressing plant leaves so as to avoid O₃ injury. *Water,*
 578 *Air, and Soil Pollution* 227, 282. doi:10.1007/s11270-016-2986-9

579 Agathokleous, E., Vanderstock, A., Kita, K., Koike, T., 2017. Stem and crown growth of
 580 Japanese larch and its hybrid F₁ grown in two soils and exposed to two free-air O₃ regimes.
 581 *Environmental Science and Pollution Research* 24, 6634–6647. doi:10.1007/s11356-017-
 582 8401-2

583 Alcántara, E., de la Guardia, M.D., Montilla, I., Romera, F.J., 2006. Interaction between ozone

584 exposure and iron nutrition in two tobacco cultivars. *Journal of Plant Nutrition* 29, 2245–
585 2255. doi:10.1080/01904160600975228

586 Ali, A.A., Abdel-Fattah, R.I., 2006. Protection of agricultural crops in Egypt against adverse
587 effects of atmospheric pollutants I. by using of ethylene diurea. *Journal of Agronomy* 5,
588 158–166.

589 Amin, S.A., Green, D.H., Hart, M.C., Küpper, F.C., Sunda, W.G., Carrano, C.J., 2009.
590 Photolysis of iron-siderophore chelates promotes bacterial-algal mutualism. *Proceedings of*
591 *the National Academy of Sciences of the United States of America* 106, 17071–17076.
592 doi:10.1073/pnas.0905512106

593 Andersen, C.P., 2003. Source-sink balance and carbon allocation below ground in plants exposed
594 to ozone. *New Phytologist* 157, 213–228. doi:10.1046/j.1469-8137.2003.00674.x

595 Ashmore, M.R., 2005. Assessing the future global impacts of ozone on vegetation. *Plant, Cell*
596 *and Environment* 28, 949–964. doi:10.1111/j.1365-3040.2005.01341.x

597 Barnes, J.D., Balaguer, L., Manrique, E., Elvira, S., Davison, A.W., 1992. A reappraisal of the
598 use of DMSO for the extraction and determination of chlorophylls *a* and *b* in lichens and
599 higher plants. *Environmental and Experimental Botany* 32, 85–100. doi:10.1016/0098-
600 8472(92)90034-Y

601 Bison, J.V., Cardoso-Gustavson, P., de Moraes, R.M., da Silva Pedrosa, G., Cruz, L.S., Freschi,
602 L., de Souza, S.R., 2018. Volatile organic compounds and nitric oxide as responses of a
603 Brazilian tropical species to ozone: the emission profile of young and mature leaves.
604 *Environmental Science and Pollution Research* 25, 3840–3848. doi:10.1007/s11356-017-

- Box, G.E.P., Cox, D.R., 1964. An Analysis of Transformations. *Journal of the Royal Statistical Society. Series B (Methodological)* 26, 211–252.
- Broberg, M.C., Feng, Z., Xin, Y., Pleijel, H., 2015. Ozone effects on wheat grain quality – A summary. *Environmental Pollution* 197, 203–213. doi:10.1016/j.envpol.2014.12.009
- Bussotti, F., Strasser, R.J., Schaub, M., 2007. Photosynthetic behavior of woody species under high ozone exposure probed with the JIP-test: a review. *Environmental Pollution* 147, 430–7. doi:10.1016/j.envpol.2006.08.036
- Calabrese, E.J., 2013. Hormetic mechanisms. *Critical Reviews in Toxicology* 43, 580–606. doi:10.3109/10408444.2013.808172
- Calabrese, E.J., 2011. Toxicology rewrites its history and rethinks its future: Giving equal focus to both harmful and beneficial effects. *Environmental Toxicology and Chemistry* 30, 2658–2673. doi:10.1002/etc.687
- Calabrese, E.J., Baldwin, L.A., 2001. U-shaped dose-responses in biology, toxicology, and public health. *Annual Review of Public Health* 22, 15–33. doi:10.1146/annurev.publhealth.22.1.15
- Cao, J., Shang, H., Chen, Z., Tian, Y., Yu, H., 2016. Effects of elevated ozone on stoichiometry and nutrient pools of *Phoebe bournei* (Hemsl.) Yang and *Phoebe zhennan* S. Lee et F. N. Wei seedlings in subtropical China. *Forests* 7, 78. doi:10.3390/f7040078
- Caregnato, F.F., Bortolin, R.C., Divan Junior, A.M., Moreira, J.C.F., 2013. Exposure to elevated ozone levels differentially affects the antioxidant capacity and the redox homeostasis of two

626 subtropical *Phaseolus vulgaris* L. varieties. Chemosphere 93, 320–330.
627 doi:10.1016/j.chemosphere.2013.04.084

628 Carnahan, J.E., Jenner, E.L., Wat, E.K.W., 1978. Prevention of ozone injury to plants by a new
629 protectant chemical. Disease Control and Pest Management 68, 1225–1229.

630 Chu, Q., Watanabe, T., Sha, Z., Osaki, M., Shinano, T., 2015. Interactions between Cs, Sr, and
631 other nutrients and trace element accumulation in *Amaranthus* shoot in response to variety
632 effect. Journal of Agricultural and Food Chemistry 63, 2355–2363. doi:10.1021/jf5058777

633 Cotrozzi, L., Remorini, D., Pellegrini, E., Landi, M., Massai, R., Nali, C., Guidi, L., Lorenzini,
634 G., 2016. Variations in physiological and biochemical traits of oak seedlings grown under
635 drought and ozone stress. Physiologia Plantarum 157, 69–84. doi:10.1111/pp1.12402

636 Craker, L.E., 1971. Ethylene production from ozone injured plants. Environmental Pollution 1,
637 299–304. doi:10.1016/0013-9327(71)90022-X

638 Craker, L.E., Starbuck, J.S., 1973. Leaf age and air pollutant susceptibility: Uptake of ozone and
639 sulfur dioxide. Environmental Research 6, 91–94. doi:10.1016/0013-9351(73)90021-2

640 Dekock, P.C., Hall, A., McDonald, M., 1960. A relation between the ratios of phosphorus to iron
641 and potassium to calcium in mustard leaves. Plant and Soil 12, 128–142.
642 doi:10.2307/42931835

643 Eckardt, N.A., Pell, E.J., 1996. Effects of ethylenediurea (EDU) on ozone-induced acceleration
644 of foliar senescence in potato (*Solanum tuberosum* L.). Environmental Pollution 92, 299–
645 306.

646 Felzer, B., Kicklighter, D., Melillo, J., Wang, C., Zhuang, Q., Prinn, R., 2004. Effects of ozone

647 on net primary production and carbon sequestration in the conterminous United States using
 648 a biogeochemistry model. *Tellus B* 56, 230–248. doi:10.1111/j.1600-0889.2004.00097.x
 649 Feng, Z.-Z., Zeng, H.-Q., Wang, X.-K., Zheng, Q.-W., Feng, Z.-W., 2008. Sensitivity of
 650 *Metasequoia glyptostroboides* to ozone stress. *Photosynthetica* 46, 463–465.
 651 doi:10.1007/s11099-008-0079-8
 652 Feng, Z., Hu, E., Wang, X., Jiang, L., Liu, X., 2015. Ground-level O₃ pollution and its impacts
 653 on food crops in China: A review. *Environmental Pollution* 199, 42–48.
 654 doi:10.1016/j.envpol.2015.01.016
 655 Feng, Z., Kobayashi, K., Ainsworth, E.A., 2008. Impact of elevated ozone concentration on
 656 growth, physiology, and yield of wheat (*Triticum aestivum* L.): a meta-analysis. *Global*
 657 *Change Biology* 14, 2696–2708. doi:10.1111/j.1365-2486.2008.01673.x
 658 Fiscus, E.L., Booker, F.L., Burkey, K.O., 2005. Crop responses to ozone: uptake, modes of
 659 action, carbon assimilation and partitioning. *Plant, Cell and Environment* 28, 997–1011.
 660 doi:10.1111/j.1365-3040.2005.01349.x
 661 Fuhrer, J., 2009. Ozone risk for crops and pastures in present and future climates.
 662 *Naturwissenschaften* 96, 173–194. doi:10.1007/s00114-008-0468-7
 663 Ganz, T., 2013. Systemic iron homeostasis. *Physiological Reviews* 93, 1721–1741.
 664 doi:10.1152/physrev.00008.2013
 665 Hayes, F., Jones, M.L.M., Mills, G., Ashmore, M., 2007. Meta-analysis of the relative sensitivity
 666 of semi-natural vegetation species to ozone. *Environmental Pollution* 146, 754–762.
 667 doi:10.1016/j.envpol.2006.06.011

668 Hendriks, C., Forsell, N., Kieseewetter, G., Schaap, M., Schöpp, W., 2016. Ozone concentrations
 669 and damage for realistic future European climate and air quality scenarios. *Atmospheric*
 670 *Environment* 144, 208–219. doi:10.1016/j.atmosenv.2016.08.026

671 Hernández, I., Munné-Bosch, S., 2015. Linking phosphorus availability with photo-oxidative
 672 stress in plants. *Journal of Experimental Botany* 66, 2889–2900. doi:10.1093/jxb/erv056

673 Hindt, M.N., Guerinot, M. Lou, 2012. Getting a sense for signals: Regulation of the plant iron
 674 deficiency response. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 1823,
 675 1521–1530. doi:10.1016/J.BBAMCR.2012.03.010

676 Iqbal, N., Trivellini, A., Masood, A., Ferrante, A., Khan, N.A., 2013. Current understanding on
 677 ethylene signaling in plants: The influence of nutrient availability. *Plant Physiology and*
 678 *Biochemistry* 73, 128–138. doi:10.1016/j.plaphy.2013.09.011

679 Jolivet, Y., Bagard, M., Cabané, M., Vaultier, M.-N., Gandin, A., Afif, D., Dizengremel, P., Le
 680 Thiec, D., 2016. Deciphering the ozone-induced changes in cellular processes: a
 681 prerequisite for ozone risk assessment at the tree and forest levels. *Annals of Forest Science*
 682 73, 923–943. doi:10.1007/s13595-016-0580-3

683 Kalabokas, P., Hjorth, J., Foret, G., Dufour, G., Eremenko, M., Siour, G., Cuesta, J., Beekmann,
 684 M., 2017. An investigation on the origin of regional springtime ozone episodes in the
 685 western Mediterranean. *Atmospheric Chemistry and Physics* 17, 3905–3928.
 686 doi:10.5194/acp-17-3905-2017

687 Karlsson, P.E., Medin, E.L., Selldén, G., Wallin, G., Ottosson, S., Pleijel, H., Skärby, L., 2002.
 688 Impact of ozone and reduced water supply on the biomass accumulation of Norway spruce

saplings. Environmental Pollution 119, 237–244. doi:10.1016/S0269-7491(01)00330-X

Knudson, L.L., Tibbitts, T.W., Edwards, G.E., 1977. Measurement of ozone injury by determination of leaf chlorophyll concentration. Plant Physiology 60, 606–608. doi:10.1104/pp.60.4.606

Kobayashi, T., Itai, R.N., Senoura, T., Oikawa, T., Ishimaru, Y., Ueda, M., Nakanishi, H., Nishizawa, N.K., 2016. Jasmonate signaling is activated in the very early stages of iron deficiency responses in rice roots. Plant Molecular Biology 91, 533–47. doi:10.1007/s11103-016-0486-3

Koike, T., 1988. Leaf structure and photosynthetic performance as related to the forest succession of deciduous broad-leaved trees. Plant Species Biology 3, 77–87. doi:10.1111/j.1442-1984.1988.tb00173.x

Li, L., Wang, X., Niu, J., Cui, J., Zhang, Q., Wan, W., Liu, B., 2017. Effects of elevated atmospheric O₃ concentrations on early and late leaf growth and elemental contents of *Acer truncatum* Bung under mild drought. Acta Ecologica Sinica 37, 31–34. doi:10.1016/J.CHNAES.2016.10.006

Lichtenthaler, H.K., 1987. Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. pp. 350–382. doi:10.1016/0076-6879(87)48036-1

Lindroth, R.L., 2010. Impacts of elevated atmospheric CO₂ and O₃ on forests: phytochemistry, trophic interactions, and ecosystem dynamics. Journal of Chemical Ecology 36, 2–21. doi:10.1007/s10886-009-9731-4

Lu, Y., Jenkins, A., Ferrier, R.C., Bailey, M., Gordon, I.J., Song, S., Huang, J., Jia, S., Zhang, F.,

710 Liu, X., Feng, Z., Zhang, Z., 2015. Addressing China's grand challenge of achieving food
 711 security while ensuring environmental sustainability. *Science Advances* 1, e1400039. doi:
 712 10.1126/sciadv.1400039

713 Manning, W., Feder, W., Vardaro, P., 1973a. Benomyl in soil and response of pinto bean
 714 plants to repeated exposures to a low level of ozone. *Phytopathological Notes* 63, 1539–
 715 1540.

716 Manning, W., Feder, W.A., Vardaro, P.M., 1973b. Reduction of chronic ozone injury on
 717 poinsettia by benomyl. *Canadian Journal of Plant Science* 53, 833–835. doi:10.4141/cjps73-
 718 162

719 Manning, W., Paoletti, E., Sandermann, H., Ernst, D., 2011a. Ethylenediurea (EDU): A research
 720 tool for assessment and verification of the effects of ground level ozone on plants under
 721 natural conditions. *Environmental Pollution* 159, 3283–3293.
 722 doi:10.1016/j.envpol.2011.07.005

723 Manning, W., Paoletti, E., Sandermann, H., Ernst, D., 2011b. Ethylenediurea (EDU): A
 724 research tool for assessment and verification of the effects of ground level ozone on plants
 725 under natural conditions. *Environmental Pollution* 159, 3283–3293.
 726 doi:10.1016/j.envpol.2011.07.005

727 Marschner, P., 2012. *Marschner's Mineral Nutrition of Higher Plants*, 3rd ed. Academic Press,
 728 London, 672pp.

729 Matyssek, R., Innes, J.L., 1999. Ozone — a risk factor for trees and forests in Europe?, in: *Forest*
 730 *Growth Responses to the Pollution Climate of the 21st Century*. Springer Netherlands,

731 Dordrecht, pp. 199–226. doi:10.1007/978-94-017-1578-2_14

732 Matyssek, R., Keller, T., Koike, T., 1993. Branch growth and leaf gas exchange of *Populus*
733 *tremula* exposed to low ozone concentrations throughout two growing seasons.
734 Environmental Pollution 79, 1–7.

735 Mills, G., Buse, A., Gimeno, B., Bermejo, V., Holland, M., Emberson, L.D., Pleijel, H., 2007. A
736 synthesis of AOT40-based response functions and critical levels of ozone for agricultural
737 and horticultural crops. Atmospheric Environment 41, 2630–2643.
738 doi:10.1016/j.atmosenv.2006.11.016

739 Moreno Pina, J., Meirelles, S.T., Maria De Moraes, R., 2017. Meteorological conditions, ozone
740 concentration and leaf age affect gas exchange in *Psidium guajava* 'Paluma'. Hoehnea 44,
741 236–245. doi:10.1590/2236-8906-50/2016

742 Munne-Bosch, S., Queval, G., Foyer, C.H., 2013. The impact of global change factors on redox
743 signaling underpinning stress tolerance. Plant Physiology 161, 5–19.
744 doi:10.1104/pp.112.205690

745 Nakaji, T., Kobayashi, T., Kuroha, M., Omori, K., Matsumoto, Y., Yonekura, T., Watanabe, K.,
746 Utriainen, J., Izuta, T., 2004. Growth and nitrogen availability of red pine seedlings under
747 high nitrogen load and elevated ozone. Water, Air, & Soil Pollution: Focus 4, 277–287.
748 doi:10.1023/B:WAFO.0000028360.61672.8d

749 Neufeld, H.S., Chappelka, A.H., Somers, G.L., Burkey, K.O., Davison, A.W., Finkelstein, P.L.,
750 2006. Visible foliar injury caused by ozone alters the relationship between SPAD meter
751 readings and chlorophyll concentrations in cutleaf coneflower. Photosynthesis Research 87,

281–286. doi:10.1007/s11120-005-9008-x

Oksanen, E., Pandey, V., Pandey, A.K., Keski-Saari, S., Kontunen-Soppela, S., Sharma, C.,
 2013. Impacts of increasing ozone on Indian plants. *Environmental Pollution* 177, 189–200.
 doi:10.1016/j.envpol.2013.02.010

Oksanen, E., Riikonen, J., Kaakinen, S., Holopainen, T., Vapaavuori, E., 2005. Structural
 characteristics and chemical composition of birch (*Betula pendula*) leaves are modified by
 increasing CO₂ and ozone. *Global Change Biology* 11, 732–748. doi:10.1111/j.1365-
 2486.2005.00938.x

Onoda, Y., Wright, I.J., Evans, J.R., Hikosaka, K., Kitajima, K., Niinemets, Ü., Poorter, H.,
 Tosens, T., Westoby, M., 2017. Physiological and structural tradeoffs underlying the leaf
 economics spectrum. *New Phytologist* 214, 1447–1463. doi:10.1111/nph.14496

Paoletti, E., Contran, N., Manning, W., Ferrara, A.M., 2009. Use of the antiozonant
 ethylenediurea (EDU) in Italy: verification of the effects of ambient ozone on crop plants
 and trees and investigation of EDU's mode of action. *Environmental Pollution* 157, 1453–
 60. doi:10.1016/j.envpol.2008.09.021

Pellegrini, E., 2014. PSII photochemistry is the primary target of oxidative stress imposed by
 ozone in *Tilia americana*. *Urban Forestry & Urban Greening* 13, 94–102.
 doi:10.1016/j.ufug.2013.10.006

Peñuelas, J., Baret, F., Filella, I., 1995. Semi-empirical indices to assess carotenoids/chlorophyll
 a ratio from leaf spectral reflectance. *Photosynthetica* 31, 221–230.

Peñuelas, J., Staudt, M., 2010. BVOCs and global change. *Trends in Plant Science* 15, 133–144.

doi:10.1016/j.tplants.2009.12.005

Pich, A., Manteuffel, R., Hillmer, S., Scholz, G., Schmidt, W., 2001. Fe homeostasis in plant cells: does nicotianamine play multiple roles in the regulation of cytoplasmic Fe concentration? *Planta* 213, 967–76. doi: 10.1007/s004250100573

Qiu, Z., Chappelka, A.H., Somers, G.L., Lockaby, B.G., Meldahl, R.S., 1993. Effects of ozone and simulated acidic precipitation on ectomycorrhizal formation on loblolly pine seedlings. *Environmental and Experimental Botany* 33, 423–431. doi:10.1016/0098-8472(93)90045-H

Rao, M. V, Lee, H.-I., Davis, K.R., 2002. Ozone-induced ethylene production is dependent on salicylic acid, and both salicylic acid and ethylene act in concert to regulate ozone-induced cell death. *The Plant Journal* 32, 447–56. doi: 10.1046/j.1365-313X.2002.01434.x

Rao, M. V, Lee, H., Creelman, R.A., Mullet, J.E., Davis, K.R., 2000. Jasmonic acid signaling modulates ozone-induced hypersensitive cell death. *The Plant Cell* 12, 1633–1646.

Rasheed, M.U., Kasurinen, A., Kivimäenpää, M., Ghimire, R., Häikiö, E., Mpmah, P., Holopainen, J.K., Holopainen, T., 2017. The responses of shoot-root-rhizosphere continuum to simultaneous fertilizer addition, warming, ozone and herbivory in young Scots pine seedlings in a high latitude field experiment. *Soil Biology and Biochemistry* 114, 279–294. doi:10.1016/j.soilbio.2017.07.024

Riikonen, J., Holopainen, T., Oksanen, E., Vapaavuori, E., 2005. Leaf photosynthetic characteristics of silver birch during three years of exposure to elevated concentrations of CO₂ and O₃ in the field. *Tree Physiology* 25, 621–632. doi:10.1093/treephys/25.5.621

Riikonen, J., Oksanen, E., Peltonen, P., Holopainen, T., Vapaavuori, E., 2003. Seasonal variation

794 in physiological characteristics of two silver birch clones in the field. Canadian Journal of
795 Forest Research 33, 2164–2176. doi:10.1139/x03-136

796 Ronen, R., Galun, M., 1984. Pigment extraction from lichens with dimethyl sulfoxide (DMSO)
797 and estimation of chlorophyll degradation. Environmental and Experimental Botany 24,
798 239–245. doi:10.1016/0098-8472(84)90004-2

799 Runeckles, V.C., Resh, H.M., 1975. Effects of cytokinins on responses of bean leaves to chronic
800 ozone treatment. Atmospheric environment 9, 749–753.

801 Ruxton, G.D., Beauchamp, G., 2008. Time for some a priori thinking about post hoc testing.
802 Behavioral Ecology 19, 690–693. doi:10.1093/beheco/arn020

803 Saitanis, C., Riga-Karandinos, A., Karandinos, M., 2001. Effects of ozone on chlorophyll and
804 quantum yield of tobacco (*Nicotiana tabacum* L.) varieties. Chemosphere 42, 945–953.
805 doi:10.1016/S0045-6535(00)00158-2

806 Saitanis, C.J., Lekkas, D. V., Agathokleous, E., Flouri, F., 2015. Screening agrochemicals as
807 potential protectants of plants against ozone phytotoxicity. Environmental Pollution 197,
808 247–255. doi:10.1016/j.envpol.2014.11.013

809 Salvatori, E., Fusaro, L., Manes, F., 2017. Effects of the antiozonant ethylenediurea (EDU) on
810 *Fraxinus ornus* L.: The role of drought. Forests 8, 320. doi:10.3390/f8090320

811 Samantaray, S., Rout, G.R., Das, P., 1998. Role of chromium on plant growth and metabolism.
812 Acta Physiol Plant 20, 201–2012.

813 Shang, B., Feng, Z., Li, P., Calatayud, V., 2018. Elevated ozone affects C, N and P ecological
814 stoichiometry and nutrient resorption of two poplar clones. Environmental Pollution 234,

815 136–144. doi:10.1016/J.ENVPOL.2017.11.056

816 Shen, J., Yuan, L., Zhang, J., Li, H., Bai, Z., Chen, X., Zhang, W., Zhang, F., 2011. Phosphorus
817 dynamics: from soil to plant. *Plant Physiology* 156, 997–1005. doi:10.1104/pp.111.175232

818 Shi, C., Kitao, M., Agathokleous, E., Watanabe, M., Tobita, H., Yazaki, K., Kitaoka, S., Koike,
819 T., 2016. Foliar chemical composition of two oak species grown in a free-air enrichment
820 system with elevated O₃ and CO₂. *Journal of Agricultural Meteorology* 72, 50–58.
821 doi:10.2480/agrmet.D-14-00018

822 Shi, C., Watanabe, T., Koike, T., 2017. Leaf stoichiometry of deciduous tree species in different
823 soils exposed to free-air O₃ enrichment over two growing seasons. *Environmental and*
824 *Experimental Botany* 138, 148–163. doi:10.1016/J.ENVEXPBOT.2017.03.012

825 Sicard, P., Anav, A., De Marco, A., Paoletti, E., 2017. Projected global tropospheric ozone
826 impacts on vegetation under different emission and climate scenarios. *Atmospheric*
827 *Chemistry and Physics Discussions* 1–34. doi:10.5194/acp-2017-74

828 Singh, A.A., Singh, S., Agrawal, M., Agrawal, S.B., 2015. Assessment of ethylene diurea-
829 induced protection in plants against ozone phytotoxicity. *Reviews of Environmental*
830 *Contamination and Toxicology* 233, 129–184. doi:10.1007/978-3-319-10479-9_4

831 Taylor, G.S., 1974. Ozone injury to tobacco in the field influenced by soil treatments with
832 benomyl and carboxin. *Phytopathology* 64, 814. doi:10.1094/Phyto-64-814

833 Tian, H., Ren, W., Tao, B., Sun, G., Chappelka, A., Wang, X., Pan, S., Yang, J., Liu, J., S. Felzer,
834 B., M. Melillo, J., Reilly, J., 2016. Climate extremes and ozone pollution: a growing threat
835 to China's food security. *Ecosystem Health and Sustainability* 2, e01203.

836 doi:10.1002/ehs2.1203

837 Trieu, T.T.N., Goto, D., Yashiro, H., Murata, R., Sudo, K., Tomita, H., Satoh, M., Nakajima, T.,
838 2017. Evaluation of summertime surface ozone in Kanto area of Japan using a semi-
839 regional model and observation. *Atmospheric Environment* 153, 163–181.
840 doi:10.1016/j.atmosenv.2017.01.030

841 Uddling, J., Karlsson, P.E., Glorvigen, A., Selldén, G., 2006. Ozone impairs autumnal resorption
842 of nitrogen from birch (*Betula pendula*) leaves, causing an increase in whole-tree nitrogen
843 loss through litter fall. *Tree physiology* 26, 113–20.

844 Vance, C.P., Uhde-Stone, C., Allan, D.L., 2003. Phosphorus acquisition and use: critical
845 adaptations by plants for securing a nonrenewable resource. *New Phytologist* 157, 423–447.
846 doi:10.1046/j.1469-8137.2003.00695.x

847 Vigani, G., Zocchi, G., Bashir, K., Philippar, K., Briat, J.F., 2013. Cellular iron homeostasis and
848 metabolism in plant. *Frontiers in Plant Science* 4, 490. doi:10.3389/fpls.2013.00490

849 Wang, S., Feng, Z., Wang, X., Gong, W., 2011. Arbuscular mycorrhizal fungi alter the response
850 of growth and nutrient uptake of snap bean (*Phaseolus vulgaris* L.) to O₃. *Journal of*
851 *Environmental Sciences* 23, 968–974. doi:10.1016/S1001-0742(10)60503-7

852 Wang, X., Agathokleous, E., Qu, L., Watanabe, M., Koike, T., 2016. Effects of CO₂ and O₃ on
853 the interaction between root of woody plants and ectomycorrhizae. *Journal of Agricultural*
854 *Meteorology* 72, 95–105. doi:10.2480/agrmet.D-14-00045

855 Whitaker, B.D., Lee, E.H., Rowland, R.A., 1990. EDU and ozone protection: Foliar
856 glycerolipids and steryl lipids in snapbean exposed to O₃. *Physiologia Plantarum* 80, 286–

293.

Wilkinson, S., Mills, G., Illidge, R., Davies, W.J., 2012. How is ozone pollution reducing our food supply? *Journal of Experimental Botany* 63, 527–536. doi:10.1093/jxb/err317

Zhuang, M., Lam, S., Li, Y., Chen, S., 2017. Elevated tropospheric ozone affects the concentration and allocation of mineral nutrients of two bamboo species. *Science of the Total Environment* 577, 231–235. doi:10.1016/j.scitotenv.2016.10.169

Table 1

Table 1 Results of statistical hypotheses testing with contrasts. Five contrasts (C_1 , C_2 , C_3 , C_4 , C_5) were applied to answer research questions, regarding comparisons, which were defined *a priori*. Chlorophyll *a* content (Chla), chlorophyll *b* content (Chlb), total chlorophyll content (TChl), total carotenoid content (TCar), chlorophyll *a* to chlorophyll *b* ratio (Chla/Chlb), optical density 435 reading to optical density 415 reading ratio (OD435/OD415), OD value at 665 nm before the acidification to OD value at 665 nm after acidification ratio (OD665/OD665a), TChl/TCar and leaf mass per area (LMA) were assessed in mature leaves of *Salix sachalinensis* plants treated five times, every nine days, with 0, 200 or 400 mg EDU L⁻¹ and exposed to background ambient ozone (AOZ) or elevated ozone (EOZ) for about a month. Hence, there were six combinational groups: AOZ×EDU0 (AOZ0), AOZ×EDU200 (AOZ200), AOZ×EDU400 (AOZ400), EOZ×EDU0 (EOZ0), EOZ×EDU200 (EOZ200), EOZ×EDU400 (EOZ400). The means are presented in Fig 1.

	1) (AOZ0) vs. (AOZ200+AOZ400)	2) AOZ200 vs. AOZ400	3) (EOZ0) vs. (EOZ200+EOZ400)	4) EOZ200 vs. EOZ400	5) AOZ0 vs. EOZ0
A) Chla	$T=-2.09$, $P=0.059$	$T=-2.23$, $P<0.05$	$T=-4.05$, $P<0.01$	$T=-2.51$, $P<0.05$	$T=3.37$, $P<0.01$
B) Chlb	$T=-2.16$, $P=0.05$	$T=-2.40$, $P<0.05$	$T=-4.35$, $P<0.001$	$T=-0.44$, $P=0.67$	$T=52.03$, $P<0.001$
C) TChl	$T=-2.22$, $P<0.05$	$T=-2.38$, $P<0.05$	$T=-4.41$, $P<0.001$	$T=-2.05$, $P=0.063$	$T=3.13$, $P<0.01$
D) TCar	$T=-0.74$, $P=0.473$	$T=-2.52$, $P<0.05$	$T=-2.08$, $P=0.059$	$T=-0.26$, $P=0.800$	$T=1.04$, $P=0.317$
E) Chla/Chlb	$T=-0.86$, $P=0.405$	$T=-0.37$, $P=0.715$	$T=-1.30$, $P=0.217$	$T=-2.71$, $P<0.05$	$T=3.08$, $P<0.01$
F) OD435/OD415	$T=0.67$, $P=0.518$	$T=0.21$, $P=0.839$	$T=-0.35$, $P=0.734$	$T=-0.97$, $P=0.339$	$T=1.71$, $P=0.113$
G) OD665/OD665a	$T=-0.16$, $P=0.876$	$T=-0.442$, $P=0.666$	$T=-0.61$, $P=0.557$	$T=-0.49$, $P=0.634$	$T=1.44$, $P=0.176$
H) TChl/TCar	$T=-1.80$, $P=0.097$	$T=1.26$, $P=0.232$	$T=-1.93$, $P=0.077$	$T=-2.13$, $P=0.054$	$T=1.76$, $P=0.104$
H) LMA	$T=-2.54$, $P=0.026$	$T=2.52$, $P=0.027$	$T=0.17$, $P=0.869$	$T=2.04$, $P=0.064$	$T=-1.36$, $P=0.199$

Table 2

Table 2 Results of statistical hypotheses testing with General Linear Model (GLM). N content, C content, and C/N ratio were assessed in shed, basal and top leaves of *Salix sachalinensis* plants. Plants were treated five times, every nine days, with 0, 200 or 400 mg EDU L⁻¹ and exposed to background ambient ozone (AOZ) or elevated ozone (EOZ) for about a month. The means are presented in Figs 2-5.

	O3	EDU	Leaf position	O3 x EDU	O3 x Leaf position	EDU x Leaf position	O3 x EDU x Leaf position
N	F=1.54, P=0.238	F=0.81, P=0.466	F=45.61, P<0.001	F=0.48, P=0.632	F=0.60, P=0.559	F=1.02, P=0.417	F=0.52, P=0.724
C	F=0.13, P=0.723	F=0.07, P=0.931	F=2.88, P=0.076	F=0.28, P=0.762	F<0.01, P=0.999	F=0.44, P=0.779	F=0.66, P=0.627
C/N	F=2.65, P=0.130	F=0.60, P=0.566	F=53.99, P<0.001	F=0.25, P=0.784	F=0.05, P=0.947	F=1.18, P=0.343	F=0.62, P=0.652
Mn	F=2.36, P=0.151	F=0.44, P=0.655	F=36.38, P<0.001	F=1.69, P=0.227	F=1.79, P=0.189	F=1.29, P=0.303	F=3.22, P<0.05
Fe	F=0.59, P=0.458	F=1.19, P=0.337	F=25.57, P<0.001	F=1.17, P=0.343	F=6.54, P<0.01	F=1.80, P=0.163	F=0.51, P=0.732
Mg	F=2.49, P=0.140	F=0.16, P=0.853	F=16.27, P<0.001	F=0.43, P=0.659	F=1.67, P=0.209	F=1.57, P=0.215	F=3.49, P<0.05
Ca	F=3.13, P=0.102	F=1.02, P=0.389	F=58.35, P<0.001	F=0.36, P=0.704	F=2.52, P=0.102	F=1.67, P=0.189	F=2.83, P<0.05
Ni	F=0.76, P=0.402	F=0.57, P=0.580	F=21.28, P<0.001	F=0.19, P=0.833	F=8.00, P<0.01	F=2.87, P<0.05	F=3.30, P<0.05
P	F=9.71, P<0.01	F=0.31, P=0.739	F=76.72, P<0.001	F=4.18, P<0.05	F=1.83, P=0.183	F=2.48, P=0.072	F=10.94, P<0.001
Al	F=0.16, P=0.700	F=0.06, P=0.939	F=34.05, P<0.001	F=0.20, P=0.822	F=0.30, P=0.743	F=0.40, P=0.809	F=0.42, P=0.792
K	F=3.28, P=0.095	F=0.50, P=0.621	F=29.18, P<0.001	F=0.95, P=0.415	F=6.37, P<0.01	F=0.71, P=0.594	F=4.26, P<0.01
Cr	F=0.30, P=0.597	F=13.29, P<0.001	F=4.24, P<0.05	F=3.30, P=0.072	F=10.93, P<0.001	F=3.88, P<0.05	F=1.36, P=0.277
K/Mg	F=3.25, P=0.097	F=0.20, P=0.820	F=40.36, P<0.001	F=0.73, P=0.505	F=0.37, P=0.693	F=1.27, P=0.310	F=2.28, P=0.091
K/Ca	F=3.20, P=0.099	F=0.07, P=0.936	F=107.44, P<0.001	F=1.07, P=0.375	F=0.27, P=0.764	F=2.41, P=0.078	F=3.25, P<0.05
Fe/Mn	F=0.38, P=0.548	F=0.05, P=0.955	F=16.01, P<0.001	F=0.02, P=0.985	F=1.98, P=0.160	F=0.22, P=0.926	F=2.09, P=0.056
Ca/Mg	F=0.43, P=0.527	F=0.29, P=0.756	F=160.12, P<0.001	F=0.11, P=0.895	F=2.39, P=0.113	F=3.60, P<0.05	F=2.25, P=0.094
P/Fe	F=6.58, P<0.05	F=1.35, P=0.295	F=226.78, P<0.001	F=9.36, P<0.01	F=0.76, P=0.478	F=3.33, P<0.05	F=12.38, P<0.001
P/Mn	F=3.71, P=0.078	F=1.56, P=0.251	F=238.21, P<0.001	F=3.04, P=0.086	F=2.76, P=0.084	F=2.38, P=0.071	F=2.46, P=0.073

Table 3

Table 3 Factor loading (A) and % contribution (B) of measured response variables to the variability of the eight principal components (F1-F8), after PCA analysis.

(A)	F1	F2	F3	F4	F5	F6	F7	F8	(B)	F1	F2	F3	F4	F5	F6	F7	F8
Chla	-0.31	0.70	-0.04	-0.40	0.13	0.00	-0.05	0.14	2.93	20.36	0.09	11.64	1.53	0.00	0.29	2.35	0.00
Chlb	0.10	0.63	0.23	-0.30	0.35	0.26	0.20	0.10	0.29	16.26	3.16	6.67	11.62	6.49	4.56	1.10	0.00
Tcar	-0.41	0.57	-0.22	-0.02	-0.32	-0.15	-0.34	0.10	5.10	13.59	2.95	0.04	9.51	2.31	13.06	1.07	0.00
OD665/OD665a	0.10	-0.16	-0.47	0.32	0.24	0.67	-0.12	0.06	0.29	1.07	13.67	7.59	5.68	44.32	1.62	0.48	0.00
N	0.09	0.73	0.10	0.38	-0.05	0.10	-0.18	-0.14	0.22	22.28	0.62	10.56	0.23	0.99	3.54	2.22	0.00
C	-0.31	0.21	0.17	0.70	0.41	-0.10	-0.11	0.06	2.87	1.82	1.71	36.04	15.76	0.88	1.45	0.42	0.00
Mn	0.56	0.33	-0.11	-0.11	0.21	-0.30	-0.37	0.19	9.22	4.64	0.76	0.85	4.29	8.79	14.76	4.21	0.00
Fe	0.27	-0.07	-0.54	0.31	-0.09	-0.40	0.14	0.28	2.23	0.20	17.74	7.15	0.81	15.82	2.23	9.56	0.00
Mg	0.78	0.00	-0.13	-0.07	0.09	0.21	-0.22	0.10	18.04	0.00	1.08	0.36	0.72	4.25	5.29	1.28	0.00
Ca	0.76	0.01	-0.37	-0.25	-0.18	0.13	-0.06	0.08	17.05	0.01	8.48	4.46	3.01	1.53	0.46	0.79	0.00
K	0.62	0.25	0.43	0.18	-0.07	-0.04	0.17	-0.18	11.58	2.63	11.43	2.51	0.42	0.19	3.27	3.70	0.00
Ni	0.47	-0.26	0.09	-0.25	0.51	-0.28	-0.14	-0.33	6.51	2.90	0.48	4.48	24.31	7.52	2.14	12.54	0.00
P	0.70	-0.07	0.33	0.29	-0.09	-0.12	-0.07	0.13	14.44	0.22	6.45	6.36	0.77	1.37	0.61	1.96	0.00
Al	0.27	0.40	-0.30	0.08	0.15	-0.10	0.65	0.15	2.21	6.65	5.29	0.51	2.03	0.92	46.16	2.67	0.00
Cr	-0.48	-0.28	-0.27	-0.07	0.45	-0.20	-0.06	0.14	6.95	3.19	4.26	0.34	19.32	3.72	0.38	2.30	0.00
LMA	-0.04	-0.32	0.60	-0.08	0.00	0.10	-0.04	0.67	0.05	4.17	21.82	0.45	0.00	0.91	0.19	53.35	0.00

Fig 1

Fig 1 Mean values $\pm$ SE ($n = 3$) of chlorophyll *a* content (Chla), chlorophyll *b* content (Chlb), total chlorophyll content (TChl), total carotenoid content (TCar), chlorophyll *a* to chlorophyll *b* ratio (Chla/Chlb), optical density 435 reading to optical density 415 reading ratio (OD435/OD415), OD value at 665 nm before the acidification to OD value at 665 nm after acidification ratio (OD665/OD665a) and TChl/TCar in mature leaves of *Salix sachalinensis* plants. Plants were treated five times, every nine days, with 0, 200 or 400 mg EDU L⁻¹ and exposed to background ambient ozone (AOZ) or elevated ozone (EOZ) for about a month. Results of statistical testing are presented in Table 1.

Fig 2

Fig 2 Mean values $\pm$ SE ($n = 3$) of N content, C content, and C/N ratio in shed, basal and top leaves of *Salix sachalinensis* plants. Plants were treated five times, every nine days, with 0, 200 or 400 mg EDU L⁻¹ and exposed to background ambient ozone (AOZ) or elevated ozone (EOZ) for about a month. Statistical analysis was conducted with General Linear Model (GLM); the only significant factor was Leaf position for N (Basal>Shed>Top) and C/N (Top>Shed>Basal). Results of statistical testing are presented in Table 2.

Fig 3

Fig 3 Mean values $\pm$ SE ($n = 3$) of P, Mn, Fe, Mg and Ca content in shed, basal and top leaves of *Salix sachalinensis* plants. Plants were treated five times, every nine days, with 0, 200 or 400 mg EDU L⁻¹ and exposed to background ambient ozone (AOZ) or elevated ozone (EOZ) for about a month. Statistical analysis was conducted with General Linear Model (GLM), and the results are presented in Table 2. Different lowercase letters above means indicate statistically significant differences in the interaction Ozone $\times$ EDU $\times$ Leaf position according to Bonferroni test ($\alpha=0.05$).

Fig 4

Fig 4 Mean values $\pm$ SE ($n = 3$) of Ni content, Al content, K content, Cr content and P/Fe ratio in shed, basal and top leaves of *Salix sachalinensis* plants. Plants were treated five times, every nine days, with 0, 200 or 400 mg EDU L^{-1} and exposed to background ambient ozone (AOZ) or elevated ozone (EOZ) for about a month. Statistical analysis was conducted with General Linear Model (GLM), and the results are presented in Table 2. Different lowercase letters above means indicate statistically significant differences in the interaction Ozone $\times$ EDU $\times$ Leaf position, whereas different lowercase below means indicate statistically significant differences in the interaction EDU $\times$ Leaf position according to Bonferroni test ($\alpha=0.05$).

Fig 5

Fig 5 Mean values $\pm$ SE ($n = 3$) of Ni content, Al content, K content, Cr content and P/Fe ratio in shed, basal and top leaves of *Salix sachalinensis* plants. Plants were treated five times, every nine days, with 0, 200 or 400 mg EDU L^{-1} and exposed to background ambient ozone (AOZ) or elevated ozone (EOZ) for about a month. Statistical analysis was conducted with General Linear Model (GLM), and the results are presented in Table 2. Different lowercase letters above means indicate statistically significant differences in the interaction Ozone $\times$ EDU $\times$ Leaf position, whereas different lowercase below means indicate statistically significant differences in the interaction EDU $\times$ Leaf position according to Bonferroni test ($\alpha=0.05$).

Fig 6

Fig 6 Biplot analysis of the main variables (shown as arrows): Mn, Fe, Mg, Ca, Ni, P, Al, K, Cr, N, C, Chla, Chlb, CarT, LMA and the OD665/OD665a. The variables EDU, Ozone, K/Mg, K/Ca, Fe/Mn, Ca/Mg, P/Fe, P/Mn, Chla/Chlb, ChlT, ChlT/CarT, N/C, and OD435/OD415 (shown as line segments) were passively only included as supplementary variables. The Centroid of the ozone treatments (Elevated Ozone, EOZ and Ambient Ozone, AOZ) as well as of the EDU treatments (0, 200 and 400 mg L⁻¹) groups of samples are also supplementary indicated by circles marked by italics fonts. Observations are shown as circle point. By the exception of EDU, Ozone and TChl, the rest of the supplementary variables, had very low contribution on the first two principal components and, although their line segments are shown on the biplot, they are not named for clarity of the figure.