Title	Serological evidence of Zika virus infection in non-human primates in Zambia
Author(s)	Wastika, Christida E.; Sasaki, Michihito; Yoshii, Kentaro; Anindita, Paulina D.; Hang'ombe, Bernard M.; Mweene, Aaron S.; Kobayashi, Shintaro; Kariwa, Hiroaki; Carr, Michael J.; Hall, William W.; Eshita, Yuki; Orba, Yasuko; Sawa, Hirofumi
Citation	Archives of virology, 164(8), 2165-2170 https://doi.org/10.1007/s00705-019-04302-0
Issue Date	2019-08
Doc URL	http://hdl.handle.net/2115/79038
Rights	The final publication is available at link.springer.com
Type	article (author version)
File Information	Archives of virology 164 (8)_ 2165-2170.pdf

[Instructions for use](#)

Serological evidence of Zika virus infection in non-human primates in Zambia

Christida E. Wastika¹, Michihito Sasaki¹, Kentaro Yoshii², Paulina D. Anindita¹, Bernard M. Hang'ombe^{3,10,11}, Aaron S. Mweene^{4,10,11}, Shintaro Kobayashi², Hiroaki Kariwa², Michael J. Carr^{6,7}, William W. Hall^{7,8,9,11}, Yuki Eshita⁵, Yasuko Orba¹, Hirofumi Sawa^{1,7,9,10,11}

¹Division of Molecular Pathobiology, Research Center for Zoonosis Control, Hokkaido University, N20, W10, Kita-Ku, Sapporo 001-0020, Japan

²Laboratory of Public Health, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo, N18, W9, Kita-ku, Sapporo 060-0818, Japan

³Department of Para-clinical Studies, School of Veterinary Medicine, University of Zambia, P.O. Box 32379, Lusaka 10101, Zambia

⁴Department of Disease Control, School of Veterinary Medicine, University of Zambia, P.O. Box 32379, Lusaka 10101, Zambia

⁵Hokudai Center for Zoonosis Control in Zambia, Research Center for Zoonosis Control, Hokkaido University, P.O. Box 32379, Lusaka 10101, Zambia

⁶National Virus Reference Laboratory, School of Medicine, University College Dublin, Dublin 4, Ireland

⁷Global Station for Zoonosis Control, Global Institution for Collaborative Research and Education (GI-CoRE), Hokkaido University, N20, W10, Kita-ku, Sapporo 001-0020, Japan

⁸Centre for Research in Infectious Diseases, School of Medicine, University College Dublin, Dublin 4, Ireland

⁹Global Virus Network, 725 West Lombard St, Baltimore, MD 21201, USA

¹⁰Global Virus Network Affiliate Centre of Excellence, University of Zambia, P.O. Box 32379, Lusaka 10101, Zambia

¹¹Africa Centre of Excellence for Infectious Diseases of Humans and Animals, P.O. Box 32379, Lusaka 10101, Zambia

Corresponding author: Hirofumi Sawa

E-mail: h-sawa@czc.hokudai.ac.jp

Telephone: +81-11-706-5185

Abstract

Zika virus (ZIKV) circulation occurs between non-human primates (NHPs) in a sylvatic transmission cycle. To investigate evidence of flavivirus infection in NHPs in Zambia, we performed plaque reduction neutralization tests (PRNT) to quantify neutralizing antibodies. PRNT revealed that sera from NHPs (African green monkeys and baboons) exhibited neutralizing activity against ZIKV (34.4%; 33/96), whereas PRNT for yellow fever virus from NHP sera showed no neutralization activity. ZIKV genomic RNA was not detected in splenic tissues from NHPs suggesting that the presence of anti-ZIKV neutralizing antibodies represented resolved infections. Our evidence suggests that ZIKV is maintained in NHP reservoirs in Zambia.

Zika virus (ZIKV) is an enveloped, single-stranded RNA virus, belonging to the family *Flaviviridae*, genus *Flavivirus*. ZIKV was serendipitously isolated from a rhesus monkey during sentinel screening for yellow fever virus in Uganda in 1947 [5]. ZIKV infection was likely underreported in the intervening decades and considered a benign infection until a large epidemic in 2007 on Yap Island in Micronesia [13]. Thereafter, the Asian lineage of ZIKV infection expanded to French Polynesia and the Americas and was associated with severe neurological disease *in utero* and in adults [20].

ZIKV is mainly transmitted by *Aedes* mosquito vectors; however, human-to-human infections have also been reported. In 2008, a scientist, who had recently returned to the United States from Senegal, sexually transmitted ZIKV to his wife [15]. Maternofetal transmission of ZIKV was first reported in Brazil and was associated with cases of microcephaly, *in utero* growth restriction and placental insufficiency [22]. Cases of ZIKV-associated microcephaly were subsequently documented in Colombia [27], Thailand and Vietnam [24], and ZIKV-related Guillain–Barré syndrome (GBS), a severe autoimmune disease, was reported in French Polynesia [6] and Colombia [31].

ZIKV infection experiments in rhesus macaque (*Macaca mulata*) and cynomolgus macaque (*Macaca fascicularis*) exhibit similar symptoms with those observed in human cases [9, 19, 30], suggesting that some NHPs are susceptible to ZIKV. Previous serological studies in humans demonstrated antibodies to ZIKV and yellow fever virus (YFV) in Zambia [2, 3]. Anti-ZIKV antibody has also been detected from baboons in Tanzania and African green monkeys from Gambia but not from baboons in Zambia using an enzyme-linked immunosorbent assay (ELISA) [4]. Hitherto, little prior knowledge exists for ZIKV infection in NHPs in Zambia. Further investigation is important to ascertain the sero-status of ZIKV and YFV in Zambian NHPs. In the present study, we have examined sera from NHPs in Zambia for the presence of neutralization antibodies against ZIKV and YFV using a plaque reduction neutralization test (PRNT).

Blood and splenic tissues were collected from Zambian malbrouck monkeys (*Chlorocebus cynosuros*; n=48), chacma baboons (*Papio ursinus*; n=25) and yellow baboons (*Papio cynocephalus*; n=23) in the Livingstone (Southern) and Mfuwe (Eastern) districts between 2009 and 2010, as previously described [33] and stored at -80°C prior to examination. NHPs were speciated by mitochondrial cytochrome b (*cytb*) gene sequencing as described by Sasaki et al. (2013) [33]. Ethical approval was obtained from the then Zambia Wildlife Authority (ZAWA), now the Department of National Parks and Wildlife, Ministry of Tourism and Arts (certificate no. 2604) [7]. PRNT was conducted using ZIKV MR-766, YFV 17D and tick-borne encephalitis virus (TBEV) Oshima 5-10 reference strains as positive controls for detection of virus-specific antibody. PRNT

against TBEV was conducted to examine cross-reactivity between flaviviruses. Prior to the experiments, all sera were thawed at 37°C, then diluted 1:5, 1:20 and 1:80 in Dulbecco's Modified Essential Medium (DMEM) for ZIKV and YFV or Minimum Essential Medium (MEM) for TBEV. The diluted sera were inactivated at 56°C for 30 minutes and thereafter incubated 1:1 with ZIKV (25 PFU/well), YFV (750 PFU/well) or TBEV (50 PFU/well) at 37°C for 30 minutes. The diluted, inactivated sera were then inoculated onto Vero cells (ZIKV and YFV) and BHK cells (TBEV) and incubated at 37°C for 1 hour. After incubation, the cells were covered by DMEM supplemented with 1.25% methylcellulose and 2% fetal bovine serum (ZIKV and YFV) or MEM supplemented with 1.5% carboxymethyl cellulose and 2% fetal bovine serum (TBEV) and incubated for 5 days (ZIKV), 4 days (TBEV) or 8 days (YFV) at 37°C. Finally, the cells were fixed using 3.7% buffered formaldehyde for 30 minutes at room temperature and stained using 0.1% (TBEV) or 1% (ZIKV and YFV) crystal violet. The plaque number was counted, and neutralizing activity was determined by a 50% reduction number of viral plaques compared to the no serum control (PRNT₅₀). Positive serum samples were judged by a 4-fold difference between flaviviruses or that, at least, showed PRNT₅₀ ≥1:40 at the final dilution. For TBEV PRNT, serum samples, which showed PRNT₅₀ ≥1:20, were considered to be positive for neutralization activity [35].

Total RNA was extracted from splenic tissues as described by Sasaki et al. (2013) [33] and employed to screen for the presence of ZIKV and YFV viral RNA. Detection of ZIKV and YFV genome were examined by real-time reverse transcriptase (RT)-PCR with an Express One-Step Super Script qRT-PCR kit (Invitrogen, Carlsbad, CA) using the StepOnePlus instrument (Applied Biosystems, Foster City, CA). The oligonucleotide primers and hydrolysis probe and final concentrations employed for detection of ZIKV were as follows: 500 nM ZIKA2 1176F (5'-GAC ATG GCT TCG GAC AG-3'), 500 nM ZIKA2 1308R (5'-CTT TGC CAA AAA GTC CAC A-3'), and 250 nM ZIKA2 1209 probe (5'-FAM-GGT GAA GCC TAC CTT GAC AAG CA-MGB-3'). Our ZIKV primers and probe were designed based on conserved regions of several ZIKV strains, MR-766 (East African, Uganda, LC002520), FSS13025 (Asian, Cambodia, KU955593), DaKAR 41525 (West African, Senegal, KU955591), and Brazil/PE243/2015 (American, Brazil, KX197192), by modifying a widely used ZIKV qRT-PCR system described by Lanciotti et al. [23] (Supplementary Figure 1). This ZIKA2 1209 qRT-PCR system detected ZIKV RNA at approximately one PFU of DaKAR 41525, MR-766, and FSS 13025 reference strains at C_T values of 32 (Supplementary Table 1). The primers sets and probe and final concentrations employed for detection of YFV genome were from a previous report by Domingo et al. (2012) [12]: 500 nM 15F (5'-GCT AAT TGA GGT GYA TTG GTC TGC-3'), 500 nM 103R (5'-CTG CTA ATC GCT

CAA MGA ACG-3') and 250 nM YFV Probe FAM41 (5'-FAM-ATC GAG TTG CTA GGC AAT AAA CAC-MGB-3'). This primer set detected YFV RNA at 1 PFU with C_T values around 33. Thermocycling conditions were 50°C for 15 minutes, 95°C for 2 minutes followed by 45 cycles 95°C for 15 seconds and 60°C for 1 minute for both ZIKV and YFV.

The PRNT revealed that 34.4% (33/96) of sera of NHPs had neutralizing antibodies against ZIKV with comparable levels of seropositivity in *Zambian malbrouck monkeys* in Livingstone (34.8%; 8/23) and in *Mfuwe* (32.0%; 8/25). Sera from *chacma baboons* in Livingstone had a relatively higher seroprevalence (48.0%; 12/25) compared to *yellow baboons* in Mfuwe (21.7%; 5/23) (Table 1, Table 2 and Supplementary Table 2). In addition, we did not detect any neutralization activity against TBEV in the NHP samples (Table 1). Although we could identify specific neutralization antibodies for ZIKV, we found one sample had both neutralization activity against both ZIKV and YFV with sera titer 160 and 40, respectively (Table 2). This phenomenon may have occurred because of either co-infection of ZIKV and YFV or because of cross-reactivity of antibodies. Flaviviruses possess similar structures and epitopes within their envelope proteins that could potentially be recognized by cross-neutralizing antibodies [34] and cross-neutralization activities among flaviviruses have been reported previously [21, 28]. Therefore, the one sample from a yellow baboon that has neutralization activities for both of ZIKV and YFV may have cross-neutralization activity. For genome screening, neither ZIKV nor YFV viral RNA were detected in splenic tissues from NHPs by real time-RT-PCR which suggested that no acute phase viremic samples were present and that the presence of anti-ZIKV neutralizing antibodies represented prior exposure to a resolved infection.

Serological studies in NHPs in Malaysia, Brazil and on the African continent revealed that some species of NHPs have been shown to have neutralization activities against ZIKV, including *Bornean orangutans*, *African green monkeys*, *baboons*, *howler monkeys*, *marmosets* and *capuchin monkeys* [4, 10, 14, 29, 37]. Along with previous reports, our study has shown seroprevalence of anti-ZIKV neutralization antibodies in *malbrouck monkeys*, *chacma baboons* and *yellow baboons* in Zambia. These studies suggest that NHPs play a role in ZIKV maintenance in nature has previously been proposed by Diallo et al. (2014) and Terzian et al. (2018) [11, 36]. Currently, neutralization assays are widely used as a method to differentiate arbovirus infections in human as reported by Fritzell et al. (2018) [16]. The viral neutralization test is the most frequently used method compared to other neutralization tests such as immunofluorescence assays. Several serological studies for ZIKV infection in animals are also used neutralization assay and, specifically PRNT [8, 10, 29, 37]. PRNT (using live viruses) was used for the determination of type-specific antibodies and differentiation of the infecting viruses. Even

though PRNT is the most specific serological diagnosis for flavivirus infection, this procedure requires tissue culture facilities and laboratories with Biosafety Level-3 (BSL-3) or BSL-4 using physical containment level 3 (P3) or P4 viruses) [25].

Serological studies of pathogens in NHPs can be applied as sentinel warning systems to assess the likelihood of human outbreaks of zoonotic disease. It has been reported that Almeida et al. succeeded in enhancing vaccine uptake prior to YFV outbreaks in humans by combined active and passive surveillance in NHPs, which resulted in vaccination being administered within a 2 km radius from the location where YFV in NHPs was detected [1]. Spillover events of flaviviruses from NHPs to humans are dependent on several factors, such as, reservoir hosts and their distribution, infectivity, pathogenicity/transmissibility of pathogens from infected hosts and survival rates of infected hosts [32]. Even though there is a paucity of information on precisely how sylvatic reservoirs with circulating ZIKV could infect humans, several urban transmissions, including mosquito-borne, sexually and maternofetal infection have been reported [17, 18, 26]. Our findings provide new insights into ZIKV circulation in Zambia and represent the first report of serologic detection of sylvatic ZIKV in Zambia. To fully understand the epidemiology and ecology of ZIKV in Zambia, further studies need to be conducted in sylvatic, urban and peri-urban settings in mosquito vectors and wildlife.

Acknowledgement

We are deeply grateful to the then Zambia Wildlife Authority (ZAWA), now the Department of National Parks and Wildlife, Ministry of Tourism and Arts, for assistance with sample collection in Zambia. ZIKV RNAs of DaKAR 41525 and FSS 13025 were kindly provided by Professor Scott C. Weaver, UTMB.

Compliance with Ethical Standards:

Funding: This study was supported by grants for Scientific Research (C), (B) from the Ministry of Education, Culture, Sports, Science and Technology, Japan (MEXT) /Japan Society for the Promotion of Science (JSPS) KAKENHI (JP15K07715, JP16H05805, JP17H03910); and grants for Scientific Research on Innovative Areas and International Group from the MEXT/JSPS KAKENHI (JP16H06431, JP16H06429, JP16K21723), and the Japan Initiative for Global Research Network of Infectious Diseases (J-GRID) from Japan Agency for Medical Research and Development (AMED) (JP18fm0108008).

Conflict of Interest: The authors declare that they have no conflicts of interest.

Ethical approval: All applicable international, national, and/or institutional guidelines for the care and use of animals were followed.

Ethical approval: This article does not contain any studies with human participants performed by any of the authors.

References

1. Almeida MA, Cardoso JaC, Dos Santos E, da Fonseca DF, Cruz LL, Faraco FJ, Bercini MA, Vettorello KC, Porto MA, Mohrdieck R, Ranieri TM, Schermann MT, Sperb AF, Paz FZ, Nunes ZM, Romano AP, Costa ZG, Gomes SL, Flannery B (2014) Surveillance for yellow Fever virus in non-human primates in southern Brazil, 2001-2011: a tool for prioritizing human populations for vaccination. *PLoS Negl Trop Dis* 8:e2741
2. Babaniyi O, Mazaba-Liwewe ML, Masaninga F, Mwaba P, Mulenga D, Songolo P, Mweene-Ndumba I, Rudatsikira E, Siziya S (2016) Prevalence of Yellow Fever in North-Western Province of Zambia. *Int. J. Public. Health. Epidemiol.*, pp 29-32
3. Babaniyi OA, Mwaba P, Songolo P, Mazaba-Liwewe ML, Mweene-Ndumba I, Masaninga F, Rudatsikira E, Siziya S (2015) Seroprevalence of Zika virus infection specific IgG in Western and North-Western Provinces of Zambia. *Int. J. Public. Health. Epidemiol.*, pp 110-114
4. Buechler CR, Bailey AL, Weiler AM, Barry GL, Breitbach ME, Stewart LM, Jasinska AJ, Freimer NB, Apetrei C, Phillips-Conroy JE, Jolly CJ, Rogers J, Friedrich TC, O'Connor DH (2017) Seroprevalence of Zika Virus in Wild African Green Monkeys and Baboons. *mSphere* 2: e00392-16
5. Bueno MG, Martinez N, Abdalla L, Duarte Dos Santos CN, Chame M (2016) Animals in the Zika Virus Life Cycle: What to Expect from Megadiverse Latin American Countries. *PLoS Negl Trop Dis* 10:e0005073
6. Cao-Lormeau VM, Blake A, Mons S, Lastere S, Roche C, Vanhomwegen J, Dub T, Baudouin L, Teissier A, Larre P, Vial AL, Decam C, Choumet V, Halstead SK, Willison HJ, Musset L, Manuguerra JC, Despres P, Fournier E, Mallet HP, Musso D, Fontanet A, Neil J, Ghawché F (2016) Guillain-Barré Syndrome outbreak associated with Zika virus infection in French Polynesia: a case-control study. *Lancet* 387:1531-1539
7. Carr M, Kawaguchi A, Sasaki M, Gonzalez G, Ito K, Thomas Y, Hang'ombe BM, Mweene AS, Zhao G, Wang D, Orba Y, Ishii A, Sawa H (2017) Isolation of a simian immunodeficiency virus from a malbrouck (*Chlorocebus cynosuros*). *Arch Virol* 162:543-548
8. Chua CL, Chan YF, Andu ESGS, Rovie-Ryan JJ, Sitam FT, Verasahib K, Sam IC (2019) Little Evidence of Zika Virus Infection in Wild Long-Tailed Macaques, Peninsular Malaysia. *Emerg Infect Dis* 25:374-376

9. Coffey LL, Pesavento PA, Keesler RI, Singapuri A, Watanabe J, Watanabe R, Yee J, Bliss-Moreau E, Cruzen C, Christe KL, Reader JR, von Morgenland W, Gibbons AM, Allen AM, Linnen J, Gao K, Delwart E, Simmons G, Stone M, Lanteri M, Bakkour S, Busch M, Morrison J, Van Rompay KK (2017) Zika Virus Tissue and Blood Compartmentalization in Acute Infection of Rhesus Macaques. *PLoS One* 12:e0171148
10. de Oliveira-Filho EF, Oliveira RAS, Ferreira DRA, Laroque PO, Pena LJ, Valença-Montenegro MM, Mota RA, Gil LHV (2018) Seroprevalence of selected flaviviruses in free-living and captive capuchin monkeys in the state of Pernambuco, Brazil. *Transbound Emerg Dis*
11. Diallo D, Sall AA, Diagne CT, Faye O, Ba Y, Hanley KA, Buenemann M, Weaver SC, Diallo M (2014) Zika virus emergence in mosquitoes in southeastern Senegal, 2011. *PLoS One* 9:e109442
12. Domingo C, Patel P, Yillah J, Weidmann M, Méndez JA, Nakouné ER, Niedrig M (2012) Advanced yellow fever virus genome detection in point-of-care facilities and reference laboratories. *J Clin Microbiol* 50:4054-4060
13. Duffy MR, Chen TH, Hancock WT, Powers AM, Kool JL, Lanciotti RS, Pretrick M, Marfel M, Holzbauer S, Dubray C, Guillaumot L, Griggs A, Bel M, Lambert AJ, Laven J, Kosoy O, Panella A, Biggerstaff BJ, Fischer M, Hayes EB (2009) Zika virus outbreak on Yap Island, Federated States of Micronesia. *N Engl J Med* 360:2536-2543
14. Favoretto S, Araujo D, Oliveira D, Duarte N, Mesquita F, Zanotto P, Durigon E (2016) First detection of Zika virus in neotropical primates in Brazil: a possible new reservoir. *bioRxiv*
15. Foy BD, Kobylinski KC, Chilson Foy JL, Blitvich BJ, Travassos da Rosa A, Haddow AD, Lanciotti RS, Tesh RB (2011) Probable non-vector-borne transmission of Zika virus, Colorado, USA. *Emerg Infect Dis* 17:880-882
16. Fritzell C, Rousset D, Adde A, Kazanji M, Van Kerkhove MD, Flamand C (2018) Current challenges and implications for dengue, chikungunya and Zika seroprevalence studies worldwide: A scoping review. *PLoS Negl Trop Dis* 12:e0006533
17. Gao D, Lou Y, He D, Porco TC, Kuang Y, Chowell G, Ruan S (2016) Prevention and Control of Zika as a Mosquito-Borne and Sexually Transmitted Disease: A Mathematical Modeling Analysis. *Sci Rep* 6:28070
18. Hallmaier-Wacker LK, Munster VJ, Knauf S (2017) Disease reservoirs: from conceptual frameworks to applicable criteria. *Emerg Microbes Infect* 6:e79

19. Hirsch AJ, Smith JL, Haese NN, Broeckel RM, Parkins CJ, Kreklywich C, DeFilippis VR, Denton M, Smith PP, Messer WB, Colgin LM, Ducore RM, Grigsby PL, Hennebold JD, Swanson T, Legasse AW, Axthelm MK, MacAllister R, Wiley CA, Nelson JA, Streblow DN (2017) Zika Virus infection of rhesus macaques leads to viral persistence in multiple tissues. *PLoS Pathog* 13:e1006219
20. Ikejezie J, Shapiro CN, Kim J, Chiu M, Almiron M, Ugarte C, Espinal MA, Aldighieri S (2017) Zika Virus Transmission-Region of the Americas, May 15, 2015-December 15, 2016. *Am J Transplant* 17:1681-1686
21. Keasey SL, Pugh CL, Jensen SM, Smith JL, Hontz RD, Durbin AP, Dudley DM, O'Connor DH, Ulrich RG (2017) Antibody Responses to Zika Virus Infections in Environments of Flavivirus Endemicity. *Clin Vaccine Immunol* 24
22. Kindhauser MK, Allen T, Frank V, Santhana RS, Dye C (2016) Zika: the origin and spread of a mosquito-borne virus. *Bull World Health Organ* 94:675-686C
23. Lanciotti RS, Kosoy OL, Laven JJ, Velez JO, Lambert AJ, Johnson AJ, Stanfield SM, Duffy MR (2008) Genetic and serologic properties of Zika virus associated with an epidemic, Yap State, Micronesia, 2007. *Emerg Infect Dis* 14:1232-1239
24. Lim SK, Lim JK, Yoon IK (2017) An Update on Zika Virus in Asia. *Infect Chemother* 49:91-100
25. Maeda A, Maeda J (2013) Review of diagnostic plaque reduction neutralization tests for flavivirus infection. *Vet J* 195:33-40
26. Marano G, Pupella S, Vaglio S, Liumbruno GM, Grazzini G (2016) Zika virus and the never-ending story of emerging pathogens and transfusion medicine. *Blood Transfus* 14:95-100
27. Mattar S, Ojeda C, Arboleda J, Arrieta G, Bosch I, Botia I, Alvis-Guzman N, Perez-Yepes C, Gerhke L, Montero G (2017) Case report: microcephaly associated with Zika virus infection, Colombia. *BMC Infect Dis* 17:423
28. Morales MA, Fabbri CM, Zunino GE, Kowalewski MM, Luppó VC, Enríe DA, Levis SC, Calderón GE (2017) Detection of the mosquito-borne flaviviruses, West Nile, Dengue, Saint Louis Encephalitis, Ilheus, Bussuquara, and Yellow Fever in free-ranging black howlers (*Alouatta caraya*) of Northeastern Argentina. *PLoS Negl Trop Dis* 11:e0005351
29. Moreira-Soto A, Carneiro IO, Fischer C, Feldmann M, Kümmerer BM, Silva NS, Santos UG, Souza BFCD, Liborio FA, Valença-Montenegro MM, Laroque PO, da Fontoura FR, Oliveira AVD, Drosten

- C, de Lamballerie X, Franke CR, Drexler JF (2018) Limited Evidence for Infection of Urban and Peri-urban Nonhuman Primates with Zika and Chikungunya Viruses in Brazil. *mSphere* 3
30. Osuna CE, Lim SY, Deleage C, Griffin BD, Stein D, Schroeder LT, Omange RW, Best K, Luo M, Hraber PT, Andersen-Elyard H, Ojeda EF, Huang S, Vanlandingham DL, Higgs S, Perelson AS, Estes JD, Safronetz D, Lewis MG, Whitney JB (2016) Zika viral dynamics and shedding in rhesus and cynomolgus macaques. *Nat Med* 22:1448-1455
 31. Parra B, Lizarazo J, Jiménez-Arango JA, Zea-Vera AF, González-Manrique G, Vargas J, Angarita JA, Zuñiga G, Lopez-Gonzalez R, Beltran CL, Rizcala KH, Morales MT, Pacheco O, Ospina ML, Kumar A, Cornblath DR, Muñoz LS, Osorio L, Barreras P, Pardo CA (2016) Guillain-Barré Syndrome Associated with Zika Virus Infection in Colombia. *N Engl J Med* 375:1513-1523
 32. Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, Lloyd-Smith JO (2017) Pathways to zoonotic spillover. *Nat Rev Microbiol* 15:502-510
 33. Sasaki M, Ishii A, Orba Y, Thomas Y, Hang'ombe BM, Moonga L, Mweene AS, Ogawa H, Nakamura I, Kimura T, Sawa H (2013) Human parainfluenza virus type 3 in wild nonhuman primates, Zambia. *Emerg Infect Dis* 19
 34. Stiasny K, Kiermayr S, Holzmann H, Heinz FX (2006) Cryptic properties of a cluster of dominant flavivirus cross-reactive antigenic sites. *J Virol* 80:9557-9568
 35. Takashima I, Morita K, Chiba M, Hayasaka D, Sato T, Takezawa C, Igarashi A, Kariwa H, Yoshimatsu K, Arikawa J, Hashimoto N (1997) A case of tick-borne encephalitis in Japan and isolation of the the virus. *J Clin Microbiol* 35:1943-1947
 36. Terzian ACB, Zini N, Sacchetto L, Rocha RF, Parra MCP, Del Sarto JL, Dias ACF, Coutinho F, Rayra J, da Silva RA, Costa VV, Fernandes NCCA, Réssio R, Díaz-Delgado J, Guerra J, Cunha MS, Catão-Dias JL, Bittar C, Reis AFN, Santos INPD, Ferreira ACM, Cruz LEAA, Rahal P, Ullmann L, Malossi C, Araújo JP, Widen S, de Rezende IM, Mello É, Pacca CC, Kroon EG, Trindade G, Drumond B, Chiaravalloti-Neto F, Vasilakis N, Teixeira MM, Nogueira ML (2018) Evidence of natural Zika virus infection in neotropical non-human primates in Brazil. *Sci Rep* 8:16034
 37. Wolfe ND, Kilbourn AM, Karesh WB, Rahman HA, Bosi EJ, Cropp BC, Andau M, Spielman A, Gubler DJ (2001) Sylvatic transmission of arboviruses among Bornean orangutans. *Am J Trop Med Hyg* 64:310-316

Table 1. Flavivirus seropositivity in Zambian non-human primates.

Collecting place	Non-human primate species	Number of sera	Not detected	PRNT (+) YFV	PRNT (+) ZIKV	PRNT (+) TBEV
Livingstone	Chacma baboons (<i>Papio ursinus</i>)	25	13	0	12 (12/25=48.0%)	0
	Zambian malbrouck monkeys (<i>Chlorocebus cynosuros</i>)	23	15	0	8 (8/23=34.8%)	0
Mfuwe	Yellow baboons (<i>Papio cynocephalus</i>)	23	18	0	5 (5/23=21.7%)	0
	Zambian malbrouck monkeys (<i>Chlorocebus cynosuros</i>)	25	17	0	8 (8/25=32.0%)	0
TOTAL		96	63	0	33 (33/96=34.4%)	0

Not detected: the antibody titer is $\leq 1:10$ for ZIKV, YFV and TBEV; PRNT (+) YFV: the antibody titer $>1:10$ for YFV or 4-fold different to ZIKV and TBEV; PRNT (+) ZIKV: the antibody titer $>1:10$ for ZIKV or 4-fold different to YFV and TBEV; PRNT (+) TBEV: the antibody titer $>1:10$ for TBEV or 4-fold different to YFV and ZIKV.

Table 2. PRNT50 values for Zika virus (ZIKV), Yellow fever virus (YFV) and Tick-borne encephalitis virus (TBEV) from non-human primate sera collected in Livingstone, Zambia.

Sample No.	Species	PRNT 50			Serological interpretation
		ZIKV	YFV	TBEV	
1	<i>Papio ursinus</i>	<10	<10	ND*	
2	<i>Papio ursinus</i>	160	10	<10	ZIKV
3	<i>Papio ursinus</i>	40	10	<10	ZIKV
4	<i>Papio ursinus</i>	<10	<10	ND	
5	<i>Papio ursinus</i>	10	<10	ND	
6	<i>Papio ursinus</i>	10	<10	ND	
7	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
8	<i>Papio ursinus</i>	160	<10	<10	ZIKV
9	<i>Papio ursinus</i>	<10	<10	ND	
10	<i>Papio ursinus</i>	40	<10	<10	ZIKV
11	<i>Papio ursinus</i>	160	40	<10	ZIKV
12	<i>Chlorocebus cynosuros</i>	10	<10	ND	
13	<i>Chlorocebus cynosuros</i>	40	<10	<10	ZIKV
14	<i>Chlorocebus cynosuros</i>	10	<10	ND	
15	<i>Papio ursinus</i>	40	<10	<10	ZIKV
16	<i>Papio ursinus</i>	<10	<10	ND	
17	<i>Papio ursinus</i>	160	<10	<10	ZIKV
18	<i>Chlorocebus cynosuros</i>	40	<10	<10	ZIKV
19	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
22	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
23	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
24	<i>Chlorocebus cynosuros</i>	40	<10	<10	ZIKV
25	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
26	<i>Chlorocebus cynosuros</i>	40	10	<10	ZIKV

27	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
28	<i>Chlorocebus cynosuros</i>	10	<10	ND	
29	<i>Chlorocebus cynosuros</i>	40	<10	<10	ZIKV
30	<i>Chlorocebus cynosuros</i>	10	<10	ND	
31	<i>Papio ursinus</i>	<10	<10	ND	
32	<i>Papio ursinus</i>	10	10	ND	
33	<i>Papio ursinus</i>	<10	<10	ND	
34	<i>Papio ursinus</i>	40	<10	<10	ZIKV
35	<i>Papio ursinus</i>	40	10	<10	ZIKV
36	<i>Papio ursinus</i>	40	<10	<10	ZIKV
37	<i>Papio ursinus</i>	10	<10	ND	
38	<i>Papio ursinus</i>	<10	<10	ND	
39	<i>Papio ursinus</i>	160	<10	<10	ZIKV
40	<i>Papio ursinus</i>	<10	<10	ND	
41	<i>Papio ursinus</i>	<10	<10	ND	
42	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
43	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
44	<i>Chlorocebus cynosuros</i>	40	<10	<10	ZIKV
45	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
46	<i>Chlorocebus cynosuros</i>	160	<10	ND	ZIKV
47	<i>Chlorocebus cynosuros</i>	160	10	<10	ZIKV
48	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
49	<i>Chlorocebus cynosuros</i>	<10	10	ND	
50	<i>Papio ursinus</i>	40	<10	<10	ZIKV

Positives are shown in bold. *ND: not determined. ZIKV: Zika virus; YFV: Yellow fever virus; TBEV: Tick-borne encephalitis virus. The numbers of PRNT50 are represented as the greatest dilution with a positive result.

Supplementary Table 1.
Cycle thresholds (C_T) values of ZIKV qRT-PCR systems.

ZIKA1107 (Lanciotti 2008)											
MR-766			DaKAR 41525				FSS 13025				
PFU	C _T mean	C _T SD	Dilution	PFU*	C _T mean	C _T SD	Dilution	PFU*	C _T mean	C _T SD	
5000	19.94	0.06									
500	23.30	0.06	1.E-01	722	29.09	0.06	1.E-01	1088	21.51	0.17	
50	26.83	0.06	1.E-02	87	32.41	0.21	1.E-02	132	24.63	0.05	
5	30.64	0.12	1.E-03	11	35.29	0.36	1.E-03	15	27.97	0.08	
0.5	33.95	0.29	1.E-04	0.9	ND		1.E-04	1.2	31.83	0.25	
0.05	37.21	0.69	1.E-05	0.1	ND		1.E-05	0.1	35.30	0.42	

ZIKA2 1209 (in this experiment)											
MR-766			DaKAR 41525				FSS 13025				
PFU	C _T mean	C _T SD	Dilution	PFU*	C _T mean	C _T SD	Dilution	PFU*	C _T mean	C _T SD	
5000	19.22	0.09									
500	22.70	0.08	1.E-01	722	22.17	0.12	1.E-01	1088	21.54	0.18	
50	26.21	0.09	1.E-02	87	25.43	0.21	1.E-02	132	24.78	0.05	
5	29.81	0.23	1.E-03	11	28.68	0.32	1.E-03	15	28.16	0.11	
0.5	33.55	0.47	1.E-04	0.9	32.46	0.33	1.E-04	1.2	32.06	0.18	
0.05	36.79	0.86	1.E-05	0.1	36.16	0.81	1.E-05	0.1	35.93	0.20	

ZIKA1107 C_T, cycle number at threshold 0.13 ΔR_n

ZIKA1209 C_T, cycle number at threshold 0.03 ΔR_n

* Estimated PFU: calculated with ZIKA1209 based on PFU of MR-766

Supplementary table 2.

Nonhuman-primate's sera (collected from Mfuwe) PRNT 50 for Zika virus, Yellow Fever virus and Tick-Borne Encephalitis virus.

Sample No.	Species	PRNT 50			Interpretation
		ZIKV	YFV	TBEV	
1	<i>Papio kindae</i>	40	<10	<10	ZIKV
2	<i>Papio kindae</i>	<10	10	ND	
3	<i>Papio cynocephalus</i>	160	10	<10	ZIKV
4	<i>Papio cynocephalus</i>	<10	<10	ND	
5	<i>Papio cynocephalus</i>	40	10	<10	ZIKV
6	<i>Papio cynocephalus</i>	<10	10	ND	
7	<i>Papio cynocephalus</i>	<10	<10	ND	
8	<i>Papio cynocephalus</i>	<10	<10	ND	
10	<i>Papio cynocephalus</i>	10	<10	ND	
11	<i>Papio cynocephalus</i>	<10	<10	ND	
12	<i>Chlorocebus cynosuros</i>	10	<10	ND	
13	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
14	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
15	<i>Chlorocebus cynosuros</i>	10	<10	ND	
16	<i>Papio cynocephalus</i>	<10	<10	ND	
17	<i>Chlorocebus cynosuros</i>	160	<10	<10	ZIKV
18	<i>Papio cynocephalus</i>	10	<10	ND	
19	<i>Papio cynocephalus</i>	40	<10	<10	ZIKV
20	<i>Papio cynocephalus</i>	<10	<10	ND	
21	<i>Papio cynocephalus</i>	<10	<10	ND	
22	<i>Papio cynocephalus</i>	160	10	<10	ZIKV
23	<i>Papio cynocephalus</i>	<10	<10	ND	
24	<i>Papio cynocephalus</i>	<10	<10	ND	
25	<i>Papio cynocephalus</i>	<10	<10	ND	
26	<i>Chlorocebus cynosuros</i>	160	10	<10	ZIKV
27	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
28	<i>Chlorocebus cynosuros</i>	10	<10	ND	
29	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
30	<i>Chlorocebus cynosuros</i>	10	<10	ND	
31	<i>Chlorocebus cynosuros</i>	40	<10	<10	ZIKV
32	<i>Chlorocebus cynosuros</i>	40	<10	<10	ZIKV
33	<i>Papio cynocephalus</i>	<10	<10	ND	
34	<i>Papio cynocephalus</i>	<10	<10	ND	
35	<i>Chlorocebus cynosuros</i>	160	<10	<10	ZIKV
36	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
37	<i>Chlorocebus cynosuros</i>	<10	<10	ND	

38	<i>Papio cynocephalus</i>	<10	<10	ND	
39	<i>Papio cynocephalus</i>	<10	<10	ND	
40	<i>Chlorocebus cynosuros</i>	40	<10	<10	ZIKV
41	<i>Chlorocebus cynosuros</i>	40	<10	<10	ZIKV
43	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
44	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
45	<i>Chlorocebus cynosuros</i>	10	<10	ND	
46	<i>Chlorocebus cynosuros</i>	10	<10	ND	
47	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
48	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
49	<i>Chlorocebus cynosuros</i>	<10	<10	ND	
50	<i>Chlorocebus cynosuros</i>	160	<10	10	ZIKV

*ND: not determined; ZIKV: Zika virus; YFV: Yellow Fever virus; TBEV: Tick-Borne Encephalitis virus. The numbers of PRNT50 are represented as the greatest dilution with a positive result.

1176F

2010FSS 13025 1141: AGGTAAGATCCTACTGCTATGAGGCATCAATATCGGACATGGCTTCGGACAGCCGCTGCC 1200

BrazilPE243 1141: AGGTAAGATCCTACTGCTATGAGGCATCAATATCAGACATGGCTTCGGACAGCCGCTGCC 1200

DaKAR 41525 1140: AGGTAAGATCCTACTGCTATGAGGCATCAATATCGGACATGGCTTCGGACAGCCGTTGTC 1199

MR766 NIID 1140: AGGTAAGATCCTATTGCTACGAGGCATCGATATCGGACATGGCTTCGGACAGTCGTTGCC 1199

1209 probe

2010FSS 13025 1201: CAACACAAGGTGAAGCCTACCTTGACAAGCAATCAGACACTCAATATGTCTGCAAAAAGAA 1260

BrazilPE243 1201: CAACACAAGGTGAAGCCTACCTTGACAAGCAATCAGACACTCAATATGTCTGCAAAAAGAA 1260

DaKAR 41525 1200: CAACACAAGGTGAAGCCTACCTTGACAAGCAGTCAGACACTCAATATGTCTGCAAGAGAA 1259

MR766 NIID 1200: CAACACAAGGTGAAGCCTACCTTGACAAGCAATCAGACACTCAATATGTCTGCAAAAAGAA 1259

1308R

2010FSS 13025 1261: CGTTAGTGGACAGAGGCTGGGGAAATGGATGTGGACTTTTTGGCAAAGGGAGCCTGGTGA 1320

BrazilPE243 1261: CGTTAGTGGACAGAGGCTGGGGAAATGGATGTGGACTTTTTGGCAAAGGGAGCTGGTGA 1320

DaKAR 41525 1260: CATTGGTGGATAGAGGTTGGGGAAATGGGTGTGGACTTTTTGGCAAAGGGAGCTGGTGA 1319

MR766 NIID 1260: CATTAGTGGACAGAGGTTGGGGAAACGGTGTGGACTTTTTGGCAAAGGGAGCTGGTGA 1319

2010FSS 13025: KU955593.1 Zika virus isolate Zika virus/H.sapiens-tc/KHM/2010/FSS13025
Brazil PE243: KX197192.1 Zika virus isolate ZIKV/H.sapiens/Brazil/PE243/2015
DaKAR 41525: KU955591.1 Zika virus isolate Zika virus/A.africanus-tc/SEN/1984/41525-DAK
MR766 NIID: LC002520.1 Zika virus genomic RNA, strain: MR766-NIID
Gray arrows show regions of ZIKA1107 primers and probe described by Lanciotti *et al* (2008).