Title	Cricket tympanal organ revisited: morphology, development and possible functions of the adult-specific chitin core beneath the anterior tympanal membrane
Author(s)	Nishino, Hiroshi; Domae, Mana; Takanashi, Takuma; Okajima, Takaharu
Citation	Cell and tissue research, 377(2), 193-214 https://doi.org/10.1007/s00441-019-03000-2
Issue Date	2019-08-14
Doc URL	http://hdl.handle.net/2115/79068
Rights	This is a post-peer-review, pre-copyedit version of an article published in Cell and Tissue Research. The final authenticated version is available online at: http://dx.doi.org/10.1007/s00441-019-03000-2
Type	article (author version)
File Information	manuscript.pdf

[Instructions for use](#)

Cricket tympanal organ revisited: morphology, development, and possible functions of the adult-specific chitin core beneath the anterior tympanal membrane

Hiroshi Nishino*¹, Mana Domae¹, Takuma Takanashi², Takaharu Okajima³

¹Research Institute for Electronic Science, Hokkaido University, Sapporo 060-0812, Japan, ²Graduate School of Information Science and Technology, Hokkaido University, Sapporo, Japan, ³Department of Forest Entomology, Forestry and Forest Products Research Institute, Matsuno-sato 1, Tsukuba, Ibaraki 305-8687, Japan

The number of text pages:

The corresponding author: Hiroshi Nishino, Research Institute for Electronic Science, Hokkaido University, Sapporo 060-0812, Japan.

Tel: +81-11-706-2596

Email: nishino@es.hokudai.ac.jp

Abstract

Vertebrates and insects are phylogenetically separated by millions of years but have commonly developed tympanal membranes for efficiently converting airborne sound to mechanical oscillation in hearing. The tympanal organ of the field cricket *Gryllus bimaculatus*, spanning 200 μm , is one of the smallest auditory organs among animals. It indirectly links to two tympana in the prothoracic tibia via tracheal vesicles. The anterior tympanal membrane is smaller and thicker than the posterior tympanal membrane, and it is thought to have minor function as a sound receiver. Using differential labeling of sensory neurons/surrounding structures and three-dimensional reconstructions, we revealed that a shell-shaped chitin mass and associated tissues are hidden behind the anterior tympanal membrane. The mass, termed the epithelial core, is progressively enlarged by discharge of cylindrical chitin from epithelial cells that start to aggregate immediately after the final molt, and it reaches a plateau in size after six days. The core, bridging between the anterior tracheal vesicle and the fluid-filled chamber containing sensory neurons, is supported by a taut membrane, suggesting the possibility that anterior displacements of the anterior tracheal vesicle are converted into fluid motion via a lever action of the core. The epithelial core did not exist in tympanal organ homologs of meso- and metathoracic legs or of nymphal legs. Taken together, the findings suggest that the epithelial core, a potential functional homolog to mammalian ossicles, underlies fine sound frequency discrimination required for adult-specific sound communications.

Key words: epithelial cells; tracheal vesicles; insect; cricket; chitin; tonotopic organization

Abbreviations: AtC: attachment cell; AN: auditory nerve; BM: basement membrane; ATM: anterior tympanal membrane; ATV: anterior tracheal vesicle; CD: ciliary dilation; CM: covering membrane; DD: dendritic dilation; DG: distal group neurons; EC: epithelial cell; ECL: epithelial cell layer; LAC: large accessory cell; La: labyrinth; Oli: olivarius organ; PG: proximal group neurons; PTM: posterior tympanal membrane; PTV:

- 1 posterior tracheal vesicle; SC: scolopale cell; ScC: scolopale cap; ScR:
- 2 scolopale rod; SGO: subgenual organ.
- 3

1 **Introduction**

2
3 In order to detect airborne sound in terrestrial environments for acoustical
4 communication, both vertebrates and insects have evolved tympanated
5 hearing organs. Since both animal groups are phylogenetically distant, this
6 is thought to be a typical example of convergent evolution (Hoy and Robert
7 1996). The emergence of tympanal membranes in prothoracic tibiae of
8 ensiferan insects dates back at least to the Eocene era, and the ensiferan
9 hearing organ is therefore one of the “oldest” tympanate ears that have
10 adapted to the terrestrial environment for more than 50 million years (Rust
11 et al., 1999; Plotnick and Smith 2012; Strauß and Lakes-Harlan 2014).
12 They are backed by an air-filled space or cavity and innervated by a
13 chordotonal sensory organ that consists of bipolar sensory neurons (Field
14 and Matheson 1999; Yack 2004).

15 Several lines of study have shown similarity between vertebrate and
16 insect hearing organs not only in external morphology but also at the
17 molecular level. As an example, mechanosensory channels and functional
18 molecules expressed in auditory sensory cells are largely shared between
19 the fruit fly chordotonal organ in the antennal base (i.e., Johnston’s organ)
20 and mouse auditory hair cells (Senthilan et al. 2012). For example, 217 of
21 274 Johnston’s-organ-associated genes have mouse or human orthologs,
22 attributable to common ancestors (Senthilan et al. 2012). Active
23 amplification of sound found in vertebrate hair cells has also been found in
24 tympanal organs of hemimetabolous insects (Kössel et al. 2008; Mhatre
25 and Robert 2013) and in Johnston’s organs of holometabolous insects (e.g.,
26 Robert and Göpfert 2002).

27 The similarity at the sensory neuronal level, however, might not be
28 surprising given that the developmental and evolutionary origins of
29 tympanal organs are derived from proprioceptive chordotonal organs that
30 retain features of ancestral mechanosensory organs (Boyan, 1993; Hoy and
31 Robert 1996; Niwa et al. 2004). Insects have independently evolved
32 tympanate hearing organs at least seventeen times, resulting in a large
33 diversity in their bodily locations, sizes, and numbers of sensory neurons
34 (Yager 1999). Thus, one might expect that “true” convergent evolution
35 rather lies in sound-transmitting apparatus adopted between animals, such as
36 humans and insects.

1 A recent study revealed that sound transmission in humans and that in
2 tree crickets share remarkable similarities in three fundamental processes
3 (Montealegre-Z et al. 2012): 1) sound detection by a tympanal
4 membrane(s), 2) converting a large displacement of vibration to a small but
5 powerful pressure by a lever system, which functionally parallels to
6 ossicles in mammals. 3) converting a mechanical pressure to fluid motion
7 that underlies the frequency discrimination of sensory neurons.

8 The tympanal organ of field crickets *Gryllus* and *Teleogryllus* (fami-
9 ly: Gryllidae; subfamily: Gryllinae) has been one of the most intensively
10 studied hearing organs in insects (Huber et al. 1989). It is the smallest in
11 animals known so far, spanning only 200 μm despite its coverage of
12 broad-band sound frequencies from 4 kHz to greater than 42 kHz (Esch et
13 al. 1980; Hutchings and Lewis 1981). Seventy sensory cells are
14 tonotopically arranged to detect higher frequency sound by more distally
15 located neurons (Oldfield et al. 1986). Field crickets have two tympanal
16 membranes of unequal sizes, in contrast to those in tettigoniids and tree
17 crickets that possess similar sizes of tympana in the proximal tibia (Ball
18 and Field 1981; Mhatre et al. 2009; Montealegre-Z et al. 2012).

19 The functions of the two tympanal membranes in field crickets have
20 been evaluated by their ablation, immobilization, and measurements of
21 mechanical properties. The posterior tympanal membrane (PTM, see Fig.
22 1a,b) is thin and six-times larger than the anterior tympanal membrane
23 (ATM, see Fig. 1c; Young and Ball 1974a). The PTM vibrates with about
24 ten-times larger amplitudes than those of the ATM (Johnstone et al. 1970).
25 Another study showed that the ATM vibrates with 20 dB larger amplitude
26 than neighboring cuticle and also about 20 dB below the vibration
27 amplitude of the PTM (Larsen, 1987). Covering the PTM with Vaseline
28 (Paton et al. 1977) or with wax (Huber et al. 1984) and detaching the PTM
29 from the leg cuticle (Nocke 1972) all significantly decreased the sensitivity
30 of auditory sensory neurons by 15-30 dB. In accordance with these findings,
31 covering the external surface of the ipsilateral PTM with water abolished
32 neural response in an auditory interneuron (Kleindienst et al., 1983). From
33 these findings, the PTM is thought to be the primary sound receiver, and
34 the motion of the PTM is crucial for hearing (Kleindienst et al. 1983).

35 In contrast, the function of the ATM has remained controversial. For
36 example, occluding the ATM with petrolatum did not affect the sensitivity

1 of auditory interneurons (Hill 1974), but removal of the ATM resulted in
2 cessation of activity of sensory neurons that tune to sound above 12 kHz
3 (Nocke 1972). Huber et al. (1984) demonstrated that crickets with their
4 ATM and PTM blocked have auditory thresholds nearly 30 dB higher than
5 those in crickets in which only the PTM is blocked, suggesting a
6 complementary function of the ATM.

7 In principle, the field cricket tympanal organ acts as a four-input system
8 in which sound reaches the ipsilateral and contralateral internal walls of
9 acoustic spiracles and the ipsilateral and contralateral surfaces of the
10 posterior tympanal membranes, although the effect of sound input from the
11 contralateral regions is small (Larsen and Michelsen 1978; Michelsen et al.
12 1994), supporting its function as a pressure-difference receiver. However,
13 how sound input to the PTM is eventually converted to mechanical
14 deformation of sensory cilia is still enigmatic. So far, cellular and
15 subcellular features of tympanal organs in field crickets have been studied
16 mainly in sectioned materials (Michel 1974; Young and Ball 1974a).

17 The aim of this study was to provide a comprehensive view of sound
18 transmission pathways including the functionally enigmatic ATM in this
19 smallest insect ear. It is scarcely understood how tympanal organ
20 morphologies affect the frequency tuning properties of sensory neurons.
21 To tackle this problem, we have developed low-invasive dissection
22 techniques that leave the connections from tympanal membranes to sensory
23 neurons intact (Nishino and Sakai 1997) but permit observations using a
24 confocal laser scanning microscope. The three-dimensional reconstructions
25 of tympanal organs allowed us to scrutinize fine anatomical links among
26 the two tympanal membranes, tracheal vesicles and the main body of the
27 tympanal organ. The results provide new insights into the function and
28 development of the ATM and related structures that should be useful for
29 future modelizing of the tympanal organ in biomechanics.

Materials and Methods

Animals

Male and female nymphal and adult field crickets, *Gryllus bimaculatus* DeGeer, at 0–15 days after the imaginal molt in a laboratory colony were used. They were maintained in a constant dark and light cycle (L:D = 12:12) at a temperature of 26°C. Over 220 individuals were used for the experiments, though the data presented here were derived from a much smaller numbers of animals. Three male and three female adult field crickets, *Teleogryllus occipitalis*, collected at Amami Islands and kept in the laboratory for 12 days were also used for inspecting the tympanal organ structures from a comparative viewpoint.

Retrograde labeling of sensory neurons

The prothoracic tympanal organ and its homologs in the meso- and metathoracic tibiae, termed tracheal organs (Young and Ball 1974b; Eibl 1978), were investigated. Animals were briefly anesthetized by carbon dioxide. A leg of interest was amputated at the trochanter and the distal section was embedded in a plastic wax with the anterior side up on a beeswax plate. The main leg nerve 5 (N5B2) and the anterior sensory nerve running along the main trachea (N5B1) were exposed by removing the anterior cuticle and were immersed in a drop of cricket saline (Nishino and Sakai 1997). The two nerves were cut with microscissors (Vannas Scissors, 501778, WPI, FL, USA) and the distal cut-ends were picked up by an electrolytically tapered tungsten rod (ID: 200 µm) with the tip slightly bent and were placed in the tip of a tapered glass electrode filled with microruby (dextran, tetramethylrhodamine and biotin, 3000 MW, ThermoFisher Scientific). Care was taken not to stretch the nerve when handling. The preparation was left in a humidified chamber at 5 °C for 24 hours. The anterior region of the epicuticle except for the ATM in the proximal tibia was carefully stripped off with fine microscissors to leave the translucent hypodermis intact (Fig. S1). The removal of the epicuticle allowed 1) the interior structure to be left nearly intact, 2) promotion of permeabilization of the fixative and fluorescent dye for counterstaining in the proximal tibia

rich with air-filled space (see below) and 3) scanning through the whole tympanal organ along the Z-axis without interference under a confocal microscope. The leg was pre-fixed in 4 % neutral paraformaldehyde solution for 15 min at room temperature.

Counterstaining and immunolabeling

Counterstaining with a fluorescent dye was used to visualize the anatomical link between the tympanal organ sensory neurons and surrounding structures such as tympanal membranes and tracheal vesicles. For differential staining, the specimens in which sensory nerves had been labeled with microruby were immersed in 0.5% Lucifer Yellow CH (Sigma Aldrich) in cricket saline (Nishino and Sakai 1997) for 40 min after pre-fixation (Nishino et al. 2016). To visualize F-actin filaments, the fixed specimens were transferred to 2.5% phalloidin solution (Acti-stain™ 488 Fluorescent Phalloidin, Cytoskeleton, Inc) for 1 h at room temperature before dehydration (Wolfrum 1990). After careful removal of the anterior tympanal membrane, chitinase from *Streptomyces* (C6137, Sigma) was applied to the epithelial core and surrounding tissues at 37°C for 3 h in phosphate buffered saline set at pH 6.0 to confirm whether these are made up from chitin. Chitinase is an extracellular enzyme complex that degrades chitin and has a molecular mass of approximately 30 kDa. Chitin is degraded to N-acetyl-D-glucosamine in two enzymatic reactions. Firstly, chitobiose units are removed from chitin by chitodextrinase-chitinase. The second reaction involves N-acetyl-glucosaminidase-chitobiase, which cleaves the disaccharide to its monomer subunits (that comprise N-acetyl-D-glucosamine). All of the specimens processed by these treatments were thoroughly rinsed with cricket saline and post-fixed in 4% neutral paraformaldehyde solution for 1-2 h at 5 °C, dehydrated in an ethanol series, and cleared in methyl salicylate.

Microscopic observation of fixed specimens and data processing

The cleared specimens stained with fluorescent dyes were observed under a confocal laser scanning microscope (LSM 5 Pascal, Zeiss) and a light microscope (Imager Z1, Zeiss). Specimens differentially labeled with

Lucifer yellow and microruby were visualized with an argon laser with a 505–530-nm bandpass filter and a helium-neon laser with a longpass filter (>560 nm), respectively. Optical sections were made at a resolution of 1024 x 1024 with 0.6–2.0- μ m intervals. Plan-Apochromat x10 0.8-NA and x20 0.8-NA (dry objectives) were used for observation of the entire tympanal organ complex, and Plan-Neofluar x40 1.3-NA (oil immersion objective) was used for observation of components of the tympanal organ. Observations were firstly made on wholemount preparations. Then transverse sections of the tibia at 100~200 μ m in thicknesses were made manually by using a twin blade razor (0.1 mm in thickness, Feather, Japan) after the specimens had been rehydrated in 70% ethanol.

The optical sections were converted to TIFF-formatted files using the software LSM Image Browser (Carl Zeiss, Jena, Germany). TIFF images were processed with image processing software (Amira ver. 3.1, Visage imaging, Berlin, Germany). The sensory neurons and surrounding structures were manually outlined in each optical section for subsequent three-dimensional representations using a volume rendering function. Three-dimensional surface models of internal structures of the legs were obtained by surface rendering.

All photographic images were processed using Adobe PhotoShop Elements 6 (San Jose, CA). In figures, two to twenty consecutive optical sections at different focal planes were overlaid as aligned stacks and flattened to one plane.

Terminology

We referred to Young and Ball (1974a) for the naming and terminology of different kinds of cells associated with the tympanal organ, to Ribeiro and Brehelin (2006) for naming and classification of hemocytes, to Nishino and Sakai (1997) for terminology of nerves, to Young and Ball (1974b) and Eibl (1978) for the tracheal organ (counterparts of the tympanal organ in the meso- and metathoracic tibiae), and to Young and Ball (1974a) for orientations of the tympanal organ and tracheal organ (see Fig. 1a). The numbers of sample sizes are shown in parentheses in Results, wherever needed.

Results

Three-dimensional structure of the tympanal organ

The tympanal organ in field crickets is an assembly of nine distinct apparata as shown in the two-dimensional optical stacks (Figs. 1d-h, 2a-f, ESM1) and three-dimensional reconstructions (Figs. 1i, j, 2 g, h) of the proximal tibia of *G. bimaculatus* (Fig. 1a-c). These apparata (Fig. 1i, j) include 1) the posterior tympanal membrane (PTM, yellow), 2) the posterior tracheal vesicle (PTV, dark blue), 3) the anterior tracheal vesicle (ATV, light blue), 4) a tent-like cellular mass (red), 5) a thin covering membrane interconnecting the lateral side of the mass and tracheal vesicles (CM, Fig. 1f), 6) sensory neurons (magenta) on the ATV, innervated by the auditory nerve (AN, Fig. 1d) diverged from N5B1, 7) the anterior tympanal membrane (ATM, light yellow), 8) the core structure hidden beneath the ATM. It attaches postero-distally to the covering membrane (green; Fig. 2 f, g, l), and 9) lipid-synthesizing organs that clog hemolymph channels proximo-distally to the cellular mass, termed the olivarius organs in primitive tettigoniids, wetas (Oli, Fig. 1d, ESM1; Lomas et al. 2013).

The PTM, which is the primary sound input site, has a hinged structure in its central to distal region (Fig. 1g; see also ESM7), reminiscent of tettigoniid tympanal membranes (Bangert et al. 1998). The inflated PTV, tightly apposed to the PTM, is located parallel to the smaller ATV (Fig. 1j, ESM2). The PTV and ATV are interconnected via two apertures (Fig. 1h, j, ESM2; Eibl 1978).

The tent-like mass is composed of two types of cells: 1) attachment cells (AtCs), each receiving insertion of a sensory neuronal cilia, and 2) large accessory cells (LACs) that primarily contribute to building up the mass (Fig. 2a; Young and Ball 1974a). The AtCs and LACs are readily identifiable by the shape of their nuclei, the former being more ellipsoidal (Figs. 2a, 3a; ESM1). All of the sensory neurons extend dendrites into single attachment cells at one-to-one correspondence via scolopale caps (SCs, Figs. 2a, c, e, 3a, ESM1). There was no indication that dendrites of proximal neurons of the distal group extended directly to modified hypodermal cells that attach to the hypodermis, as suggested in *T. commodus* (Young and Ball 1974a). The cellular mass is attached to the

dorsal cuticular ceiling via a bundle of approximately 30 ligaments originating from hypodermal cells (inset, Fig. 1e). The attachment site is thinner than surrounding cuticles (arrow, Fig. 1e). The bundle of ligaments radiate ventrally and cover the whole cellular mass (Fig. 1e; ESM1).

The covering membrane (CM) is thin, approximately 0.5 μm in thicknesses (Fig. 1f). The membrane is secreted by unidentified cells with diameters of 5-7 μm that line the inside of the membrane (inset, Fig. 1f). The membrane descends along the lateral edge of the cellular mass to attach to the anterior region of the dorsal membrane of the PTV and the anterior edge of the dorsal membrane of the ATV (ESM1). The plasma serum is often coagulated inside the covering membrane through a fixation process (e.g., see ESM6), suggesting that the sensory dendrites are immersed in a lipidic fluid (Huber et al. 1989). In all of the fixed specimens with interior structures left intact, no hemocytes were observed around sensory dendrites (Figs. 1f; 2a; ESM1), though aggregation of a few hemocytes was often observed in the proximo-posterior region of the fluid chamber (Fig. 2f; ESM1).

The sensory neurons are classified into a proximal group (PG, Fig. 2a-d, g) and a distal group (DG, Fig. 2b, e, f, h) based on whether dendrites are attached to the cellular mass externally or immersed in fluid. Based on this criterion, only proximal group A defined by Young and Ball (1974a) was categorized into the proximal group and other neurons were categorized into the distal group. Neurons in the proximal group, a group of neurons that is a possible functional homolog of the intermediate organ in wetas (Ball and Field 1981; Nishino and Field 2003; Strauß et al. 2017), extend dendrites along the antero-dorsal to postero-ventral region of the cellular mass (Fig. 2d). Proximal group neurons attach to the AtCs at their ciliated region of dendrites (Fig. 3b), where mechanosensory channels are distributed (review: Li et al. 2018).

Neurons in the distal group are arranged proximo-distally on the dorsal membrane of the ATV (Fig. 2b, e, f), of which thicknesses are approximately 1-2 μm (Fig. 2e, f; ESM1). As shown in the dorsal view, somata and dendrites of the distal group of neurons are arranged diagonally at about 45° against the longitudinal axis of the tibia, in which the more distal cells occupy the more anterior region (Fig. 2c, h). This arrangement allows maximal use of the space in the limited dimension of the tympanal

organ. Dendrites of the distal group of neurons bridge between the dorsal membrane of the ATV and the bottom of the cellular mass (Fig. 2e-g, j). The dendrites of the distal 20 neurons of the distal group extend to AtCs elongated from the bottom of the cellular mass (Fig. 2a; ESM1). Their dendritic arrangements and orientations are topographically complex. The array of sensory dendrites winds like an S shape (broken lines in Fig. 2i, k), in which dendrites of more proximal neurons extend into the attachment cells at smaller angles due to the narrower spaces between the ATV and the cellular mass in the proximal direction (Fig. 2e, f; ESM1). This is in reverse sequence for the tettigoniid *crista acustica*, in which distances between the attachment cells and the dorsal membrane gradually become larger in the proximal direction (Hummel et al. 2017). Dendrites located in the middle region of the distal group neurons tended to be denser than those in the proximal and distal ends (Fig. 2c, i).

Being consistent with our previous findings (Nishino et al. 2016), phalloidin stainings showed that scolopale rods of sensory neurons are rich in actin filaments (Fig. 3b-d), and peripheral regions of large accessory cells showed weak immunoreactivity to actin filaments (Fig. 3a). Subcellular structures, such as the scolopale cap (ScC), ciliary dilation (CD), and dendritic dilation (DD), typical of scolopidia participating in auditory/vibratory detection (Field and Matheson 1998; Yack 2004), were detectable in dendritic cilia (Fig. 3b, d).

Structures of meso- and metathoracic tracheal organs

Counterparts of the prothoracic tympanal organ, termed tracheal organs in the meso and metathoracic tibiae (Young and Ball 1974b; Eibl et al. 1978), resemble the premature tympanal organ in nymphs (Ball and Young 1974). Tracheal organs are not equipped with tympanal membranes or core structures (Fig. 4). As shown by Young and Ball (1974b) and by Eibl (1978), only sensory neurons corresponding to proximal group A are present (Fig. 4c, d). The non-inflated tracheae corresponding to the ATV and PTV run in parallel but are not interconnected by two apertures (Fig. 4a, b). Olivarius organs adjacent to the cellular mass are present in meso- and metathoracic tracheal organs (Fig. 4a, b). Closer observation revealed that unlike the tympanal organs in phalate adults (Fig. 4e-g) and 6th instar

nymphs (Fig. 4h), which possess a well-developed covering membrane, there were no covering membranes in tracheal organs (Fig. 4c, d). Instead, the cellular mass was suspended to the anterior cuticular wall via fine fibers (Fig. 4c; ESM3), resembling those seen in the *crista acustica* homologs in the atympanate ground weta *Hemiandrus pallitarsis* (Strauß et al. 2017). Since hemolymph passage is permitted beneath the cellular mass, a number of hemocytes were detectable beneath the cellular mass of meso and matathoracic tracheal organs (Fig. 4c, d; ESM3).

Epithelial core and associated tissues in the mature adult

The epithelial core, previously termed the “tracheal body” (Friedman 1972a, b; Huber et al. 1989), exists in *G. bimaculatus* and *T. occipitalis* but only in mature adults (Figs. 2g and 5). Mature cores are shell-shaped and are approximately 140 µm x 90 µm square and 60 µm in thickness in *G. bimaculatus* (Fig. 5a-e) and approximately 110 µm x 60 µm square and 25 µm in thickness in *T. occipitalis* (Fig. 5f-h). The designation of tracheal body is, however, inappropriate because the core is neither part of the trachea nor a product secreted by tracheoblast cells (Wigglesworth 1954; Affolter and Causinius 2008). Instead, the core is built up by external scaffolding of epithelial cells (also referred to as epidermal cells, Wigglesworth 1959) to the dorso-anterior edge of the ATV. Since the core is adhered to a mechanically unstable position and protrudes anteriorly like a lever (Fig. 6i-n), the core needs to be supported by surrounding tissues for keeping its original position. As shown in three-dimensional reconstructions of the core region (Fig. 5) and transverse sections (Fig. 6), the following six features were detectable beneath the ATM.

- 1) The basement membrane (BM) that extends from the dorsal rim of the ATM approaches the core and attaches nearly perpendicularly to the fringe of the antero-dorsal surface of the core (Fig. 5a, b, d; ESMs4, 5), the site being slightly more ventral than that of the covering membrane on the opposite side (ESMs4, 5). The BM is composed of a layer of single cells and is 4-5 µm in thickness (ESM5). The width of the membrane becomes smaller in the distal direction, as shown in the dorsal view (Fig. 5b). The antero-proximal region of the BM is firmly attached to the ATM (ESM5).

- 2) The attachment site of the BM to the core is wedged by epithelial cell layers (ECLs, Fig. 6l-n) that run in parallel along the anteriormost surface of the core (ESMs4, 5). The ECLs are embedded in a sheet of whitish connective tissues (see Fig. 9a). The thicknesses of ECLs are approximately 7 μm near the core and 20 μm in the ATM rim area (ESM5). The complex of BM/ECLs is taut and not flaccid, literally like a tympanal membrane.
- 3) The space between the complex of BM/ECLs and the ATM is a chamber immersed in hemolymph that contains a few hemocytes and epithelial cells (see Fig. 8e, f; ESM5).
- 4) There are labyrinth-like frameworks that loosely support the proximo-ventral side of the core (ESMs4, 5). A bundle of fibers mechanically bridges the ECLs and the anterior wall of the ATV/ventral region of the core (Figs. 5c, 6b-f). The cavities made by labyrinth are immersed in hemolymph that contains numerous plasmatocytes (ESMs4, 5).
- 5) The posterior surface of the core is attached to the fluid-filled chamber at its distal two thirds (Figs. 2l, 6j, k, m, n; ESMs6, 7). Contact of the core with the fluid is not direct but via a thin covering membrane. The covering membrane is often detached from the core in the proximal half region in *T. occipitalis* (e.g., Fig. 6i; ESM6), while it attaches to the central region of the proximal edge of the core in *G. bimaculatus* (Fig. 8f; ESM5).
- 6) As in the PTV supported ventrally by the cuticular ledge (Fig. 6a-h), the ATV on which the main body of the tympanal organ lies is supported ventrally by a thick cuticular ledge that protrudes posteriorly from the rim of the ATM (Fig. 6c-f; ESM7; Michel 1974; Eibl 1978).

The epithelial core is hard, but surrounding tissues are elastic

The epithelial core itself is a shell-shaped, hard, and translucent mass (Fig. 7a). The core was degraded by placing a small amount of chitinase powder onto it and subsequent incubation at 37 °C for 3 h (Fig. 7b), suggesting that the core is made up by N-acetylglucosamine, which is the primary monomeric unit of the polymer chitin. The ligaments connecting the

cellular mass to the hypodermis were also degraded by chitinase, resulting in collapse of the cellular mass onto the ATV (Fig. 7d). The covering membrane and tent-like cellular mass maintained their structures after application of the chitinase (Fig. 7c), suggesting that they have a composition other than chitin.

The elasticity of the basement membrane and that of epithelial cell layers were assessed in a fresh prothoracic leg with the epicuticle of the ATM and the surrounding epicuticle carefully removed (ESM8). This treatment leaves the structures underlying the ATM intact. Pressure applied to the exocuticle of the distal femur caused a volumetric increase in the ATV and PTV because tracheal systems are closed in the tibia and, in turn, air in the extracellular space surrounding the main body of the tympanal organ pushes the basement membrane away in an outward (anterior) direction (n=8, ESM8), suggesting that the melanized basement membrane is elastic but sufficiently hard to push away the epithelial cell layers ventrally. The movement is larger in the proximal region of the basement membrane than in the distal region (ESM8). Such movements were never observed in intact specimens, suggesting that the movement of the basement membrane is normally hindered by its suspension to the ATM in the proximal region.

Progressive development of the epithelial core and associated tissues

In the nymphal and pharate adult of *G. bimaculatus*, the covering membrane is attached to the interior wall of the anterior cuticle (Fig. 4e, f). Therefore, the space between the cellular mass and the anterior cuticle is narrow (ca. 60 μm). Moreover, the ATV and the PTV are not directly attached but separated by tracheoblast cells (arrow, Fig. 4f; Wigglesworth 1954).

The core and labyrinth are concurrently formed between the ATM and ATV. Immediately after the final molt, epithelial cells with diameters of 4-5 μm aggregate beneath the ATM, forming cell layers (Figs. 8a, 9a). The layers are especially thick in the ventro-distal region beneath the ATM (ESMs4, 5). The epithelial cells produce thin filopodium-like processes laterally to connect themselves to each other (Fig. 8b). Some of the epithelial cells invaginate by elongation of thick filopodia (Fig. 8c),

1 allowing their migration toward the future core region. The epithelial cells
2 remaining just beneath the ATM secrete chitin outward (anteriorly), which
3 contributes to thickening of the ATM from $8.13 \pm 0.58 \mu\text{m}$ immediately
4 after the final molt ($n = 3$, Fig. 8c) to $19.78 \pm 0.79 \mu\text{m}$ in a mature adult
5 (measured by Schneider et al., 2017).

6 A sheet of single cells with diameters of 6-8 μm , probably plasmatocytes,
7 is static from the final molt and becomes the basement membrane (Fig. 8d,
8 e, Wigglesworth 1973). The intercellular space appears to be filled with
9 connective tissues discharged by plasmatocytes (Wigglesworth 1956). The
10 basement membrane is gradually melanized over a period of one week (see
11 Fig. 9a, b). The small space between the basement membrane and the ATM
12 is maintained throughout the adult stage (Fig. 8e, f, ESM5;
13 Friedman1972a).

14 In contrast, plasmatocytes that aggregated ventral to the future core
15 region are active. The plasmatocytes phagocytize filopodium processes of
16 epithelial cells and shape them into frameworks in the labyrinth, except for
17 the anterior-most epithelial cell layers (Fig. 8g; ESMs4, 5). Debris of the
18 filopodia of epithelial cells is continually removed away by phagocytosis
19 by plasmatocytes that stay beneath the ATM through adult life (ESMs4, 5).

20 The core augmentation occurs progressively. At the end of day 1,
21 invaginated epithelial cells are scaffolded onto the ATV and start to secrete
22 chitin inward (Fig. 9c). This is the beginning of core formation. On day 2,
23 individual epithelial cells start to discharge chitin material (Fig. 8h-j).
24 Three-dimensional reconstructions of tractable chitin materials in the core
25 (Fig. 8i, j) showed their cylindrical appearances, corresponding to those
26 found by Friedman (1972b) in *G. assimilis*. The chitin cylinders
27 collectively form a small fan-shaped core (Figs. 9d). From day 3,
28 concomitant with inward recruitment of more epithelial cells, the core
29 grows accretionally (Fig. 9e-h) and enlarges toward dorso-anteriorly until
30 the size reaches a plateau on day 6 (Fig. 9i). The enlarged core gradually
31 separates the ATV from the ATM, making the cellular mass 120 μm away
32 from the ATM in the mature adult (Fig. 6l, ESM5).

33 The epithelial cell nuclei lining the posterior surface of the core are
34 orderly arranged like a fan (Fig. 8k, l). The early epithelial cells scaffolded
35 onto the ATV undergo apoptosis and become themselves part of the core.
36 Scaffolding of epithelial cells, chitin secretion, and attachment of new

1 epithelial cells to the cell bodies of old cells should be precisely regulated
2 spatio-temporally so that accretionary growth of the shell-shaped mass
3 occurs (Fig. 8m).

4 5 **Formation of the epithelial core is independent of innervation of** 6 **peripheral nerves**

7
8 The formation of the epithelial core is a robust process that is unaffected by
9 denervation after the imaginal molt. For example, decerebration (including
10 the corpus allatum) and denervation of N5B1 (innervating the tympanal
11 organ) and N5B2 (innervating part of the SGO) in the proximal tibia
12 immediately after the imaginal molt resulted in normal core formation and
13 associated structures (n = 5). Cutting all of the peripheral nerves
14 unilaterally close to the prothoracic ganglion immediately after the final
15 molt also resulted in normal formation of the core and associated structures.
16 Thus, the core augmentation proceeds without control of sensory neurons
17 or efferent neurons in the adult stage. Keeping animals in constant dark
18 from the final instar to adult stage also resulted in normal formation of the
19 core (n = 7), suggesting that core formation is not affected by light
20 impinging through the translucent ATM or by light-entrained circadian
21 rhythms.

22 23 **Discussion**

24 25 **Overview**

26
27 Studies on tympanal organs in field crickets have a long history dating back
28 to Schwabe's anatomical study (1906) and Autrum's physiological study
29 (1941). In contrast to detailed descriptions about frequency tuning of
30 auditory sensory neurons (Oldfield, 1986; Pollack and Imaizumi 1999;
31 Imaizumi and Pollack 1999, 2005), the key structure transmitting the
32 mechanical energy to the sensory neurons has remained enigmatic. The
33 epithelial core, first discovered by Michel (1974), was only briefly referred
34 to in a monograph (Huber et al., 1989). The tissue embedding the core,
35 termed the suspensorium (Schwabe 1906; Michel 1974), has been thought
36 to be a mechanical damper (Larsen and Michelsen 1978).

1 Our study revealed, for the first time, that the ATM has a bilayering
2 structure: a thick outer membrane and an underlying thin membrane
3 complex composed of a basement membrane and epithelial cell layers. The
4 epithelial core, bridging between the ATV and the fluid-filled sensory
5 chamber, is supported by a taut membrane complex. There are no
6 energy-absorbing structures around the core, suggesting more active
7 functions other than a passive mechanical damper. On the other hand, the
8 outer membrane of the ATM is 20 μm in thickness and has no mechanical
9 links with the internal structure. Thus it is reasonable that immobilizing the
10 outer membrane had no effect on the physiological properties of sensory
11 neurons in *T. commodus* (Hill 1974) or on acoustic orientation in *T.*
12 *oceanicus* (Bailey and Thomson 1977). All sound transmission apparatus
13 including the trachea, the epithelial core, and tympanal membranes that are
14 made up by chitin or materials polymerized with chitin will be important
15 for impedance matching on sound transmission between their junctions.

16 The meso- and metathoracic counterparts of the prothoracic tympanal
17 organ, tracheal organs, retain evolutionary conserved features (Yager 1999;
18 Strauß and Lakes-Harlan 2014). These organs lack an epithelial core,
19 covering membrane, and distal group neurons (that tune to high-frequency
20 sounds), suggesting an intimate functional link of the core with
21 high-frequency sound detection in the fluid-filled environment.

22 The core development is precisely regulated but independent of
23 peripheral neuronal control. The hearing system in the cricket, therefore,
24 offers not only a model for understanding a miniature ear achieving a broad
25 audible range but also a model for dynamic cell migration based on
26 self-assembly (Fig. 8m), nowadays extensively studied in developmental
27 biology (e.g., Ma et al. 2017; Matsubaysahi et al. 2017).

28 Our histological procedures permitted observation in almost intact
29 specimens. Generally, tissues connected to the exoskeleton cause
30 inadvertent deformation of the structure through formaldehyde fixation.
31 Moreover, embedding agents such as epons hardly penetrate into the cuticle,
32 resulting in deformation or even loss of internal tissues in sectioned
33 materials. Young and Ball (1974a) failed to find the epithelial core in *T.*
34 *commodus*, presumably due to the loss of the core during sectioning.
35 Removal of the exocuticle and two-step fixation allowed wholemount
36 confocal observations of semi-intact specimens at sufficiently high

1 resolutions (Fig. 3), which are beyond the resolutions of micro CT (> 3
2 μm).

3 4 **Possible sound transmission pathways**

5
6 Several lines of studies in field crickets have shown that the PTM is crucial
7 for hearing (Johnstone et al. 1970; Paton et al. 1977; Huber et al. 1984) and
8 that it acts as a pressure-difference receiver that compares external sound
9 impinging onto the PTM with internal sound impinging to the tracheal
10 opening in the prothorax and transmitting to the internal vesicle (Larsen
11 and Michelsen 1978; Larsen 1987; Michelsen et al. 1994). The PTM is
12 broadly tuned with an optimum well above the calling song frequency
13 (Lankheet et al. 2017; Schneider et al. 2017). Its tuning is not sharp enough
14 to account for the tuning of individual sensory neurons, which is generally
15 more selective (Oldfield et al. 1986; Hutching and Lewis 1981; Imaizumi
16 and Pollack 1999). These findings suggested the existence of a second filter
17 somewhere in the tibia (Larsen et al. 1989). In this context, our findings
18 provide insights into how sound energy is mapped in a
19 frequency-dependent manner for individual sensory neurons.

20 21 *Structures surrounding the core*

22 In general, sensory neurons beneath the cellular mass, i.e., the distal group
23 of neurons, are mechanically isolated from structures other than tracheal
24 vesicles. The regions proximal and distal to the tympanal organ are
25 protected by olivarius organs, the thick ATM, and cuticular ledges. These
26 structures will prevent the disturbance of hemolymph passage and the
27 disturbance of muscular movements to the tympanal organ. Moreover, the
28 ATV is largely surrounded by incompressible fluid, that is, the fluid-filled
29 sensory chamber, the ventral hemolymph channel, and labyrinth, except for
30 the antero-distal edge to which the core is attached. We assume that the
31 labyrinth has two functions: 1) frameworks, made from filopodia of
32 epithelial cells, mechanically support the ventral side of the core loosely. 2)
33 the cavities in the labyrinth encloses hemolymph, creating a static
34 environment immersed in fluid.

35 In such a protected environment, the sound impinging to the PTM
36 could be transmitted effectively between solid media, from the large PTV

1 to the small ATV, and to the core, all of which are appositioned in parallel
2 (Fig. 10c, d). Two sound transmission pathways are conceivable from the
3 anatomy of these pathways. First, the thin region of the PTV can resonate
4 with the PTM vibration and transmit sound energy to the ATV via its
5 junction. Since the thin dorsal membrane of the PTV is attached directly to
6 the fluid-filled chamber, longitudinal waves of sound components may
7 enter the fluid and affect the sensitivity of sensory neurons (Greenfield,
8 2002), although there are no specialized structures that effectively
9 compress the fluid laterally in the junction between the PTV and ATV (Fig.
10 10c, d). Second, compressional waves of sounds can lead to instantaneous
11 displacements of the ATV. Airflows through the two apertures potentially
12 result in a volumetric increase of the ATV and push the interior wall
13 outward. Although it is not known whether either of these or both
14 contribute to the stimulation of sensory neurons, instantaneous
15 displacement of the ATV would be more effective for core displacement.

16
17 *Core - is it designed to convert mechanical displacement to fluid motion?*

18 Neurons of the distal group are immersed in hemocyte-free fluid sealed by
19 the covering membrane. The dendrites of these neurons are arranged with
20 dendrites of distal neurons oriented progressively more perpendicularly
21 against the dorsal membrane and occupying the wider fluid space between
22 the dorsal membrane and the cellular mass. Neurophysiological studies
23 have shown that sensory neurons are tonotopically organized so that more
24 distal neurons in the distal group tune to sound with higher frequencies
25 (Oldfield et al. 1986). The fluid-filled chamber is attached to the core in the
26 distal region via the thin covering membrane (Fig. 10a, b). From these
27 findings, it is tempting to speculate that the core is ideally situated for
28 converting its mechanical displacement to fluid motion, which is the direct
29 source of mechanical deformation of the cilia of the neurons in the distal
30 group.

31 The core has topographically different connectivities with surrounding
32 tissues (Fig. 10a-d). The proximal region of the core, inflated in *G.*
33 *bimaculatus*, is supported anteriorly by complex of the elastic basement
34 membrane/epithelial cell layers, but only loosely supported ventrally by
35 labyrinth-like frameworks (Fig. 10b, c). The distal region of the core is
36 firmly supported by thick epithelial cell layers (dark blue, Fig. 10d). Along

1 the dorso-ventral axis, the ventral side of the core is adhered to the ATV in
2 a large area (yellow line, Fig. 10a, b), but the attachment site of the core to
3 the covering membrane is limited to the distal two thirds (Fig. 10b).

4 Given these features, we propose that a class 1 lever action operates on
5 the core (Fig. 10e, f). The membrane complex is taut like a tympanal
6 membrane and is backed by a small chamber immersed in incompressible
7 hemolymph (Fig. 10e). The membrane complex attaches to the core at
8 anteriormost areas so that the core in the stationary state is immobilized by
9 a delicate balance between the fluid-filled chamber, the labyrinth, and the
10 membrane complex (Fig. 10e). Therefore, when the core attachment to the
11 ATV is displaced outward, i.e. anteriorly (thin arrow, Fig. 10e), a small but
12 powerful inward pressure will be generated at the attachment site to the
13 fluid chamber (thick arrow, Fig. 10e). Moreover, the proximal inflated
14 region of the core is attached to the anterior wall of the ATV from the
15 lateral direction (Fig. 10c), and the distal thin region of the core is attached
16 to the corner between the dorsal membrane and the anterior wall of the
17 ATV (Fig. 10d). Due to this geometry, the outward pressure to the anterior
18 wall of the ATV can displace the proximal region of the core anteriorly but
19 displace the distal region of the core more dorso-posteriorly. This
20 complementary lever operating proximo-distally might help to generate a
21 diagonal fluid motion from the disto-anterior direction to the
22 proximo-posterior direction and subserve frequency discrimination of
23 sensory neurons that are also arranged diagonally along the proximo-distal
24 axis (Fig. 10g). The presumed core action in field crickets is, in principle,
25 parallel to the action of the tympanal plate of the tettigoniid *Copiphora*
26 *gorgonensis*, which is attached to the fluid chamber containing sensory
27 neurons and that vibrates 180° out of phase against the thin tympanal
28 membranes (Monteraegre-Z and Robert 2015).

29 Further physiological investigations are needed to determine the extent
30 to which the fluid motion driven by the core contributes to generation of
31 traveling wave and frequency discrimination. The only available data are
32 results of a study showing that there was no electrical response of the
33 auditory nerve to tones above 12 kHz after ablating the ATM in *G.*
34 *campestris* (Nocke 1972). Nevertheless, this is in good accordance with the
35 finding that the most distal 20 neurons that tune to sounds higher than 15
36 kHz (Huber et al. 1989) are ideally situated for receiving a fluid current.

1 Since high-frequency sounds tend to attenuate during transmission in the
2 medium, distal neurons might need assistance from fluid motion for sharp
3 frequency tuning. Systematic disruption of the core integrity is needed for
4 elucidating concrete core functions in the future.

5 6 **Functional aspects of two neural groups**

7
8 The cricket tympanal organ is composed of two anatomically defined
9 neuronal groups; the proximal group and the distal group. While the somata
10 of the proximal group are firmly suspended to the anterior cuticular wall,
11 their dendritic tips are inserted to the cellular mass beyond the apertured
12 area. This aperture possibly limits propagation of vibration transmitted
13 through the longitudinal ATV and permits detection of the displacement of
14 the mass against the soma attachment site. Adequate stimulus of the
15 proximal group, therefore, would be displacements of the mass along the
16 anterior-posterior axis (Fig. 10a), a scheme similar to the subgenual organ
17 in which sensory dendrites attach to the ellipsoidal cellular mass (Nishino
18 and Field 2003; Strauß et al. 2012). However, the primary sources of
19 vibration that evoke displacement of the mass are still enigmatic. The
20 ligaments connecting the cellular mass to the thinned cuticle are one
21 possible route for transmission of cuticular vibration to proximal group
22 neurons.

23 The stimulus situation in distal group neurons is also complex even
24 though their dendrites are immersed in fluid. The two apertures appear to
25 limit transmission of sound energy to the ATV region between them (Fig.
26 10a). However, tensile stress could be loaded on the cilia not only directly
27 by the fluid motion but also indirectly by displacement of the cellular mass
28 and/or the dorsal membrane of tracheal vesicles (Bangert et al. 1998).

29 Physiological data imply the presence of such complex interplay
30 between the cellular mass and fluid-filled chamber. A striking characteristic
31 of cricket sensory neurons equivalent to the distal group is the occurrence
32 of additional sensitivity peaks at frequencies other than characteristic
33 frequencies (Esch et al. 1980; Hutching and Lewis 1981; Imaizumi and
34 Pollack 1999). Additional sensitivity peaks of auditory receptors often
35 emerge when sounds with higher intensities are applied (Imaizumi and
36 Pollack 1999). Therefore, such complex responses of single sensory

neurons may be attributable to the proximo-distal gradation of connectivity strength with surrounding tissues; i.e., distal neurons may be more sensitive to the core motion than to the cellular mass motion and in reverse sequence for proximal neurons.

Our assumption does not exclude the importance of frequency filtering based on intrinsic properties of scolopidial sensilla (Oldfield et al. 1986). Since morphological specializations of dendrites, scolopale caps, and attachment cells are evident among individual neurons of the distal group (Young and Ball 1974a), the stiffness gradient between dendritic cilia and attachment cells might confer additional frequency filtering to sensory neurons, as suggested in *crista acustica* of Tettigoniidae (Hummel et al. 2017). In fact, tuning properties of sensory neurons are invariable from the last instar that has no epithelial cores, although the auditory thresholds of sensory neurons are 30-45 dB higher in last instars than in adults (Ball and Hill 1978; Staudacher 2009).

The developmental data suggested the importance of the distal group of neurons in adult life. In contrast to the proximal group of neurons, the number of which is invariable from the fourth instar, the number of neurons in the distal group increases progressively from the sixth instar to the adult stage (Ball and Young 1974). Sound-transmitting apparatus linked to the distal group of neurons, such as tracheal vesicles and the epithelial core, progressively mature after the imaginal molt. Maturation of the core, which takes six days after the imaginal molt, is important for transmitting sufficiently powerful pressure into the fluid. The thresholds of acoustic orientation behaviors to the calling song with carrier frequency 5 kHz in female crickets gradually decrease by 20-30 dB in about 11 days after the imaginal molt (Sergejeva and Popov 1994). The improvement in sensitivity after the imaginal molt may be, at least partly, attributable to the formation of the epithelial core.

Evolutionary perspective

The tympanal organ of the field cricket (Gryllinae) differs greatly from the *crista acustica* of tettigoniidae in that it has unequal-sized tympanal membranes and no clear differences in sensory neuronal sizes. Despite its translucent appearance like a specialized membrane, the ATM is

1 acoustically less sensitive than the PTM (Larsen, 1987; Huber et al. 1989;
2 Schneider et al., 2017). Given that tympanal membranes in Gryllidae are
3 morphologically diverse (Huber et al., 1989; Schneider et al. 2017), one
4 might speculate that this system reflects habitats and behavioral ecology
5 unique to field crickets.

6 In a “tree-thinking” perspective, the subfamily Eneopterinae species are
7 phylogenetically close to Gryllinae but have adapted to tropical rainforests
8 (ter Hofstede et al. 2015; Chintauan-Marquier et al. 2016). The tribe
9 Lebinthini *L. bitaeniatus* has an ATM covered with a cuticular fold, in
10 addition to the PTM (Schneider et al. 2017). The ATM is smaller than the
11 PTM and the structure of the tympanal organ resembles that of field
12 crickets, retaining conserved features between Gryllinae and Eneopterinae.
13 However, the ATM of *L. bitaeniatus* is thinner than the PTV, which results
14 in sensitivity reversal of the PTM and ATM between *G. bimaculatus* and *L.*
15 *bitaeniatus* (Schneider et al. 2017). On the other hand, another tribe,
16 Eurepini, that has adapted to ground and shrubs in Australia (Grandcolas et
17 al. 2010), has a PTM but no ATM (Schneider et al. 2017). These results
18 indicate the possibility that their habitats, in addition to their unique
19 acoustic communications (ter Hofstede et al. 2015), affect ear morphology
20 in these cricket species.

21 In field crickets, 1) adaptation to ground habitats and 2) ultrasound
22 detection for avoiding bats could be reflected in morphologies of the
23 tympanal organ. Field crickets, in general, have habits that dig burrows in
24 the ground for shelter (Michelsen 1998; Gawarek et al. 2014). Eibl (1978)
25 found long sensory hairs close to the PTM that presumably detect
26 burrowing movements in field crickets. Their habitats that have many hard
27 obstacles such as stones and their burrowing habits raise the potential for
28 damage to the thin part of the cuticle, i.e., tympanal membranes. Damage to
29 the ATM leads to immediate dysfunctioning of the tympanal organ because
30 it is in close proximity to the fluid-filled sensory neurons sealed by the thin
31 covering membrane. This situation differs from that of the PTM, which is
32 indirectly attached to the tympanal organ via the PTV. Since sound
33 impinges not only to the PTV but also to the tracheal opening in the
34 prothorax (Larsen and Michelsen, 1978; Poulet and Hedwig, 2001), sound
35 transmission through the PTV to the tympanal organ is still functional
36 under condition of damage of the PTM (Kleindiest et al., 1983; Schmitz et

al., 1983). Therefore, strong selective pressure could be exerted on the ATM rather than on the PTV.

Sensitivity to ultrasound is thought to be acquired later than acquirement of an intraspecific communication system during an evolutionary process (Stumpner and von Helversen 2001; Strauß and Stumpner 2014). Since avoidance of ultrasound emitted by bats provides high survival value, modification of the pre-existing tympanal organ must have been needed for ultrasound detection at high sensitivity. Therefore, the thickening of the ATM and the core formation might be an evolutionary solution for protecting the distal group of neurons from mechanical damage as well as enabling high-frequency sound detection. Thickening of the ATM and external core attachment are indeed achieved by a small modification of the pre-existing structure.

The core size and shape differ even between two field cricket species, *G. bimaculatus* and *T. occipitalis*, the former being larger and more inflated proximally. Comparative neuroanatomy between crickets equipped with a thick anterior membrane (Gryllinae), thin membrane (Lebinthini), and no membrane (Eurepini) will be useful for assessing functions of the cores and evolution of sound transmission pathways.

In conclusion, compared to tettigonnid ears in which bilateral compression of two equal-sized tympanal membranes produces fluid motion (Montealegre-Z and Robert 2015), the field cricket tympanal organ is a more perturbation-resistant ear, in which vibrations of the single tympanal membrane are likely converted to unilateral fluid motion via a lever action (Fig. 10g). In this context, the cricket ear systematically resembles mammalian ears that enclose ossicles deep in the middle ear (Fig. 10h). Measurements of mechanical properties of sound transmission pathways will be the priority toward application of this miniature ear for industrial use, as directed by a beautiful biomimetics example of the ear of the parasitoid fly *Orimia ochracea* to a hearing aid (Mason et al. 2001; Miles and Hoy 2006).

Acknowledgments

This work was supported by Innovative Materials Engineering Based on Biological Diversity in Grant-in-Aid for Scientific Research on Innovative Areas (project number: 80332477) to HN and TT. HN thanks Dr. Masnori

1 Ochiai (Hokkaido University) for expertized comments on identification of
2 blood cells and epithelial cells and Dr. Larry Field (University of
3 Canterbury) for valuable discussions from comparative viewpoints.

4 Conflict of Interest

5 The authors declare that they have no conflict of interest.

6

References

- Affolter M, Caussinus E (2008) Tracheal branching morphogenesis in *Drosophila*: new insights into cell behavior and organ architecture. *Development* 135:2055-2064
- Autrum HJ (1941) Über Gehör- und Erschütterungssinn bei Locustiden. *Z Vergl Physiol* 28:580-637
- Ball EE, Hill KG (1978) Functional development of the auditory system of the cricket, *Teleogryllus commodus*. *J Comp Physiol* 127:131-138
- Ball EE, Young D (1974) Structure and development of the auditory system in the prothoracic leg of the cricket *Teleogryllus commodus* (Walker) II. Postembryonic development. *Z Zellforsch* 147:313-324
- Ball EE, Field LH. (1981) Structure of auditory system of the weta *Hemideina Crassidens* (Blanchard, 1851) (Orthoptera, Ensifera, Gryllacridoidea, Stenopelmatidae). 1. Morphology and histology. *Cell Tissue Res* 217:321–343.
- Bailey WJ, Thomson P (1977) Acoustic orientation in the cricket *Teleogryllus oceanicus* (Le Guillou). *J Exp Biol* 67:61-75.
- Bangert M, Kalmring K, Sickmann T, Stephen R, Jatho M, Lakes-Harlan R (1998) Stimulus transmission in the auditory receptor organs of the foreleg of bushcrickets (Tettigoniidae) I. The role of the tympana. *Hearing Res* 115:27-38
- Boyan GS (1993) Another look at insect audition: the tympanic receptors as an evolutionary specialization of the chordotonal system. *J Insect Physiol* 39:187-200.
- Chintauan-Marquier IO, Legendre F, Hugel S, Robillard T, Grandcolas F, Nel A, Zuccon D, Desutter-Grandcolas L (2016) Laying the foundations of evolutionary and systematic studies in crickets (Insecta, Orthoptera): a

1 multilocus phylogenetic analysis. Cladistics 32:54-81.

2
3 Eibl E (1978). Morphology of the sense organs in the proximal parts of the
4 tibiae of *Gryllus campestris* L. and *Gryllus bimaculatus* DeGeer (Insecta,
5 Ensifera). Zoomorphol 89:185-205

6
7 Esch HF, Huber F, Wohlers DW (1980) Primary auditory neurons in
8 crickets: physiology and central projections. J Comp Physiol 137:27-38

9
10 Field LH, Matheson T (1998) Chordotonal organs of insects. Adv Insect
11 Physiol 27:1–228

12
13 Friedman MH (1972a) A light and electron microscopic study of sensory
14 organs and associated structures in the foreleg tibia of the cricket, *Gryllus*
15 *assimilis*. J Morphol 138:263-327

16
17 Friedman MH (1972b) An electron microscopic study of the tympanal
18 organ and associated structures in the foreleg tibia of the cricket, *Gryllus*
19 *assimilis*. J Morphol 138:329-347

20
21 Gawalek M, Dudek K, Ekner-Grzyb A, Kwieciski Z, Sliwowska JH (2014)
22 Ecology of the field cricket (Gryllidae: Orthoptera) in farmland: the
23 importance of livestock grazing. North-West J Zool 10:325-332

24
25 Göpfert MC, Robert D (2002) Motion generation by *Drosophila*
26 mechanosensory neurons. Proc Natl Acad Sci USA 100:5514-5519

27
28 Grandcolas-D L, Blanchet E, Robillard T, Magal C, Vannier F, Dangles O
29 (2010) Evolution of the cercal sensory system in a tropical cricket clade
30 (Orthoptera: Grylloidea: Eneopterinae): a phylogenetic approach. Biol J
31 Linn Soc 99:614-631

32
33 Greenfield, M. D. (2002) Signalers and Receivers. Oxford University Press,
34 Oxford.414pp

35
36 Hill KG (1974) Carrier frequency as a factor in phonotactic behavior of

- female crickets (*Teleogryllus commodus*). J Comp Physiol 93:7-18.
- Ter Hofstede HM, Sconich S, Robillard T, Hedwig B (2015) Evolution of communication system by sensory adaptation of startle behavior. Curr Biol 25:3245-3252
- Hoy RR, Robert D (1996) Tympanal hearing in insects. Annu Rev Entmol 41:433-450
- Huber F, Kleindienst HU, Weber T, Thorson J (1984) Auditory behavior of the cricket. III. Tracking of male calling song by surgically and developmentally one-eared females, and the curious role of the anterior tympanum. J Comp Physiol 155:725-738
- Huber F, Moore TE, Loher W (1989) Cricket behavior and neurobiology. Cornell University Press
- Hummel J, Kössl M, Nowotony M (2017) Morphological basis for a tonotopic design of an insect ear. J Comp Neurol 525:2443-2455
- Hutchings M, Lewis DB (1981) Response properties of primary auditory fibers in the cricket *Teleogryllus oceanicus* (Le Guillou). J Comp Physiol 143:129-134
- Imaizumi K, Pollack GS (1999) Neural coding of sound frequency by cricket auditory receptors. J Neurosci 19:1508-1506
- Imaizumi K, Pollack GS (2005) Central projections of auditory receptor neurons of crickets. J Comp Neurol 493:439-447
- Johnstone BM, Saunders JC, Johnstone JR (1970) Tympanic membrane response in the cricket. Nature 227:625-626
- Kleindienst HU, Wohlers DW, Larsen ON (1983) Tympanal membrane motion is necessary for hearing in crickets. J Comp Physiol 151:397-400

- 1 Kössl M, Möckel D, Weber M, Seyfarth E-A (2008) Otoacoustic emissions
2 from insect ears: evidence of active hearing? J Comp Physiol A
3 194:597-609
4
- 5 Lankheet MJ, Cerkenik U., Larsen ON, van Leeuwen JL (2017)
6 Frequency tuning and directional sensitivity of tympanal vibrations in the
7 field cricket *Gryllus bimaculatus*. J R Soc Interface 14:20170035
8
- 9 Larsen ON (1987) The cricket's anterior tympanum revisited. The Science
10 of Nature 74:92-94
11
- 12 Larsen ON, Michelsen A (1978) Biophysics of the ensiferan ear. III. The
13 cricket ear as a four-input system. J Comp Physiol 123:217-227
14
- 15 Larsen ON, Kleindienst H-U, Michelsen A (1989) Biophysical aspects of
16 sound reception. In: Huber F, Moore T, Loher W (eds) Cricket behavior and
17 neurobiology. Cornell Univ Press, Ithaca, pp 364-390
18
- 19 Li T, Bellen HJ, Groves AK (2018) Using *Drosophila* to study mechanisms
20 of hereditary hearing loss. Dis Model Mech dmm 031492
21
- 22 Lomas KF, Greenwood DR, Windmill JFC, Jackson JC, Corfield J, Parsons
23 S (2013) Discovery of a lipid synthesizing organ in the auditory system of
24 an insect. Plos One 7:e51486
25
- 26 Ma M, Cao X, Dai J, Pastor-Pareja JC (2017) Basement membrane
27 manipulation in *Drosophila* wing discs affects Dpp retention but not
28 growth mechanoregulation. Development 42:97-106
29
- 30 Matsubayashi Y, Louani A, Dragu A, Sánchez-Sánchez BJ, Serna-Morales
31 E, Yolland L, Gyoergy A, Vizcay G, Fleck RA, Heddleston JM, Chew TL,
32 Siekhaus DE, Stramer BM (2017) A moving source of matrix components
33 is essential for *de novo* basement membrane formation. Curr Biol 27,
34 3526-3534
35
- 36 Mason AC, Oshinsky ML, Hoy RR (2001) Hyperacute directional hearing

1 in a microscale auditory system. Nature 410:486-490

2
3 Mhatre N, Montealegre-Z F, Balakrishnan R, Robert D (2009) Mechanical
4 response of the tympanal membranes of the tree cricket *Oecanthus henryi*.
5 J Comp Physiol A 195:453-462

6
7 Mhatre N, Robert D (2013) A tympanal insect ear exploits a critical
8 oscillator for active amplification and tuning. Curr Biol 23:1952-1957

9
10 Michel K 1974. Das Tympanalorgan von *Gryllus bimaculatus* DeGeer
11 (Saltatoria, Gryllidae). Z Morphol Tiere 77:285-315

12
13 Michelsen A, Povov AV, Lewis B (1994) Physics of directional hearing in
14 the cricket *Gryllus bimaculatus*. J Comp Physiol 175:153-164

15
16 Michelsen A (1998) The tuned cricket. New Physiol Sci 13:32-38

17
18 Miles RN, Hoy RR (2006) The development of a biologically-inspired
19 directional microphone for hearing aids. Audit Neurotol 11:86-94

20
21 Montealegre-Z F, Jonsson T, Robson-Brown KT, Postles M, Robert D.
22 (2012) Convergent evolution between insect and mammalian audition.
23 Science 338:968-971

24
25 Montealegre-Z F, Robert D (2015) Biomechanics of hearing in katydids. J
26 Comp Physiol A 201:5-18

27
28 Nishino H, Sakai M (1997) Three neural groups in the femoral chordotonal
29 organ of the cricket *Gryllus bimaculatus*: central projections and soma
30 arrangement and displacement during joint flexion. J Exp Biol
31 200:2583-2595

32
33 Nishino H, Field LH (2003) Somatotopic mapping of chordotonal organ
34 neurons in a primitive ensiferan, the New Zealand weta *Hemideina*
35 *femorata*: II. Complex tibial organ. J Comp Neurol 464:327-342

- 1 Nishino H, Mukai H, Takanashi T (2016) Chordotonal organs in
2 hemipteran insects: unique peripheral structures but conserved central
3 organization revealed by comparative neuroanatomy. *Cell Tissue Res*
4 366:549-572
- 5
- 6 Niwa N, Hiromi Y, Okabe M (2004) A conserved developmental program
7 for sensory organ formation in *Drosophila melanogaster*. *Nat Genet*
8 36:293-297
- 9
- 10 Nocke H (1972) Physiological aspects of sound communication in crickets
11 (*Gryllus campestris* L.). *J Comp Physiol* 80:141-162
- 12
- 13 Oldfield BP, Kleindienst HU, Huber F (1986) Physiology and tonotopic
14 organization of auditory receptors in the cricket *Gryllus bimaculatus*
15 DeGeer. *J Comp Physiol* 159:457-464
- 16
- 17 Paton JA, Capranica RR, Dragsten PR, Webb WW (1977) Physiological
18 basis for auditory frequency analysis in field crickets. *J Comp Physiol*
19 119:221-240
- 20
- 21 Plotnick RE, Smith DM (2012) Exceptionally preserved fossil insect ears
22 from the Eocene green river formation of Colorado. *J Paleontol* 86:19-24
- 23
- 24 Pollack GS, Imaizumi K (1999) Neural analysis of sound frequency in
25 insects. *BioEssays* 21:295-303
- 26
- 27 Poulet JFA, Hedwig B (2001) Tympanic membrane oscillation and auditory
28 receptor activity in the stridulating cricket *Gryllus bimaculatus*. *J Exp Biol*
29 204:1281-1293
- 30
- 31 Ribeiro C, Brehelin M (2006) Insect haemocytes: what type of cell is that?
32 *J Insect Physiol* 52:417-429
- 33
- 34 Robert D, Göpfert MC (2002) Novel schemes for hearing and orientation in
35 insects. *Curr Opin Neurobiol.* 12:715-720
- 36

1 Rust J, Stumpner A, Gottwald J (1999) Singing and hearing in a Tertiary
2 bushcricket. *Nature* 399:650.

3
4 Schmitz B, Scharstein H, Wendler G (1983) Phonotaxis in *Gryllus*
5 *campestris* L. (Orthoptera, Gryllidae) H. Acoustic orientation of female
6 crickets after occlusion of single sound entrances. *J Comp Physiol*
7 152:257-264.

8
9 Schneider ES, Römer H, Robillard T, Schimdt AKD (2017) Hearing with
10 exceptionally thin tympana: ear morphology and tympanal membrane
11 vibrations in eneopterine crickets. *Sci Rep* 7:15266

12
13 Schwabe, J. (1906) Beiträge zur Morphologie und Histologie der
14 tympanalen Sinnesapparate der Orthopteren. *Zoologica* 50:1–154.

15
16 Senthilan PR, Piepenbrock D, Overzmyradov G, Nadrowski B, Bechstedt S,
17 Pauls S, Winker M, Möbius W, Howard J, Göpfert MC (2012) *Drosophila*
18 auditory organ genes and genetic hearing defects. *Cell* 150:1042-1054

19
20 Sergejeva MV, Popov AV (1994) Ontogeny of positive phonotaxis in
21 female crickets *Gryllus bimaculatus* De Geer: dynamics of sensitivity,
22 frequency-intensity domain, and selectivity to temporal pattern of the male
23 calling song. *J Comp Physiol A* 174:381-389

24
25 Staudacher EM (2009) The auditory system of last instars in *Gryllus*
26 *bimaculatus* DeGeer. *Physiol Entomol* 34:18-29

27
28 Strauß J, Lehmann GUC, Lehmann AW, Lakes-Harlan R (2012) Spatial
29 organization of tettigoniid auditory receptors: insights from neuronal
30 tracing. *J Morphol* 273:1280-1290

31
32 Strauß J, Lakes-Harlan R (2014) Evolutionary and Phylogenetic Origins of
33 Tympanal Hearing Organs in Insects. B. Hedwig (ed.), *Insect Hearing and*
34 *Acoustic Communication, Animal Signals and Communication*. Vol. 1. pp.
35 5-28. Springer-Verlag Berlin Heidelberg

- 1 Strauß J, Lomas K, Field LH (2017) The complex tibial organ of the New
2 Zealand ground weta: sensory adaptations for vibrational signal detection.
3 Sci Rep 7:2031
4
- 5 Stumpner A, von Helversen D (2001) Evolution and function of auditory
6 systems in insects. Naturwissenschaften 88:159-170.
7
- 8 Yack JE (2004) The structure and function of an auditory chordotonal
9 organs in insects. Microsc Res Tech 63:315-337
10
- 11 Yager DD (1999) Structure, development, and evolution of insect auditory
12 systems. Microsc Res Tech 47:380-400
13
- 14 Young D, Ball E (1974a) Structure and development of the auditory system
15 in the prothoracic leg of the cricket *Teleogryllus commodus* (Walker). I.
16 Adult structure. Z Zellforsch 147:293-312
17
- 18 Young D, Ball E (1974b) Structure and development of the tracheal organ
19 in the mesothoracic leg of the cricket *Teleogryllus commodus* (Walker). Z
20 Zellforsch 147:325-334
21
- 22 Wigglesworth VB (1954) Growth and regeneration in the tracheal system
23 of an insect, *Rhodnius prolixus* (Hemiptera). J Cell Sci 95: 115-137.
24
- 25 Wigglesworth VB (1956) The Haemocytes and Connective Tissue
26 Formation in an Insect, *Rhodnius Prolixus* (Hemiptera). J Cell Sci s3-97:
27 89-98
28
- 29 Wigglesworth VB (1959) The role of the epidermal cells in the migration of
30 tracheoles in *Rhodnius prolixus* (Hemiptera). J Exp Biol 36:632-640
31
- 32 Wigglesworth VB (1973) Haemocytes and basement membrane formation
33 in *Rhodnius*. J Insect Physiol 19:831-844
34
- 35 Wolfrum U (1990) Actin filaments: the main components of the scolopale
36 in insect sensilla. Cell Tissue Res 261:85–96
37

Figure Legends

Fig. 1. Prothoracic leg tympanal organ and surrounding structures in minimally dissected specimens. (a-c) Posterior tympanal membrane (PTM, b) and anterior tympanal membrane (ATM, c) in the prothoracic tibia (a). (d-j) Two-dimensional optical stacks (d-h) and three-dimensional reconstructions (i, j) of the tympanal organ in which sensory neurons are labeled by microruby and surrounding structures are counterstained with Lucifer Yellow. From posterior to anterior, the PTM, PTV (posterior tracheal vesicle), ATV (anterior tracheal vesicle), core, and ATM are appositioned in parallel (see also ESMs 1,2). The PTV and ATV are interconnected by two apertures (h, j, see also ESM2). Sensory neurons located between the cellular mass and the dorsal membrane of the ATV are immersed in lipidic fluid sealed by the covering membrane (CM). The CM is secreted by cells lining the membrane (inset in f). The cellular mass is suspended by a bundle of ligaments (inset in e) to the thinned cuticle (arrows in e). The distal two thirds of the PTM has a hinge-like structure, as shown in a transverse section (g). AN, auditory nerve; Oli, olivarius organ; SGO: subgenual organ. *Scale bars:* (a) 1 mm; (b-d) 500 μm ; (e-g, inset in e) 50 μm ; (inset in f) 10 μm ; (h) 100 μm ; (i, j) 200 μm .

Fig. 2 The main body of the tympanal organ. (a-h) The cellular mass is composed of small attachment cells (AtCs) and large accessory cells (LACs). Sensory neurons, viewed dorsally (b, c) and proximally (d-f), are grouped into the proximal group (PG) and distal group (DG). The numbers of cells in g and h are seven for the PG and 60 for the DG. The PG dendrites externally attach to the proximal edge of the cellular mass (d). The DG dendrites (b) and scolopale caps (ScCs, c) are orderly arranged proximo-distally. DG somata are firmly attached to the dorsal membrane of the ATV (e, f). Note that most distal neurons extend dendrites to attachment cells elongated from the cellular mass (a, f). (g, h) Three-dimensional reconstructions of sensory neurons, AtCs and LACs, showing that sensory dendrites bridging between the dorsal membrane of the ATV and the bottom of the cellular mass, gradually rise more perpendicularly to the membrane from proximal to distal. (i-k) Topographically complex arrangements of the dendritic cilia of the DG, viewed dorsally (i) and

distally (j, k). The orientations of dendrites and ScCs are schematized (i, k).
 (l) An optical section close to the covering membrane showing that the core
 attaches to the covering membrane at its' dorso-distal region. The outline of
 the core (broken line) is overlaid. SC: scolopale cell. Scale bars: (a-h, l) 50
 μm ; (i-k) 20 μm .

Fig. 3 Subcellular structures of sensory neurons revealed by phalloidin
 staining. (a) The LAC and AtC are readily identifiable by their spherical
 and ellipsoidal shapes of nuclei, respectively. (b-d) Different magnified
 images of the ciliated region of sensory dendrites showing that scolopale
 rods (ScR) rich in actin filaments highlighten an orderly arrangement of
 sensory dendrites and specialized structures such as the scolopale cap
 (ScC), ciliary dilation (CD), and dendritic dilation (DD). SC: scolopale cell.
 Scale bars: (a, c) 50 μm ; (b, d) 10 μm .

Fig. 4 Meso and metathoracic tracheal organs (a-d) and tympanal organs in
 nymphs (e-h). (a-d) Meso and metathoracic tracheal organs have a cellular
 mass structurally resembling that in the prothoracic tympanal organ but
 lacking inflated tracheae interconnected by two apertures (a, b), distal
 group sensory neurons, covering membrane, and core (c, d). The cellular
 mass is suspended to the ATV via fine fibers (c, see also ESM3). Olivarius
 organs (Oli) are stuffed in the space distal and proximal to the tracheal
 organ and the subgenual organ (SGO). Note hemocytes aggregating
 beneath the cellular mass (c, d). (e-h) Tympanal organs in a phalate adult
 (e-g) and 6th instar (h) showing that the PTV and ATV are spaced by
 tracheoblast cells (arrow in f), but the covering membrane has already
 developed (g, h). The cellular mass is located closely to the anterior cuticle
 since the core has not developed yet (e, f). Scale bars: (a, b) 500 μm ; (c, d,
 f-h) 50 μm ; (e) 100 μm .

Fig. 5 Three-dimensional reconstructions (volume rendering) of the
 epithelial core of *G. bimaculatus* (a-e) and a smaller field cricket, *T.*
occipitalis (f-h), in which the epicuticle of the ATM has been removed. The
 basement membrane (BM) that extends from the rim of the anterior
 tympanal membrane is attached to the antero-distal edge of the core (d, see
 also Fig. 9b). The width of the BM becomes smaller distally (b), resulting

in the distal region of the core being closer to the ATM (e). A smaller core is present in the field cricket *T. occipitalis* (f-h). ECL: epithelial cell layer; La: labyrinth. Scale bars = 50 μ m.

Fig. 6 Transverse sections of the prothoracic tibia in *T. occipitalis* (a-k) and *G. bimaculatus* (l-n) showing varied association of the core with surrounding tissues proximo-distally. The apparatus comprising the tympanal organ are overlaid on the optical sections (a-h). Numbers in a-h are distances from the proximal tip of the cellular mass. The attachment fashion between the PTV and ATV is varied from proximal to distal (a-h). The frameworks in the labyrinth (La) support the proximal region of the core from its ventral aspect (c, d). Contact of the core with the lipidic fluid is not direct but via the covering membrane (CM) (i, j), and only the distal region of the core attaches to the covering membrane (d-f, k, m, n). The basement membrane (BM) is attached posteriorly to the dorsal region of the core (j, k, l-n). Transverse sections distal to the cellular mass (i-n, see Fig. 2d-f for proximal regions) show that the CM keeps its tautness (l-n). Scale bars: (a-h) 100 μ m; (i-n) 50 μ m.

Fig. 7 Application of chitinase to the epithelial core. (a) The epithelial core was cleared in methyl salicylate and viewed under differential interference contrast. (b-d) Chitinase treatment resulted in breakdown of the core (b) and the ligaments (d) connecting the cellular mass to the dorsal cuticle, suggesting that these are made up by chitin. The cellular mass and covering membrane were not affected by application of chitinase (c). Scale bars: 100 μ m.

Fig. 8 Cellular contribution to the core formation in *G. bimaculatus*. (a-c) After the final molt, epithelial cells (ECs) with diameters of 4-5 μ m aggregate beneath the ATM (a) and interconnect laterally by thin filopodia (b). Some of epithelial cells (ECs) start to migrate posteriorly to attach to the ATV by producing thick filopodia posteriorly (c). (d, e) The basement membrane (BM) comprising a sheet of single cells with diameters of 7-8 μ m (d) attaches to the periphery of the core anlage (e). (f) The proximal edge of the core attaches to the CM in its central region. The epicuticle of the ATM is removed. (g) The epithelial cell layers (ECLs) ventral to the

core region are shaped into the labyrinth (La) by phagocytosis of thin filopodia by plasmatocytes. (h-j) The ECs scaffolding on the ATV start to secrete chitin inside (h) at day 1 (see Fig. 9c). Three-dimensional reconstructions of tractable chitin cylinders show that epithelial cells secrete chitin inside and participate in enlargement of the chitin mass (i, j). (k, l) Mature core in which epithelial cells are radiated like a fan. New epithelial cells attach to the cell bodies of old cells and secrete chitin (arrowheads in l). (m) Working hypotheses of core development. The core is formed by repetition of 1) epithelial cell scaffolding to the ATV, 2) secretion of chitin, and 3) new cells scaffolding to old cells. Note that epithelial cells located laterally tend to secrete thicker but shorter chitin compared to the central region, collectively contributing to accretionary growth of the fan-shaped core. Scale bars: (a, d-g, k, l) 50 μm ; (b, c) 10 μm ; (h-j) 20 μm .

Fig. 9 Epithelial core development. (a, b) Intact ATM viewed under an objective microscope in adults at different ages, showing that epithelial cells embedded in whitish connective tissues aggregate beneath the ATM at day 1 (a) and the melanized basement membrane (BM) attaches to the edge of the core at day 6, demarcating the core (outlined by a broken line in b). (c-h) Progressive enlargement of the core from day 1 to day 6. (i) Volumetric measurement of the core during development showing that the core size reaches a plateau at day 6, although some inter-individual differences are evident ($n=5$ for each day). Scale bars: (a, b) 100 μm ; (c-h) 50 μm .

Fig. 10 Sound transmission pathways in the tympanal organ of field crickets. (a) Schematized representation of sound transmission apparatus viewed anteriorly. (b) Schematic representation of elaborated structures underlying the ATM. The three attachment sites to the core and the BM to ATM attachment site are shown in red. (c, d) Different connectivities of the proximal region of the core (e) and the distal region of the core (d) with the ATV and the membrane complex. (e, f) Possible lever action to the core. Outward inflation of the ATV results in inward displacement of the dorsal core due to the support of the membrane complex functioning as a fulcrum. This is class 1 lever action (f). (g, h) The core action (g) is presumably

functionally parallel to that of mammalian ossicles (h) in that they both convert large mechanical displacements to small but powerful pressures in the fluid. See text for abbreviations. Scale bars: (a) 500 μm ; (b-e) 100 μm .

Legends of Supplementary Materials

SM1 Procedures for low-invasive observation of the interior structure of the tympanal organ in *G. bimaculatus*.

ESM1 Adult tympanal organ in the left prothoracic tibia of *G. bimaculatus*, viewed anteriorly, showing sensory neurons (magenta) and associated structures (green) including the epithelial core.

ESM2 Tympanal organ of the left prothoracic tibia of *G. bimaculatus*, viewed dorsally, showing parallel appositions between the PTM, PTV, ATV, core and ATM. Note two apertures interconnecting the PTV and ATV. See text for abbreviations.

ESM3 Tracheal organ in the left mesothoracic tibia of *G. bimaculatus*, viewed anteriorly, showing no epithelial core, no covering membrane and no distal group neurons. The cellular mass is suspended to the anterior cuticle wall and the dorsal membrane of the ATV via thin fibers. Note hemocytes aggregating beneath the cellular mass. SGO: subgenual organ.

ESM4 ATM of the left prothoracic tibia of *G. bimaculatus*, viewed anteriorly, showing elaborate structures associated with the core. The epicuticle of the ATM was carefully removed in this specimen. See text for abbreviations.

ESM5 Tympanal organ in the left prothoracic tibia of *G. bimaculatus*, viewed proximally, showing different connectivities of the core with surrounding structures. The epicuticle of the ATM was removed in this specimen. Note that a small air-filled space between the ATV and the complex of the basement membrane/epithelial cell layers becomes larger distally. The labyrinth-like frameworks, originating from filopodia of epithelial cells, bridge between the epithelial cell layers and ATV and core.

1 The central region of the proximal tip of the core attaches to the covering
2 membrane. The epicuticle of the ATM and that of the surrounding
3 exoskeleton were removed in this specimen. See text for abbreviations.

4
5 **ESM6** Tympanal organ in the left prothoracic tibia of *T. occipitalis*, viewed
6 proximally, showing the covering membrane attached to the distal region of
7 the core. Note the lipidic fluid coagulated inside the covering membrane.
8 See text for abbreviations.

9
10 **ESM7** Low-magnified image of the tympanal organ in the left prothoracic
11 tibia of *T. occipitalis*, viewed proximally, showing different attachments
12 between the PTV and ATV proximo-distally. Note thick ledges protrude
13 from the ATM to support the ATV ventrally. See text for abbreviations.

14
15 **ESM8** Movement of the basement membrane and the underlying core. The
16 slow pressure-on to the exocuticle of the distal femur results in movement
17 the basement membrane toward ventrally while the pressure-off results in
18 return of the basement membrane to its original position. Note that the
19 displacement of the distal core region is smaller than that of the proximal
20 region.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Fig. 9

Fig. 10