Title	Singular Bogoliubov transformations and inequivalent representations of canonical commutation relations
Author(s)	Arai, Asao
Citation	Reviews in Mathematical Physics, 31(08), 1950026 https://doi.org/10.1142/S0129055X19500260
Issue Date	2019-09
Doc URL	http://hdl.handle.net/2115/79187
Rights	Electronic version of an article published as Reviews in Mathematical Physics, Vol. 31, No. 08, 1950026 (2019), https://doi.org/10.1142/S0129055X19500260 ©copyright World Scientific Publishing Company, https://www.worldscientific.com/worldscinet/rmp
Type	article (author version)
File Information	RMP31-08_1950026.pdf

[Instructions for use](#)

Singular Bogoliubov Transformations and Inequivalent Representations of Canonical Commutation Relations

Asao Arai

Department of Mathematics, Hokkaido University

Sapporo, Hokkaido 060-0810

Japan

E-mail: arai@math.sci.hokudai.ac.jp

Abstract

We introduce a concept of singular Bogoliubov transformation on the abstract boson Fock space and construct a representation of canonical commutation relations (CCRs) which is inequivalent to any direct sum of the Fock representation. Sufficient conditions for the representation to be irreducible are formulated. Moreover, an example of such representations of CCRs is given.

Keywords: singular Bogoliubov transformation, canonical commutation relations, inequivalent representation, boson Fock space.

Mathematics Subject Classification 2010: 47L60.

1 Introduction

In this paper we introduce a concept of singular Bogoliubov transformation on the abstract boson Fock space over a Hilbert space $\mathcal{H}$. A singular Bogoliubov transformation is induced by a pair (S, T) of *unbounded* linear operators on $\mathcal{H}$ obeying suitable constraints on their domain of definition and some consistency conditions (see (4.1) and (4.2)). As in the case of the standard bosonic Bogoliubov transformations (e.g., [11, 12, 13]), a singular Bogoliubov transformation also yields a representation $\pi_B(\mathcal{D})$ of the canonical commutation relations (CCRs) over a dense subspace $\mathcal{D}$ of $\mathcal{H}$. We are interested in conditions under which the representation $\pi_B(\mathcal{D})$ is inequivalent to any direct sum representation of the Fock representation $\pi_F(\mathcal{D})$ of the CCRs over $\mathcal{D}$. We formulate also sufficient conditions for $\pi_B(\mathcal{D})$ to be irreducible.

One of the motivations for this work comes from representation theoretic considerations on the Casimir effect [3, 7, 14]. In the paper [5], there is a suggestion (a heuristic argument) that a representation of CCRs inequivalent to the Fock one may be associated with the Casimir effect and one may use this representation to explain the Casimir effect

without invoking zero-point energy. But, in this paper, we do not discuss the Casimir effect.

The outline of the present paper is as follows. In Section 2, we recall the definition of representation of the CCRs over a complex inner product space and state a proposition on irreducibility of a representation of CCRs. In Section 3, we review the Fock representation of CCRs and prove some related facts. In Section 4, we introduce a singular Bogoliubov transformation. As mentioned above, it yields a representation $\pi_B(\mathcal{D})$ of the CCRs over $\mathcal{D}$. We prove that $\pi_B(\mathcal{D})$ is inequivalent to any direct sum representation of the Fock representation $\pi_F(\mathcal{D})$ (Theorem 4.5). We formulate also sufficient conditions for $\pi_B(\mathcal{D})$ to be irreducible (Theorems 4.7 and 4.8). In Section 5, we show that, for each pair (K, L) of injective symmetric operators with suitable properties, there may exist a singular Bogoliubov transformation and a representation $\pi_{K,L}(\mathcal{D})$ of the CCRs over $\mathcal{D}$, which is inequivalent to any direct sum representation of $\pi_F(\mathcal{D})$ (Theorem 5.3). We show that, under an additional condition, $\pi_{K,L}(\mathcal{D})$ is irreducible (Theorem 5.4). In the last section, we discuss an example of $\pi_{K,L}(\mathcal{D})$ in the case $\mathcal{H} = L^2(\Lambda)$ with Λ being a finite box in the d -dimensional Euclidean space $\mathbb{R}^d$ ($d \geq 2$). This example may be related to a Casimir effect which may be caused by a quantum scalar field on Λ . But, in the present paper, we do not discuss this aspect.

2 Preliminary

For a linear operator A on a Hilbert space $\mathcal{H}$, we denote by $D(A)$ the domain of A . If A is densely defined, then A has its adjoint; we denote it by A^* .

Let $\mathcal{F}$ be a complex Hilbert space and $\mathcal{D}$ be a dense subspace of $\mathcal{F}$. Let $\mathcal{V}$ be a complex inner product space with inner product $\langle \cdot, \cdot \rangle_{\mathcal{V}}$ (linear in the second variable and anti-linear in the first) and norm $\|\cdot\|_{\mathcal{V}}$ (we sometimes omit the subscript $\mathcal{V}$ in $\langle \cdot, \cdot \rangle_{\mathcal{V}}$ and $\|\cdot\|_{\mathcal{V}}$ if there is no danger of confusion). Suppose that, for each $f \in \mathcal{V}$, a densely defined closed linear operator $C(f)$ on $\mathcal{F}$ is given. Then the triple $(\mathcal{F}, \mathcal{D}, \{C(f), C(f)^* | f \in \mathcal{V}\})$ is called a representation of the CCRs over $\mathcal{V}$ if the following (i)–(iii) hold:

- (i) (domain invariance) For all $f \in \mathcal{V}$, $\mathcal{D} \subset D(C(f)) \cap D(C(f)^*)$, $C(f)\mathcal{D} \subset \mathcal{D}$, $C(f)^*\mathcal{D} \subset \mathcal{D}$.
- (ii) (anti-linearity in test vectors) For all $f, g \in \mathcal{V}$ and $\alpha, \beta \in \mathbb{C}$ (the set of complex numbers), $C(\alpha f + \beta g) = \alpha^* C(f) + \beta^* C(g)$ on $\mathcal{D}$, where, for $z \in \mathbb{C}$, z^* denotes the complex conjugate of z .
- (iii) (CCRs over $\mathcal{V}$) For all $f, g \in \mathcal{V}$,

$$[C(f), C(g)^*] = \langle f, g \rangle_{\mathcal{V}}, \quad [C(f), C(g)] = 0 \quad \text{on } \mathcal{D},$$

where, for linear operators A and B on a Hilbert space, $[A, B] := AB - BA$.

Two representations $(\mathcal{F}, \mathcal{D}, \{C(f), C(f)^* | f \in \mathcal{V}\})$ and $(\mathcal{F}', \mathcal{D}', \{C'(f), C'(f)^* | f \in \mathcal{V}\})$ of the CCRs over $\mathcal{V}$ are said to be equivalent if there exists a unitary operator $U : \mathcal{F} \rightarrow \mathcal{F}'$ such that, for all $f \in \mathcal{V}$, $UC(f)U^{-1} = C'(f)$.

For two linear operators A and B on a Hilbert space $\mathcal{X}$, the symbol $A \subset B$ means that B is an extension of A , i.e., $D(A) \subset D(B)$ and $A\Psi = B\Psi$, $\forall \Psi \in D(A)$.

As usual, we denote by $\mathfrak{B}(\mathcal{X})$ the Banach space of everywhere defined bounded linear operators on $\mathcal{X}$.

For a set $\mathfrak{A}$ of linear operators on a Hilbert space $\mathcal{X}$, the set

$$\mathfrak{A}' := \{T \in \mathfrak{B}(\mathcal{X}) | TA \subset AT, \forall A \in \mathfrak{A}\} \subset \mathfrak{B}(\mathcal{X})$$

is called the strong commutant of $\mathfrak{A}$.

The set $\mathfrak{A}$ is said to be reducible if there is a non-trivial closed subspace $\mathcal{M}$ of $\mathcal{X}$ ($\mathcal{M} \neq \{0\}$, $\mathcal{X}$) such that every $A \in \mathfrak{A}$ is reduced by $\mathcal{M}$ (i.e., $P_{\mathcal{M}}A \subset AP_{\mathcal{M}}$, where $P_{\mathcal{M}}$ is the orthogonal projection onto $\mathcal{M}$). The set $\mathfrak{A}$ is said to be irreducible if it is not reducible.

A representation $(\mathcal{F}, \mathcal{D}, \{C(f), C(f)^* | f \in \mathcal{V}\})$ of the CCRs over $\mathcal{V}$ is said to be reducible (resp. irreducible) if the set $\{C(f), C(f)^* | f \in \mathcal{V}\}$ is reducible (resp. irreducible).

The following fact is well known (see, e.g., [2, Proposition 5.9]):

Lemma 2.1 *Let $\mathfrak{A}$ be a set of linear operators on $\mathcal{X}$.*

- (i) *If $\mathfrak{A}' = \mathbb{C}I := \{\alpha I | \alpha \in \mathbb{C}\}$ (I denotes identity), then $\mathfrak{A}$ is irreducible.*
- (ii) *If $\mathfrak{A}$ is an irreducible set of densely defined linear operators on $\mathcal{X}$ and $*$ -invariant (i.e., $A \in \mathfrak{A} \implies A^* \in \mathfrak{A}$), then $\mathfrak{A}' = \mathbb{C}I$.*

3 Fock Representation of CCRs and Related Facts

Let $\mathcal{H}$ be a complex Hilbert space and, for each $n \in \mathbb{N}$ (the set of natural numbers), $\otimes_s^n \mathcal{H}$ be the n -fold symmetric tensor product Hilbert space of $\mathcal{H}$ with convention $\otimes_s^0 \mathcal{H} := \mathbb{C}$. Then the boson Fock space over $\mathcal{H}$ is defined to be the direct sum Hilbert space of $\{\otimes_s^n \mathcal{H}\}_{n=0}^\infty$:

$$\begin{aligned} \mathcal{F}_b(\mathcal{H}) &:= \oplus_{n=0}^\infty \otimes_s^n \mathcal{H} \\ &= \left\{ \Psi = \{\Psi^{(n)}\}_{n=0}^\infty | \Psi^{(n)} \in \otimes_s^n \mathcal{H}, n \geq 0, \sum_{n=0}^\infty \|\Psi^{(n)}\|^2 < \infty \right\}. \end{aligned}$$

The subspace

$$\begin{aligned} \mathcal{F}_0(\mathcal{H}) &:= \{\Psi = \{\Psi^{(n)}\}_{n=0}^\infty | \Psi^{(n)} \in \otimes_s^n \mathcal{H}, n \geq 0 \text{ and there exists an } n_0 \in \mathbb{N} \text{ such that} \\ &\quad \text{for all } n \geq n_0, \Psi^{(n)} = 0\} \end{aligned}$$

is dense in $\mathcal{F}_b(\mathcal{H})$. This subspace is called the finite particle subspace of $\mathcal{F}_b(\mathcal{H})$.

We denote by $A(f)$ the annihilation operator with test vector $f \in \mathcal{H}$ on $\mathcal{F}_b(\mathcal{H})$, which is the unique densely defined closed linear operator on $\mathcal{F}_b(\mathcal{H})$ such that its adjoint $A(f)^*$, called the creation operator with test vector f , takes the following form: for all $\Psi \in D(A(f)^*)$, $(A(f)^*\Psi)^{(0)} = 0$ and

$$(A(f)^*\Psi)^{(n)} = \sqrt{n}S_n(f \otimes \Psi^{(n-1)}), \quad n \geq 1,$$

where S_n is the symmetrization operator on the n -fold tensor product $\otimes^n \mathcal{H}$ of $\mathcal{H}$. For all $f \in \mathcal{H}$, $\mathcal{F}_0(\mathcal{H}) \subset D(A(f)) \cap D(A(f)^*)$ and $A(f)$ and $A(f)^*$ leave $\mathcal{F}_0(\mathcal{H})$ invariant. Moreover, $\{A(f), A(f)^* | f \in \mathcal{H}\}$ satisfies the CCRs over $\mathcal{H}$:

$$[A(f), A(g)^*] = \langle f, g \rangle_{\mathcal{H}}, \quad [A(f), A(g)] = 0, \quad [A(f)^*, A(g)^*] = 0 \quad (f, g \in \mathcal{H}) \quad (3.1)$$

on $\mathcal{F}_0(\mathcal{H})$.

The following fact is well known (see, e.g., [2, Theorem 5.14]):

Theorem 3.1 *Let $\mathcal{D}$ be a dense subspace of $\mathcal{H}$. Then $\{A(f), A(f)^* | f \in \mathcal{D}\}' = \mathbb{C}I$ and*

$$\pi_F(\mathcal{D}) := (\mathcal{F}_b(\mathcal{H}), \mathcal{F}_0(\mathcal{H}), \{A(f), A(f)^* | f \in \mathcal{D}\}) \quad (3.2)$$

is an irreducible representation of the CCRs over $\mathcal{D}$.

The representation $\pi_F(\mathcal{D})$ is called the Fock representation of the CCRs over $\mathcal{D}$.

An anti-linear mapping $J : \mathcal{H} \rightarrow \mathcal{H}$ is called a conjugation on $\mathcal{H}$ if $J^2 = I$ and $\|Jf\| = \|f\|$, $f \in \mathcal{H}$.

The next proposition plays a basic role in the theory of representations of CCRs in boson Fock spaces:

Proposition 3.2 *Assume that $\mathcal{H}$ is separable. Let S and T be (not necessarily bounded) linear operators on $\mathcal{H}$ with the following property: $D(S) \cap D(T)$ includes a dense subspace $\mathcal{D}$ of $\mathcal{H}$ such that $T\mathcal{D}$ is dense in $\mathcal{H}$ and $T_{\mathcal{D}} := T \upharpoonright \mathcal{D}$, the restriction of T to $\mathcal{D}$, is injective. Let J be a conjugation on $\mathcal{H}$ and suppose that there exists a non-zero vector $\Omega \in \mathcal{F}_b(\mathcal{H})$ such that, for all $\Psi \in \mathcal{F}_0(\mathcal{H})$ and $f \in \mathcal{D}$,*

$$\langle A(Tf)^* \Psi, \Omega \rangle = -\langle A(JSf) \Psi, \Omega \rangle. \quad (3.3)$$

Then $ST_{\mathcal{D}}^{-1}$ is a densely defined closable operator and its closure $\overline{ST_{\mathcal{D}}^{-1}}$ is Hilbert-Schmidt.

Proof. Let $\mathcal{D}_T := T\mathcal{D}$. Then, by the present assumption, $\mathcal{D}_T$ is dense in $\mathcal{H}$. Hence $T_{\mathcal{D}}^{-1}$ is densely defined with $D(T_{\mathcal{D}}^{-1}) = \mathcal{D}_T$. By (3.3), we have $\langle A(g)^* \Psi, \Omega \rangle = -\langle A(JST_{\mathcal{D}}^{-1}g) \Psi, \Omega \rangle$ for all $g \in \mathcal{D}_T$. Hence it follows from [1, Proposition 3.3]¹ that $ST_{\mathcal{D}}^{-1}$ is bounded (hence closable) and $\overline{ST_{\mathcal{D}}^{-1}}$ is Hilbert-Schmidt. ■

Lemma 3.3 *Let X and Y be (not necessarily bounded) linear operators on $\mathcal{H}$ such that there exists a dense subspace $\mathcal{D} \subset D(X) \cap D(Y)$ and the following equation holds:*

$$\langle Xf, Xg \rangle - \langle Yf, Yg \rangle = \langle f, g \rangle, \quad f, g \in \mathcal{D}. \quad (3.4)$$

Let $X_{\mathcal{D}} := X \upharpoonright \mathcal{D}$. Then $X_{\mathcal{D}}$ is injective and $X_{\mathcal{D}}^{-1}$ is bounded with $\|X_{\mathcal{D}}^{-1}\| \leq 1$.

¹In the cited proposition, it is assumed that an operator L satisfying $\langle A(g)^* \Psi, \Omega \rangle = \langle A(JLg) \Psi, \Omega \rangle$ is bounded. But, as is seen from the proof, it is not necessary.

Proof. It follows from (3.4) that, for all $f \in \mathcal{D}$, $\|X_{\mathcal{D}}f\|^2 = \|f\|^2 + \|Yf\|^2 \geq \|f\|^2$, which implies the desired result. $\blacksquare$

The following proposition is new (to the author's best knowledge) and plays a crucial role in the present paper.

Proposition 3.4 *Assume that $\mathcal{H}$ is separable. Let X and Y be as in Lemma 3.3 and suppose that X is unbounded and $X\mathcal{D}$ is dense in $\mathcal{H}$. Let Ω be a vector in $\mathcal{F}_b(\mathcal{H})$ satisfying*

$$\langle A(Xf)^*\Psi, \Omega \rangle = -\langle A(JYf)\Psi, \Omega \rangle, \quad \Psi \in \mathcal{F}_0(\mathcal{H}), f \in \mathcal{D}. \quad (3.5)$$

Then $\Omega = 0$.

Proof. We prove the proposition by reductio ad absurdum. Suppose that there existed a non-zero vector $\Omega \in \mathcal{F}_b(\mathcal{H})$ such that (3.5) holds. Since $\text{Ran } X_{\mathcal{D}}$ is dense by the present assumption and $X_{\mathcal{D}}$ is injective by Lemma 3.3, it follows from Proposition 3.2 that $K := \overline{YX_{\mathcal{D}}^{-1}}$ is Hilbert-Schmidt. Hence K^*K is trace class. Using (3.4) and a limiting argument, one can show that

$$K^*K = I - L^*L, \quad (3.6)$$

where $L := \overline{X_{\mathcal{D}}^{-1}}$. For a linear operator A , we denote by $\sigma(A)$ the spectrum of A . Since K^*K is a non-negative trace class operator, it is a non-negative compact operator. Hence $\sigma(K^*K) \setminus \{0\} = \{\lambda_n\}_{n=1}^N$, where $N < \infty$ or $N = \infty$ and λ_n is a positive eigenvalue of K^*K with a finite multiplicity; if $N = \infty$, then $\lim_{n \rightarrow \infty} \lambda_n = 0 \cdots (*)$. Hence $\sigma(L^*L) \setminus \{1\} = \{1 - \lambda_n\}_{n=1}^N$.

Suppose that there existed a constant $\gamma > 0$ such that $L^*L \geq \gamma$. Then, for all $g \in \text{Ran } X_{\mathcal{D}}$, $\|X_{\mathcal{D}}^{-1}g\|^2 \geq \gamma\|g\|^2$. Hence $\|X_{\mathcal{D}}f\|^2 \leq \gamma^{-1}\|f\|^2$, $f \in \mathcal{D}$. But this contradicts the unboundedness of X . Hence $\inf \sigma(L^*L) = 0$. Hence $N = \infty$ and there exists a subsequence $\{\lambda_{n_k}\}_{k=1}^{\infty}$ such that $\lim_{k \rightarrow \infty} (1 - \lambda_{n_k}) = 0$, i.e., $\lim_{k \rightarrow \infty} \lambda_{n_k} = 1$. But this contradicts (*). Thus we arrive at a contradiction. $\blacksquare$

In concluding this section, we state a fact which may be used to prove the irreducibility of representations of CCRs. For each $f \in \mathcal{H}$, one has a self-adjoint operator

$$\Phi(f) := \frac{1}{\sqrt{2}} \overline{(A(f)^* + A(f))},$$

called the Segal field operator with test vector f . It is easy to see that $\Phi(f)$ leaves $\mathcal{F}_0(\mathcal{H})$ invariant and, for all $f, g \in \mathcal{H}$,

$$[\Phi(f), \Phi(g)] = i\text{Im} \langle f, g \rangle \quad \text{on } \mathcal{F}_0(\mathcal{H}), \quad (3.7)$$

where, for a complex number $z \in \mathbb{C}$, $\text{Im } z$ denotes the imaginary part of z .

Let J be a conjugation on $\mathcal{H}$. Then the set

$$\mathcal{H}_J := \{f \in \mathcal{H} | Jf = f\} \quad (3.8)$$

is a real Hilbert space with the inner product of $\mathcal{H}$. It follows that

$$\mathcal{H} = \{f + ig | f, g \in \mathcal{H}_J\}.$$

Let A and B be densely defined linear operators on $\mathcal{H}$ with the following properties (i) and (ii):

(i) $JA \subset AJ, JB \subset BJ$.

(ii) There exists a subspace $\mathcal{E}$ in $\mathcal{H}_J$ such that $\mathcal{E} \subset D(A) \cap D(B)$ and, for all $f, g \in \mathcal{E}$, $\langle Af, Bg \rangle = \langle f, g \rangle$.

Property (i) means that, for all $\Psi \in D(A)$, $J\Psi \in D(A)$ and $AJ\Psi = JA\Psi$. Hence, in particular, $A(D(A) \cap \mathcal{H}_J) \subset \mathcal{H}_J$. Similarly one has that $B(D(B) \cap \mathcal{H}_J) \subset \mathcal{H}_J$.

Let

$$\phi_A(f) := \Phi(Af), \quad \pi_B(f) := \Phi(iBg), \quad f \in D(A) \cap \mathcal{H}_J, g \in D(B) \cap \mathcal{H}_J.$$

Then it follows from (3.7) that $\phi_A(f)$ and $\pi_B(g)$ obey the Heisenberg commutation relations over $\mathcal{E}$ on $\mathcal{F}_0(\mathcal{H})$: for all $f, g \in \mathcal{E}$,

$$[\phi_A(f), \pi_B(g)] = i \langle f, g \rangle, \quad [\phi_A(f), \phi_A(g)] = 0, \quad [\pi_B(f), \pi_B(g)] = 0$$

on $\mathcal{F}_0(\mathcal{H})$. In fact, $e^{i\phi_A(f)}$ and $e^{i\pi_B(g)}$ obey the Weyl relations (see, e.g., [2, Theorem 5.25(ii)]):

$$e^{i\phi_A(f)} e^{i\pi_B(g)} = e^{-i \langle f, g \rangle} e^{i\pi_B(g)} e^{i\phi_A(f)}, \quad f, g \in \mathcal{E}.$$

Lemma 3.5 *Let $C := A + iB$. Suppose that $C\mathcal{E}$ is dense in $\mathcal{H}$. Then $\{e^{i\phi_A(f)}, e^{i\pi_B(f)} | f \in \mathcal{E}\}' = \mathbb{C}I$. In particular, $\{e^{i\phi_A(f)}, e^{i\pi_B(f)} | f \in \mathcal{E}\}$ is irreducible.*

Proof. Let $T \in \{e^{i\phi_A(f)}, e^{i\pi_B(f)} | f \in \mathcal{E}\}'$. Then, for all $f \in \mathcal{E}$, $Te^{i\phi_A(f)} = e^{i\phi_A(f)}T$ and $Te^{i\pi_B(f)} = e^{i\pi_B(f)}T$. By a general formula for the Weyl operator $e^{i\Phi(u)}$ ($u \in \mathcal{H}$) (see, e.g., [2, Theorem 5.25(ii)]), we have

$$e^{i\Phi(Cf)} = e^{i\|f\|^2/2} e^{i\phi_A(f)} e^{i\pi_B(f)}, \quad f \in \mathcal{E}.$$

Hence it follows that $Te^{i\Phi(Cf)} = e^{i\Phi(Cf)}T$, $f \in \mathcal{E}$. Recall that, for any dense subspace $\mathcal{D}$ of $\mathcal{H}$, $\{e^{i\Phi(u)} | u \in \mathcal{D}\}$ is irreducible (e.g., [2, Theorem 5.25(iv)]). Note that $\{e^{i\Phi(u)} | u \in \mathcal{D}\}$ is $*$ -invariant. Hence, by Lemma 2.1(ii), $\{e^{i\Phi(u)} | u \in \mathcal{D}\}' = \mathbb{C}I$. Therefore $\{e^{i\Phi(Cf)} | f \in \mathcal{E}\}' = \mathbb{C}I$. Hence $T = \alpha I$ for some $\alpha \in \mathbb{C}$. $\blacksquare$

4 Singular Bogoliubov Transformations and Representations of CCRs

4.1 Definitions

Let T and S be densely defined (not necessarily bounded) linear operators on $\mathcal{H}$ such that there exists a dense subspace $\mathcal{D} \subset D(T) \cap D(S)$ and the following equations hold:

$$\langle Tf, Tg \rangle - \langle Sf, Sg \rangle = \langle f, g \rangle, \quad (4.1)$$

$$\langle Tf, JSg \rangle = \langle Sf, JTg \rangle, \quad f, g \in \mathcal{D}, \quad (4.2)$$

where J is a conjugation on $\mathcal{H}$.

For each $f \in \mathcal{D}$, the operator

$$B_0(f) := A(Tf) + A(JSf)^*$$

is a densely defined linear operator with $D(B_0(f)) \supset \mathcal{F}_0(\mathcal{H})$ and $B_0(f)^*$ is densely defined with $D(B_0(f)^*) \supset \mathcal{F}_0(\mathcal{H})$. Hence $B_0(f)$ is closable. Therefore one can define a densely defined closed operator $B(f)$ on $\mathcal{F}_b(\mathcal{H})$ by

$$B(f) := \overline{B_0(f)}. \quad (4.3)$$

It is obvious that $B(f)$ and $B(f)^*$ leave $\mathcal{F}_0(\mathcal{H})$ invariant. Moreover, using (4.1) and (4.2), one can easily show that $B(\cdot)$ and $B(\cdot)^*$ satisfy the CCRs over $\mathcal{D}$ on $\mathcal{F}_0(\mathcal{H})$: for all $f, g \in \mathcal{D}$,

$$[B(f), B(g)^*] = \langle f, g \rangle, \quad [B(f), B(g)] = 0, \quad [B(f)^*, B(g)^*] = 0 \quad \text{on } \mathcal{F}_0(\mathcal{H}). \quad (4.4)$$

Therefore

$$\pi_B(\mathcal{D}) := (\mathcal{F}_b(\mathcal{H}), \mathcal{F}_0(\mathcal{H}), \{B(f), B(f)^* | f \in \mathcal{D}\}) \quad (4.5)$$

is a representation of the CCRs over $\mathcal{D}$.

The correspondence $T_B: (A(\cdot), A(\cdot)^*) \mapsto (B(\cdot), B(\cdot)^*)$ is a generalization of the standard bosonic Bogoliubov transformation (see, e.g., [11, 12, 13], [9, p.316], [6, §2] and references therein) in the sense that S or T may be unbounded and T_B is not necessarily invertible. Equation (4.1) implies that T is bounded if and only if S is bounded. Based on this property, we say that the Bogoliubov transformation T_B is *singular* if T or S is unbounded (then both T and S are unbounded).

4.2 Inequivalence to any direct sum representation of the Fock representation $\pi_F(\mathcal{D})$

We want to find conditions under which $\pi_B(\mathcal{D})$ is inequivalent to any direct sum representation of the Fock representation $\pi_F(\mathcal{D})$.

Lemma 4.1 *Let $\mathcal{D}$ be a dense subspace of $\mathcal{H}$ and $\pi_C(\mathcal{D}) := (\mathcal{F}, \mathcal{D}, \{C(f), C(f)^* | f \in \mathcal{D}\})$ be a representation of the CCRs over $\mathcal{D}$. Suppose that $\pi_C(\mathcal{D})$ is equivalent to a direct sum representation $\oplus_{n=1}^N \pi_F(\mathcal{D})$ of $\pi_F(\mathcal{D})$ with $N < \infty$ or $N = \infty$. Then there exists a non-zero vector $\Omega \in \cap_{f \in \mathcal{D}} D(C(f))$ such that $C(f)\Omega = 0$, $f \in \mathcal{D}$.*

Proof. By the present assumption, there exists a unitary operator U from $\mathcal{F}_b(\mathcal{H})$ to $\mathcal{G}_N := \oplus_{n=1}^N \mathcal{F}_b(\mathcal{H})$ such that $UC(f)U^{-1} = \oplus_{n=1}^N A(f)$, $f \in \mathcal{D}$. Let $\Omega_{\mathcal{H}}$ be the Fock vacuum in $\mathcal{F}_b(\mathcal{H})$:

$$\Omega_{\mathcal{H}}^{(0)} = 1, \quad \Omega_{\mathcal{H}}^{(n)} = 0, \quad n \geq 1.$$

Then one has

$$A(f)\Omega_{\mathcal{H}} = 0, \quad f \in \mathcal{H}. \quad (4.6)$$

It is easy to see that $\Psi_0 := \bigoplus_{n=1}^N n^{-1} \Omega_{\mathcal{H}}$ is a non-zero vector in $\mathcal{G}_N$ (note that, in the case $N = \infty$, $\sum_{n=1}^{\infty} \|n^{-1} \Omega_{\mathcal{H}}\|^2 = \sum_{n=1}^{\infty} n^{-2} < \infty$). By (4.6), $\Psi_0 \in D(\bigoplus_{n=1}^N A(f))$, $f \in \mathcal{D}$ and $(\bigoplus_{n=1}^N A(f))\Psi_0 = 0$. Hence, putting $\Omega := U^{-1}\Psi_0$, one sees that $\Omega \in \bigcap_{f \in \mathcal{D}} D(C(f))$ and $C(f)\Omega = 0$, $f \in \mathcal{D}$. ■

To state and prove results on the above mentioned inequivalence, we consider two cases separately: the one is the case where T is bounded and the other is the case where T is unbounded.

Lemma 4.2 *Assume that $\mathcal{H}$ is separable. Suppose that T is bounded with $T\mathcal{D}$ being dense in $\mathcal{H}$ and $\bar{S}$ is not Hilbert-Schmidt. Then there exist no non-zero vectors $\Omega \in \bigcap_{f \in \mathcal{D}} D(B(f))$ such that $B(f)\Omega = 0$, $f \in \mathcal{D}$.*

Proof. Suppose that there existed a non-zero vector $\Omega \in \bigcap_{f \in \mathcal{D}} D(B(f))$ that $B(f)\Omega = 0$, $f \in \mathcal{D}$. Then, for all $\Psi \in \mathcal{F}_0(\mathcal{H})$ and $f \in \mathcal{D}$, (3.3) holds. Hence, by Proposition 3.2, $\overline{ST_{\mathcal{D}}^{-1}}$ is Hilbert-Schmidt. By the present assumption, $\overline{T_{\mathcal{D}}} = \bar{T}$ is bounded. Hence $U := \overline{ST_{\mathcal{D}}^{-1}T_{\mathcal{D}}}$ is Hilbert-Schmidt. One has $U = \bar{S}$. Hence $\bar{S}$ is Hilbert-Schmidt. But this contradicts the assumption that $\bar{S}$ is not Hilbert-Schmidt. ■

Theorem 4.3 *Assume that $\mathcal{H}$ is separable. Suppose that T is bounded with $T\mathcal{D}$ being dense in $\mathcal{H}$ and $\bar{S}$ is not Hilbert-Schmidt. Then $\pi_B(\mathcal{D})$ is inequivalent to any direct sum representations of the Fock representation $\pi_F(\mathcal{D})$. In particular, if $\pi_B(\mathcal{D})$ is irreducible, then $\pi_B(\mathcal{D})$ is inequivalent to $\pi_F(\mathcal{D})$.*

Proof. Suppose that $\pi_B(\mathcal{D})$ were equivalent to a direct sum representation of $\pi_F(\mathcal{D})$. Then, by Lemma 4.1, there exists a non-zero vector $\Omega \in \bigcap_{f \in \mathcal{D}} D(B(f))$ such that $B(f)\Omega = 0$, $f \in \mathcal{D}$. But this contradicts Lemma 4.2. ■

We next consider the case where T is unbounded.

Lemma 4.4 *Assume that $\mathcal{H}$ is separable. Suppose that T is unbounded and $T\mathcal{D}$ is dense in $\mathcal{H}$. Then there exist no non-zero vectors $\Omega \in \bigcap_{f \in \mathcal{D}} D(B(f))$ such that $B(f)\Omega = 0$, $f \in \mathcal{D}$.*

Proof. Suppose that there existed a non-zero vector $\Omega \in \bigcap_{f \in \mathcal{D}} D(B(f))$ such that $B(f)\Omega = 0$, $f \in \mathcal{D}$. Then (3.5) holds with $X = T$ and $Y = S$. But this contradicts Proposition 3.4. ■

The following theorem is one of the main results in the abstract framework of the present paper.

Theorem 4.5 *Assume that $\mathcal{H}$ is separable. Suppose that T is unbounded and $T\mathcal{D}$ is dense in $\mathcal{H}$. Then $\pi_B(\mathcal{D})$ is inequivalent to any direct sum representation of the Fock representation $\pi_F(\mathcal{D})$. In particular, if $\pi_B(\mathcal{D})$ is irreducible, then $\pi_B(\mathcal{D})$ is inequivalent to $\pi_F(\mathcal{D})$.*

Proof. Suppose that $\pi_B(\mathcal{D})$ were equivalent to a direct sum representation of $\pi_F(\mathcal{D})$. Then, by Lemma 4.1, there exists a non-zero vector $\Omega \in \bigcap_{f \in \mathcal{D}} D(B(f))$ such that $B(f)\Omega = 0$, $f \in \mathcal{D}$. But this contradicts Lemma 4.4. ■

4.3 Irreducibility

In this subsection, we formulate sufficient conditions for $\pi_B(\mathcal{D})$ to be irreducible. To be as general as possible, we consider two cases. The first one uses the following condition:

Assumption (I) There exists a dense subspace $\mathcal{D}_1$ of $\mathcal{H}$ such that the following (i) and (ii) hold:

$$(i) \quad \mathcal{D}_1 \subset D(TT^*) \cap D(JSS^*J) \cap D(ST^*) \cap D(JTS^*J).$$

$$(ii) \quad T^*\mathcal{D}_1 \subset \mathcal{D}, (S^*J)\mathcal{D}_1 \subset \mathcal{D} \text{ and}$$

$$TT^* - JSS^*J = I, \quad ST^*J = JTS^* \quad \text{on } \mathcal{D}_1.$$

Lemma 4.6 Suppose that Assumption (I) holds. Then, for all $f \in \mathcal{D}_1$,

$$A(f) = B(T^*f) - B(S^*Jf)^* \quad \text{on } \mathcal{F}_0(\mathcal{H}). \quad (4.7)$$

Proof. One needs only to compute the right hand side of (4.7) using (4.3) and Assumption (I). $\blacksquare$

Theorem 4.7 Suppose in addition to (4.1) and (4.2) that Assumption (I) holds. Then $\pi_B(\mathcal{D})$ is irreducible.

Proof. Let $Z \in \{B(f), B(f)^* | f \in \mathcal{D}\}'$. Then $Z \in \mathfrak{B}(\mathcal{F}_b(\mathcal{H}))$ and for all $f \in \mathcal{D}$, $ZB(f) \subset B(f)Z$, $ZB(f)^* \subset B(f)^*Z$. By using (4.7), one can show that, for all $\Phi, \Psi \in \mathcal{F}_0(\mathcal{H})$ and $f \in \mathcal{D}_1$,

$$\langle \Phi, ZA(f)\Psi \rangle = \langle A(f)^*\Phi, Z\Psi \rangle.$$

Since $\mathcal{F}_0(\mathcal{H})$ is a core for $A(f)^*$, it follows that $Z\Psi \in D(A(f)^{**}) = D(A(f))$ and $A(f)Z\Psi = ZA(f)\Psi$. Since $\mathcal{F}_0(\mathcal{H})$ is a core for $A(f)$, this implies that $ZA(f) \subset A(f)Z$. Similarly one has $ZA(f)^* \subset A(f)^*Z$ for all $f \in \mathcal{D}_1$. Hence $Z \in \{A(f), A(f)^* | f \in \mathcal{D}_1\}'$. By Theorem 3.1, $\{A(f), A(f)^* | f \in \mathcal{D}_1\}' = \mathbb{C}I$. Hence $\{B(f), B(f)^* | f \in \mathcal{D}\}' = \mathbb{C}I$. Therefore, by Lemma 2.1(i), $\{B(f), B(f)^* | f \in \mathcal{D}\}$ is irreducible. $\blacksquare$

Another criterion for the irreducibility of $\pi_B(\mathcal{D})$ is given as follows:

Theorem 4.8 Suppose in addition to (4.1) and (4.2) that

$$JS \subset SJ, \quad JT \subset TJ.$$

Let $\mathcal{D}_J := \mathcal{D} \cap \mathcal{H}_J$, $R := (1+i)T + (1-i)S$ and suppose that $R\mathcal{D}_J$ is dense in $\mathcal{H}$. Then $\pi_B(\mathcal{D})$ is irreducible.

Proof. Let $Z \in \{B(f), B(f)^* | f \in \mathcal{D}\}'$. Then it is easy to see that, for all $f \in \mathcal{D}_J$,

$$\frac{1}{\sqrt{2}}(B(f)^* + B(f)) = \phi_{T+S}(f), \quad \frac{i}{\sqrt{2}}(B(f)^* - B(f)) = \pi_{T-S}(f)$$

on $\mathcal{F}_0(\mathcal{H})$. Hence, in the same way as in the proof of Theorem 4.8, one can show that $Z\phi_{T+S}(f) \subset \phi_{T+S}(f)Z$ and $Z\pi_{T-S}(f) \subset \pi_{T-S}(f)Z$. Since $\phi_{T+S}(f)$ and $\pi_{T-S}(f)$ are self-adjoint, it follows that $Ze^{i\phi_{T+S}(f)} = e^{i\phi_{T+S}(f)}Z$ and $Ze^{i\pi_{T-S}(f)} = e^{i\pi_{T-S}(f)}Z$. Note that $T + S + i(T - S) = R$. Hence, by Lemma 3.5, $Z = \alpha I$ for some $\alpha \in \mathbb{C}$. Thus $\pi_B(\mathcal{D})$ is irreducible. $\blacksquare$

5 A General Class of Singular Bogoliubov Transformations

In this section, we introduce a general class of singular Bogoliubov transformations. Let K and L be injective (not necessarily bounded) symmetric operators on a Hilbert space $\mathcal{H}$ such that

$$\mathcal{D}_{K,L} := D(K^{-1}L) \cap D(KL^{-1}) \quad (5.1)$$

is dense in $\mathcal{H}$ and, for a conjugation J on $\mathcal{H}$,

$$JK \subset KJ, \quad JL \subset LJ. \quad (5.2)$$

Then one can define densely defined linear operators

$$T_{\pm} := \frac{1}{2}(K^{-1}L \pm KL^{-1}) \quad (5.3)$$

with $D(T_{\pm}) = \mathcal{D}_{K,L}$. It follows from (5.2) that

$$JT_{\pm} \subset T_{\pm}J.$$

Lemma 5.1 *For all $f, g \in \mathcal{D}_{K,L}$,*

$$\langle T_+f, T_+g \rangle - \langle T_-f, T_-g \rangle = \langle f, g \rangle, \quad (5.4)$$

$$\langle T_+f, JT_-g \rangle = \langle T_-f, JT_+g \rangle. \quad (5.5)$$

Proof. This follows from direct computations, where the symmetry of K and L are used. ■

Remark 5.2 Operators of type $T_{\pm}$ with K and L being injective self-adjoint operators such that $K^{-1}L$ and KL^{-1} are *bounded* are introduced in [1, Lemma 5.8]. But here we are mainly interested in the case where $T_{\pm}$ are *unbounded*.

Let $\mathcal{D}$ be a dense subspace of $\mathcal{H}$ such that $\mathcal{D} \subset \mathcal{D}_{K,L}$. Then, as in Section 4, one can define a densely defined closed linear operator

$$B_{K,L}(f) := \overline{A(T_+f) + A(JT_-f)^*}, \quad f \in \mathcal{D} \quad (5.6)$$

and see that

$$\pi_{K,L}(\mathcal{D}) := (\mathcal{F}_b(\mathcal{H}), \mathcal{F}_0(\mathcal{H}), \{B_{K,L}(f), B_{K,L}(f)^* | f \in \mathcal{D}\}) \quad (5.7)$$

is a representation of the CCRs over $\mathcal{D}$.

Note that, if T_+ or T_- is unbounded, then the correspondence: $(A(\cdot), A(\cdot)^*) \mapsto (B_{K,L}(\cdot), B_{K,L}(\cdot)^*)$ is a singular Bogoliubov transformation. Thus, for each pair (K, L) of injective symmetric operators with additional properties, there exists a singular Bogoliubov transformation.

A simple application of Lemma 4.4 and Theorem 4.5 yields the following theorem:

Theorem 5.3 *Let $\mathcal{H}$ be separable. Suppose that T_+ is unbounded and $T_+\mathcal{D}$ is dense. Then:*

- (i) *There exists no non-zero vector $\Omega \in \cap_{f \in \mathcal{D}} D(B_{K,L}(f))$ such that $B_{K,L}(f)\Omega = 0$, $f \in \mathcal{D}$.*
- (ii) *The representation $\pi_{K,L}(\mathcal{D})$ is inequivalent to any direct sum representation of the Fock representation $\pi_F(\mathcal{D})$. In particular, if $\pi_{K,L}(\mathcal{D})$ is irreducible, then $\pi_{K,L}(\mathcal{D})$ is inequivalent to $\pi_F(\mathcal{D})$.*

With regard to irreducibility of $\pi_{K,L}(\mathcal{D})$, we have the following result.

Theorem 5.4 *Suppose that $(K^{-1}L + iKL^{-1})(\mathcal{D} \cap \mathcal{H}_J)$ is dense in $\mathcal{H}$. Then $\pi_{K,L}(\mathcal{D})$ is irreducible.*

Proof. We need only to apply Theorem 4.8 to the case where $T = T_+$, $S = T_-$. In this case $R = K^{-1}L + iKL^{-1}$. ■

6 An Example

In this section we show by an explicit construction that there exists a pair (K, L) of symmetric operators and a conjugation J , satisfying conditions described in Section 5 so that the conclusions of Theorems 5.3 and 5.4 hold.

The example given below is related to a construction of two quantum scalar fields on the finite box

$$\Lambda := (0, L)^{d-1} \times (0, L_d) = \{\mathbf{x} = (x_1, \dots, x_d) \mid x_1, \dots, x_{d-1} \in (0, L), x_d \in (0, L_d)\} \quad (6.1)$$

in the d -dimensional Euclidean space $\mathbb{R}^d$ with $d \geq 2$, $L > 0$ and $L_d > 0$.

Remark 6.1 There may be another choice of Λ . For example $\Lambda = \mathbb{R}^{d-1} \times (0, L_d)$. But, in this paper, we restrict ourselves to the case given above.

We denote by Δ_D the Dirichlet Laplacian for Λ on $L^2(\Lambda)$ (see, e.g., [10, p.263] for the definition of Δ_D).² The operator $-\Delta_D$ is a non-negative self-adjoint operator. Let

$$\Gamma := \left\{ \mathbf{k} = (k_1, \dots, k_d) \mid k_j \in \frac{\pi}{L}\mathbb{N}, j = 1, \dots, d-1, k_d \in \frac{\pi}{L_d}\mathbb{N} \right\}$$

where $\alpha\mathbb{N} := \{\alpha n \mid n \in \mathbb{N}\}$ ($\alpha \in \mathbb{R}$). It is well known that the spectrum $\sigma(-\Delta_D)$ of $-\Delta_D$ is purely discrete with

$$\sigma(-\Delta_D) = \{\mathbf{k}^2 \mid \mathbf{k} \in \Gamma\}.$$

²One can consider also Laplacians with other boundary conditions. But, in the present paper, we treat only the Dirichlet Laplacian for simplicity.

An eigenvector of $-\Delta_D$ with eigenvalue $\mathbf{k}^2$ is given by

$$\varphi_{\mathbf{k}}(\mathbf{x}) := \prod_{j=1}^d \varphi_{k_j}(x_j), \quad \mathbf{x} = (x_1, \dots, x_d) \in \Lambda \quad (6.2)$$

with

$$\varphi_{k_j}(x_j) := \sqrt{\frac{2}{L}} \sin(k_j x_j), \quad x_j \in (0, L), \quad j = 1, \dots, d-1, \quad (6.3)$$

$$\varphi_{k_d}(x_d) := \sqrt{\frac{2}{L_d}} \sin(k_d x_d), \quad x_d \in (0, L_d). \quad (6.4)$$

It is well known that $\{\varphi_{\mathbf{k}} | \mathbf{k} \in \Gamma\}$ is a complete orthonormal system (CONS) of $L^2(\Lambda)$.

Let $m \geq 0$ be a constant and

$$h := (-\Delta_D + m^2)^{1/2}, \quad (6.5)$$

Then we have

$$h\varphi_{\mathbf{k}} = \omega(\mathbf{k})\varphi_{\mathbf{k}}, \quad \mathbf{k} \in \Gamma, \quad (6.6)$$

with

$$\omega(\mathbf{k}) := \sqrt{\mathbf{k}^2 + m^2}. \quad (6.7)$$

Hence the spectrum of h is purely discrete with

$$\sigma(h) = \{\omega(\mathbf{k}) | \mathbf{k} \in \Gamma\}. \quad (6.8)$$

Note that

$$0 < \omega(\mathbf{k})^{-1} \leq \frac{1}{\sqrt{\left(\frac{\pi}{L}\right)^2 (d-1) + \left(\frac{\pi}{L_d}\right)^2}}, \quad \mathbf{k} \in \Gamma. \quad (6.9)$$

Hence $h^{-1} \in \mathfrak{B}(L^2(\Lambda))$.

We set

$$L_{\text{real}}^2(\Lambda) := \{f \in L^2(\Lambda) | f^* = f\},$$

the real Hilbert space of real elements in $L^2(\Lambda)$. For all $\alpha \in \mathbb{R}$ and $f \in D(h^\alpha)$, we have

$$h^\alpha f = \sum_{\mathbf{k} \in \Gamma} \omega(\mathbf{k})^\alpha \langle \varphi_{\mathbf{k}}, f \rangle \varphi_{\mathbf{k}}$$

in the sense of $L^2(\Lambda)$ -convergence. Hence

$$h^\alpha D(h^\alpha) \cap L_{\text{real}}^2(\Lambda) \subset L_{\text{real}}^2(\Lambda). \quad (6.10)$$

We next consider the case where a “wall”

$$W_a := \{\mathbf{x} \in \Lambda | x_d = a\} \quad (6.11)$$

perpendicular to the x_d -axis is placed in Λ with $0 < a < L_d$. In this case Λ is decomposed as follows:

$$\Lambda = \Lambda_1 \cup W_a \cup \Lambda_2, \quad (6.12)$$

where

$$\Lambda_1 := \{\mathbf{x} \in \Lambda | 0 < x_d < a\}, \quad \Lambda_2 := \{\mathbf{x} \in \Lambda | a < x_d < L_d\}. \quad (6.13)$$

This decomposition of the box Λ induces the orthogonal decomposition

$$L^2(\Lambda) = L^2(\Lambda_1) \oplus L^2(\Lambda_2) \quad (6.14)$$

of the Hilbert space $L^2(\Lambda)$. But one should note that the Dirichlet Laplacian Δ_D cannot be reduced by $L^2(\Lambda_\ell)$ ($\ell = 1, 2$).

In what follows, we assume the following:

Assumption (a) $L^2/L_d^2 \in \mathbb{Q}$ (the set of rational numbers) and $a^2/L_d^2 \notin \mathbb{Q}$

Let Λ_1 and Λ_2 be given by (6.13) and Δ_ℓ be the Dirichlet Laplacian for Λ_ℓ ($\ell = 1, 2$). For each $\mathbf{k} \in \Gamma$ and $\ell = 1, 2$, we define functions $\psi_{\mathbf{k}}^{(\ell)}$ on Λ_ℓ as follows:

$$\psi_{\mathbf{k}}^{(1)}(\mathbf{x}) := \left(\prod_{j=1}^{d-1} \varphi_{k_j}(x_j) \right) \psi_{k_d}^{(1)}(x_d), \quad \mathbf{x} \in \Lambda_1, \quad (6.15)$$

$$\psi_{\mathbf{k}}^{(2)}(\mathbf{x}) := \left(\prod_{j=1}^{d-1} \varphi_{k_j}(x_j) \right) \psi_{k_d}^{(2)}(x_d), \quad \mathbf{x} \in \Lambda_2, \quad (6.16)$$

where

$$\psi_{k_d}^{(1)}(x_d) := \sqrt{\frac{2}{a}} \sin \frac{L_d k_d x_d}{a}, \quad x_d \in (0, a), \quad (6.17)$$

$$\psi_{k_d}^{(2)}(x_d) := \sqrt{\frac{2}{L_d - a}} \sin \frac{L_d k_d (x_d - a)}{L_d - a}, \quad x_d \in (a, L_d). \quad (6.18)$$

The set $\{\psi_{\mathbf{k}}^{(\ell)}\}_{\mathbf{k} \in \Gamma}$ is a CONS of $L^2(\Lambda_\ell)$.

Using the Dirichlet Laplacian Δ_ℓ for Λ_ℓ ($\ell = 1, 2$), we introduce a non-negative self-adjoint operator

$$h_{a,\ell} := (-\Delta_\ell + m^2)^{1/2} \quad (6.19)$$

on $L^2(\Lambda_\ell)$ and define

$$h_a := h_{a,1} \oplus h_{a,2} \quad (6.20)$$

relative to the decomposition (6.14).

Let

$$\begin{aligned}\omega_1(\mathbf{k}) &:= \sqrt{k_1^2 + \cdots + k_{d-1}^2 + (L_d k_d/a)^2 + m^2}, \\ \omega_2(\mathbf{k}) &:= \sqrt{k_1^2 + \cdots + k_{d-1}^2 + (L_d k_d/(L_d - a))^2 + m^2}, \quad \mathbf{k} \in \Gamma.\end{aligned}$$

Then we have

$$h_{a,\ell}\psi_{\mathbf{k}}^{(\ell)} = \omega_{\ell}(\mathbf{k})\psi_{\mathbf{k}}^{(\ell)}, \quad \mathbf{k} \in \Gamma. \quad (6.21)$$

By (6.14), each $f \in L^2(\Lambda)$ is written as

$$f = f^{(1)} + f^{(2)}$$

or $f = (f^{(1)}, f^{(2)})$, where

$$f^{(\ell)} := \chi_{\Lambda_{\ell}} f \in L^2(\Lambda_{\ell})$$

with $\chi_{\Lambda_{\ell}}$ being the characteristic function of Λ_{ℓ} .

The direct sum operator

$$-\Delta_{12} := (-\Delta_1) \oplus (-\Delta_2)$$

is a non-negative self-adjoint operator on $L^2(\Lambda)$. By functional calculus, we have

$$(-\Delta_{12})^{1/2} = (-\Delta_1)^{1/2} \oplus (-\Delta_2)^{1/2}. \quad (6.22)$$

Lemma 6.2 $D((-\Delta_{12})^{1/2}) \subset D((-\Delta_D)^{1/2})$ and, for all $f \in D((-\Delta_{12})^{1/2})$,

$$\|(-\Delta_D)^{1/2} f\| = \|(-\Delta_{12})^{1/2} f\|. \quad (6.23)$$

In particular, $(-\Delta_D)^{1/2}(-\Delta_{12})^{-1/2}$ is in $\mathfrak{B}(L^2(\Lambda))$.

Proof. Let $f = f^{(1)} + f^{(2)} \in D((-\Delta_{12})^{1/2})$. Then, by the definition of Δ_{ℓ} , there exists a sequence $\{f_n^{(\ell)}\}_{n=1}^{\infty}$ in $C_0^{\infty}(\Lambda_{\ell})$ such that $f_n^{(\ell)} \rightarrow f^{(\ell)}$ ($n \rightarrow \infty$) and

$$\|(-\Delta_{\ell})^{1/2} f^{(\ell)}\|^2 = \lim_{n \rightarrow \infty} \sum_{j=1}^d \|\partial_j f_n^{(\ell)}\|^2,$$

where $\partial_j := \partial/\partial x_j$. Hence, letting $f_n := f_n^{(1)} + f_n^{(2)}$, we see that $f_n \in C_0^{\infty}(\Lambda)$ and $f_n \rightarrow f$ ($n \rightarrow \infty$),

$$\lim_{n \rightarrow \infty} \sum_{j=1}^d \|\partial_j f_n\|^2 = \|(-\Delta_1)^{1/2} f^{(1)}\|^2 + \|(-\Delta_2)^{1/2} f^{(2)}\|^2 = \|(-\Delta_{12})^{1/2} f\|^2.$$

Therefore $f \in D((-\Delta_D)^{1/2})$ (hence $D((-\Delta_{12})^{1/2}) \subset D((-\Delta_D)^{1/2})$) and (6.23) holds. $\blacksquare$

We next study relations between h and h_a .

Lemma 6.3

- (i) $D(h_a) \subset D(h)$ and hh_a^{-1} is in $\mathfrak{B}(L^2(\Lambda))$.
- (ii) $D(h_a^{1/2}) \subset D(h^{1/2})$ and $h^{1/2}h_a^{-1/2}$ is in $\mathfrak{B}(L^2(\Lambda))$.

Proof. (i) It follows from functional calculus that

$$D(h) = D((-\Delta_D)^{1/2}), \quad D(h_a) = D((-\Delta_{12})^{1/2}).$$

Hence, by Lemma 6.2, $D(h_a) \subset D(h)$. Hence $D(hh_a^{-1}) = L^2(\Lambda)$ and $hh_a^{-1} \in \mathfrak{B}(L^2(\Lambda))$.³

(ii) By part (i), h is h_a -bounded. Since h and h_a are non-negative self-adjoint operators, it follows from a general theorem ([8, Theorem X.18(a)]) that $h^{1/2}$ is $h_a^{1/2}$ -bounded. Hence $D(h_a^{1/2}) \subset D(h^{1/2})$ and $h^{1/2}h_a^{-1/2}$ is bounded with $D(h^{1/2}h_a^{-1/2}) = L^2(\Lambda)$. ■

For convenience, we extend each eigenfunction $\psi_{\mathbf{k}}^{(\ell)}$ of $h_{a,\ell}$ ($\mathbf{k} \in \Gamma$) to a function on Λ in the following way:

$$\tilde{\psi}_{\mathbf{k}}^{(\ell)}(\mathbf{x}) := \begin{cases} \psi_{\mathbf{k}}^{(\ell)}(\mathbf{x}) & \text{if } \mathbf{x} \in \Lambda_\ell \\ 0 & \text{if } \mathbf{x} \in \Lambda \setminus \Lambda_\ell \end{cases}. \quad (6.24)$$

Similarly we denote by $\tilde{\psi}_{k_d}^{(\ell)}$ ($\ell = 1, 2$) the extensions of the functions $\psi_{k_d}^{(1)}$ and $\psi_{k_d}^{(2)}$ to $(0, L_d)$.

By functional calculus, we have for all $\alpha > 0$

$$h_a^\alpha \tilde{\psi}_{\mathbf{k}}^{(\ell)} = \omega_\ell(\mathbf{k})^\alpha \tilde{\psi}_{\mathbf{k}}^{(\ell)}, \quad \ell = 1, 2, \mathbf{k} \in \Gamma. \quad (6.25)$$

For each $(\mathbf{k}, \mathbf{p}) \in \Gamma \times \Gamma$ and $\ell = 1, 2$, we define

$$\gamma_{\mathbf{k}\mathbf{p}}^{(\ell)} := \langle \varphi_{\mathbf{k}}, \tilde{\psi}_{\mathbf{p}}^{(\ell)} \rangle. \quad (6.26)$$

It follows from (6.2), (6.15) and (6.16) that

$$\gamma_{\mathbf{k}\mathbf{p}}^{(\ell)} = \left(\prod_{j=1}^{d-1} \delta_{k_j p_j} \right) \langle \varphi_{k_d}, \tilde{\psi}_{p_d}^{(\ell)} \rangle. \quad (6.27)$$

We set

$$c_1 := \frac{L_d}{a}, \quad c_2 := \frac{L_d}{L_d - a}. \quad (6.28)$$

Lemma 6.4 For all $k_d, p_d \in (\pi/L_d)\mathbb{N}$,

$$\langle \varphi_{k_d}, \tilde{\psi}_{p_d}^{(1)} \rangle = \frac{2}{\sqrt{L_d a}} \frac{(-1)^{L_d p_d} c_1 p_d \sin(a k_d)}{k_d^2 - c_1^2 p_d^2}, \quad (6.29)$$

$$\langle \varphi_{k_d}, \tilde{\psi}_{p_d}^{(2)} \rangle = -\frac{2}{\sqrt{L_d(L_d - a)}} \frac{c_2 p_d \sin(a k_d)}{k_d^2 - c_2^2 p_d^2}. \quad (6.30)$$

³The following fact is well known (which follows from the closed graph theorem): let A and B be closed linear operators on a Banach space $\mathcal{X}$. Suppose that $D(A) \subset D(B)$ and A is bijective. Then BA^{-1} is in $\mathfrak{B}(\mathcal{X})$.

Proof. By (6.4) and (6.17), we have

$$\left\langle \varphi_{k_d}, \tilde{\psi}_{p_d}^{(1)} \right\rangle = \frac{2}{\sqrt{L_d a}} \int_0^a \sin(k_d x_d) \sin(c_1 p_d x_d) dx_d.$$

Then, by direct computations, we obtain (6.29). Similarly one can prove (6.30). $\blacksquare$

Lemma 6.5 $D(h_a^{1/2}) \subsetneq D(h^{1/2})$.

Proof. By functional calculus, for all $\mathbf{k} \in \Gamma$, $\varphi_{\mathbf{k}} \in D(h^{1/2})$ with

$$h^{1/2} \varphi_{\mathbf{k}} = \omega(\mathbf{k})^{1/2} \varphi_{\mathbf{k}}.$$

It is sufficient to show that, for some $\mathbf{k}_0 \in \Gamma$, $\varphi_{\mathbf{k}_0} \notin D(h_a^{1/2})$. Let $\mathbf{k} \in \Gamma$ be such that $\varphi_{\mathbf{k}} \in D(h_a^{1/2})$. Then, by the fact that $\{\tilde{\psi}_{\mathbf{p}}^{(\ell)} | \ell = 1, 2, \mathbf{p} \in \Gamma\}$ is a CONS of $L^2(\Lambda)$, we have

$$s_{\mathbf{k}} := \sum_{\ell=1}^2 \sum_{\mathbf{p} \in \Gamma} \left| \left\langle \tilde{\psi}_{\mathbf{p}}^{(\ell)}, h_a^{1/2} \varphi_{\mathbf{k}} \right\rangle \right|^2 = \|h_a^{1/2} \varphi_{\mathbf{k}}\|^2 < \infty.$$

By the symmetry of $h_a^{1/2}$ and (6.25), we have

$$\left\langle \tilde{\psi}_{\mathbf{p}}^{(\ell)}, h_a^{1/2} \varphi_{\mathbf{k}} \right\rangle = \left\langle h_a^{1/2} \tilde{\psi}_{\mathbf{p}}^{(\ell)}, \varphi_{\mathbf{k}} \right\rangle = \omega_{\ell}(\mathbf{p})^{1/2} \gamma_{\mathbf{k}\mathbf{p}}^{(\ell)}.$$

Hence

$$s_{\mathbf{k}} = \sum_{\ell=1}^2 \sum_{\mathbf{p} \in \Gamma} \omega_{\ell}(\mathbf{p}) \left| \gamma_{\mathbf{k}\mathbf{p}}^{(\ell)} \right|^2.$$

Using (6.27), we have

$$s_{\mathbf{k}} = \sum_{\ell=1}^2 \sum_{p_d \in (\pi/L_d)\mathbb{N}} \omega_{\ell}(\tilde{\mathbf{k}}, p_d) \left| \left\langle \varphi_{k_d}, \tilde{\psi}_{p_d}^{(\ell)} \right\rangle \right|^2,$$

where

$$\tilde{\mathbf{k}} := (k_1, \dots, k_{d-1}) \in \left(\frac{\pi}{L} \mathbb{N} \right)^{d-1}.$$

By Lemma 6.4, we have

$$\lim_{p_d \rightarrow \infty} \left| p_d \left\langle \varphi_{k_d}, \tilde{\psi}_{p_d}^{(1)} \right\rangle \right|^2 = \frac{4 \sin^2(ak_d)}{L_d a c_1^2}. \quad \lim_{p_d \rightarrow \infty} \left| p_d \left\langle \varphi_{k_d}, \tilde{\psi}_{p_d}^{(2)} \right\rangle \right|^2 = \frac{4 \sin^2(ak_d)}{L_d (L_d - a) c_2^2}.$$

Now let $\mathbf{k}_0 := (\tilde{\mathbf{k}}, k_d)$ with $\tilde{\mathbf{k}} \in ((\pi/L)\mathbb{N})^{d-1}$ arbitrary and $k_d = \pi/L_d$. Then $0 < ak_d < \pi$. Hence $\sin ak_d \neq 0$. Therefore there exists a constant $C > 0$ such that

$$\left| \left\langle \varphi_{k_d}, \tilde{\psi}_{p_d}^{(\ell)} \right\rangle \right|^2 \geq \frac{C}{c_{\ell} p_d^2}.$$

Note that

$$\omega_\ell(\tilde{\mathbf{k}}, p_d) \geq c_\ell p_d.$$

Hence

$$s_{\mathbf{k}_0} \geq C \sum_{p_d \in (\pi/L_d)\mathbb{N}} \frac{1}{p_d} = \infty.$$

Thus $\varphi_{\mathbf{k}_0} \notin D(h_a^{1/2})$. ■

The following fact shows a singular nature of the pair (h, h_a) of one-particle Hamiltonians:

Lemma 6.6 *The operator $h^{-1/2}h_a^{1/2}$ is unbounded.*

Proof. Let $T := h^{-1/2}h_a^{1/2}$. We prove the unboundedness of T by reductio ad absurdum. Suppose that T were bounded. Since $D(T) = D(h_a^{1/2})$, T is densely defined. Hence it is closable and the closure $\bar{T}$ is in $\mathfrak{B}(L^2(\Lambda))$. We have $T^* = (\bar{T})^*$. Hence $D(T^*) = L^2(\Lambda)$. Since $h^{-1/2}$ is bounded with $D(h^{-1/2}) = L^2(\Lambda)$, it follows that $T^* = (h_a^{1/2})^*(h^{-1/2})^* = h_a^{1/2}h^{-1/2}$. This implies that $D(h^{1/2}) \subset D(h_a^{1/2})$ and $D(T^*) = h^{1/2}D(h_a^{1/2})$. But this contradicts Lemma 6.5. ■

Lemma 6.7 *Let*

$$S_\pm := \frac{1}{2}(h^{-1/2}h_a^{1/2} \pm h^{1/2}h_a^{-1/2}). \quad (6.31)$$

Then $S_\pm$ are unbounded.

Proof. This follows from Lemma 6.3(ii) and Lemma 6.6. ■

Note that

$$\mathcal{D}_a := D(S_+) = D(S_-) = D(h_a^{1/2}). \quad (6.32)$$

We denote by C_Λ the complex conjugation on $L^2(\Lambda)$:

$$C_\Lambda f := f^*, \quad f \in L^2(\Lambda).$$

Then it is easy to see that, for all $\alpha \in \mathbb{R}$,

$$C_\Lambda h^\alpha \subset h^\alpha C_\Lambda, \quad C_\Lambda h_a^\alpha \subset h_a^\alpha C_\Lambda.$$

Hence

$$C_\Lambda S_\pm \subset S_\pm C_\Lambda. \quad (6.33)$$

The next lemma follows from Lemma 5.1 and (6.33):

Lemma 6.8 For all $f, g \in \mathcal{D}_a$, the following equations hold:

$$\langle S_+ f, S_+ g \rangle - \langle S_- f, S_- g \rangle = \langle f, g \rangle, \quad (6.34)$$

$$\langle S_+ f, S_- g \rangle = \langle S_- f, S_+ g \rangle. \quad (6.35)$$

Lemma 6.9 The range $\text{Ran } S_+$ of S_+ is dense in $L^2(\Lambda)$.

Proof. It is sufficient to show that the orthogonal complement $(\text{Ran } S_+)^\perp$ of $\text{Ran } S_+$ is $\{0\}$. Let $g \in (\text{Ran } S_+)^\perp$. Then, for all $f \in D(h_a^{1/2})$, $\langle S_+ f, g \rangle = 0$, which implies that $\langle h_a^{1/2} f, h^{-1/2} g \rangle = -\langle G f, g \rangle$ with $G := h^{1/2} h_a^{-1/2}$. By Lemma 6.3(ii), G is bounded. Hence $\langle G f, g \rangle = \langle f, G^* g \rangle$. Hence $h^{-1/2} g \in D(h_a^{1/2})$ and $h_a^{1/2} h^{-1/2} g = -G^* g$. Hence $g = -G G^* g$, which implies that $\|g\|^2 = -\|G^* g\|^2 \leq 0$. Therefore $g = 0$. Thus $(\text{Ran } S_+)^\perp = \{0\}$. ■

Lemma 6.10 $(h^{-1/2} h_a^{1/2} + i h^{1/2} h_a^{-1/2})(\mathcal{D}_a \cap L_{\text{real}}^2(\Lambda))$ is dense in $L^2(\Lambda)$.

Proof. Suppose that there exists an element $g \in L^2(\Lambda)$ such that, for all $f \in \mathcal{D}_a \cap L_{\text{real}}(\Lambda)$, $\langle g, (h^{-1/2} h_a^{1/2} + i h^{1/2} h_a^{-1/2}) f \rangle = 0 \dots (*)$. Since $C_\Lambda h_a^{1/2} \subset h_a^{1/2} C_\Lambda$, it follows that, each $f \in \mathcal{D}_a$ is written $f = f_1 + i f_2$ with f_1 and f_2 being in $\mathcal{D}_a \cap L_{\text{real}}(\Lambda)$. Hence $(*)$ extends to all $f \in \mathcal{D}_a$. Let $G := h^{1/2} h_a^{-1/2}$. Then, by Lemma 6.3(ii), G is bounded. Hence it follows that, for all $f \in \mathcal{D}_a = D(h_a^{1/2})$, $\langle h^{-1/2} g, h_a^{1/2} f \rangle = \langle i G^* g, f \rangle$. This implies that $h^{-1/2} g \in D(h_a^{1/2})$ and $h_a^{1/2} h^{-1/2} g = i G^* g$. Hence $g = i G G^* g$, which implies that $\|g\|^2 = i \|G^* g\|^2$. Hence $\|g\|^2 = 0$, implying that $g = 0$. Thus the desired result follows. ■

We now define

$$b(f) := \overline{a(S_+ f) + a(C_\Lambda S_- f)^*}, \quad f \in \mathcal{D}_a. \quad (6.36)$$

Theorem 6.11 The triple

$$\pi_b(\mathcal{D}_a) := (\mathcal{F}_b(L^2(\Lambda)), \mathcal{F}_0(L^2(\Lambda)), \{b(f), b(f)^* | f \in \mathcal{D}_a\}).$$

is a representation of the CCRs over $\mathcal{D}_a$. Moreover, the following (i) and (ii) hold:

- (i) There exists no non-zero vector $\Omega \in \cap_{f \in \mathcal{D}_a} D(b(f))$ such that $b(f)\Omega = 0$, $f \in \mathcal{D}_a$.
- (ii) The representation $\pi_b(\mathcal{D}_a)$ is irreducible and inequivalent to the Fock representation $\pi_F(\mathcal{D}_a)$.

Proof. (i) By Lemma 6.9, we can apply Theorem 5.3(i) to the case where $K = h^{1/2}$, $L = h_a^{1/2}$, $J = C_\Lambda$ and $T_+ = S_+$. Then the desired result follows.

(ii) The irreducibility of $\pi_b(\mathcal{D}_a)$ follows from Lemma 6.10 and an application of Theorem 5.4. The inequivalence of $\pi_b(\mathcal{D}_a)$ to $\pi_F(\mathcal{D}_a)$ is due to an application of Theorem 5.3 (ii). ■

Acknowledgment

The author thanks the referee for constructive comments on the original manuscript, which have been helpful in writing this revised version.

References

- [1] A. Arai, A family of inequivalent Weyl representations of canonical commutation relations with applications to quantum field theory. *Rev. Math. Phys.* **28** (2016), 1650007 (26 pages).
- [2] A. Arai, *Analysis on Fock Spaces and Mathematical Theory of Quantum Fields*, World Scientific, Singapore, 2018.
- [3] H. B. G. Casimir, On the attraction between two perfectly conducting plates, *Proc. Koninklijke Nederlandse Akademie van Wetenschappen* **51** (1948), 793–795.
- [4] C. Dappiaggi, G. Nosari and N. Pinamonti, The Casimir effect from the point of view of algebraic quantum field theory, *Math. Phys. Anal. Geom.* **19** (2016), 12 (44 pages).
- [5] H. Ezawa, K. Nakamura and K. Watanabe, The Casimir force from Lorentz’s, in *Frontiers in Quantum Physics* (S. C. Lim, R. Abd-Shukor, K. H. Kwek eds.), Springer, Singapore, 1998, 160–169.
- [6] F. Hiroshima, I. Sasaki, H. Spohn and A. Suzuki, *Enhanced Binding in Quantum Field Theory*, COE Lecture Note Vol. 38, Institute of Mathematics for Industry, Kyushu University, 2012.
- [7] S. K. Lamoreaux, Demonstration of the Casimir force in the 0.6 to 6 μm range, *Phys. Rev. Lett.* **78** (1998), 5–8; Erratum *Phys. Rev. Lett.* **81** (1998), 5475–5476.
- [8] M. Reed and B. Simon, *Methods of Modern Mathematical Physics II: Fourier Analysis, Self-Adjointness*, Academic Press, New York, 1975.
- [9] M. Reed and B. Simon, *Methods of Modern Mathematical Physics III: Scattering Theory*, Academic Press, New York, 1979.
- [10] M. Reed and B. Simon, *Methods of Modern Mathematical Physics IV: Analysis of Operators*, Academic Press, New York, 1978.
- [11] S. N. M. Ruijsenaars, On Bogoliubov transformations for systems of relativistic charged particles, *J. Math. Phys.* **18** (1977), 517–526.
- [12] S. N. M. Ruijsenaars, On Bogoliubov transformations. II. The general case, *Ann. of Phys.* **116** (1978), 105–134.
- [13] D. Shale, Linear symmetries of free boson fields, *Trans. Amer. Math. Soc.* **103** (1962), 149–167.

- [14] M. J. Sparnaay, Measurements of attractive forces between flat plates, *Physica* **24** (1958), 751–764.