



Title	Impact of QnrB19, a pentapeptide repeat protein mimicking double stranded DNA, on the quinolone resistance in Salmonella Typhimurium. [an abstract of dissertation and a summary of dissertation review]
Author(s)	Ruttana, PACHANON
Citation	北海道大学. 博士(獣医学) 甲第14276号
Issue Date	2020-09-25
Doc URL	<a href="http://hdl.handle.net/2115/79721">http://hdl.handle.net/2115/79721</a>
Rights(URL)	<a href="https://creativecommons.org/licenses/by/4.0/">https://creativecommons.org/licenses/by/4.0/</a>
Type	theses (doctoral - abstract and summary of review)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Pachanon_Ruttana_abstract.pdf (論文内容の要旨)



[Instructions for use](#)

学位論文内容の要旨  
Abstract of the dissertation

博士の専攻分野の名称：博士（獣医学）

氏名： Pachanon Ruttana  
Name

学位論文題名  
The title of the doctoral dissertation

Impact of QnrB19, a pentapeptide repeat protein mimicking double stranded  
DNA, on the quinolone resistance in *Salmonella* Typhimurium

（二重鎖DNAを模倣するペントペプチド繰返し蛋白QnrB19の  
ネズミチフス菌キノロン耐性化への影響）

Fluoroquinolones have been used for the drugs of choice to treat non-typhoidal *Salmonella* (NTS) infection in humans and animals. However, plasmid-mediated quinolone resistance (PMQR) mechanism has emerged in Enterobacteriaceae. Plasmid-encoded quinolone resistance protein Qnr is an important factor among PMQR in bacterial resistance to quinolones. This protein interacts with DNA gyrase and reduces susceptibility to quinolones.

However, as there are limited information of the predominant Qnr types in *Salmonella* isolates, I investigated the prevalence of Qnr in 692 reports published from 2008 to 2017 and found that 4,459 Enterobacteriaceae isolates had Qnrs (1,917 QnrB, 1,545 QnrS, 498 QnrA, 459 QnrD, 27 QnrVC, 12 QnrC and 1 QnrE). And QnrB19 was found in the highest number of isolates among Qnr reported in *Salmonella* isolates.

In Chapter I, *in vitro* assays for wild type *S. Typhimurium* DNA gyrases, QnrB19 and quinolones were performed. The IC<sub>50</sub>s of norfloxacin and ciprofloxacin against DNA gyrases were increased around 3-fold by the addition of QnrB19 and the results exhibited that the contribution of QnrB19 did not associate with the difference between R1 group of these two quinolones. The IC<sub>50</sub> of nalidixic acid was 59- and 110-folds higher than norfloxacin and of ciprofloxacin in the absence of QnrB19, respectively, and this was similar to the that in the presence of 18 nM QnrB19. These results showed that the fluorine at R6 and/or piperazine at R7 and/or hydrogen at R8 group might associate with the ability

of QnrB19 to increase IC<sub>50</sub>s. QnrB19 was shown for the first time *in vitro* to have ability to grant non-classical quinolone resistance to *S. Typhimurium* DNA gyrase in connection with the structure at R6 and/or R7 and/or R8.

In Chapter II, I identified the activity of novel fluoroquinolones with the potentiality attributed by unique R1 group, 6-amino-3,5-difluoropyridine-2-yl, by *in vitro* assay. I compared IC<sub>50</sub> of WQ-3810, WQ-3334 and WQ-4065 with specific features of quinolones at positions R1, R6, R7 or R8. WQ-3810 and WQ-3334 (6-amino-3,5-difluoropyridine-2-yl at the R1 group) showed stronger inhibitory activity against *S. Typhimurium* DNA gyrases with a lower IC<sub>50</sub>s than WQ-4065 (6-ethylamino-3,5-difluoropyridine-2-yl at the R1). Moreover, WQ-3810 and WQ-3334 showed greater inhibitory activity against *S. Typhimurium* DNA gyrases even in the presence of QnrB19. The results suggested that novel fluoroquinolones, WQ-3810 and WQ-3334 could be the good candidate drug for wild type *Salmonella* and *Salmonella* carrying QnrB19.

Comparison of conventional quinolones (nalidixic acid, norfloxacin and ciprofloxacin) in Chapter I and novel fluoroquinolones (WQ-3810, WQ-3334 and WQ-4065) in Chapter II showed that 6-amino-3,5-difluoropyridine-2-yl group at the R1 in WQ-3810 and WQ-3334 added a strong inhibitory activity against *S. Typhimurium* DNA gyrases to quinolones with very low IC<sub>50</sub>. IC<sub>50</sub>s of WQ-3810 and WQ-3334 were greatly increased in the presence of QnrB19, however, similar to those of ciprofloxacin and norfloxacin. Structural at R1 may cause the different impact by QnrB19 on the inhibitory activities of quinolones against DNA gyrase. The knowledge obtained in my study can be applied to design new compounds against bacteria carrying QnrB19 or other pentapeptide repeat proteins.