Title	Monitoring changes in land use and distribution of mangroves in the southeastern part of the Mekong River Delta, Vietnam
Author(s)	Hong, Huynh Thi Cam; Avtar, Ram; Fujii, Masahiko
Citation	Tropical ecology, 60(4), 552-565 https://doi.org/10.1007/s42965-020-00053-1
Issue Date	2019-12
Doc URL	http://hdl.handle.net/2115/79825
Rights	This is a post-peer-review, pre-copyedit version of an article published in Tropical ecology. The final authenticated version is available online at: http://dx.doi.org/10.1007/s42965-020-00053-1
Type	article (author version)
File Information	Manuscript_Tropical Ecology_MF_RA-06122019_clean.pdf

[Instructions for use](#)

Monitoring Changes in Land Use and Distribution of Mangroves in the Southeastern Part of the Mekong River Delta, Vietnam

Huynh Thi Cam Hong ¹, Ram Avtar ^{1,2*}, and Masahiko Fujii ^{1,2}

¹ Graduate School of Environmental Science, Hokkaido University, Sapporo 060-0810,
Japan;

² Faculty of Environmental Earth Science, Hokkaido University, Sapporo 060-0810,
Japan; htchong@ees.hokudai.ac.jp; ram@ees.hokudai.ac.jp; mfujii@ees.hokudai.ac.jp

* Correspondence: ram@ees.hokudai.ac.jp; Tel.: +81-011-706-2261

Abstract: Mangroves are one of the most valuable and productive coastal ecosystems. Previous studies show a severe loss of mangroves around the world over the last several decades due to natural and anthropogenic activities. Mangroves located in the Southeastern part of the Mekong River Delta (MRD) are also affected by these activities. Shrimp farming is considered as one of the main drivers causing the rapid loss and degradation of mangroves. The goal of this study is to assess the spatiotemporal changes in land use and distribution of mangroves in the Soc Trang and Bac Lieu provinces of the southeastern part of the MRD. Multi-temporal Landsat data were used for land use and land cover (LULC) classification using the maximum-likelihood classification (MLC) algorithm. The changes in mangrove forest areas were monitored using medium spatial resolution (Landsat-5 Thematic Mapper (TM) and Landsat-8 Operational Land Imager (OLI)) satellite imageries from 1988 to 2018. In the study area, there were seven major LULC types namely, dense mangroves, sparse mangroves, aquaculture farms, arable land with crop cover, arable land without crop cover, settlements, and water bodies. The overall accuracies of the LULC maps in 1988, 1998, 2008, and 2018 were 81.2%, 83.3%, 78.3%, and 81.9%, respectively. This study reveals that dense and sparse mangrove forests have decreased by 90% from 5,495 hectares (ha) to 515 ha and by 55% from 14,105 ha to 6,289 ha, respectively from 1988 to

2018. On the other hand, the aquaculture farm has increased at the rate of 5,024 ha/year for a period of 30 years. This rapid growth of aquaculture farming activities caused the rapid loss and degradation of mangroves in the MRD. Quantitative information about mangrove change obtained by this study is considered to be useful for future coastal management and relevant policies in the MRD.

Keywords: Mangrove forests, aquaculture, Landsat.

1. Introduction

Mangrove forests are generally referred to as tidal wetland ecosystems found in the intertidal zones between marine and terrestrial ecosystems (Saenger et al., 1983). Mangrove forests are restricted to the tropics and subtropics and considered highly productive ecosystems that provide various ecosystem services as well as climate change mitigation and adaptation (CCMA) options (Duncan et al., 2016). They serve as an enormous carbon sink, and at the same time provide climate and disturbance regulation services (Wylie et al., 2016). Besides, mangroves also play an important role in stabilizing the shoreline, preventing coastal erosion and well recognized in reducing the risks of tsunamis or storms (Blankespoor et al., 2014; Danielsen et al., 2005; Spalding et al., 2014; Van Coppenolle et al., 2018; T. Quoc Vo et al., 2013). Moreover, they also provide food, clean water, and other useful materials for local people (Phan et al., 2015). In addition, mangroves provide various benefits, which include aquatic resources, entertainment, ecotourism, biofiltration and so on (Locatelli et al., 2014; UNEP, 2014).

The total mangrove area accounts for 0.7% of the total tropical forests in the world (Donato et al., 2011). In the Southeast Asian region, the area of mangrove forests cover was 37,019 km² and 35,694 km² in 2000 and 2012, respectively (Hamilton et al., 2016). The largest extension of mangrove forests in 2010 was found in Asia with 42% of mangrove cover, followed by Africa with 20% (Long et al., 2011). In Vietnam, the area of mangroves

was estimated to be about 400,000 ha in the 1960s. The area declined dramatically to 73,000 ha in 1990 due to the use of herbicides during the Indochina War, and the expansion of agriculture and aquaculture areas (FAO, 2015). In Vietnam, the total mangrove area accounted for 270,000 ha in 2015 (FAO, 2015).

In the Vietnamese part of the Mekong River Delta (MRD), mangrove forest ecosystems are recognized as one of the most productive and biologically complex ecosystems. These mangrove forest ecosystems have been mostly distributed along with the MRD's coastal provinces, including Tra Vinh, Ben Tre, Long An, Kien Giang, Ca Mau, and especially Soc Trang and Bac Lieu (Cosslett, 2014). In recent decades, mangrove forest ecosystems have played vital roles not only in the biogeochemical cycle but also in providing a livelihood to the local people by aquaculture and fishing. Previous studies suggested that mangrove forests used to cover the the delta by more than 250,000 ha (Phan Nguyen Hong, 1993; Thu et al., 2007). The Indochina War, forest fires, collection of fuelwood, and other human activities have resulted in the reduction of mangrove forests (Feller et al., 2017), thus increasing the potential risks of flooding and land subsidence (Bakker, S.A, 2017; Truong et al., 2018). Especially, since the early 2000s, mangrove forests have been cleared for shrimp farming in many areas (Phan Nguyen Hong, 1993; Hong P., 1995; Van Hao, 1999). In addition, climate change has caused the sea level rise leading to saline water intrusion, which also causes a change in land use and mangrove distribution (Nguyen, 2016, Dasgupta et al., 2017; Tran et al., 2019).

Among many factors that have affected the mangroves of the MRD, the most important factor contributing to mangrove destruction is the aquaculture activities (Thu et al., 2007). Previous studies have demonstrated a complex relationship between mangroves and shrimp farming. Shrimp farming in the MRD has resulted in thousands of hectares of mangrove forests being converted to shrimp ponds. The pattern of land use and land cover (LULC) in

the MRD has been changing significantly over the decades, consequently affecting both the economic growth and environmental sustainability in the region. There are a large number of previous studies that have been conducted, focusing on mangrove restoration and development in recent decades (Ellison, 2000; Kaly et al., 1998; Lee et al., 2019; Nguyen, Van Tam et al., 2016). However, these studies have reported that mangrove restoration was a complex issue without sound and evidence-based restoration policies (Lee et al., 2019). Therefore, geospatial data-based information can play a significant role in providing evidence-based information about changes in mangrove forest areas and various drivers such as the expansion of aquaculture and its impacts on mangrove health. This analysis can help in understanding and identifying the complex relationship between mangrove status and aquaculture farming activities (Thu et al., 2007).

The LULC classification is an integral part of extracting thematic information about various classes from the satellite data. It is useful for land resource management and planning. Multi-temporal LULC data are valuable sources of information to know the trajectory of various land use and cover types and change pattern. Monitoring the changes in LULC based on remote sensing was conducted in previous studies with the aim to monitor the cropping pattern (Minh et al., 2019; Nguyen et al., 2016), as well as detecting the distribution of mangroves and urban areas at the local and global scales (Avtar et al., 2013; Avtar et al., 2016; C. Singh et al., 2010; Vo et al., 2013). The applications of geospatial data to study mangrove cover changes have been successfully demonstrated through previous studies (e.g. Avtar et al., 2018; Chandra Giri et al., 2007; T.-M. Lee et al., 2009; Muttitanon et al., 2005; I. J. Singh et al., 2004; Tuan Quoc Vo et al., 2015). Moreover, many studies have indicated that remote sensing has advantages over the traditional field-based investigation in the monitoring of mangrove forests because of the synoptic view and large area of coverage (Pham et al., 2019; Avtar et al., 2016; Kuenzer et al., 2011; C. Singh et al., 2010). The use of multi-spatiotemporal geospatial data can provide useful information about mangrove forests

globally. Pham et al. (2019) have reviewed the role of multispectral, Synthetic Aperture Radar (SAR), as well as hyperspectral data for estimating various biophysical parameters of mangroves. Their study summarized the most commonly used methodologies globally for estimating the biophysical parameters of mangroves. Among these methodologies, the pixel-based classification methods are considered the most commonly used in mangrove classification (Avtar et al., 2016; T. N. K. D. Binh et al., 2005; Kamal et al., 2011; Thu et al., 2007; Tong et al., 2004). Supervised classification algorithms were effectively used for mangrove classification using traditional satellite data (Conchedda et al., 2008; C. Giri et al., 2011; Kuenzer et al., 2011).

Although there are many studies on the impacts of shrimp farming on mangrove change, most of them focused on limited provinces of MRD, such as Kien Giang, Ca Mau, Tra Vinh, and Ben Tre (Tan Phong Nguyen et al., 2019; Thu et al., 2007; Van Cuong et al., 2015; Tuan Quoc Vo et al., 2015). In Soc Trang and Bac Lieu provinces, the sea dyke system has been built to protect the residential area for the purpose of economic development in the area. At present, coastal erosion along with deforestation for aquaculture has not been adequately controlled. Therefore, local authorities have a plan to protect the mangroves and rehabilitate the mangroves outside the sea dyke, which is considered essential. In addition, local authorities have co-operated with villagers to co-manage mangrove forests inside sea dykes through the benefits of mangrove-shrimp farming (DARD, 2017).

Therefore, due to the limited comprehensive data on temporal changes in the mangrove ecosystems of Soc Trang and Bac Lieu provinces, this study provides a comprehensive assessment of long-term changes in mangroves distribution and its relationship with aquaculture activities. This study investigates the potential application of Landsat data using the maximum-likelihood classification (MLC) algorithm to monitor land cover changes and mangroves distribution in the Soc Trang and Bac Lieu provinces from 1988 to 2018. Change detection methods were also used to have better visualization and understanding of changing

trends of mangrove forests. This study can support the local government in the decision-making process for land use planning and management practices in the future such as mangrove plantation plans for sea dyke's protection, policies of mangroves development along with aquaculture.

2. Materials and Methods

2.1. Study area

In the study, two coastal provinces in the Southeastern region of the MRD, namely Soc Trang (Lat. 9°12'N -9°56'N, Long. 105°33'E–106°23'E) and Bac Lieu (Lat. 9°00'–9°38'9"N, Long. 105°14'15"–105°51'54"E) were selected (Figure 1). The total coastline is about 128 km in length, of which Soc Trang and Bac Lieu provinces cover 72 km and 56 km, respectively (DARD, 2017). In terms of climatic features, Soc Trang and Bac Lieu provinces are located in the tropical climate zone with high monthly mean atmospheric temperature (Evers et al., 2009; Ha et al., 2018; Huu, 2011). These provinces have two distinct seasons, i.e. wet season (from May to November) and dry season (from December to April). The average annual rainfall in Soc Trang and Bac Lieu varies from 2,000 to 2,300 mm (DARD, 2017). Noticeably, in the dry season, rainfall accounts for only 10% of the total annual rainfall, with almost no rain (often triggering droughts) in January through March, while the rainy season has continuous rain occasionally that can last for 3-5 days causing serious flooding (DARD, 2011). The annual mean atmospheric temperature is 26⁰C, with a maximum temperature of 30⁰C in April and May and a minimum temperature of 20⁰C in December and January. The average annual sunshine hours are 2,500-2,600 hours (DARD, 2017).

According to the Department of Agriculture and Rural Development (DARD), the average humidity during the dry season is around 80% (DARD, 2017). Historical data from the Meteorological Department (MD) shows that the increase in average humidity in Soc

Trang and Bac Lieu provinces is up to 85% during the wet season (DARD, 2017). Noticeably, although Soc Trang and Bac Lieu provinces are warm coastal provinces, they are less affected by storms, and tropical cyclones. However, they are under strong impacts of the semi-diurnal system of the East Sea tides and a part of the diurnal system of the West Sea. The monthly average water level of high and low tides in Soc Trang and Bac Lieu provinces varies between -1.9 m and 1.9 m (DARD, 2017). Climate change in recent years is expected to increase the risk in the study area, due to flooding from the Mekong River as well as droughts and seawater intrusion (Hak et al., 2016; Toan, 2014).

Figure 1. The location of Soc Trang and Bac Lieu provinces in the MRD, Vietnam, and distribution of field survey points.

2.2. Field data

In this study, two field surveys were conducted from June 1 to July 30, 2017 and from March 15 to April 15, 2018, to collect ground truth data necessary for LULC classification

and validation of results. In the field surveys, 170 and 125 field sites were randomly selected in the Soc Trang and Bac Lieu provinces, respectively. Global Positioning System (GPS) device (Gamin 60x) was used to mark the locations of various LULC types. The GPS photos were also collected for various LULC types. The geo-tagged photos are highly effective in improving the accuracy of the LULC classification (Oba et al., 2014). These photos have clearly shown the difference in the vegetation vigor and density of mangrove forests in the Soc Trang and Bac Lieu provinces. Figure 2 shows the distribution of the ground truth data collected during the field survey. Figure B1 shows dense mangrove forests defined as the area in which the average mangrove density is more than 5,000 trees per ha (The People's Committee of Bac Lieu Province, 2017). The mangroves are mainly located along the national sea dyke systems (Figures B1a and B1b). A part of the mangroves located outside the sea dyke systems was affected by high waves and was gradually swept into the sea water (Figure B1c).

In Figure B2, the sparse mangrove forests are shown with an average mangrove density of less than or equal to 5,000 trees per ha (The People's Committee of Bac Lieu Province, 2017). At some field sites, the mangrove forests' density and age were directly observed according to information by DARD (2017). In addition, the total percent of mangroves cover in each aquaculture farm associated with aquaculture production were also investigated. This information and data were used as training samples for the MLC of Landsat data and validation. Each training site was delineated on the images using polygons that cover multiple pixels in the homogeneous areas. The ground data were collected based on the homogeneity of the area/class to cover the representative sample of Landsat data. The status of mangrove forests and aquaculture farming areas were also recorded as pieces of evidence for the validation of results. Besides ground truth data, high spatial resolution images on Google Earth (GE) and administrative land use maps at the provincial-level (Bac Lieu and Soc Trang) were used for validation of the classification results.

Figure 2. Distribution of ground truth data collected in Soc Trang and Bac Lieu provinces.

Xu et al. (2017) demonstrated the importance of field photographs in identifying the various LULC types and landscapes in the study area. Figure B1-B5 show a number of geo-tagged photographs with the density of mangrove forests, aquaculture farms, as well as arable land with and without crop cover. The selection of ground truth data was based on the number of classes for LULC classification. Figures B3, B4, and B5 show the aquaculture area, arable land with crop cover, and arable land without crop cover, respectively.

2.3. Satellite data

In this study, Landsat-5 TM and Landsat 8 OLI data of 1988, 1998, 2008, and 2018 were downloaded from the United States Geological Survey (USGS) server (<https://earthexplorer.usgs.gov/>) for land cover classification. The Landsat data with less than 10% of cloud cover were selected from Landsat archived. Table 1 summarizes the detailed information of the satellite data that was used in this study in every 10-years period. Landsat data were only collected in the dry season to eliminate not only the effects of clouds and shadows, but also the change in surface reflectance in different seasons.

Table 1. Detailed information of satellite data collected from the USGS server.

Satellite	Sensor	Path/Row	Spatial resolution (m)	Date of acquisition
Landsat 5	TM	125/53	30	30 January
		125/54		1988
Landsat 5	TM	125/53	30	30 January
		125/54		1998
Landsat 5	TM	125/53	30	9 March 2008
		125/54		
Landsat 8	OLI	125/53	30	5 March 2018
		125/54		

211

212 2.3.1. Landsat data processing

213 Figure 3 shows the methodology adopted in this study. Information about various objects
214 on the earth's surface was generated from the Landsat data using reflectance information in
215 the form of digital number (DN) (Avtar et al., 2013). Visual image interpretation techniques
216 were also used to visualize the quality of data as well as various land cover types. Different
217 forms of distortion or shift were noticed and considered during the visual interpretation
218 process. Therefore, the geometric rectification, including image-to-image registration was
219 applied. The image enhancement and histogram matching techniques were also used to
220 enhance the quality of the image (Sari et al., 2016; Moreira et al., 2014; Tokola et al., 2001).
221 In the study, the Normalized Difference Vegetation Index (NDVI) was also generated to
222 monitor the spectral reflectance properties of vegetation. The NDVI was calculated as
223 expressed in Equation (1):

$$224 \quad \text{NDVI} = (\text{NIR} - \text{Red}) / (\text{NIR} + \text{Red}) \quad (1)$$

where NIR is the reflectance at the near infra-red band, and

Red is the reflectance at the red band.

NDVI has been widely used to identify various vegetation types (Alatorre et al., 2016; Avtar et al., 2011; Mohajane et al., 2018; Rouse Jr et al., 1974; Ruiz-Luna et al., 2010). The dense and sparse mangrove forests, agriculture areas, and aquaculture areas can be differentiated using NDVI values. Therefore, we further used NDVI in MLC classification.

Figure 3. Flowchart of the methodology of the study.

2.3.2. Image classification

There are various classification algorithms for LULC classification. Among these algorithms, the MLC is widely used due to its successful applications (Dan, 2017; Hall et al., 1995; Lillesand et al., 2015; Townshend, 1992). The MLC is a parametric classification algorithm that assumes a classed signature in a normal distribution (Gaussian statistical distribution) and statistically calculates the probability that a particular pixel belongs to a

particular class (Ramsey III et al., 1996). Each pixel is assigned to the class that has the highest probability (Dan et al. 2017; Hall et al., 1995). The ArcGIS 10.4 and ENVI 5.2 software were used for classification and change detection. It was also a useful tool to determine the relationship between the LULC class and spectral reflectance. Each LULC map was classified into seven classes, namely, dense mangroves, sparse mangroves, aquaculture farms, arable land with crop cover, arable land without crop cover, settlements, and water bodies (Table 2). The selection of classes for classification was based on the fieldwork and other existing maps.

Table 2. Land-cover classified in this study.

Land-cover class	Description
Water bodies	Rivers, canals
Aquaculture farms	Shrimp farms
Settlements	Residence, commercial, industrial, transportation
Arable land without crop cover	No vegetation covered
Arable land with crop cover	Rice fields, fruit gardens
Sparse mangroves	Mangrove coverage of less than 50%
Dense mangroves	Mangrove coverage of more than 50%

2.3.3. Accuracy assessment

In this study, LULC maps were generated using Landsat data and MLC algorithm. An accuracy assessment of classified maps was done using the confusion matrix. The ground truth data collected during field surveys and Google Earth (GE) images were used for accuracy assessment of classified maps (Congalton, 1991; Islam et al., 2018). The sampling points were randomly collected during the field surveys. These points were converted to GE images to see the coverage and distribution in different land cover categories. These points were compared with locations on the GE images to cross-validate their locations' accuracy.

267 In the process of LULC classification, the maps of mangrove classification were compared
268 with the available mangrove maps from official mapping sources to avoid misclassification
269 between mangroves and other crops. The overall accuracy and kappa coefficients were also
270 calculated to evaluate the agreement between reference and prediction data (Congalton,
271 1991; Liu et al., 2008; Long & Giri, 2011).

272 **3. Results**

273 *3.1. Land use and land cover maps*

274 The spatiotemporal pattern of various LULC classes for 1988, 1998, 2008, and 2018
275 Landsat data using MLC classifier are shown in Figure 4. The agricultural area was mostly
276 located in the upper region of the study area. The mangrove forests were commonly
277 distributed in the coastal regions, while most of the aquaculture farms were located between
278 agricultural lands and mangrove forests.

Figure 4. The LULC classification maps of Soc Trang and Bac Lieu provinces of the years 1988, 1998, 2008, and 2018, based on MLC of Landsat 5 TM (1988, 1998 and 2008) and Landsat 8 OLI (2018) images.

The temporal changes in the LULC classes from 1988 to 2018 were illustrated in Figure 5. The arable land with crop cover was the most dominant land cover type, amounting to 463,376 ha (80.9%) of the total area in 1988. The areas of dense and sparse mangrove forests were only 5,495 ha (1.0%) and 14,105 ha (2.5%), respectively, in 1988. The aquaculture area covered 9,673 ha (1.7%) in 1988. Noticeably, the aquaculture farms have significantly increased, reaching 160,393 ha (28.0%) in 2018. The opposite trend was seen in the total area of arable land with and without crop cover. The total area of agricultural land gradually decreased over the 30 years from 526,221 ha in 1988 to 387,102 ha in 2018. In terms of the settlement area, this area had increased from 95 ha in 1988 to 2,660 ha in 2018. The areas under water bodies showed a slight decrease, from 17,017 ha in 1988 to 15,646 ha in 2018, but the change may be affected by seasonal variations in water coverage in the study area.

The results showed that mangrove forests were extensively distributed in the coastal regions of Soc Trang and Bac Lieu provinces and along the rivers interlinked with sea water. The decrease in the areas of dense and sparse mangroves from 1988 to 2008 was faster than that from 2008 to 2018. The annual rates of change in dense mangrove forests were 243 ha/year and 13 ha/year from 1988 to 2008 and 2008 to 2018, respectively. The annual change

rates of sparse mangroves were 352 ha/year and 79 ha/year from 1988 to 2008 and 2008 to 2018, respectively. The study reveals that the annual deforestation rate of sparse mangroves was more than that of dense mangroves. Mangrove deforestation in the study area was caused by various drivers in the MRD. Indeed, previous studies pointed out that these drivers include shrimp farming (De Graaf & Xuan, 1998; T. T. P. Ha, van Dijk, & Visser, 2014; Johnston, Trong, Tien, & Xuan, 2000), the Indochina war, and human activities such as cutting down trees for fuelwood or agricultural area expansion (Phan Nguyen Hong, 1993; Orchard et al., 2016). This low rate was mainly due to the mangrove conservation projects and programmes in the MRD in recent years. One thing to note is that the annual rate of change in the aquaculture farms from 1988 to 2018 (5,024 ha/year) was higher than that in previous periods. As mentioned earlier, the expansion of aquaculture farms was mainly occurred in between mangroves and agricultural areas, implying prominent land-use change from mangroves to agricultural areas (Figure 4).

Figure 5. Changing pattern of LULC classes in the study area from 1988 to 2018.

Generally, the results of the LULC classification show the decline in the mangrove forest area, while the calculated NDVI value indicates the degradation of the mangroves quality. Therefore, the NDVI-based analysis can be used to monitor mangrove density as reported by previous studies (Adi et al., 2016; Sari et al, 2016). Figure 6 shows the temporal variation in the mean NDVI of dense and sparse mangrove forests. The mean NDVI values of dense mangroves were always higher than that of sparse mangroves. As can be seen from the figure, the mean NVDI values of dense mangroves show a decreasing trend. This trend shows that together with the dense mangrove forest area, the quality of dense mangrove forests has also declined during the 1988 to 2018. In the case of sparse mangroves, the mean NVDI values have increased from 1988 to 1998, then have decreased gradually from 1998 to 2018.

Figure 6. The NDVI mean values of dense and sparse mangroves in the years 1988, 1998, 2008, and 2018 (standard deviation values in table A5)

3.2. Accuracy assessment

In terms of the validation of classified images, 420 sampling points were randomly created. The accuracy assessment results are shown in Appendix A of LULC maps of 1988, 1998, 2008, and 2018. Table 3 illustrates the overall accuracy of LULC maps and kappa coefficient based on a confusion matrix. The overall accuracy of LULC maps for the year 1988, 1998, 2008, and 2018 were 81.2%, 83.3%, 78.3%, and 81.9%, respectively, and the kappa coefficient were 0.78, 0.81, 0.75, and 0.79, respectively.

Table 3. The accuracy assessment of the land cover maps from 1988 to 2018.

Class	1988		1998		2008		2018	
	PA (%)	UA (%)	PA (%)	UA (%)	PA (%)	UA (%)	PA (%)	UA (%)
C1	95.0	86.4	98.3	98.3	91.7	96.5	80.0	90.6
C2	75.0	80.4	78.3	78.3	71.7	82.7	76.7	82.1
C3	81.7	79.0	80.0	98.0	71.7	81.1	85.0	81.0
C4	78.3	72.3	81.7	60.5	75.0	57.7	85.0	75.0
C5	78.3	79.7	75.0	83.3	76.7	62.2	83.3	73.5
C6	75.0	86.5	78.3	92.2	76.7	76.7	80.0	84.2
C7	85.0	85.0	91.7	84.6	66.7	87.0	83.3	90.9
Overall								
accuracy	81.2		83.3		78.3		81.9	
(%)								
Overall								
kappa	0.78		0.81		0.75		0.79	
coefficient								

PA: Producer's accuracy, UA: User's accuracy; C1: Water bodies, C2: Aquaculture farms, C3: Settlements, C4: Arable land without crop cover, C5: Arable land with crop cover, C6: Sparse mangroves, C7: Dense mangroves.

3.3. Change detection

Table 4 shows clear changes in LULC area in every 10-year period between 1988 and 2018. This study reveals that mangrove forests have decreased by 12,796 ha from 1988 to 2018 in the Soc Trang and Bac Lieu provinces. In the first 10-year period (1988-1998), the area of sparse mangroves had rapidly decreased by 6,602 ha and continued to decline by 7,816 ha during the last 30 years. The decrease was mainly due to the expansion of aquaculture farms in the coastal regions of the MRD.

Table 4. The changes in LULC area in every 10-year period between 1988 and 2018.

Land cover	1988-1998		1998-2008		2008-2018		1988-2018	
	ha	%	ha	%	ha	%	ha	%
C1	-1,004	-0.18	597	0.10	-964	-0.17	-1,371	-0.24
C2	27,894	4.87	98,484	17.20	24,342	4.25	150,720	26.32
C3	597	0.10	1,251	0.22	717	0.13	2,565	0.45
C4	191,630	33.47	-45,894	-8.01	-10,254	-1.79	135,482	23.66
C5	-211,760	-36.98	-49,912	-8.72	-12,928	-2.26	274,600	-47.96
C6	-6,602	-1.15	-429	-0.07	-785	-0.14	-7,816	-1.36

C7	-755	-0.13	-4,096	-0.72	-129	-0.02	-4,980	-0.87
----	------	-------	--------	-------	------	-------	--------	-------

C1: Water bodies, C2: Aquaculture farms, C3: Settlements, C4: Arable land without crop cover, C5: Arable land with crop cover, C6: Sparse mangroves, C7: Dense mangroves.

The magnitude of change in different LULC classes from 1988 to 2018 is shown in Table 5. It is clearly seen that the aquaculture farms have increased, amounting to approximately 152,839 ha during the last 30 years. The LULC classes of arable land with crop cover, arable land without crop cover, sparse mangroves, and dense mangroves converted into aquaculture farms by 27.1%, 29.1 %, 45.7%, and 35.4%, respectively. One thing to note is that the settlements and water bodies accounted for the lowest percentages of the total LULC categories. The total unchanged areas of these two classes represented 80% and 78% during the last 30 years, respectively.

Table 5. The post-classification change detection matrix indicating the direction of change between 1988 and 2018.

LULC classes changes	From							
		C1	C2	C3	C4	C5	C6	C7
To C1		13,103.73*	53.37	10.89	75.69	1,986.39	77.49	29.25
(ha) C2		396.99	7,964.50*	-	13,899.60	137,993.94	6,560.19	1,952.46
C3		74.34	36.72	60.03*	91.89	1,811.52	14.13	6.39
C4		624.24	1,125.18	14.85	21,499.83*	162,931.14	3,135.42	599.22
C5		675.27	403.29	-	15,447.15	168,181.38*	2,868.48	1,817.82

	C6	1,849.32	421.83	-	200.97	1,590.39	1,654.20*	918.28
	C7	166.95	13.32	-	2.88	7.83	61.83	190.53*
To	C1	77.58*	0.53	12.70	0.15	0.42	0.54	0.53
(%)	C2	2.35	79.50*	-	27.14	29.08	45.65	35.41
	C3	0.44	0.37	69.99*	0.18	0.38	0.10	0.12
	C4	3.70	11.23	17.31	41.98*	34.34	21.82	10.87
	C5	4.00	4.03	-	30.16	35.44*	19.96	32.97
	C6	10.95	4.21	-	0.39	0.34	11.51*	16.65
	C7	0.99	0.13	-	0.01	-	0.43	3.46*

363 C1: Water bodies, C2: Aquaculture farms, C3: Settlements, C4: Arable land without crop
364 cover, C5: Arable land with crop cover, C6: Sparse mangroves, C7: Dense mangroves.
365 Astarisks (*) show unchanged areas (the areas that were not converted to any other
366 classes).

367 Figure 7 shows the geospatial distribution of LULC changed and unchanged areas from
368 1988 to 2018. During the period 1988-2018, the total area of regional land-use change
369 accounted for 62.9%, while the total unchanged area was 37.1%. The annual rate of change
370 was 11,998 ha/year. The results show that the major LULC change was due to the expansion
371 of aquaculture farms. The aquaculture farms accounted for approximately 160,406 ha and
372 increased from 1.7% in 1988 to 28.0X% in 2018. During that period, dense and sparse
373 mangroves were converted into aquaculture farms by 35.4% and 45.7%, respectively.
374 Besides, arable land with crop cover was converted into aquaculture farms areas by 27.1%
375 (Table 5). Mangrove areas in the Soc Trang and Bac Lieu provinces have been continuously
376 decreasing over the years. In 2018, the total mangrove forest area in these two provinces was

377 6,804 ha. More than 90% of dense mangrove areas and 55% of sparse mangrove areas had
378 reduced during the study period of 30 years. It can be indicated that settlements continued to
379 increase, while both arable land with crop cover and mangrove forests decreased. In fact,
380 mangrove forests were converted into not only aquaculture farms but also agricultural areas.

(a)

(b)

(c)

Figure 7. Change detection of mangrove forests (a) from 1988 to 1998, (b) from 1998 to 2008 and (c) from 2008 to 2018.

4. Discussion

4.1. Spatio-temporal changes of mangrove forests

The multi-temporal LULC maps show the changing trends of the LULC classes in a simplified way (Figure 4). These maps illustrate the changing pattern of various LULC classes. For instance, the aquaculture farming areas were mainly expanded in between the agriculture areas and mangrove forests, as mangrove forests serve as nurseries and food-supplies for marine and brackish water animals, and therefore, have high potentials to function as aquaculture farming areas as well once converted. The mangroves also absorb wastes generated by shrimp farming (Gautier et al., 2001; Wösten et al., 2003). The key findings of the study show that the areas of agriculture, dense and sparse mangroves, and water bodies in Soc Trang and Bac Lieu provinces have decreasing trends of LULC change, while the areas of aquaculture farms and settlements show increasing trends. Findings from this study are similar to those from previous studies (Binh et al., 2008; Tong et al., 2004). The previous studies have reported that a large number of mangrove and agricultural lands in the MRD have been converted into aquaculture farms during the last decades. Consequently, it is clear that the rapid increase in the aquaculture farms in the MRD have negative impacts on mangroves. One of the recent studies (Truong et al., 2018) also reported a decrease in the mangrove forests in the MRD.

The aquaculture area in the study site has increased rapidly from 1988 to 2018. It is believed that aquaculture has brought more economic benefits than maintaining and conserving mangrove forests (Vo et al., 2013). From 2008 to 2018, both dense and sparse

mangrove forests only slightly decreased, which was due to a number of national and international mangrove forest conservation projects and programs. Mangroves, mainly *S. caseolaris*, were planted in the mudflats along the coastal lines in the MRD, especially in Cu Lao Dung district, Soc Trang province (DARD, 2017). The mangrove replantation in the MRD usually occurred after the sediment deposition in the shallow seabed. In Bac Lieu province, however, the replantation as well as conservation of mangrove forests have been limited in recent years due to the serious erosion in the coastal lines.

Based on the analysis of NDVI, it is clear that the quality of mangroves (especially density of mangroves) has also changed. A series of results of field observation data, Landsat data and NDVI-based analysis suggest that dense mangroves have been damaged severely compared to sparse mangroves between 1988 and 2018. The NDVI-based analysis shows that there was a fluctuation in the density of sparse mangroves during the study period. It shows an increasing pattern for the period 1988-1998 and a decreasing pattern for the period 1998-2018. Noticeably, the decreasing pattern for the period 1998-2018 was mainly due to a number of mangrove forest management policies in the regional master plan that was approved by the Prime Minister in the 1996-1999 and 2000-2006 periods (FAO 2016). Under these policies, an households or organization can convert up to 40% of mangroves area to other uses such as housing and aquaculture (mainly shrimp farms – i.e. shrimp farmers in protection and production mangrove forests must maintain at least 60% mangrove canopy cover) (Decision 186/2006/QD-TTg). According to this plan, the mangrove areas in the MRD, including Soc Trang and Bac Lieu provinces, were divided into three zones such as full protection zone, buffer zone, and economic zone. In these zones, 60% of the land area was used for mangrove forest development, and the rest (40%) was used for aquaculture (mainly shrimp farming), agriculture, and other utilizations (Minh et al., 2001).

Together with other Asian countries such as India and Thailand, the mangrove forest areas in the MRD have a decreasing trend in recent decades as a result of the economic benefits (Liu et al., 2018). The recovery of the mangrove forests as well as ecosystem protection in the MRD are major challenges for the Vietnamese government. However, recovery and conservation of the mangrove forests are crucial not only for the ecosystem protection in Soc Trang and Bac Lieu provinces but also for sustainable development in the MRD. Therefore, the application of advanced techniques for the assessment of LULC change is very essential for better mangrove forest management. In this study, the use of moderate resolution (30 m) Landsat data has enabled the monitoring of spatio-temporal changes in mangroves in the study area because of low-cost or free data, long temporal coverage, and wider spatial coverage of Landsat data (Aziz et al., 2015; Chandra Giri et al., 2015; Ottinger et al., 2016; T. Vo et al., 2013). However, these data still have limitations of clouds and shadows which can affect the accuracy of classification (Avtar et al., 2013; Avtar et al., 2016). Therefore, the use of high-resolution satellite data along with in-situ information can improve the accuracy of classification. Use of SAR data can help to overcome the limitation of clouds and shadows (Avtar et al., 2018; Minh et al., 2019; Nguyen et al., 2016; Vo et al., 2013) and can also provide more information about biophysical parameters of mangroves, such as biomass, density and height. For example, (e.g. Avtar et al. (2013, 2016) had successfully applied SAR satellite images to detect forest area and biomass. Pham et al. (2018) have applied SAR data to observe the mangrove forest changes in the MRD. Thus, such applications could be widely used in the MRD in the future. SAR data were not used in this study because of the unavailability of historical SAR data. In the future, we can study about the role of distance from the coastal areas in mangroves degradation and impacts of socio-economic factors on change in mangroves areas.

4.2. Driving factors of mangrove degradation

Mangrove forest dynamics are undergoing constant changes due to both natural and anthropogenic forces. The decrease in mangrove areas was mainly due to an expansion of aquaculture farms. Besides aquaculture farms, mangrove areas have been lost due to fuelwood collection and cutting of mangrove trees for house construction materials (Thu & Populus, 2007). Moreover, the rapid urbanization along the coastal regions had been noticed through the field surveys in the study area. Baumgartner et al. (2016) found that the degradation of mangroves in the MRD was also caused by the short-term mangrove exploitation for immediate economic benefits by local communities rather than long-term sustainable use of mangroves. Local farmers have individually expanded their aquaculture land without land-use planning. Besides, there are small new segments of the mangroves that were planted along the coast, especially in the mudflats area outside the sea dyke systems (Albers et al., 2015; Joffre et al., 2018).

Recently, the Vietnamese government has allowed local people to join mangrove protection along with aquaculture farming activities under a model of “land use allocation for forestry production purposes”. For example, DARD (2017) showed that the Nha Mat district of Bac Lieu province was one of the regions that allowed local farmers to join the above model. In this model, the local people can use 30% of the mangrove area for their aquaculture production for a 5-years period (Truong et al., 2018). However, a large number of local people had gradually cut down the roots of mangroves. As a result, these mangroves become weak or dead. The information collected from local people through two field surveys revealed that the mangrove forest coverage has affected the aquaculture production. In fact, local authorities allow households who have contracted co-management of mangrove forest to convert up to 40% of mangrove area where aquaculture is combined. This was consistent with Truong et al. (2018) that revealed that if mangroves coverage is more than 60% then it had a negative impacts on shrimp productivity. Therefore, the use of accurate mangrove coverage area maps can be helpful for local stakeholders including policy makers. The results

of this study may assist in the decision-making process for conserving mangroves, preventing environmental degradation, and improving the ecosystem management and planning in Bac Lieu and Soc Trang provinces in the MRD.

5. Conclusions

Mangrove forests are important to human beings as they provide various ecosystem services to support human livelihood activities. Detecting changes in the mangrove forest areas, especially in the coastal regions, is essential for ecosystem conservation as well as coastline protection. In this study, Landsat data were used to evaluate the LULC changes caused by development activities and policies over the 30 years. Results show a significant disappearance of the mangroves over the study period. Remote sensing-based analysis could identify low- and high-density mangrove forests based on the spectral signature. Besides, the NDVI data could also show the changes in the density and health condition of mangrove forests. This study has revealed that the expansion of aquaculture farms has been one of the major factors leading to the considerable decrease in the areas as well as the quality (density and health condition) of dense and sparse mangroves. Nevertheless, mangroves plantation activities did not compensate sufficiently for the loss of mangrove area caused by aquaculture expansion. Therefore, there is an urgent need for mangrove research and management policies and strategies, as well as plans for the sustainable development of mangroves in the MRD.

Acknowledgments: Authors would like to thank USGS for providing Landsat data and Can Tho University for providing support during field-work. We would also like to thank JICA-JICE for their continuous support and Hokkaido University for providing facilities. This study was also supported by the “Integrated Research Program for Advancing Climate Models (TOUGOU program)” from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan. The authors are also thankful to Stanley Anak Suab, Huynh

Vuong Thu Minh, Drs. La Ode Muhammad Yasir Haya and Amit Kumar Batar for suggestions and encouragement to draft the manuscript. The authors also would like to thank five students in Can Tho University for helping in the fieldwork.

Appendix A

Tables A1 through A4 show the results of the accuracy assessment of the four LULC maps in 1988, 1998, 2008 and 2018, respectively. The overall accuracy varied from 83.3 % to 78.3% and the kappa coefficient varied from 0.81 to 0.75.

Table A1. Confusion matrix of L8 OLI (1988) images representing classification accuracy of supervised classification.

Class	C1	C2	C3	C4	C5	C6	C7	Total (Producer's)
C1	57	2	1	0	0	0	0	60
C2	7	45	4	3	1	0	0	60
C3	1	1	49	8	1	0	0	60
C4	0	1	3	47	9	0	0	60
C5	1	4	1	7	47	0	0	60
C6	0	1	4	0	1	45	9	60
C7	0	2	0	0	0	7	51	60
Total (User's)	66	56	62	65	59	52	60	420

C1: Water bodies, C2: Aquaculture farms, C3: Settlements, C4: Arable land without crop cover, C5: Arable land with crop cover, C6: Sparse mangroves, C7: Dense mangroves.

Table A2. Same as in Table A1 but for LS5 TM (1998) images.

Class	C1	C2	C3	C4	C5	C6	C7	Total (Producer's)
C1	59	0	0	1	0	0	0	60
C2	1	47	0	12	0	0	0	60
C3	0	4	48	7	1	0	0	60
C4	0	2	1	49	8	0	0	60
C5	0	7	0	8	45	0	0	60
C6	0	0	0	3	0	47	10	60
C7	0	0	0	1	0	4	55	60
Total (User's)	60	60	49	81	54	51	65	420

Table A3. Same as in Table A1 but for LS5 TM (2008) images.

Class	C1	C2	C3	C4	C5	C6	C7	Total (Producer's)
C1	55	5	0	0	0	0	0	60
C2	0	43	5	7	5	0	0	60
C3	2	1	43	5	9	0	0	60
C4	0	1	2	45	12	0	0	60
C5	0	0	2	12	46	0	0	60
C6	0	1	0	5	2	46	6	60

C7	0	1	1	4	0	14	40	60
Total								
(User's)	57	52	53	78	74	60	46	420

526

527 **Table A4.** Same as in Table A1 but for LS5 TM (2018) images.

Class	C1	C2	C3	C4	C5	C6	C7	Total (Producer's)
C1	48	7	2	2	1	0	0	60
C2	5	46	4	3	2	0	0	60
C3	0	2	51	7	0	0	0	60
C4	0	1	2	51	4	1	1	60
C5	0	0	4	4	50	2	0	60
C6	0	0	0	0	8	48	4	60
C7	0	0	0	1	3	6	50	60
Total (User's)	53	56	63	68	68	67	55	420

528

529 **Table A5.** NDVI value of dense and sparse mangroves

Year	Mangrove class	Minimum value	Maximum value	Standard deviation	Mean value
1988	DM	0.655	0.909	0.059	0.748
	SM	0.046	0.734	0.111	0.463
1998	DM	0.660	0.859	0.042	0.732
	SM	0.278	0.662	0.074	0.563
2008	DM	0.560	0.695	0.021	0.590
	SM	0.235	0.630	0.087	0.417
2018	DM	0.455	0.558	0.017	0.474
	SM	0.337	0.455	0.032	0.393

DM: Dense mangroves, SM: Sparse mangroves

Figure B1. Photos of dense mangroves.

534 **Figure B2.** Photos of sparse mangroves.

535 **Figure B3.** Photos of aquaculture farms.

536

537

538

539

9°29'40".82N -105°15'41".33E

9°30'11".52N -105°31'25".98E

9°31'3".50N -105°53'45".50E

(a)

(b)

(c)

Figure B5. Photos of agriculture lands without crops.

References

- Adi, W., & Sari, S. P. (2016). Detection of mangrove distribution in Pongok Island. *Procedia Environmental Sciences*, 33, 253–257.
- Alatorre, L. C., Sánchez-Carrillo, S., Miramontes-Beltrán, S., Medina, R. J., Torres-Olave, M. E., Bravo, L. C., ... Uc, M. (2016). Temporal changes of NDVI for qualitative environmental assessment of mangroves: Shrimp farming impact on the health decline of the arid mangroves in the Gulf of California (1990–2010). *Journal of Arid Environments*, 125, 98–109. <https://doi.org/10.1016/j.jaridenv.2015.10.010>
- Albers, T., & Schmitt, K. (2015). Dyke design, floodplain restoration and mangrove co-management as parts of an area coastal protection strategy for the mud coasts of the Mekong Delta, Vietnam. *Wetlands Ecology and Management*, 23(6), 991–1004.
- Avtar, R., Herath, S., Saito, O., Gera, W., Singh, G., Mishra, B., & Takeuchi, K. (2013). *Application of remote sensing techniques toward the role of traditional water bodies with respect to vegetation conditions* (Vol. 16). <https://doi.org/10.1007/s10668-013-9507-4>
- Avtar, R., Kumar, P., Oono, A., Saraswat, C., Dorji, S., & Hlaing, Z. (2016). *Potential application of remote sensing in monitoring ecosystem services of forests, mangroves and urban areas* (Vol. 32). <https://doi.org/10.1080/10106049.2016.1206974>

559 Avtar, R., Mukherjee, S., Abayakoon, S. B. S., Sophal, C., & Thapa, R. (2018). *Integrating*
560 *ALOS-PALSAR and ground based observations for forest biomass estimation for*
561 *REDD+ in Cambodia* (Vol. 8). <https://doi.org/10.30852/sb.2018.414>

562 Avtar, R., Singh, C. K., Shashtri, S., & Mukherjee, S. (2011). Identification of erosional and
563 inundation hazard zones in Ken–Betwa river linking area, India, using remote sensing
564 and GIS. *Environmental Monitoring and Assessment*, 182(1–4), 341–360.

565 Avtar, R., Suzuki, R., Takeuchi, W., & Sawada, H. (2013). PALSAR 50 m Mosaic Data
566 Based National Level Biomass Estimation in Cambodia for Implementation of
567 REDD+ Mechanism. *PLOS ONE*, 8(10), e74807.
568 <https://doi.org/10.1371/journal.pone.0074807>

569 Aziz, A. A., Phinn, S., & Dargusch, P. (2015). Investigating the decline of ecosystem services
570 in a production mangrove forest using Landsat and object-based image analysis.
571 *Estuarine, Coastal and Shelf Science*, 164, 353–366.

572 Bakker, S.A. (2017, May). *Coastal protection in the Mekong Delta: Investigation of shore*
573 *nourishment and mangroves as Building with Nature solutions*.

574 Baumgartner, U., Kell, S., & Nguyen, T. H. (2016). Arbitrary mangrove-to-water ratios
575 imposed on shrimp farmers in Vietnam contradict with the aims of sustainable forest
576 management. *SpringerPlus*, 5, 438–438. <https://doi.org/10.1186/s40064-016-2070-3>

577 Binh, C. T., Phillips, M., & Demaine, H. (2008). *Integrated shrimp-mangrove farming*
578 *systems in the Mekong Delta of Vietnam* (Vol. 28). [https://doi.org/10.1046/j.1365-](https://doi.org/10.1046/j.1365-2109.1997.00901.x)
579 [2109.1997.00901.x](https://doi.org/10.1046/j.1365-2109.1997.00901.x)

580 Binh, T. N. K. D., Vromant, N., Hung, N. T., Hens, L., & Boon, E. K. (2005). Land Cover
581 Changes Between 1968 and 2003 In Cai Nuoc, Ca Mau Peninsula, Vietnam.
582 *Environment, Development and Sustainability*, 7(4), 519–536.
583 <https://doi.org/10.1007/s10668-004-6001-z>

584 Blankespoor, B., Dasgupta, S., & Laplante, B. (2014). Sea-level rise and coastal wetlands.
585 *Ambio*, 43(8), 996–1005. <https://doi.org/10.1007/s13280-014-0500-4>

586 Conchedda, G., Durieux, L., & Mayaux, P. (2008). An object-based method for mapping and
587 change analysis in mangrove ecosystems. *Theme Issue: Remote Sensing of the*
588 *Coastal Ecosystems*, 63(5), 578–589. <https://doi.org/10.1016/j.isprsjprs.2008.04.002>

589 Congalton, R. G. (1991). A review of assessing the accuracy of classifications of remotely
590 sensed data. *Remote Sensing of Environment*, 37(1), 35–46.
591 [https://doi.org/10.1016/0034-4257\(91\)90048-B](https://doi.org/10.1016/0034-4257(91)90048-B)

592 Cosslett, T. L., C., P. D. (2014). *Water resources and food security in the Vietnam Mekong*
593 *Delta. Switzerland: Springer International Publishing.*

594 Dan, T. T. (2017). *The Change of Vegetation in Mekong Delta and Its Effect for the*
595 *Hydrological Mechanism Using GIS and Remote Sensing.*

596 Danielsen, F., K Sørensen, M., Olwig, M., Selvam, V., Parish, F., Burgess, N., ...
597 Suryadiputra, N. (2005). The Asian Tsunami: A Protective Role for Coastal
598 Vegetation. *Science (New York, N.Y.)*, 310, 643.
599 <https://doi.org/10.1126/science.1118387>

600 DARD. (2017). *Department of Agriculture and Rural Development.* Soc Trang and Bac Lieu.

601 Dasgupta, S., Sobhan, I., & Wheeler, D. (2017). The impact of climate change and aquatic
602 salinization on mangrove species in the Bangladesh Sundarbans. *Ambio*, 46(6), 680–
603 694. <https://doi.org/10.1007/s13280-017-0911-0>

604 De Graaf, G., & Xuan, T. (1998). Extensive shrimp farming, mangrove clearance and marine
605 fisheries in the southern provinces of Vietnam. *Mangroves and Salt Marshes*, 2(3),
606 159–166.

607 Donato, D. C., Kauffman, J. B., Murdiyarso, D., Kurnianto, S., Stidham, M., & Kanninen,
608 M. (2011). Mangroves among the most carbon-rich forests in the tropics. *Nature*
609 *Geoscience*, 4, 293.

610 Duncan, C., Primavera, J. H., Pettorelli, N., Thompson, J. R., Loma, R. J. A., & Koldewey,
611 H. J. (2016). Rehabilitating mangrove ecosystem services: A case study on the
612 relative benefits of abandoned pond reversion from Panay Island, Philippines.
613 *Turning the Tide on Mangrove Loss*, 109(2), 772–782.
614 <https://doi.org/10.1016/j.marpolbul.2016.05.049>

615 Ellison, A. M. (2000). Mangrove restoration: Do we know enough? *Restoration Ecology*,
616 8(3), 219–229.

617 Evers, H.-D., & Benedikter, S. (2009). *Strategic group formation in the Mekong Delta: The*
618 *development of a modern hydraulic society*. ZEF Working Paper Series.

619 FAO. (2015). *Mangrove-related policy and institutional framework in Vietnam*. Viet Nam.

620 Feller, I. C., Friess, D. A., Krauss, K. W., & Lewis, R. R. (2017). The state of the world's
621 mangroves in the 21st century under climate change. *Hydrobiologia*, 803(1), 1–12.
622 <https://doi.org/10.1007/s10750-017-3331-z>

623 Gautier, D., Amador, J., & Newmark, F. (2001). The use of mangrove wetland as a biofilter
624 to treat shrimp pond effluents: Preliminary results of an experiment on the Caribbean
625 coast of Colombia. *Aquaculture Research*, 32(10), 787–799.

626 Giri, C., Ochieng, E., Tieszen, L. L., Zhu, Z., Singh, A., Loveland, T., ... Duke, N. (2011).
627 Status and distribution of mangrove forests of the world using earth observation
628 satellite data. *Global Ecology and Biogeography*, 20(1), 154–159.
629 <https://doi.org/10.1111/j.1466-8238.2010.00584.x>

630 Giri, Chandra, Long, J., Abbas, S., Murali, R. M., Qamer, F. M., Pengra, B., & Thau, D.
631 (2015). Distribution and dynamics of mangrove forests of South Asia. *Land*
632 *Cover/Land Use Change (LC/LUC) and Environmental Impacts in South Asia*, 148,
633 101–111. <https://doi.org/10.1016/j.jenvman.2014.01.020>

634 Giri, Chandra, Pengra, B., Zhu, Z., Singh, A., & Tieszen, L. L. (2007). Monitoring mangrove
635 forest dynamics of the Sundarbans in Bangladesh and India using multi-temporal
636 satellite data from 1973 to 2000. *Estuarine, Coastal and Shelf Science*, 73(1), 91–
637 100. <https://doi.org/10.1016/j.ecss.2006.12.019>

638 H. Oba, M. Hirota, R. Chbeir, H. Ishikawa, & S. Yokoyama. (2014). Towards Better Land
639 Cover Classification Using Geo-tagged Photographs. *2014 IEEE International*
640 *Symposium on Multimedia*, 320–327. <https://doi.org/10.1109/ISM.2014.78>

641 Ha, T., Dieperink, C., Otter, H. S., & Hoekstra, P. (2018). Governance conditions for adaptive
642 freshwater management in the Vietnamese Mekong Delta. *Journal of Hydrology*, 557,
643 116–127.

644 Ha, T. T. P., van Dijk, H., & Visser, L. (2014). Impacts of changes in mangrove forest
 645 management practices on forest accessibility and livelihood: A case study in
 646 mangrove-shrimp farming system in Ca Mau Province, Mekong Delta, Vietnam.
 647 *Land Use Policy*, 36, 89–101. <https://doi.org/10.1016/j.landusepol.2013.07.002>
 648 Hak, D., Nadaoka, K., Patrick Bernado, L., Le Phu, V., Hong Quan, N., Quang Toan, T., ...
 649 Pham Dang Tri, V. (2016). Spatio-temporal variations of sea level around the Mekong
 650 Delta: Their causes and consequences on the coastal environment. *Hydrological*
 651 *Research Letters*, 10(2), 60–66. <https://doi.org/10.3178/hr1.10.60>
 652 Hall, F. G., Townshend, J. R., & Engman, E. T. (1995). Status of remote sensing algorithms
 653 for estimation of land surface state parameters. *Remote Sensing of Environment*,
 654 51(1), 138–156.
 655 Hamilton, S. E., & Casey, D. (2016). Creation of a high spatio-temporal resolution global
 656 database of continuous mangrove forest cover for the 21st century (CGMFC-21).
 657 *Global Ecology and Biogeography*, 25(6), 729–738.
 658 <https://doi.org/10.1111/geb.12449>
 659 Hong, P. (1995). Tác động của việc nuôi quảng canh tôm đến môi trường và tài nguyên đa
 660 dạng sinh học vùng đất ngập nước ven biển (Impact of shrimp culture extension on
 661 environment and biodiversity resources in marsh of coastal zone). *Journal of*
 662 *Fisheries*, 25, 6–9.
 663 <https://earthexplorer.usgs.gov/>. (n.d.).
 664 [http://www.fao.org/fileadmin/templates/rap/files/meeting/2016/161220_05_Vietnam_polic](http://www.fao.org/fileadmin/templates/rap/files/meeting/2016/161220_05_Vietnam_policy_presentation.pdf)
 665 [y_presentation.pdf](http://www.fao.org/fileadmin/templates/rap/files/meeting/2016/161220_05_Vietnam_policy_presentation.pdf). (n.d.).
 666 Huu, P. C. (2011). Planning and implementation of the dyke systems in the Mekong Delta,
 667 Vietnam. *Unpublished PhD Thesis. Faculty of Mathematics and Natural Sciences,*
 668 *Rheinischen Friedrich-Wilhelms, University of Bonn, Bonn.*
 669 Islam, M. M., Borgqvist, H., & Kumar, L. (2018). *Monitoring Mangrove Forest Landcover*
 670 *Changes in the Coastline of Bangladesh from 1976 to 2015.*
 671 <https://doi.org/10.1080/10106049.2018.1489423>

672 Joffe, O., Poortvliet, M., & Klerkx, L. (2018). Are shrimp farmers actual gamblers? An
673 analysis of risk perception and risk management behaviors among shrimp farmers in
674 the Mekong Delta. *Aquaculture*, 495.
675 <https://doi.org/10.1016/j.aquaculture.2018.06.012>

676 Johnston, D., Trong, N. V., Tien, D. V., & Xuan, T. T. (2000). Shrimp yields and harvest
677 characteristics of mixed shrimp–mangrove forestry farms in southern Vietnam:
678 Factors affecting production. *Aquaculture*, 188(3), 263–284.
679 [https://doi.org/10.1016/S0044-8486\(00\)00348-3](https://doi.org/10.1016/S0044-8486(00)00348-3)

680 Kaly, U. L., & Jones, G. P. (1998). *Mangrove Restoration: A Potential Tool for Coastal*
681 *Management in Tropical Developing Countries*. 27(8), 6.

682 Kamal, M., & Phinn, S. (2011). Hyperspectral Data for Mangrove Species Mapping: A
683 Comparison of Pixel-Based and Object-Based Approach. *Remote Sensing*, 3(10).
684 <https://doi.org/10.3390/rs3102222>

685 Kuenzer, C., Bluemel, A., Gebhardt, S., Quoc, T. V., & Dech, S. (2011). Remote Sensing of
686 Mangrove Ecosystems: A Review. *Remote Sensing*, 3(5).
687 <https://doi.org/10.3390/rs3050878>

688 Lee, S. Y., Hamilton, S., Barbier, E. B., Primavera, J., & Lewis, R. R. (2019). Better
689 restoration policies are needed to conserve mangrove ecosystems. *Nature Ecology &*
690 *Evolution*, 3(6), 870–872. <https://doi.org/10.1038/s41559-019-0861-y>

691 Lee, T.-M., & Yeh, H.-C. (2009). Applying remote sensing techniques to monitor shifting
692 wetland vegetation: A case study of Danshui River estuary mangrove communities,
693 Taiwan. *Wetland Restoration and Ecological Engineering*, 35(4), 487–496.
694 <https://doi.org/10.1016/j.ecoleng.2008.01.007>

695 Lillesand, T., Kiefer, R. W., & Chipman, J. (2015). *Remote sensing and image interpretation*.
696 John Wiley & Sons.

697 Liu, K., Li, X., Shi, X., & Wang, S. (2008). Monitoring mangrove forest changes using
698 remote sensing and GIS data with decision-tree learning. *Wetlands*, 28(2), 336.
699 <https://doi.org/10.1672/06-91.1>

700 Locatelli, T., Binet, T., Kairo, J. G., King, L., Madden, S., Patenaude, G., ... Huxham, M.
 701 (2014). Turning the tide: How blue carbon and payments for ecosystem services
 702 (PES) might help save mangrove forests. *Ambio*, 43(8), 981–995.
 703 <https://doi.org/10.1007/s13280-014-0530-y>
 704 Long, J. B., & Giri, C. (2011). Mapping the Philippines' Mangrove Forests Using Landsat
 705 Imagery. *Sensors*, 11(3). <https://doi.org/10.3390/s110302972>
 706 MDP. (2011). *Towards a Mekong Delta Plan, Vietnam-Netherlands Cooperation, Water*
 707 *Sector Synthesis, Hanoi, Vietnam.*
 708 Minh, H. V. T., Avtar, R., Mohan, G., Misra, P., & Kurasaki, M. (2019). Monitoring and
 709 Mapping of Rice Cropping Pattern in Flooding Area in the Vietnamese Mekong Delta
 710 Using Sentinel-1A Data: A Case of An Giang Province. *ISPRS International Journal*
 711 *of Geo-Information*, 8(5). <https://doi.org/10.3390/ijgi8050211>
 712 Minh, T., Yakupitiyage, A., & Macintosh, D. (2001). *Management of the integrated*
 713 *mangrove-aquaculture farming systems in the Mekong delta of Vietnam.* Integrated
 714 Tropical Coastal Zone Management, School of Environment
 715 Mohajane, M., Essahlaoui, A., Oudija, F., El Hafyani, M., Hmadi, A. E., El Ouali, A., ...
 716 Teodoro, A. C. (2018). Land Use/Land Cover (LULC) Using Landsat Data Series
 717 (MSS, TM, ETM+ and OLI) in Azrou Forest, in the Central Middle Atlas of Morocco.
 718 *Environments*, 5(12), 131.
 719 Moreira, E., & Valeriano, M. (2014). Application and evaluation of topographic correction
 720 methods to improve land cover mapping using object-based classification.
 721 *International Journal of Applied Earth Observation and Geoinformation*, 32, 208–
 722 217. <https://doi.org/10.1016/j.jag.2014.04.006>
 723 Muttitanon, W., & Tripathi, N. (2005). *Land use/land cover changes in the coastal zone of*
 724 *Ban Don Bay, Thailand using Landsat 5 TM data* (Vol. 26).
 725 <https://doi.org/10.1080/0143116051233132666>
 726 Nguyen, T. N. (2016). Assessment of salt water intrusion in Kien Giang Province, Viet Nam,
 727 Master's thesis, Graduate School of Environmental Science, Hokkaido University,
 728 68pp.

729 Nguyen, D. B., Gruber, A., & Wagner, W. (2016). Mapping rice extent and cropping scheme
730 in the Mekong Delta using Sentinel-1A data. *Remote Sensing Letters*, 7(12), 1209–
731 1218. <https://doi.org/10.1080/2150704X.2016.1225172>

732 Nguyen, Tan Phong, & Parnell, K. E. (2019). Coastal land use planning in Ben Tre, Vietnam:
733 Constraints and recommendations. *Heliyon*, 5(4), e01487.

734 Nguyen, T.P., Van Tam, N., Quoi, L. P., & Parnell, K. E. (2016). Community perspectives
735 on an internationally funded mangrove restoration project: Kien Giang province,
736 Vietnam. *Ocean & Coastal Management*, 119, 146–154.
737 <https://doi.org/10.1016/j.ocecoaman.2015.10.008>

738 Orchard, S. E., Stringer, L. C., & Quinn, C. H. (2016). Mangrove system dynamics in
739 Southeast Asia: Linking livelihoods and ecosystem services in Vietnam. *Regional*
740 *Environmental Change*, 16(3), 865–879.

741 Ottinger, M., Clauss, K., & Kuenzer, C. (2016). *Aquaculture: Relevance, Distribution,*
742 *Impacts and Spatial Assessments – A Review* (Vol. 119).
743 <https://doi.org/10.1016/j.ocecoaman.2015.10.015>

744 Pham, L. T. H., Vo, T. Q., Dang, T. D., & Nguyen, U. T. N. (2019). Monitoring mangrove
745 association changes in the Can Gio biosphere reserve and implications for
746 management. *Remote Sensing Applications: Society and Environment*, 13, 298–305.
747 <https://doi.org/10.1016/j.rsase.2018.11.009>

748 Pham, T. D., Yoshino, K., Le, N. N., & Bui, D. T. (2018). Estimating aboveground biomass
749 of a mangrove plantation on the Northern coast of Vietnam using machine learning
750 techniques with an integration of ALOS-2 PALSAR-2 and Sentinel-2A data.
751 *International Journal of Remote Sensing*, 39(22), 7761–7788.
752 <https://doi.org/10.1080/01431161.2018.1471544>

753 Phan, L. K., van Thiel de Vries, J. S. M., & Stive, M. J. F. (2015). Coastal Mangrove Squeeze
754 in the Mekong Delta. *Journal of Coastal Research*, 233–243.
755 <https://doi.org/10.2112/JCOASTRES-D-14-00049.1>

756 Phan Nguyen Hong, H. T. S. (1993). *Mangroves of Viet Nam*.

- 757 Ramsey III, E., & R. Jensen, J. (1996). *Remote Sensing of Mangrove Wetlands: Relating*
758 *Canopy Spectra to Site-Specific Data* (Vol. 62).
- 759 Richards John, A., & Xiuping, J. (1999). *Remote sensing digital image analysis: An*
760 *introduction*.
- 761 Rouse Jr, J., Haas, R., Schell, J., & Deering, D. (1974). *Monitoring vegetation systems in the*
762 *Great Plains with ERTS*.
- 763 Ruiz-Luna, A., Escobar, A. C., & Berlanga-Robles, C. (2010). Assessing distribution
764 patterns, extent, and current condition of northwest Mexico mangroves. *Wetlands*,
765 30(4), 717–723.
- 766 Saenger, P., Hegerl, E.J., D., J. D. S. (1983). *Global status of mangrove ecosystems. The*
767 *Environmentalist*, 3 (Supplement 3): 1-88.
- 768 Sari, S. P., & Rosalina, D. (2016). Mapping and Monitoring of Mangrove Density Changes
769 on tin Mining Area. *The 2nd International Symposium on LAPAN-IPB Satellite*
770 *(LISAT) for Food Security and Environmental Monitoring*, 33, 436–442.
771 <https://doi.org/10.1016/j.proenv.2016.03.094>
- 772 Satyanarayana, B., M Muslim, A., Amira Izzaty Horsali, N., Mat Zauki, N. A., Otero, V.,
773 Nadzri, M., ... Dahdouh-Guebas, F. (2018). *Status of the undisturbed mangroves at*
774 *Brunei Bay, East Malaysia: A preliminary assessment based on remote sensing and*
775 *ground-truth observations* (Vol. 6). <https://doi.org/10.7717/peerj.4397>
- 776 Singh, C., Shashtri, S., Mukherjee, S., Avtar, R., & Singh, S. (2010). *Monitoring change in*
777 *land use and land cover in Rupnagar district of Punjab, India using Landsat and IRS*
778 *LISS III satellite data* (Vol. 13). <https://doi.org/10.12775/v10090-010-0018-8>
- 779 Singh, I. J., Singh, S. K., Kushwaha, S. P. S., Ashutosh, S., & Singh, R. K. (2004).
780 Assessment and monitoring of estuarine mangrove forests of Goa using satellite
781 remote sensing. *Journal of the Indian Society of Remote Sensing*, 32(2), 167–174.
782 <https://doi.org/10.1007/BF03030873>
- 783 Spalding, M., McIvor, A., Tonneijck, F., Tol, S., & Eijk, P. van. (2014). *Mangroves for*
784 *coastal defence*.

785 The People's Committee of Bac Lieu Province. (2017). *Forest status data in 2016 in Bac*
786 *Lieu province*. Bac Lieu, Viet Nam.

787 Thu, P. M., & Populus, J. (2007). Status and changes of mangrove forest in Mekong Delta:
788 Case study in Tra Vinh, Vietnam. *Sedimentological and Ecohydrological Processes*
789 *of Asian Deltas: The Yangtze and the Mekong*, 71(1), 98–109.
790 <https://doi.org/10.1016/j.ecss.2006.08.007>

791 Toan, T. (2014). Climate Change and Sea Level Rise in the Mekong Delta: Flood, Tidal
792 Inundation, Salinity Intrusion, and Irrigation Adaptation Methods. *Coastal Disasters*
793 *and Climate Change in Vietnam: Engineering and Planning Perspectives*, 199–217.

794 Tokola, T., Sarkeala, J., & Lin, M. (2001). Use of topographic correction in Landsat TM-
795 based forest interpretation in Nepal. *International Journal of Remote Sensing*, 22,
796 551–563. <https://doi.org/10.1080/01431160050505856>

797 Tong, P. H. S., Auda, Y., Populus, J., Aizpuru, M., Habshi, A. A., & Blasco, F. (2004).
798 Assessment from space of mangroves evolution in the Mekong Delta, in relation to
799 extensive shrimp farming. *International Journal of Remote Sensing*, 25(21), 4795–
800 4812. <https://doi.org/10.1080/01431160412331270858>

801 Townshend, J. G. (1992). Land cover. *International Journal of Remote Sensing*, 13(6–7),
802 1319–1328.

803 Tran, T. V., Tran, D. X., Myint, S. W., Huang, C., Pham, H. V., Luu, T. H., & Vo, T. M. T.
804 (2019). Examining spatiotemporal salinity dynamics in the Mekong River Delta using
805 Landsat time series imagery and a spatial regression approach. *Science of The Total*
806 *Environment*, 687, 1087–1097. <https://doi.org/10.1016/j.scitotenv.2019.06.056>

807 Truong, T. D., & Do, L. H. (2018). Mangrove forests and aquaculture in the Mekong river
808 delta. *Land Use Policy*, 73, 20–28. <https://doi.org/10.1016/j.landusepol.2018.01.029>

809 UNEP. (2014). *THE IMPORTANCE OF MANGROVES TO PEOPLE: A CALL TO ACTION*.

810 Van Coppenolle, R., Schwarz, C., & Temmerman, S. (2018). Contribution of Mangroves and
811 Salt Marshes to Nature-Based Mitigation of Coastal Flood Risks in Major Deltas of
812 the World. *Estuaries and Coasts*, 41(6), 1699–1711. [https://doi.org/10.1007/s12237-](https://doi.org/10.1007/s12237-018-0394-7)
813 [018-0394-7](https://doi.org/10.1007/s12237-018-0394-7)

814 Van Cuong, C., Brown, S., To, H. H., & Hockings, M. (2015). Using Melaleuca fences as
815 soft coastal engineering for mangrove restoration in Kien Giang, Vietnam. *Ecological*
816 *Engineering*, 81, 256–265. <https://doi.org/10.1016/j.ecoleng.2015.04.031>

817 Van Hao, N. (1999). *Shrimp Health Research in Vietnam, Including Current and Planned*
818 *Activities*. 94–97. ACIAR; 1998.

819 Vo, T., Oppelt, N., Leinenkugel, P., & Kuenzer, C. (2013). *Remote Sensing in Mapping*
820 *Mangrove Ecosystems—An Object-Based Approach* (Vol. 5).
821 <https://doi.org/10.3390/rs5010183>

822 Vo, T. Quoc, Oppelt, N., Leinenkugel, P., & Kuenzer, C. (2013). Remote Sensing in Mapping
823 Mangrove Ecosystems—An Object-Based Approach. *Remote Sensing*, 5(1).
824 <https://doi.org/10.3390/rs5010183>

825 Vo, Tuan Quoc, Kuenzer, C., & Oppelt, N. (2015). How remote sensing supports mangrove
826 ecosystem service valuation: A case study in Ca Mau province, Vietnam. *Ecosystem*
827 *Services*, 14, 67–75.

828 Wösten, J., De Willigen, P., Tri, N., Lien, T., & Smith, S. (2003). Nutrient dynamics in
829 mangrove areas of the Red River Estuary in Vietnam. *Estuarine, Coastal and Shelf*
830 *Science*, 57(1–2), 65–72.

831 Wylie, L., Sutton-Grier, A. E., & Moore, A. (2016). Keys to successful blue carbon projects:
832 Lessons learned from global case studies. *Marine Policy*, 65, 76–84.
833 <https://doi.org/10.1016/j.marpol.2015.12.020>

834 Xu, G., Zhu, X., Fu, D., Dong, J., & Xiao, X. (2017). Automatic land cover classification of
835 geo-tagged field photos by deep learning. *Environmental Modelling & Software*, 91,
836 127–134. <https://doi.org/10.1016/j.envsoft.2017.02.004>
837