Title	Histamine Production Behaviors of a Psychrotolerant Histamine-Producer, <i>Morganella psychrotolerans</i> , in Various Environmental Conditions
Author(s)	Wang, Di; Yamaki, Shogo; Kawai, Yuji; Yamazaki, Koji
Citation	Current microbiology, 77, 460-467 https://doi.org/10.1007/s00284-019-01853-y
Issue Date	2020-01-02
Doc URL	http://hdl.handle.net/2115/80113
Rights	This is a post-peer-review, pre-copyedit version of an article published in Current Microbiology. The final authenticated version is available online at: https://doi.org/10.1007/s00284-019-01853-y
Type	article (author version)
File Information	Wang_Blind revised.2.pdf

[Instructions for use](#)

Histamine production behaviors of a psychrotolerant histamine-producer,
***Morganella psychrotolerans*, in various environmental conditions**

Di Wang, Shogo Yamaki, Yuji Kawai, and Koji Yamazaki*

Laboratory of Marine Food Science and Technology, Faculty of Fisheries Sciences, Hokkaido
University, Minato, Hakodate, Hokkaido, 041-6811, Japan

*Corresponding author. Tel./fax: +81 138 40 5574, E-mail: yamasaki@fish.hokudai.ac.jp

Acknowledgements

This work was supported by JSPS KAKENHI (Grant Number JP17K07931).

Compliance with Ethical Standards

Conflict of Interest

The authors declare that they have no conflict of interest.

Abstract

Histamine food poisoning is a major safety concern related to seafood consumption worldwide. *Morganella psychrotolerans* is a novel psychrotolerant histamine-producer. In this study, the histamine production behaviors of *M. psychrotolerans* and two other major histamine-producers, mesophilic *Morganella morganii* and psychrotrophic *Photobacterium phosphoreum*, were compared in seafood products, and histamine accumulation by *M. psychrotolerans* was characterized at various pH and temperature levels in culture broth. The growth of *M. psychrotolerans* and *P. phosphoreum* increased similarly at 4°C in canned tuna, but *M. psychrotolerans* produced much higher levels of histamine than *P. phosphoreum*. Histamine accumulation by *M. psychrotolerans* was induced at lower environmental pH condition at 4 and 20 °C. The optimal temperature and pH for producing histamine by crude histidine decarboxylase of *M. psychrotolerans* was 30 °C and pH 7, respectively. The activity of the crude HDC extracted from *M. psychrotolerans* cells at 10 °C retained 45% of the activity at 30°C. Histidine decarboxylase gene expression of *M. psychrotolerans* was induced by low pH conditions. These results suggest that *M. psychrotolerans* are also a very important histamine-producer leading to histamine poisoning associated with seafood below the refrigeration temperature.

Histamine production behaviors of a psychrotolerant histamine-producer, *Morganella psychrotolerans*, in various environmental conditions

Introduction

Histamine fish poisoning is a foodborne toxicity caused by ingesting high levels of histamine [1]. The U.S. Food and Drug Administration (FDA) guideline for safe levels of histamine concentration in seafood is 50 mg/kg, while histamine concentration greater than 500 mg/kg is hazardous to humans [2]. Poisoning symptoms include rash, headache, nausea, diarrhea, and so on [3]. Although many cases show mild symptoms, rare cases have shown a potential to threaten human life [4, 5]. Histamine accumulation in food is caused by histamine-producing bacteria that produce histidine decarboxylase (HDC), an enzyme that decarboxylates free histidine to histamine [6]. Once histamine is formed in food, it cannot be destroyed by common food processing treatments such as heating and freezing [7].

Histamine-producing bacteria are categorized into low and high histamine-producers. The bacteria producing more than 1,000 mg/L of histamine in culture broth are classified as high histamine-producing bacteria [8]. Generally, mesophilic bacteria such as *Morganella morganii*, *Photobacterium damsela*, and *Klebsiella pneumoniae* are well documented as high histamine-producing bacteria in fishery products. These mesophilic histamine-producers will prevent forming histamine by controlling the storage temperature and time during the transports, processing [9]. However, psychrotrophic histamine-producers such as some *Photobacterium* spp. capable of produce histamine to a harmful level in seafood at

refrigeration temperature, and normal cold chain cannot completely suppress histamine production by them. *P. kishitanii* and *P. aquimaris* were recently isolated from fish [10], and *P. phosphoreum* caused histamine food poisoning incident in Japan [11].

M. psychrotolerans is a novel psychrotolerant histamine-producer, that can grow at 0–2 °C. Its biochemical characteristics are very similar to those of *M. morganii* [12]. Previous studies have documented the poisoning incidents caused by *M. psychrotolerans* [13, 14, 15]. In our previous study, we suggested that *M. psychrotolerans* is commonly distributed through retail seafood and accumulates histamine in culture broth during low-temperature incubation [16].

In this study, we compared the histamine production abilities of *M. psychrotolerans*, *P. phosphoreum*, and *M. morganii* in canned tuna and investigated the effect of environmental factors (pH and temperature) on histamine production by *M. psychrotolerans* in culture broth. The activities of crude HDC and *hdc* gene expression levels were also characterized.

Materials and methods

Bacterial strains and growth conditions

M. psychrotolerans JCM 16473^T and *M. morganii* NBRC 3848^T were inoculated in tryptic soy broth (TSB, BD, Franklin Lakes, NJ) supplemented with 1% L-histidine (Wako Pure Chemical Industries, Osaka, Japan) at 25 and 30 °C (TSBH, pH 6.0), respectively. *P. phosphoreum* NBRC 13896 was cultured in TSBH supplemented with 1.5% NaCl (Final

NaCl concentration in medium: 2%) at 25 °C. The cultures were incubated for 24 h for the experiments described below.

Inoculation with bacteria to tuna samples

Canned tuna (Hagoromo Food co. Ltd., Shizuoka, Japan) was purchased from a local supermarket. A 100 g of tuna was aseptically put into a stomacher bag (Seward Ltd., Worthing, U.K.). The precultured bacteria were centrifuged (10,000×g, 2 min, 4 °C) and washed twice with phosphate buffered saline (PBS, pH 7.2). After serial dilution by PBS, each bacterial suspension was inoculated to tuna samples (populations about 3 log CFU/g), and the samples were stored at 4 or 20 °C. A 5 g of the sample was aseptically weighted in a stomacher bag, homogenized in 45 mL PBS at 200 rpm for 1 min, and used to determine total plate counts and histamine accumulation. The homogenized solutions were serially diluted with PBS and spread onto plates. *M. psychrotolerans* or *M. morganii* colonies were counted after incubating for 48 h at 25 or 30 °C on tryptic soy agar (TSA, BD, Franklin Lakes, NJ), respectively. *P. phosphoreum* colonies was counted after incubating for 48 h at 25 °C on TSA-SC agar (TSA supplemented with 1.5% NaCl). Histamine contents were determined with the histamine test kit (Kikkoman Biochemifa Company, Tokyo, Japan), according to the manufacturer's instructions.

***M. psychrotolerans* growth and histamine production under different environmental conditions**

Pre-cultured *M. psychrotolerans* was washed and diluted with PBS. Then, 100 μ L of the bacteria suspension was mixed with 9.9 mL TSB supplemented with 1% L-histidine (TSBH, pH 5 - 8, adjusted with 1M NaOH or HCl). The working culture (approximately about 5.5 log CFU/mL of an initial inoculum size) was incubated at 4 or 20 °C. At each sampling time, the bacterial population and the histamine concentration in the culture medium were determined.

Activity of Cell-free crude HDC under different environmental conditions

The 5 mL of precultured *M. psychrotolerans* cells were harvested by centrifugation (10,000 $\times$ g, 5 min, 4 °C), and washed twice with McIlvaine buffer (0.01M citrate-0.02 M phosphate buffer pH6.2). The bacterial cells were resuspended in 1 mL of the same buffer (approximately 9.51 log CFU/mL in working buffer), and disrupted using zirconia beads (zircon prep mini, Nippon Genetics Co. Ltd., Tokyo, Japan) and a Bead Beater (2,500 rpm, 15 min, CD-1000, EYELA, Tokyo, Japan). A cell-free supernatant was collected as a crude HDC solution by centrifugation (10,000 $\times$ g, 15 min, 4 °C). The crude HDC solution (0.2 mL) was mixed with the reaction buffer (1.8 mL, 0.1M citrate-0.2 M phosphate buffer) supplemented with 1% L-histidine. After 1 h incubation, the reaction terminated by heating at 100°C for 5 min. The activity of crude HDC was determined by checking the concentration of histamine in the solution at conditions ranging from pH 4-8 (30 °C) and 10-50 °C (pH 6.2). The crude HDC solution was directly used for the analysis of extraction.

***M. psychrotolerans* hdc gene expression by quantitative RT-PCR**

Gene expression levels of *hdc* at different pH (5 - 8) and temperature levels (4 or 20 °C) were quantified by quantitative RT-PCR (qRT-PCR). The *hdc* and reference gene (16S rDNA) primers are shown in Table 1. Each primer was designed according to previously reported nucleotide sequences [6, 17]. RNA extraction was extracted using NucleoSpin RNA extraction kit (Macherey-Nagel GmbH, Germany) from *M. psychrotolerans* cells cultured at 4 °C for 6 d or 20 °C for 24 h. The concentration and purity (A_{260}/A_{280}) ratio of all RNA samples were checked with a spectrophotometer (GeneQuant pro, UK). The RNA was stored at - 70 °C and subsequently used for cDNA synthesis. The cDNA was synthesized using RevertAid First strand cDNA Synthesis Kit (Fermentas, Thermo Fisher Scientific Inc. USA). The contamination of DNA residues was determined in RNA sample by a control reaction that had a cDNA synthesis without reverse transcriptase enzyme. A 20 µL reaction mixture for qRT-PCR contained 10 µL SsoAdvanced™ Universal SYBR® Green Supermix (Bio-Rad Laboratory Inc., Hercules, CA, USA), 1 µL cDNA, 3 µL primer mix (final concentration: 300 nM of each primer), and 6 µL water. The PCR conditions were an initial cycle at 95 °C for 45 s, which was followed by 40 cycles at 95 °C for 15 s, and 61 °C for 1 min in an ABI 7500 Real-Time PCR system (Applied Biosystems, Foster City, CA, USA). For each target gene, the real-time PCR efficiency was calculated according to the equation $E = 10^{(-1/\text{slope})}$. The gene expression level was calculated by the Pfaffl method [18].

Statistical analysis

All the experiments were carried out in triplicate. The mean values and standard deviation were calculated from the triplicate results. The statistical analyses were performed using Tukey's *post hoc* range method. Significance level was based on 5% ($P < 0.05$). All statistical analyses were done using RStudio Desktop (RStudio Desktop, Inc., Boston, MA).

Results

Histamine accumulation in canned tuna

To evaluate the histamine production capabilities of *M. psychrotolerans* in canned tuna, we compared them with major mesophilic and psychrophilic histamine-producers, *M. morganii* and *P. phosphoreum*. Changes in viable cell counts and histamine concentration at 4 and 20 °C are shown in Fig. 1. At 20 °C, the growth curves of the three bacteria showed similar trends. The numbers of the viable cell count increased to more than 6 log CFU/g after 12 h from the initial counts, and exceeded 8 log CFU/g after 36 h (Fig. 1-A). Histamine concentrations in the samples inoculated with *M. psychrotolerans* and *M. morganii* exceeded 1,000 mg/kg after 36 h, and were 3,000 mg/kg after 60 h. However, histamine concentration in samples with *P. phosphoreum* was less than 400 mg/kg after 60 h (Fig. 1-B). When the tuna samples inoculated with the three bacteria were stored at 4 °C, the viable cell counts of *M. psychrotolerans* and *P. phosphoreum* were detected to be more than 6 log CFU/g after 4 d, and reached to about 8 log CFU/g after 8 d. However, *M. morganii* did not grow at 4 °C (Fig. 1-C). Histamine production by *M. psychrotolerans* and *P. phosphoreum* were 1,467, 2,611

and 556, 883 mg/kg in canned tuna after 8 d and 10 d at 4 °C, respectively (Fig. 1-D).

Effect of pH and temperature on *M. psychrotolerans* growth and histamine accumulation in culture broth

The *M. psychrotolerans* growth and histamine accumulation at different pH and temperature levels are shown in Fig. 2. At 20 °C, the cell populations rapidly increased to approximately 9 log CFU/mL at pH 6, 7, and 8 after 24 h. The viable cells also reached 9 log CFU/mL at pH 5 after 36 h (Fig. 2-A). High levels of histamine (> 2,600 mg/L) were produced by *M. psychrotolerans* under all pH conditions after 36 h (Fig. 2-B). At 4 °C, the cell populations showed a rapid increase at pH 6 - 8, and the viable cell counts reached to 9 log CFU/mL after 6 d (Fig. 2-C). Significant histamine accumulations (> 1,700 mg/L) were observed after 6 d at pH 5, 6, and 7 (Fig. 2-D). The histamine concentrations increased with decreasing pH at both pH conditions (after 48 h and 8 d incubated at 20 °C and 4 °C, respectively).

Effect of temperature and pH on cell-free crude HDC activity

Effects of temperature and pH on crude HDC extracted from *M. psychrotolerans* cells are shown in Fig. 3. The results showed that the crude HDC exhibited the most active at 30 °C, and it showed about two-fold higher compared with those at 10, 20, and 40 °C. However, the HDC activity at 50 °C was 30% of the activity at 30 °C (Fig. 3-A). Also, crude HDC activity was greatly affected by the environmental pH. The maximum activity was shown at pH 7.

The HDC activities at pH 4, 5, 6, and 8 were 16%, 49%, 76% and 48%, respectively, compared with that at pH 7 (Fig. 3-B).

Effect of pH and temperature on *hdc* gene expression levels

Changes of *hdc* gene expression at different pH and temperature levels are shown in Fig. 4. The *hdc* gene expressions were also affected by the environmental pH. Expression levels decreased with increasing pH, and the expression levels at pH 5 were 9.1-fold and 14.7-fold higher at 20 and 4 °C, respectively, than the levels at pH 8.

Discussion

M. morganii and some of *Photobacterium* spp. like *P. phosphoreum*, *P. kishitanii* and *P. aquimaris* are recognized as major mesophilic and psychrophilic histamine-producing bacteria [11, 19], respectively. The effects of environmental factors on histamine accumulation by psychrotolerant *M. psychrotolerans* have not been studied well yet. Therefore, to avoid the histamine poisoning associated with seafood, in-depth studies on the relations between environmental factors and the histamine accumulation by *M. psychrotolerans* are needed. In this study, *M. psychrotolerans* and *M. morganii* produced similar amounts of histamine at room temperature (20 °C) during the incubation. In addition, *M. psychrotolerans* produced greater amount of histamine than *P. phosphoreum* at 20 °C (Fig. 1-B). Moreover, *M. psychrotolerans* produced higher amount of histamine (more than 1,000 mg/kg at 8 d) than *P. phosphoreum* (about 556 mg/kg) at low temperature (4 °C) (Fig. 1-D). *P.*

phosphoreum NBRC 13896 has reported that be classified as *P. kishitanii*, a more prolific histamine producing bacteria at low temperatures [8]. And Bjornsdottir-Butler [8] reported that *P. kishitanii* produced ≤ 1545 mg/L histamine in LSW-70 with 1% L-histidine at 20 °C after 48 h. However, *M. psychrotolerans* produced ≥ 3700 mg/L histamine in TSBH at 20 °C after 48 h in our study (Fig. 2-B). *Photobacterium spp.* as the marine bacteria, it requires salt for growing, and the optimum activity for producing histamine was shown at 1-4% NaCl concentration [20, 21]. On the contrary, Emborg *et al* [12] have reported *M. psychrotolerans* can grow at pH 4.6 - 9.2, temperature 2 - 35 °C, and does not require salt for its growth. The NaCl concentration of the canned tuna used in this study was about $0.14 \pm 0.01\%$ (Data not shown). The salt concentration of common raw fish muscle is about 0.2-0.4% [22, 23, 24]. This could be a reason that *M. psychrotolerans* showed higher ability for producing histamine than *P. phosphoreum* NBRC 13896 in this study. In our previous study, *M. psychrotolerans* has been found from 40.6% retail seafood in Japan [16]. *M. morganii* and *M. psychrotolerans* exhibit very similar biochemical characteristics, and cannot be distinguished with the API50CH-E kit (BioMérieux, France), commonly used for rapid bacterial identification [12]. Therefore, some of the isolates identified as *M. morganii* for a causative microorganism on histamine outbreaks associated with seafood have been reasonably suspected to be *M. psychrotolerans*.

Although no big differences were observed in the growth curves of *M. psychrotolerans* under various pH conditions, the histamine accumulations were greatly affected by environmental pH at both temperatures (Fig. 2). Namely, *M. psychrotolerans* produced high

amount of histamine at low environmental pH conditions (Fig. 2-B and D). These findings were consistent with the results found by Torres et al [25], who showed that the histamine production by *M. morganii* was reduced at higher pH than 6.6. Morii et al [21] have also reported the optimum culture pH for histamine production by *P. phosphoreum* to be 5.5 - 6.5.

Maximum crude HDC activity was observed at pH 7 (Fig. 3-B). Similarly, the HDC activity of other gram-negative histamine-producers, *M. morganii* [26] and *E. aerogenes* [27], was optimal at pH 6.5. The crude HDC activity of *M. psychrotolerans* at pH 4 was 18% of the activity at pH 7 (Fig. 3-B). However, the expression level of the *hdc* gene in *M. psychrotolerans* cells at low pH condition was much higher than that at neutral pH condition (Fig. 4). The decarboxylation of histidine by histamine-producers requires *hdc* gene expression and HDC enzyme synthesis in the cells. Moriii [21] reported the crude HDC from *P. phosphoreum* cells exhibited no activity at pH 4. Although *M. psychrotolerans* did not grow at pH 4 in culture broth, its crude HCD still showed an activity (Fig. 3-B). Hence, it indicates that the histamine-producing ability of *M. psychrotolerans* under low environmental pH is higher than that of *P. phosphoreum*. The optimum temperature for the *M. psychrotolerans* crude HDC was at 30 °C, and the crude HDC maintained 45% of its optimum activity at 10 °C (Fig. 3-A). Bjornsdottir-Butler et al. [20] reported that the optimum activity of the HDC of *P. kishitanii* and *P. angustum* were at 30 °C. And Moriii [21] reported that the crude HDC from *P. phosphoreum* at 10 °C retained 10 % activity of the optimum temperature. The pH of fish muscle is 5.5 - 6.5 [25, 28]. Also, it is often adjusted with organic acids to inhibit the growth of microorganisms. From these findings, it can be argued that *M. psychrotolerans* is

more important than other psychrophilic histamine producers as a risk factor for histamine poisoning associated with seafood during low-temperature storage. However, the crude HDC activity of *M. psychrotolerans* decreased at alkaline pH (pH 8) (Fig. 3-B). Corresponding with previous studies reported that the HDC activity from *M. morganii* [26], *P. phosphoreum* [29], and *Streptococcus thermophilus* [30] were greatly reduced at pH 8. Bjornsdottir-Butler et al [28] have shown an excellent strategy for reducing histamine accumulation in fish muscle: pH elevation by trisodium phosphate treatment. From the results of this study, pH elevation can be one of the effective methods to suppress the histamine production by *M. psychrotolerans* in seafood as well.

The *hdc* gene expression of *M. psychrotolerant* was greatly induced at acidic conditions at both 4 and 20 °C (Fig. 4). This phenomenon in *M. psychrotolerans* agrees with other histamine-producing bacteria. Many reports indicate that low pH is a necessary factor for inducing *hdc* gene expression in *M. morganii* [31, 32], *P. iliopiscarium* [1], *P. damsela* subsp. *damsela* [32], and *S. thermophiles* [33]. Histidine decarboxylase cluster of *M. morganii* is composed of *hdcT1* (putative histidine/histamine antiporters), *hdc*, *hdcT2* (putative histidine/histamine antiporters), and *hisRS* (histidyl-tRNA synthetase), suggesting that bacterial cells incorporate extracellular histidine and excrete histamine to extracellular environment [31, 32]. The initial acidic to neutral pH enhances the *hdc* gene expression, forming HDC and histamine, which is excreted from the cells. The bacterial cells sense the acidic pH conditions, which can turn the histidine over to histamine and CO₂. Thus, a large of H⁺ were eliminate and the bacteria could escape from acid stress through the decarboxylation

reaction of the amine as a survival strategy [31, 32, 34, 35, 36].

This study is the first report on the histamine production behaviors of *M. psychrotolerans* in various environmental conditions and characterization of the crude HDC activity and *hdc* gene expression level of *M. psychrotolerans*. Our results indicate that *M. psychrotolerans* has a higher capacity to produce histamine toxicologically in seafood under low temperature than *P. phosphoreum*. Results of this study could be useful for improving seafood safety and developing effective strategies to reduce a risk of histamine poisoning by *M. psychrotolerans*. However, further studies are needed to construct suitable strategies, especially for disinfection and suppression of *M. psychrotolerans* growth in seafood.

References

1. Takahashi H, Ogai M, Miya S, Kuda T, Kimura B (2015) Effects of environmental factors on histamine production in the psychrophilic histamine-producing bacterium *Photobacterium iliopiscarium*. Food Control 52:39–42.
<https://doi.org/10.4315/0362-028X.JFP-18-022>
2. U.S. Food and Drug Administration. (2011) Chapter 7. Scombrotoxin (histamine) formation. In fish and fishery products hazards and controls guidance (4rd ed.). (pp. 113-152).
3. Taylor SL (1986) Histamine food poisoning: toxicology and clinical aspects. Crit Rev Toxicol 17:91–128. <https://doi.org/0.3109/10408448609023767>

- 251 4. D'Aloia A, Vizzardi E, Pina Della P, Bugatti S, Del Magro F, Raddino R, Curnis A,
252 Dei Cas L (2011) A scombroid poisoning causing a life-threatening acute pulmonary
253 edema and coronary syndrome in a young healthy patient. *Cardiovasc Toxicolo*
254 11:280–283. <https://doi.org/10.1007/s12012-011-9115-1>
- 255 5. SanchezGuerrero IM, Vidal JB, Escudero AI (1997) Scombroid fish poisoning: a
256 potentially life-threatening allergic-like reaction. *J Allergy Clin Immunol* 100:433–
257 434. [https://doi.org/10.1016/S0091-6749\(97\)70263-X](https://doi.org/10.1016/S0091-6749(97)70263-X)
- 258 6. Ferrario C, Ricci G, Borgo F, Fortina MG (2012) Species-specific DNA probe and
259 development of a quantitative PCR assay for the detection of *Morganella morganii*.
260 *Letters in Applied Microbiology* 54:292–298.
261 <https://doi.org/10.1111/j.1472-765X.2012.03206.x>
- 262 7. Etkind P, Wilson ME, Gallagher K, Cournoyer J (1987) Bluefish-associated
263 scombroid poisoning. An example of the expanding spectrum of food poisoning from
264 seafood. *JAMA* 258:3409–3410. <https://doi.org/10.1001/jama.1987.03400230069034>
- 265 8. Bjornsdottir-Butler K, Abraham A, Harper A, Dunlap PV, Benner RA Jr. (2018)
266 Biogenic amine production by and phylogenetic analysis of 23 *Photobacterium*
267 species. *Journal of Food Protection* 81:1264–1274.
268 <https://doi.org/10.4315/0362-028X.JFP-18-022>
- 269 9. Bjornsdottir-Butler K, Bencsath FA, McCarthy S, Benner RA Jr. (2017) Heat

- 270 resistance of histidine decarboxylase from gram-negative histamine-producing
271 bacteria in seafood. Journal of Food Protection 80:1273–1279.
272 <https://doi.org/10.4315/0362-028X.JFP-17-008>
- 273 10. Torido Y, Ohshima C, Takahashi H, Miya S, Iwakawa A, Kuda T, Kimura B (2014)
274 Distribution of psychrophilic and mesophilic histamine-producing bacteria in retailed
275 fish in Japan. Food Control 46:338–342.
276 <https://doi.org/10.1016/j.foodcont.2014.05.045>
- 277 11. Kanki M, Yoda T, Ishibashi M, Tsukamoto T (2004) *Photobacterium phosphoreum*
278 caused a histamine fish poisoning incident. International Journal of Food
279 Microbiology 92:79–87. <https://doi.org/10.1016/j.ijfoodmicro.2003.08.019>
- 280 12. Emborg J (2006) *Morganella psychrotolerans* sp. nov., a histamine-producing
281 bacterium isolated from various seafoods. International Journal of Systematic and
282 Evolutionary Microbiology 56:2473–2479. <https://doi.org/10.1099/ijs.0.64357-0>
- 283 13. Emborg J, Dalgaard P (2006) Formation of histamine and biogenic amines in
284 cold-smoked tuna: an investigation of psychrotolerant bacteria from samples
285 implicated in cases of histamine fish poisoning. Journal of Food Protection 69:897–
286 906. <https://doi.org/10.4315/0362-028X-69.4.897>
- 287 14. Emborg J, Laursen BG, Dalgaard P (2005) Significant histamine formation in tuna
288 (*Thunnus albacares*) at 2 °C—effect of vacuum- and modified atmosphere-packaging

- 289 on psychrotolerant bacteria. International Journal of Food Microbiology 101:263–279.
290 <https://doi.org/10.1016/j.ijfoodmicro.2004.12.001>
- 291 15. Torger Børresen. (2008) Improving seafood products for the consumer (pp. 292–324).
292 Abington, England: Woodhead Publishing Ltd.
- 293 16. Kato R, Wang D, Yamaki S, Kawai Y, Yamazaki K (2017) Distribution on fishery
294 products and histamine producing ability of *Morganella psychrotolerans*, a
295 psychrotrophic histamine-producing bacterium. Jpn J Food Microbiol 34:158–165 (in
296 Japanese). <https://doi.org/10.5803/jsfm.34.158>
- 297 17. Bjornsdottir-Butler K, Leon MS, Benner RA Jr. (2016) Draft genome sequences of
298 histamine-producing *Morganella psychrotolerans* strains. Genome Announc 4:2473–1.
299 <https://doi.org/10.1128/genomeA.01001-16>
- 300 18. Pfaffl MW (2001) A new mathematical model for relative quantification in real-time
301 RT-PCR. Nucleic Acids Res. 29. <https://doi.org/10.1093/nar/29.9.e45>
- 302 19. Kim SH, An H, Wei CI, Visessanguan W, Benjakul S, Morrissey MT, Su YC, Pitta TP
303 (2003) Molecular detection of a histamine former, *Morganella morganii*, in albacore,
304 mackerel, sardine, and a processing plant. Journal of Food Science 68:453–457.
305 <https://doi.org/10.1111/j.1365-2621.2003.tb05693.x>
- 306 20. Bjornsdottir-Butler K, McCarthy SA, Dunlap PV, Benner RA Jr. (2016)
307 *Photobacterium angustum* and *Photobacterium kishitanii*, Psychrotrophic high-level

308 histamine-producing bacteria indigenous to tuna. Applied and Environmental
309 Microbiology 82:2167–2176. <https://doi.org/10.1128/AEM.02833-15>

310 21. Morii H, Izumi Y, Kasama K, Ishimoto R (1994) Factors influencing histamine
311 formation by psychrotrophic luminous bacteria *Photobacterium phosphoreum*. Nippon
312 Suisan Gakkaishi 60:773–777. <https://doi.org/10.2331/suisan.60.773>

313 22. Geirsdottir M, Hlynisdottir H, Thorkelsson G, Sigurgisladdottir S (2007) Solubility and
314 viscosity of herring (*Clupea harengus*) proteins as affected by freezing and frozen
315 storage. Journal of Food Science 72:C376–C380.
316 <https://doi.org/10.1111/j.1750-3841.2007.00471.x>

317 23. Manthey-Karl M, Lehmann I, Ostermeyer U, Schröder U (2016) Natural chemical
318 composition of commercial fish species: Characterisation of pangasius, wild and
319 farmed turbot and barramundi. Foods 5:58–14. <https://doi.org/10.3390/foods5030058>

320 24. Thorarinsdottir KA, Arason S, Bogason SG, Kristbergsson K (2004) The effects of
321 various salt concentrations during brine curing of cod (*Gadus morhua*). International
322 Journal of Food Science and Technology 39:79–89.
323 <https://doi.org/10.1046/j.0950-5423.2003.00757.x>

324 25. Torres S, Roeckel M, Marti MC (2002) Histamine formation by *Morganella morganii*
325 isolated from *Trachurus murphyi* (chilean mackerel). Latin American Applied
326 Research 32:209–214.

- 327 26. Tanase S, Guirard BM, Snell EE (1985) Purification and properties of a pyridoxal
328 5'-phosphate-dependent histidine decarboxylase from *Morganella morganii* AM-15. J
329 Biol Chem 260:6738–6746. <https://doi.org/10.1128/jb.169.9.3963-3968.1987>
- 330 27. Zou Y, Hu W, Jiang A, Tian M (2014) Partial purification and characterization of a
331 novel histidine decarboxylase from *Enterobacter aerogenes* DL-1. Preparative
332 Biochemistry and Biotechnology 45:605–615.
333 <https://doi.org/10.1080/10826068.2014.940541>
- 334 28. Bjornsdottir-Butler K, Green DP, Bolton GE, McClellan-Green PD (2015) Control of
335 histamine-producing bacteria and histamine formation in fish muscle by trisodium
336 phosphate. Journal of Food Science 80:M1253–M1258.
337 <https://doi.org/10.1111/1750-3841.12875>
- 338 29. Morii H, Kasama K (2004) Activity of two histidine decarboxylases from
339 *Photobacterium phosphoreum* at different temperatures, pHs, and NaCl concentrations.
340 Journal of Food Protection 67:1736–1742.
341 <https://doi.org/10.4315/0362-028X-67.8.1736>
- 342 30. Tabanelli G, Torriani S, Rossi F, Rizzotti L, Gardini F (2012) Effect of
343 chemico-physical parameters on the histidine decarboxylase (HdcA) enzymatic
344 activity in *Streptococcus thermophilus* PRI60. Journal of Food Science 77:M231–
345 M237. <https://doi.org/10.1111/j.1750-3841.2012.02628.x>

- 346 31. Ferrario C, Borgo F, Las Rivas De B, Muñoz R, Ricci G, Fortina MG (2013)
347 Sequencing, characterization, and gene expression analysis of the histidine
348 decarboxylase gene cluster of *Morganella morganii*. Curr Microbiol 68:404–411.
349 <https://doi.org/10.1007/s00284-013-0490-7>
- 350 32. Kimura B, Takahashi H, Hokimoto S, Tanaka Y, Fujii T (2009) Induction of the
351 histidine decarboxylase genes of *Photobacterium damsela* subsp. *damsela* (formally
352 *P. histaminum*) at low pH. J Appl Microbiol 107:485–497.
353 <https://doi.org/10.1111/j.1365-2672.2009.04223.x>
- 354 33. Calles-Enriquez M, Eriksen BH, Andersen PS, Rattray FP, Johansen AH, Fernandez
355 M, Ladero V, Alvarez MA (2010) Sequencing and transcriptional analysis of the
356 *Streptococcus thermophilus* histamine biosynthesis gene cluster: factors that affect
357 differential *hdcA* expression. Applied and Environmental Microbiology 76:6231–6238.
358 <https://doi.org/10.1128/AEM.00827-10>
- 359 34. Landete JM, Las Rivas De B, Marcobal A, Muñoz R (2008) Updated molecular
360 knowledge about histamine biosynthesis by bacteria. Critical Reviews in Food Science
361 and Nutrition 48:697–714. <https://doi.org/10.1080/10408390701639041>
- 362 35. Merrell DS, Camilli A (2000) Regulation of *Vibrio cholerae* genes required for acid
363 tolerance by a member of the “ToxR-Like” family of transcriptional regulators.
364 Journal of Bacteriology 182:5342–5350.
365 <https://doi.org/10.1128/JB.182.19.5342-5350.2000>

366 36. Tanaka Y, Kimura B, Takahashi H, Watanabe T, Obata H, Kai A, Morozumi S, Fujii
367 T (2008) Lysine decarboxylase of *Vibrio parahaemolyticus*: kinetics of transcription
368 and role in acid resistance. J Appl Microbiol 104:1283–1293.
369 <https://doi.org/10.1111/j.1365-2672.2007.03652.x>

370

371

372

373

374

375

376

377

378

379

380

381

TABLE AND FIGURE LEGENDS

Table 1. Primer sequences used for qRT-PCR experiments

Fig. 1. The growth curve (A, C) and histamine accumulation (B, D) in canned tuna at 20 °C (A, B) and 4 °C (C, D). Symbols represent *Morganella psychrotolerans* (○), *M. morganii* (□), and *Photobacterium phosphoreum* (△) for viable counts. Bars represent *M. psychrotolerans* (gray), *P. phosphoreum* (white), and *M. morganii* (black) for histamine concentration. Dagger (†) indicates histamine concentration less than 4 mg/kg. Means with different letters within the same incubation time indicate significantly difference ($p < 0.05$).

Fig. 2. The growth curve (A, C) and histamine production (B, D) by *M. psychrotolerans* at 20 °C (A, B) and 4 °C (C, D). Symbols represent pH 5 (○), pH 6 (□), pH 7 (△), and pH 8 (◇) for viable counts. Bars represent pH 5 (white), pH 6 (gray), pH 7 (black), and pH 8 (slash) for histamine concentration. Dagger (†) means less than 4 mg/kg of histamine concentration. Means with different letters within the same incubation time indicate significantly difference ($p < 0.05$).

Fig. 3. The effects of temperature (A) and pH (B) on histamine accumulation by the cell-free crude histidine decarboxylase of *M. psychrotolerans*. Means with different letters within the same temperature and pH indicate significantly difference ($p < 0.05$).

403

404 Fig. 4. Relative expression of the histidine decarboxylase gene of *M. psychrotolerans* under various pH
405 conditions. Bars represent pH 5 (white), pH 6 (gray), pH 7 (black), and pH 8 (slash). Means with
406 different letters within the same temperature indicate significantly difference ($p < 0.05$).

407

Fig. 1 Wang et al.

Fig. 2 Wang et al.

Fig. 3 Wang et al.

Fig. 4 Wang et al.

Table. 1 Wang et al.

	Primer	Sequences (5'-3')
hdc	Hdc-MP-f	ATTCAATTGGTGTTTCCGGC
	Hdc-MP-r	TATGACCATTACGGGAACCG
16s rDNA	Mor-Ref-f	TTTCAGTCGGGAGGAAGGTG
	Mor-Ref-r	GGGGATTTCACATCTGACTYA