Title	Fine root dynamics and partitioning of root respiration into growth and maintenance components in cool temperate deciduous and evergreen forests
Author(s)	Sun, Lifei; Hirano, Takashi; Yazaki, Tomotsugu; Teramoto, Munemasa; Liang, Naishen
Citation	Plant and soil, 446(1-2), 471-486 https://doi.org/10.1007/s11104-019-04343-z
Issue Date	2020-01
Doc URL	http://hdl.handle.net/2115/80115
Rights	This is a post-peer-review, pre-copyedit version of an article published in Plant and Soil. The final authenticated version is available online at: http://dx.doi.org/10.1007/s11104-019-04343-z
Type	article (author version)
File Information	Sun et al., Plant and Soil.pdf

[Instructions for use](#)

Title: Fine root dynamics and partitioning of root respiration into growth and maintenance components in cool temperate deciduous and evergreen forests

Lifei Sun^a, Takashi Hirano^{a,*}, Tomotsugu Yazaki^{a,**}, Munemasa Teramoto^b and Naishen Liang^b

^a Research Faculty of Agriculture, Hokkaido University

^b Center for Global Environmental Research, National Institute for Environmental Studies

*Corresponding author: Takashi Hirano

Research Faculty of Agriculture, Hokkaido University

N9 W9, Kita-ku, Sapporo 060-8589, Japan

Tel & Fax +81-11-706-3689, Email hirano@env.agr.hokudai.ac.jp

**Present address: School of Agriculture, Meiji University

Running head: Growth and maintenance respirations of tree fine roots

Abstract

Aims: We aim to show the seasonality of fine root dynamics and examine the relationship between root respiration (R_r) and fine root dynamics. In addition, we try partitioning R_r into growth (R_g) and maintenance (R_m) components.

Methods: Soil respiration (R_s), fine root biomass (B), and fine root production (P) were measured simultaneously over a growing season in adjoining deciduous (DF) and evergreen (EF) forests. The R_r was separated from R_s by the trenching method, and R_r was partitioned into R_g and R_m using an empirical model.

Results: The seasonality of P was almost the same in both forests, though that of B was different. The R_r showed a positive correlation with P in both sites. Annual R_r was estimated to be 610 (DF) and 393 (EF) g C m⁻² year⁻¹. Annual R_g and R_m were 121 and 166 (DF), and 86 and 182 (EF) g C m⁻² year⁻¹, respectively.

27 *Conclusions:* We found a clear seasonal pattern in P and a positive linearity between R_r and P . Despite considerable
28 uncertainty due to the small sample size, presence of larger roots, and measurement uncertainty, the results suggest
29 that our approach is capable of partitioning R_r .

30

31 Key words: biomass, chamber, ingrowth core, production, soil respiration, trenching.

32

Introduction

Terrestrial ecosystems have sequestered carbon dioxide (CO₂) at a rate of 3.2 ± 0.8 Pg C year⁻¹ during the last decade until 2016, accounting for 30% of total CO₂ emissions from fossil fuels, industry, and land-use change (Le Quéré et al. 2018). Forests, one of the carbon-richest ecosystems, covers about 30% of the land surface (Bonan 2008; Keenan et al. 2015) and was estimated to be a large CO₂ sink of 2.4 ± 0.4 Pg C year⁻¹ from 1990 to 2007 (Pan et al. 2011). Therefore, forests are crucial ecosystems for mitigating climate change mostly owing to capturing atmospheric CO₂ during growth.

Belowground biological processes, such as root respiration and microbial decomposition of soil organic matter, greatly contribute to the carbon balance of forest ecosystems (Davidson et al. 2006; Janssens et al. 2001). In particular, fine roots, which are commonly defined as thin roots less than 2 mm in diameter, are known for their vital role in global biogeochemical cycles (McCormack et al. 2015). Fine roots function to absorb nutrients and water from the soil and have a shorter life cycle through production, mortality, and littering with average turnover rates of about one year in temperate forests (Brunner et al. 2013; Finér et al. 2011b). In addition, despite their small biomass the contribution of fine roots to the net primary production (NPP) of forests is relatively high at 32% for boreal forest (Yuan and Chen 2010), and 22% globally (McCormack et al. 2015). Also, fine root litter is a major carbon source in forest soils (Richter et al. 1999). Therefore, fine root dynamics are key processes that govern forest productivity and biogeochemical cycles.

Fine root phenology strongly influences belowground carbon dynamics and allocation, whereas seasonality and variability in the production and respiration of fine roots are not well understood among forest types (Abramoff and Finzi 2015; McCormack et al. 2014; Radville et al. 2016). Moreover, the underlying mechanisms controlling fine root dynamics remain limited because of the lack of experimental evidence (Liu et al. 2018; Radville et al. 2016). In particular, the dynamic functions of fine roots, such as respiration, are not quantitatively understood, which has restrained the improvement of terrestrial biosphere models to predict the responses of forest ecosystems to environmental changes (McCormack et al. 2015; Warren et al. 2015). Therefore, field experiments which simultaneously measure the production and respiration of fine roots in different forest types over a growing season will help improve our qualitative understanding of the role of fine roots in forest carbon cycles.

Empirical models exist that partition plant respiration into growth and maintenance requirements (Amthor 2000; McCree 1974; Penning de Vries 1974; Thornley 1970), plus the respiratory cost of ion uptake by fine roots (Chapin III et al. 2011; Johnson 1990; Lambers et al. 2008). In these models, growth and maintenance respirations correlate linearly with production (growth) and biomass, respectively. Problems with plant respiration models occur in the quantification of respiration partitioning (Gifford 2003), which lack a mechanistic basis (Sweetlove et al. 2013). Nevertheless, laboratory experiments suggested that the models were useful for understanding the carbon balance of plants, though the partitioning would differ by plant species and environmental conditions (Lambers et al. 2008). Also, the partitioning of respiration into these functional components will help in the understanding of the ecological controls over plant respiration (Chapin III et al. 2011). These models have been adopted by terrestrial biosphere models, such as Biome-BGC (Thornton and Rosenbloom 2005), to explicitly represent autotrophic respiration as the sum of growth and maintenance respirations. However, the responses of the models' rate coefficients to abiotic and biotic factors have not been well understood (Hopkins et al. 2013) owing to the lack of field experiments (Amthor 2000). So far, there have been no comparisons of model predictions to field conditions, except for (George et al. 2003) partitioning fine root respiration in field conditions. They separately estimated fine root respiration for growth, maintenance, and ion uptake (only nitrogen) in evergreen and deciduous forests using short-term chamber measurements and literature information.

In this study, root respiration and fine root dynamics (production and biomass) were measured simultaneously in adjoining deciduous and evergreen forests in a cool temperate climate over a growing season in order to: 1) quantify root respiration with reliability, 2) compare fine root phenology between the different forest types, and 3) examine the relationship between root respiration and fine root dynamics. In addition, we try partitioning the root respiration into growth and maintenance components on an annual basis using an empirical model. After the growth and maintenance respirations of fine roots are quantified in the two forests, the model's rate coefficients are compared between the forests and to coefficients reported from laboratory and greenhouse experiments.

Materials and method

Study site

The study was conducted in two adjacent sites (deciduous and evergreen forests) in southern Hokkaido, northern Japan (42° 44' N, 141° 31' E, 125 m above sea level). The soil type was volcanogenous regosol with a high water permeability. In both sites, below the leaf litter and a 4 cm thick root mat, there is a 10 cm thick surface soil layer (A horizon) and an unweathered pumice layer (C horizon); a B horizon is lacking (Hirano et al. 2003a).

The deciduous forest (DF) was established in 1957 as a Japanese larch (*Larix kaempferi*) plantation. However, it was severely devastated by typhoon Songda in 2004. The typhoon blew down about 90% of the trees (Sano et al. 2010). In 2005 the fallen tree stems were removed; afterwards the plantation has not been managed (Hirano et al. 2017). In 2015 the dominant tree species was still Japanese larch, followed by *Ulmus davidiana* var. *japonica* (Japanese elm), *Acer pictum* subsp. *dissectum* (maple), and *Quercus crispula* (oak), with a density of 2360 stems ha⁻¹ for trees taller than 2 m. The forest floor is covered with *Dryopteridaceae* sp. (fern) and shrub species, including *Rubus idaeus* (raspberry) (Yazaki et al. 2016).

The evergreen forest (EF) was a Japanese spruce (*Picea glehnii*) plantation established next to the deciduous forest in 1979. Spruce saplings were planted in a double row 1.3 m wide at intervals approximately 6 m between rows. Almost no trees fell during typhoon Songda. Tree density was 2250 stems ha⁻¹ in 2015. Understory vegetation was sparse.

Decadal mean annual air temperature and precipitation from 2008 to 2017 were 8.3 ± 0.3°C and 1312 ± 172 mm (mean ± 1 standard deviation (SD)), respectively, at a meteorological station (Tomakomai) 14 km from the study site. The lowest and highest monthly mean air temperatures were -3.9°C in January and 20.8°C in August, respectively. About 85% of the annual precipitation falls in the snow-free period from April to November. Snow usually covers the ground from early December through early April.

Experimental design

Field experiments were conducted from September 2014 to November 2015. In September 2014, to measure soil CO₂ efflux, a set of four 0.5 m × 0.5 m aluminum collars were installed, where one collar was used for a control, another for root sampling, the third for litter removal, and the fourth for trenching. Five replications were established in each site for partitioning total soil respiration (R_s) into leaf litter decomposition (R_l), belowground

heterotrophic respiration (R_h), and root respiration (R_r). To keep root density similar, four collars in each set were placed on a circle with a 2 m radius around larch trees in the DF site, or were aligned 2 m away from a tree row in the EF site (Fig. 1). Collar spacing was 20 cm, except for the trenching collars where it was 40 cm to ease the trenching effect on the next collar. Collars were inserted only 1 to 2 cm into the soil to minimize damaging surface roots (Wang et al. 2005). To avoid lateral air leak under the collars, each collar was surrounded with a bank of soil. For control collars (CC), no treatment was applied. In sampling collars (SC), the fine root experiments (described later) were conducted. In litter removal collars (LC), 1 mm meshed screens were set between the litter layer and soil surface in early November 2014. The mesh was temporarily removed with the overlying litter, and then soil CO_2 efflux was measured without leaf litter. After the measurement, the mesh and litter were immediately put back on the collar. The trenching method was applied to remove R_r from R_s . For trenching collars (TC), four PVC boards 4 mm thick were vertically inserted to a depth of 0.3 m along a square collar in November 2014 to prevent roots from intruding. The decomposition of dead roots (R_d) left in TC was estimated by the root litter bag method (described later). R_l and R_r were calculated using the following equations:

$$R_l = R_s - R_{LC} \quad (1)$$

$$R_r = R_s - R_l - R_h = R_s - (R_{TC} - R_d) \quad (2)$$

where R_s is soil CO_2 efflux measured in CC or SC, and R_{LC} and R_{TC} are CO_2 effluxes from LC and TC, respectively.

Soil CO_2 efflux

Soil CO_2 efflux was measured once or twice a month from September 2014 to November 2015, except during the snow season. Before the treatments, CO_2 efflux was also measured in September and October 2014 to check for initial spatial variations among the CC, SC, LC, and TC collars. The measurements were conducted manually on each collar between 10:00 and 16:00 using a closed chamber system with two 50 cm tall cubic chambers made of transparent acrylic plastic (Sun et al. 2017). CO_2 concentration and air temperature inside the chamber were measured every 5 s with an infrared gas analyzer (LI820; Li-Cor Inc., Lincoln, NB, USA) and a thermocouple probe (MHP; Omega Engineering, Stamford, CA, USA). Outputs were recorded using a data logger (CR1000; Campbell Scientific Inc., Logan, UT, USA). Each chamber was closed for 180 s. Despite the short closing, soil

CO₂ efflux might have been underestimated by about 5% because of lateral diffusion in relatively porous root mats (Hutchinson and Livingston 2001). Immediately after measurement, soil temperature (T_s) at a depth of 5 cm and volumetric soil water content (SWC) of the top 5 cm layer were measured, respectively, in each collar with a thermocouple probe (MHP; Omega Engineering) and a soil moisture sensor (SM150; Delta-T Devices Ltd., Cambridge, UK). Sprouts in each collar were picked carefully before each measurement. In addition, T_s at a depth of 5 cm were monitored hourly in two CC at each site using button-type temperature loggers (Thermochron SL type, KN laboratories, Osaka, Japan). SWC at a depth of 3 cm were also monitored every half-hour in four replications at a nearby re-growing forest site using TDR sensors (CS615; Campbell Scientific, Inc.) along with air temperature and precipitation (Hirano et al. 2017).

The relationship between soil CO₂ efflux (R , $\mu\text{mol m}^{-2} \text{s}^{-1}$) and T_s ($^{\circ}\text{C}$) was analyzed for each collar using the following exponential equation:

$$R = a \cdot \exp(b \cdot T_s) \quad (3)$$

where a and b are fitting parameters. To analyze the relationship with SWC, temperature-normalized soil CO₂ efflux (R_b , $\mu\text{mol m}^{-2} \text{s}^{-1}$) was calculated using the following equation (Hirano et al. 2003b):

$$R_b = R \cdot \exp\{b \cdot (T_b - T_s)\} \quad (4)$$

where T_b is base temperature set as the mean T_s at each site. Also, bivariate models combining Eq. 3 for T_s with various equations for SWC (Reichstein and Janssens 2009) were applied to estimate soil CO₂ efflux. CO₂ efflux was calculated hourly from the monitoring data of T_s and SWC using a best-fit model and converted into annual CO₂ emission between November 2014 and November 2015.

Dead root decomposition in trenched collars

The root litter bag method was applied to assess CO₂ emissions (R_d) through the decomposition of dead roots left in TC (Epron et al. 1999; Gholz et al. 2000; Silver and Miya 2001). To determine the initial dry weight of dead roots (X_0 , g DM m⁻²), live roots were extracted in the same manner as in the fine root experiment (described later) from 30 cm thick surface soil samples in a 0.3 m $\times$ 0.3 m area near each TC ($n = 5$) at each site in early November 2014. The roots were separated into three classes by diameter: fine (< 2 mm), medium (2–10 mm), and coarse

roots (> 10 mm). The root samples were air-dried, and parts of the samples were oven-dried at 70°C for 48 hours to determine the water content of each root class. From the water content, the dry weights of the remaining air-dried samples were determined. Air-dried samples were put into 2 mm meshed bags that were 10 cm × 10 cm; samples were 1, 3, and 5 g in dry weight, respectively, for fine, medium, and coarse roots. In each site, 25 bags for each class were buried in the A horizon near TC in late November 2014. Five bags were collected from each class five times until May 2017, and the dead roots remaining in the bags were oven-dried. The decomposition constant k (year⁻¹) was determined using the following equation (Wieder and Lang 1982):

$$Y = Y_0 \cdot \exp(-k \cdot t) \quad (5)$$

where Y is the dry weight of remaining dead roots (g DM bag⁻¹) at time t (year), and Y_0 is the initial dry weight of root samples (g DM bag⁻¹). From k and X_0 , annual R_d (g C m⁻² year⁻¹) from November 2014 was calculated for each root class using a conventional conversion factor of 0.5 from root dry matter to carbon (Chapin III et al. 2011; Ravindranath and Ostwald 2008).

$$R_d = 0.5 \cdot X_0 \{1 - \exp(-k)\} \quad (6)$$

Fine root experiment

The sequential soil core method was applied to measure fine root biomass (B ; g DM m⁻²) and its temporal variation (ΔB ; g DM m⁻² period⁻¹). Three soil cores were sampled from each SC in mid-November 2014, early May, mid-June, early August, late September, and mid-November 2015. To confirm the trenching effect on fine root existence, soil cores were also sampled similarly from TC in early May and mid-June 2015. Using a perforated board for positioning, core sampling was made in each SC at three points, which were randomly selected from a grid with 8 cm spacing. Soil cores down to 14 cm were taken using a stainless-steel edged tube with an inner diameter of 2.4 cm. The total sampling area was 13.6 cm² for each collar each time, which accounted for 0.5% of the collar area (0.25 m²). The collected core samples were first divided into the root mat (top 4 cm) and A horizon (lower 10 cm), and then washed several times with tap water to remove soil particles and fragmented litter. Next, living fine roots were extracted and were visually separated into tree and herbaceous roots in the DF site, where herbaceous plant roots were not negligible. The fine roots were dried for 48 hours at 70°C and weighed. Tree fine roots originated

mainly from larch trees and shrub species in the DF site, but were almost entirely from spruce trees in the EF site.

The ingrowth core method (e.g. (Vogt et al. 1998)) was applied to measure fine root production (P ; g DM m⁻² period⁻¹). Ingrowth cores were made of plastic cylindrical frames wrapped in a 2 mm meshed sheet. The cores were 14 cm long and had a diameter of 2.3 cm. Soils were taken from the A horizon in September 2014, and roots were removed using a 2 mm mesh after air drying. The cores were filled with root-free, air-dried soil and inserted vertically into three pits after the soil core sampling at each SC. The ingrowth cores were replaced simultaneously with the sequential core sampling. Soil samples in the collected ingrowth cores were first divided into the top 4 cm layer (root mat) and below 10 cm (A horizon), and washed to extract living fine roots. The fine roots were weighed after drying for 48 hours at 70°C. The dry matter is equivalent to P through each sampling interval, and annual P was calculated as the sum of periodic P over a year. Using a conventional factor of 0.5, dry matter was converted to carbon. Annual mortality (M ; g DM m⁻² year⁻¹) was calculated as a difference between annual P and annual ΔB . In addition, fine-root turnover rates were calculated as the ratio of annual P and mean B (Brunner et al. 2013).

Growth and maintenance respirations

The simple equation of growth (R_g) and maintenance respiration (R_m) partitioning follows (Amthor 2000) and (Thornley 1970):

$$R_r = R_g + R_m = g_R \cdot P + m_R \cdot B \quad (7)$$

where g_R is the growth respiration coefficient, and m_R is the maintenance respiration coefficient. To adopt the equation for field data, it was modified into:

$$R_r = R_g + R_m + h = c \cdot P + d \cdot \exp(f \cdot T_s) \cdot B + h \quad (8)$$

where c is g_R (g C g DM⁻¹), d is base respiration rate per unit biomass at 0°C (g C g DM⁻¹ day⁻¹), f is a temperature coefficient (°C⁻¹), T_s is soil temperature at a depth of 5 cm (°C), and h is residual (g C m⁻² day⁻¹), which includes the respiration of ion uptake, herbaceous roots, and thicker roots. $d \cdot \exp(f \cdot T_s)$ is equivalent to m_R (g C g DM⁻¹ day⁻¹) at T_s . The parameters of c , d , f , and h were determined by curve fitting to the data set of R_r (g C m⁻² day⁻¹), P (g DM m⁻² day⁻¹), B (g DM m⁻²), and T_s , which were calculated at the same intervals. The R_r and T_s were averaged

over each interval of P measurement, and B was the mean of two measurements at the beginning and end of an interval. Next, R_r , P , and B were spatially averaged for each site ($n = 5$). Thus, the number of the data set for the four variables (R_r , T_s , P , and B) was equivalent to the frequency of P measurement ($n = 5$).

Statistical analysis

ANOVA and post hoc Tukey's HSD were applied to compare environmental factors, fine root biomass, fine root production, and soil CO₂ effluxes among treatments. t -test was applied to compare mean values between the sites. Parameters in the equations were determined by curve fitting to the data set ($n = 5$) for each forest. In addition, uncertainties (± 1 SD) of annual R_g and R_m were estimated from those of the parameters and measurements using the law of error propagation. Data analyses were conducted with a software package (Origin Pro 2015J; Origin Lab Corporation, Northampton, MA, USA).

Results

Environmental conditions

Mean air temperature at 1.5 m aboveground and total precipitation measured in the nearby re-growing forest site during the growing season from May to October 2015 were 14.0°C and 822 mm, respectively, which are within the range of 1 SD from their decadal means of $14.4 \pm 0.5^\circ\text{C}$ and 949 ± 188 mm between 2006 and 2015. Daily mean T_s peaked in August (Fig. 2a). Mean T_s was 8.8°C in the DF site and 7.5°C in the EF site from November 2014 to November 2015. The higher T_s in the DF site was mainly caused by more incident solar radiation on the forest floor through the defoliated canopy in spring. In 2015, SWC showed a seasonal variation with a minimum in August, which was opposite to T_s (Fig. 2b).

Mean T_s was not significantly different among four treatments at both sites; however, it was significantly lower in the EF site (Table 1). In contrast, mean SWC significantly increased after trenching (TC), though no significant difference was found between the sites.

Decomposition of dead roots

Dead roots decayed exponentially. The decomposition constant k was higher in the DF site than in the EF site and was highest for fine roots, followed by medium and coarse roots in both sites (Table 2). Using Eq. 6, annual R_d was calculated for each root class using the k values and initial root biomass and totaled 131 and 89 g C m⁻² year⁻¹, respectively, in the DF and EF sites (Table 2), accounting for 20% and 16% of the initial carbon content. The contribution of fine root decomposition to the total R_d was 34% and 52%, respectively, in the DF and EF sites.

Soil CO₂ efflux

Soil CO₂ efflux was measured before the treatment and showed no significant difference among four collars in all five replications in each site (ANOVA, $P > 0.05$), which indicates that correction for spatial variation in CO₂ efflux was unnecessary. Also, no significant difference was found between CO₂ effluxes from CC and SC collars in 2015 even after the start of soil core sampling (paired t -test, $P > 0.05$); mean CO₂ effluxes (± 1 SD) were 4.97 ± 4.30 (CC) and 4.77 ± 3.34 (SC) $\mu\text{mol m}^{-2} \text{s}^{-1}$ in the DF site, 3.91 ± 2.76 (CC) and 4.02 ± 2.69 (SC) $\mu\text{mol m}^{-2} \text{s}^{-1}$ in the EF site. These results indicate that the effect of core sampling on CO₂ efflux was negligible. Thus, CO₂ efflux from SC was used as R_s to minimize spatial discrepancy between flux measurement and root sampling.

Soil CO₂ efflux showed a similar seasonal variation in both sites with a peak in late July (Fig. 3). Soil CO₂ efflux was significantly correlated to T_s ($P < 0.001$) in an exponential manner (Eq. 3) on every collar ($R^2 > 0.55$). Figure 4 shows the mean relationships for SC and TC. The Q_{10} values (mean ± 1 SD, $n = 5$) calculated using the fitting parameter (b) were 2.7 ± 0.5 (SC) and 3.0 ± 0.5 (TC) in the DF site, and 2.4 ± 0.2 (SC) and 2.4 ± 0.2 (TC) in the EF site. The results were significantly higher ($P < 0.05$) in the DF site but not significantly different ($P > 0.05$) between SC and TC. Soil CO₂ efflux was normalized for mean T_s (R_b) according to Eq. 4 using T_b of 14°C and 12°C, respectively, in the DF and EF sites (Table 1). In both sites, R_b had no significant relationship with SWC ranging between 0.1 and 0.6 m³ m⁻³ (Fig. 5). Also, adding SWC did not improve adjusted R^2 than T_s alone. Thus, using Eq. 3, hourly CO₂ efflux was estimated from T_s monitoring data for each collar and summed up over an annual period starting in November 2014. The annual values ($n = 5$) were 1255 ± 401 (SC), 1044 ± 334 (LC), and 776 ± 83 (TC) g C m⁻² year⁻¹ in the DF site, and 1139 ± 386 (SC), 1027 ± 323 (LC), and 835 ± 172 (TC) g C m⁻² year⁻¹ in the EF site. The SD was lower in TC than in SC, because spatial variations in root respiration were

excluded in TC. Annual R_s , R_l , R_h , and R_r were calculated from the annual soil CO₂ effluxes and R_d (Table 3). Here, R_r includes not only the respiration of tree fine roots but also those of tree medium and coarse roots, and herbaceous roots. The contributions of R_r to R_s were 49% and 35%, respectively, in the DF and EF sites. Also, R_d accounted for 10% and 8% of R_s , respectively, in the DF and EF sites.

Fine root dynamics

Tree fine root biomass (B) showed a clear seasonal variation with a peak in September in the DF site (Fig. 6a), whereas seasonality was indeterminable in herbaceous fine root biomass. In the EF site, B had less seasonality (Fig. 6c). No herbaceous fine root biomass was found in the EF site. Peak B was 482 and 559 g DM m⁻², respectively, in the DF and EF sites (Figs. 6b and 6c). On average, 56% and 49% of B existed in the root mat in the DF and EF sites, respectively. In TC, B in May and June 2015 were 32.0 ± 6.6 and 11.6 ± 1.4 g DM m⁻², respectively, in the DF site ($n = 5$), and 41.8 ± 10.2 and 16.4 ± 3.0 g DM m⁻², respectively, in the EF site ($n = 5$). B decreased significantly between May and June ($P < 0.001$) in both sites. In June 2015, B in the trenched collars (TC) decreased to 3% in comparison with the non-trenched collars (SC) in both sites.

No herbaceous fine root was found in the ingrowth cores even in the DF site, probably because herbaceous plants were picked out periodically. Tree fine root production (P) showed a distinct seasonality with a peak between 2.3 to 2.5 g DM m⁻² day⁻¹ in summer (mid-June to early August) in both sites (Figs. 6d and 6e). The seasonal variations of B and P were similar in the DF site (Figs. 6b and 6d). Even during winter and early spring between November 21, 2014 and May 1, 2015, P was measured at 0.14 ± 0.08 and 0.22 ± 0.14 g DM m⁻² day⁻¹ in the DF and EF sites, respectively; P was not significantly different between the sites ($P = 0.26$). The P was positively correlated ($P < 0.05$) with mean T_s during core-sampling intervals (Fig. 7) but showed no significant relationship with SWC ($P > 0.05$, data not shown). In both sites, R_r was significantly correlated with P (Fig. 8, $P < 0.05$), but not with B ($P > 0.05$, data not shown).

Annual P was not significantly different between the DF and EF sites (Table 4). Although the root mat was only 4 cm thick, P in the root mat accounted for 45 to 48% of total P . Mean B was significantly larger in the EF site ($P < 0.01$), although the difference was not significant if herbaceous fine root biomass was included ($P > 0.05$). In

the DF site, herbaceous fine root biomass accounted for 25% (110 g DM m⁻²) of total *B*. In November 2014, fine root accounted for 20% and 33% of total tree root biomass, respectively, in the DF and EF sites (Table 2). The tree fine root turnover rate was $1.08 \pm 0.36 \text{ year}^{-1}$ in the DF site, which was significantly larger than $0.68 \pm 0.12 \text{ year}^{-1}$ in the EF site ($P < 0.05$).

Partitioning root respiration into growth and maintenance components

R_r was partitioned into R_g and R_m using Eq. 8. In both sites, no significant correlation was found between P and B ($P > 0.05$). Despite a relatively high R^2 (≥ 0.87), the fitting was not significant ($P \geq 0.065$) because of the small number of samples ($n = 5$) for both sites (Table 5). All four parameters were not significantly different between the sites ($P \geq 0.55$). The Q_{10} of R_m calculated from f (temperature coefficient) were 3.3 and 2.6, respectively, for DF and EF. Using these parameters, annual R_g and R_m were calculated (Table 6). The annual totals ($R_g + R_m + h$) were 638 and 396 g C m⁻² year⁻¹, respectively, in the DF and EF sites. These totals exceeded R_r by 28 and 3 g C m⁻² year⁻¹, respectively, in the DF and EF sites. The contribution of R_g to the total was almost the same (19 vs. 22%), though R_m was larger in the EF site (26 vs. 46%). As a result, h contributed more in the DF site.

Discussion

Soil respiration and root respiration

Root respiration (R_r) was calculated as the residual of total soil respiration (R_s) after subtracting leaf litter decomposition (R_l) and soil heterotrophic respiration (R_h) (Eq. 2). Thus, R_r corresponds to mycorrhizosphere respiration, which consists of not only respiration of living root tissue but also rhizomicrobial and mycorrhizal respiration (Moyano et al. 2009). There was no significant difference in soil CO₂ efflux between control (CT) and sampling collars (SC) even after soil sampling. Thus, soil CO₂ efflux on SC was treated as R_s , because it enabled us to directly compare soil CO₂ efflux and fine root dynamics in the same collar.

Trenching is a commonly used method to estimate soil heterotrophic respiration, although it potentially causes biases owing to dead root decomposition, lack of water uptake by roots, and lack of root litter input (Epron et al. 1999; Hanson et al. 2000; Subke et al. 2006). In this study, dead root decomposition (R_d) was quantified by the

litter bag method. Soil water content (SWC) was significantly higher in TC (Table 1), but no relationship with soil CO₂ efflux was found using bivariate models and temperature-normalized CO₂ efflux (Fig. 5), suggesting that an increase in SWC minimally affected soil CO₂ efflux. Tree fine root biomass (*B*) in TC was only 3% in comparison with SC in June 2015, which indicates that trenching worked effectively. In contrast, belowground microbial respiration was probably underestimated to a small extent due to the lack of root litter input (Subke et al. 2006). Also, trenching killed ectomycorrhizal fungi and consequently might have enhanced the decomposition of soil organic matter by saprotrophic fungi (Gadgil effect) (Fernandez and Kennedy 2016).

The annual R_s of 1255 and 1139 g C m⁻² year⁻¹ (Table 3) estimated from hourly T_s , were relatively large but within the range of reported values from temperate forests (Subke et al. 2006), which were 828 ± 267 g C m⁻² year⁻¹ for coniferous forests and 976 ± 514 g C m⁻² year⁻¹ for deciduous forests, respectively (mean $\pm$ 1 SD, $n = 28$). Our results were also larger than the annual R_s of 760 ± 40 g C m⁻² year⁻¹ measured for 8 years in a Japanese larch plantation in central Japan (Teramoto et al. 2017), which had an annual air temperature (8.6°C) similar to our site (8.3°C). Before the typhoon disturbance, annual R_s was 934 g C m⁻² year⁻¹ in the DF site in 2003 (Liang et al. 2010), which was smaller by 321 g C m⁻² year⁻¹ than in 2015 despite having similar annual mean soil temperatures (8.6°C in 2003 and 8.8°C in 2015). The larger R_s in 2015 was partly due to higher soil temperature during the summer. In 2015, soil temperature in July and August were higher on average by 2.4°C because of a sparser canopy. The annual contributions of R_r to R_s (R_r / R_s) were 0.49 and 0.35, respectively, in the DF and EF sites (Table 3). These ratios were compatible with results from a meta-analysis (Subke et al. 2006), where R_r / R_s was calculated to be 0.43 ± 0.21 for deciduous forests and 0.49 ± 0.14 for coniferous forests ($n = 28$, respectively).

The decomposition of dead tree roots was well explained with the rate constant of k (Table 2). The k was higher in the DF site than in the EF site (0.16–0.42 vs. 0.10–0.29 year⁻¹) and highest for fine roots. A study conducted in a beech forest in France showed k values of 0.38 year⁻¹ for fine roots and 0.22 year⁻¹ for coarse roots (> 2 mm) (Epron et al. 1999), which is similar to the result in the DF site. Also, a meta-analysis reported that mean k values for fine roots were 0.44 and 0.17 year⁻¹, respectively, for broadleaf and coniferous forests, and k values became lower with root diameter (Silver and Miya 2001). Annual R_d were 131 and 89 g C m⁻² year⁻¹, respectively, for DF and EF sites (Table 3), which accounted for 10% and 8% of annual R_s , respectively. If the R_d was neglected in Eq.

2, R_r would be calculated to be 479 and 304 g C m⁻² year⁻¹, respectively, for DF and EF sites, which would underestimate R_r by 21% and 23%. In a beech forest, (Epron et al. 1999) reported that R_r was 40% underestimated by annual R_d of 160 g C m⁻² year⁻¹.

Fine root dynamics

We applied the sequential coring method for tree fine root biomass (B) and the ingrowth core method for tree fine root production (P) for a total soil depth of 14 cm. In this area 83–85% of fine roots existed in the surface 15 cm of the soil layer (Sakai et al. 2007), suggesting underestimation of B and P in this study. The ingrowth core method has been widely applied mainly because of its easy applicability (Addo-Danso et al. 2016). However, the ingrowth core method tends to underestimate P in comparison with the minirhizotron method, which yields more reliable estimates (Addo-Danso et al. 2016; Finér et al. 2011b; Hendricks et al. 2006; Majdi et al. 2005). Thus, our P results could be underestimated.

In both sites, P showed a clear seasonality with a peak in summer and was found to be 0.14–0.22 g DM m⁻² day⁻¹ even during a period from winter through early spring (Fig. 6). The period ranged from November 21, 2014 to May 1, 2015, in which albedo indicated that snow covered the ground from December 2 to April 4 in a nearby re-growing forest site (Hirano et al. 2017). Mean soil temperature in the period was 1.4 and 0.4°C, respectively, in the DF and EF sites. A delay in temperature rise in April in the EF site (Fig. 2) was caused by delayed thaw due to a dense canopy, indicating that snow covered the ground until April 20 in the EF site. In addition, larch trees were still leafless on May 1, 2015. In this area larch trees usually begin to leaf out in early May (Hirano et al. 2003a). Therefore, despite including early spring after thaw the measured P suggests that tree fine roots grew during winter (Radville et al. 2016).

The seasonal P was well explained by soil temperature (T_s) (Fig. 7). Similar seasonal variations were reported from many boreal and temperate forests (Brassard et al. 2009; Fukuzawa et al. 2013; Noguchi et al. 2005; Steinaker et al. 2010; Tierney et al. 2003). Also, B increased in summer in the DF site, whereas no clear seasonal variation was found in the EF site (Fig. 6). Although summer B was comparable, winter B was greater in the EF site. The discrepancy in seasonality was probably attributed to the life span of fine roots. Mean life span, as the reciprocal

of a turnover rate, was 0.93 and 1.47 year, respectively, in the DF and EF sites (Table 4). In the EF site, the longer life span may not have decreased B even in winter. In boreal and temperate forests, some studies reported seasonal variations similar to the DF site (Brassard et al. 2009; Coleman et al. 2000), but other studies found no seasonality (Fukuzawa et al. 2013; Makkonen and Helmisaari 1998; Noguchi et al. 2005).

Annual mean B was 340 and 516 g DM m⁻², respectively, in the DF and EF sites (Table 4). These values are smaller than mean values of 505 g DM m⁻² in deciduous broadleaf forests and 607 g DM m⁻² in evergreen conifer forests in temperate regions (Finér et al. 2011a). The reasons for the smaller B are possibly because of the thin surface soil in both sites, tree thinning by the typhoon in the DF site, and possibly limited sampling depth. Annual P in the two sites were almost the same (368 vs. 352 g DM m⁻² year⁻¹) despite the large difference of B . As a result, turnover rates, which are defined as the ratio of annual P and mean B , were significantly higher in the DF site (1.09 year⁻¹) than in the EF site (0.68 year⁻¹). If the maximum B is used instead of mean B (Brunner et al. 2013), turnover rates of the two sites are almost identical (0.30 vs. 0.31 year⁻¹). Although the ingrowth core method was used in this study, the annual P was similar to the mean P of temperate forests (337 g DM m⁻² year⁻¹), which was measured by various methods, especially using the sequential coring method (Finér et al. 2011b). Also, the P of this study was slightly larger than that measured by the minirhizotron method from a cedar plantation (320 g DM m⁻² year⁻¹) in central Japan (Noguchi et al. 2005). A review paper showed that the turnover rates from mean B were 1.10 year⁻¹ for both deciduous *Fagus sylvatica* and evergreen *Picea abies* (Brunner et al. 2013). These turnover rates are almost identical with the DF turnover rate. The turnover rate of the EF site was smaller than that of *Picea abies*.

Growth and maintenance respiration

R_r was partitioned into growth (R_g) and maintenance respirations (R_m) of tree fine roots using a model (Eq. 8) by multiple regression. The defects of the model include the lack of respiration due to root-derived compounds (rhizomicrobial and mycorrhizal respiration) and ion uptake. In addition, respiration except tree fine roots were not included in the analyses. Moreover, B and P were probably underestimated because of the limited sampling depth of 14 cm and the use of the ingrowth core method. These facts certainly resulted in considerable uncertainty. Rhizomicrobial and mycorrhizal respiration may be related to P and B , because they depend on rhizodeposits and

carbohydrates derived from roots (Moyano et al. 2009). Ion uptake respiration is expected to correlate to tree photosynthesis (Lambers et al. 2008). Although we did not consider ion uptake respiration, gross primary production (GPP) of the larch forest growing in the DF site before the typhoon disturbance peaked in June-July (Hirano et al. 2003a; Hirata et al. 2007). Also, in an evergreen conifer forest of *Pinus resinosa* in Japan, GPP peaked in June-August (Mizoguchi et al. 2012). These seasonal variations of GPP are like those of P with a peak in summer (Fig. 6). Thus, ion uptake respiration was probably partly included in R_g . The residual (h) was much larger in the DF site (Table 6), which is consistent with the fact that B accounted for only 20% of total tree root biomass (Table 2), and herbaceous root biomass accounted for 24% of total fine root biomass (Table 4).

The sums of R_g and R_m , which were rough estimates of tree fine root respiration, were similar between the DF and EF sites (287 vs. 268 g C m⁻² year⁻¹; Table 6), although their contributions to total root respiration were different (45 vs. 68%). (George et al. 2003) used short-term chamber measurements and published information to partition fine root respiration into growth, maintenance, and ion uptake (only nitrogen) components in deciduous (*Liquidambar styraciflua*) and evergreen (*Pinus taeda*) forests. They reported that annual fine root respiration in the deciduous and evergreen forests were 245 and 639 g C m⁻² year⁻¹, respectively, and maintenance respiration accounted for 86 and 98% of the total fine root respiration, respectively. Although annual P (345 g DM m⁻² year⁻¹) of the deciduous forest was almost identical to that (368 g DM m⁻² year⁻¹) of the DF site (Table 4), growth respiration (24 g C m⁻² year⁻¹) was only 20% of the DF's R_g (Table 6). Their ion uptake respiration was less than 4% of total fine root respiration. When limited to the growing season between May and November, tree fine root respiration (R_g plus R_m) were calculated to be 260 and 230 g C m⁻², respectively, in the DF and EF sites. These values are similar to those estimated from chamber measurements for *Quercus rubra* (229 g C m⁻²), *Tsuga canadensis* (242 g C m⁻²), and *Fraxinus alba* (270 g C m⁻²) in the Harvard Forest (Abramoff and Finzi 2016).

The R_g accounted for 42 and 32% of the total fine root respiration in the DF and EF sites, respectively (Table 6). The higher contribution in the DF site might be due to its higher turnover rate (Lambers et al. 2008). Growth (g_R) and maintenance (m_R) coefficients (Eq. 7) of whole root respiration for some species have been determined from growth chamber or greenhouse experiments. Using the units of mmol O₂ (g DM)⁻¹ for g_R and nmol O₂ (g DM)⁻¹ s⁻¹ for m_R , and explicitly including the respiration of ion uptake in g_R , whole root g_R and m_R , respectively,

were 11 and 26 for *Dactylis glomerata*, 19 and 21 for *Festuca ovina*, 12 and 6 for *Quercus suber*, and 18 and 22 for *Triticum aestivum* (Lambers et al. 2008). For *Eucalyptus* sp. cuttings, g_R and m_R of whole root respiration were 5.2 mmol CO₂ (g DM)⁻¹ and 9.7 nmol CO₂ (g DM)⁻¹ s⁻¹, respectively, at 22°C (Thongo M'Bou et al. 2010), in which ion uptake respiration was not separated from R_g . As for fine roots in field conditions (George et al. 2003), g_R coefficients were 5.1 (*P. taeda*) and 5.8 mmol CO₂ (g DM)⁻¹ (*L. styraciflua*), and m_R coefficients were 8.9 (*P. taeda*) and 10.2 nmol CO₂ (g DM)⁻¹ s⁻¹ (*L. styraciflua*) at 25°C. In our study for tree fine roots, g_R ($= c$) were 26.7 and 20.0 mmol CO₂ (g DM)⁻¹, respectively, in the DF and EF sites. In addition, m_R ($= d \cdot \exp(f \cdot T_s)$) were 5.5 (DF) and 2.7 (EF) nmol CO₂ (g DM)⁻¹ s⁻¹ at 22°C, and 7.9 (DF) and 3.6 (EF) nmol CO₂ (g DM)⁻¹ s⁻¹ at 25°C. The m_R increases with T_s according to Q_{10} of 3.3 and 2.6, respectively, in the DF and EF sites. Although the respiratory quotient should be considered, g_R of our study was relatively higher than those of the other species. However, our study's m_R were lower. Most of the differences in the coefficients possibly arose from differences in chemical composition of the roots, different rates of alternative pathway respiration, and different methods used (Lambers et al. 2008). In addition, differences between whole root respiration and fine root respiration would be another reason for the different coefficients.

Conclusions

We conducted a year-round field experiment to measure the production and biomass of tree fine roots, and soil respiration simultaneously in adjoining deciduous and evergreen forests on the same soil type. The periodic measurement of the three items was made in the same collar to minimize inconsistency among items by spatial positions. Using the field data, we partitioned tree fine root respiration into its growth and maintenance components on an annual basis by multiple regression using an empirical model. Growth (g_R) and maintenance (m_R) coefficients of the model were not significantly different between the forests. In comparison with reported values from laboratory or greenhouse experiments, in our study g_R was higher, but m_R was lower. Considerable uncertainty in the partitioning arose from a small sample size, the existence of thicker roots, and uncertainty in field measurement. However, the results suggest the availability of the approach using the model in field. To obtain more reliable results, ion uptake respiration should be incorporated into the model.

449

450 **Acknowledgements**

451 This study was supported by JSPS KAKENHI (nos. 25241002 and 17K20037) and the Environment Research and
452 Technology Development Fund (2-1705) of the Environmental Restoration and Conservation Agency. We thank
453 the Hokkaido Regional Office of the Forestry Agency for allowing us to use the study site, and N. Saigusa, R.
454 Hirata and the staff of CGER for managing the site, and K. Fukuzawa and K. Takagi for teaching us how to measure
455 fine roots.

456

References

- Abramoff RZ, Finzi A (2016) Seasonality and partitioning of root allocation to rhizosphere soils in a midlatitude forest. *Ecosphere* 7: 1-19. doi: e01547. 10.1002/ecs2.1547.
- Abramoff RZ, Finzi AC (2015) Are above- and below-ground phenology in sync? *New Phytologist* 205: 1054-1061. doi: 10.1111/nph.13111.
- Addo-Danso SD, Prescott CE, Smith AR (2016) Methods for estimating root biomass and production in forest and woodland ecosystem carbon studies: A review. *Forest Ecol Manag* 359: 332-351. doi: 10.1016/j.foreco.2015.08.015.
- Amthor J, S. (2000) The McCree-de Wit-Penning de Vries-Thornly respiration paradigms: 30 years later. *Ann Bot-London* 86: 1-20.
- Bonan G (2008) Forests and climate change: forcings, feedbacks, and the climate benefits of forests. *Science* 320: 1444-1449.
- Brassard BW, Chen HYH, Bergeron Y (2009) Influence of Environmental Variability on Root Dynamics in Northern Forests. *Crit Rev Plant Sci* 28: 179-197. doi: 10.1080/07352680902776572.
- Brunner I, Bakker MR, Björk RG, Hirano Y, Lukac M, Aranda X, Børja I, Eldhuset TD, Helmisaari HS, Jourdan C, Konôpka B, López BC, Miguel Pérez C, Persson H, Ostonen I (2013) Fine-root turnover rates of European forests revisited: an analysis of data from sequential coring and ingrowth cores. *Plant Soil* 362: 357-372. doi: 10.1007/s11104-012-1313-5.
- Chapin III FS, Matson PA, Vitousek PM (2011) *Plant Respiration. Principles of Terrestrial Ecosystem Ecology*. Second Edition edn. Springer, New York.
- Coleman MD, Dickson RE, Isebrands JG (2000) Contrasting fine-root production, survival and soil CO₂ efflux in pine and poplar plantations. *Plant Soil* 225: 129-139.
- Davidson EA, Richardson AD, Savage KE, Hollinger D (2006) A distinct seasonal pattern of the ratio of soil respiration to total ecosystem respiration in a spruce-dominated forest. *Global Change Biol* 12: 230-239. doi: 10.1111/j.1365-2486.2005.01062.x.
- Epron D, Farque L, Lucot E, Badot PM (1999) Soil CO₂ efflux in a beech forest: the contribution of root respiration. *Annals of Forest Science* 56: 289-295.
- Fernandez CW, Kennedy PG (2016) Revisiting the 'Gadgil effect': do interguild fungal interactions control carbon cycling in forest soils? *New Phytol* 209: 1382-1394. doi: 10.1111/nph.13648.
- Finér L, Ohashi M, Noguchi K, Hirano Y (2011a) Factors causing variation in fine root biomass in forest ecosystems. *Forest Ecol Manag* 261: 265-277. doi: 10.1016/j.foreco.2010.10.016.
- Finér L, Ohashi M, Noguchi K, Hirano Y (2011b) Fine root production and turnover in forest ecosystems in relation to stand and environmental characteristics. *Forest Ecol Manag* 262: 2008-2023. doi: 10.1016/j.foreco.2011.08.042.
- Fukuzawa K, Shibata H, Takagi K, Satoh F, Koike T, Sasa K (2013) Temporal variation in fine-root

biomass, production and mortality in a cool temperate forest covered with dense understory vegetation in northern Japan. *Forest Ecol Manag* 310: 700-710. doi: 10.1016/j.foreco.2013.09.015.

George K, Norby RJ, Hamilton JG, DeLucia EH (2003) Fine-root respiration in a loblolly pine and sweetgum forest growing in elevated CO₂. *New Phytologist* 160: 511-522. doi: 10.1046/j.1469-8137.2003.00911.x.

Gholz HL, Wein DA, Smitherman SM, Harmon ME, Parton WJ (2000) Long-term dynamics of pine and hardwood litter in contrasting environments: toward a global model of decomposition. *Global Change Biol* 6: 751-765.

Gifford RM (2003) Plant respiration in productivity models: conceptualisation, representation and issues for global terrestrial carbon-cycle research. *Functional Plant Biology* 30: 171. doi: 10.1071/fp02083.

Hanson PJ, Edwards NT, Garten CT, Andrews JA (2000) Separation root and soil microbial contributions to soil respiration: A review of methods and observations. *Biogeochemistry* 48: 115-146.

Hendricks JJ, Hendrick RL, Wilson CA, Mitchell RJ, Pecot SD, Guo D (2006) Assessing the patterns and controls of fine root dynamics: an empirical test and methodological review. *J Ecol* 94: 40-57. doi: 10.1111/j.1365-2745.2005.01067.x.

Hirano T, Hirata R, Fujinuma Y, Saigusa N, Yamamoto S, Harazono Y, Takada M, Inukai K, Inoue G (2003a) CO₂ and water vapor exchange of a larch forest in northern Japan. *Tellus B* 55: 244-257. doi: 10.1034/j.1600-0889.2003.00063.x.

Hirano T, Kim H, Tanaka Y (2003b) Long-term half-hourly measurement of soil CO₂ concentration and soil respiration in a temperate deciduous forest. *J Geophys Res-Atmos* 108. doi: 10.1029/2003JD003766.

Hirano T, Suzuki K, Hirata R (2017) Energy balance and evapotranspiration changes in a larch forest caused by severe disturbance during an early secondary succession. *Agr Forest Meteorol* 232: 457-468. doi: 10.1016/j.agrformet.2016.10.003.

Hirata R, Hirano T, Saigusa N, Fujinuma Y, Inukai K, Kitamori Y, Takahashi Y, Yamamoto S (2007) Seasonal and interannual variations in carbon dioxide exchange of a temperate larch forest. *Agr Forest Meteorol* 147: 110-124. doi: 10.1016/j.agrformet.2007.07.005.

Hopkins F, Gonzalez-Meler MA, Flower CE, Lynch DJ, Czimczik C, Tang J, Subke J-A (2013) Ecosystem-level controls on root-rhizosphere respiration. *New Phytologist* 199: 339-351. doi: 10.1111/nph.12271.

Hutchinson GL, Livingston GP (2001) Vents and seals in non-steady-state chambers used for measuring gas exchange between soil and the atmosphere. *Eur J Soil Sci* 52: 675-682.

Janssens IA, Lankreijer H, Matteucci G, Kowalski AS, Buchmann N, Epron D, Pilegaard K, Kutsch W, Longdoz B, Grunwald T, Montagnani L, Dore S, Rebmann C, Moors EJ, Grelle A, Rannik U, Morgenstern K, Oltchev S, Clement R, Gudmundsson J, Minerbi S, Berbigier P, Ibrom

- A, Moncrieff J, Aubinet M, Bernhofer C, Jensen NO, Vesala T, Granier A, Schulze ED, Lindroth A, Dolman AJ, Jarvis PG, Ceulemans R, Valentini R (2001) Productivity overshadows temperature in determining soil and ecosystem respiration across European forests. *Global Change Biol* 7: 269-278.
- Johnson IR (1990) Plant respiration in relation to growth, maintenance, ion uptake and nitrogen assimilation. *Plant Cell Environ* 13: 319-328.
- Keenan RJ, Reams GA, Achard F, de Freitas JV, Grainger A, Lindquist E (2015) Dynamics of global forest area: Results from the FAO Global Forest Resources Assessment 2015. *Forest Ecol Manag* 352: 9-20. doi: 10.1016/j.foreco.2015.06.014.
- Lambers H, Chapin III FS, Pons TL (2008) The Role of Respiration in Plant Carbon Balance. *Plant Physiological Ecology*. Second Edition edn. Springer, New York.
- Le Quéré C, Andrew RM, Friedlingstein P, Sitch S, Hauck J, Pongratz J, Pickers PA, Korsbakken JI, Peters GP, Canadell JG, Arneeth A, Arora VK, Barbero L, Bastos A, Bopp L, Chevallier F, Chini LP, Ciais P, Doney SC, Gkritzalis T, Goll DS, Harris I, Haverd V, Hoffman FM, Hoppema M, Houghton RA, Hurtt G, Ilyina T, Jain AK, Johannessen T, Jones CD, Kato E, Keeling RF, Goldewijk KK, Landschützer P, Lefèvre N, Lienert S, Liu Z, Lombardozzi D, Metzl N, Munro DR, Nabel JEMS, Nakaoka S-i, Neill C, Olsen A, Ono T, Patra P, Peregon A, Peters W, Peylin P, Pfeil B, Pierrot D, Poulter B, Rehder G, Resplandy L, Robertson E, Rocher M, Rödenbeck C, Schuster U, Schwinger J, Séférian R, Skjelvan I, Steinhoff T, Sutton A, Tans PP, Tian H, Tilbrook B, Tubiello FN, van der Laan-Luijkx IT, van der Werf GR, Viovy N, Walker AP, Wiltshire AJ, Wright R, Zaehle S, Zheng B (2018) Global Carbon Budget 2018. *Earth System Science Data* 10: 2141-2194. doi: 10.5194/essd-10-2141-2018.
- Liang N, Hirano T, Zheng ZM, Tang J, Fujinuma Y (2010) Soil CO₂ efflux of a larch forest in northern Japan. *Biogeosciences* 7: 3447-3457. doi: 10.5194/bg-7-3447-2010.
- Liu S, Luo D, Yang H, Shi Z, Liu Q, Zhang L, Kang Y (2018) Fine Root Dynamics in Three Forest Types with Different Origins in a Subalpine Region of the Eastern Qinghai-Tibetan Plateau. *Forests* 9: 517. doi: 10.3390/f9090517.
- Majdi H, Pregitzer K, Morén A-S, Nylund J-E, I. Ågren G (2005) Measuring Fine Root Turnover in Forest Ecosystems. *Plant Soil* 276: 1-8. doi: 10.1007/s11104-005-3104-8.
- Makkonen K, Helmisaari HS (1998) Seasonal and yearly variations of fine-root biomass and necromass in a Scots pine (*Pinus sylvestris* L.) stand. *Forest Ecol Manag* 102: 283-290.
- McCormack ML, Adams TS, Smithwick AH, Eissenstat DM (2014) Variability in root production, phenology, and turnover rate among 12 temperate tree species. *Ecology* 95: 2224-2235.
- McCormack ML, Dickie IA, Eissenstat DM, Fahey TJ, Fernandez CW, Guo D, Helmisaari HS, Hobbie EA, Iversen CM, Jackson RB, Leppalammi-Kujansuu J, Norby RJ, Phillips RP, Pregitzer KS, Pritchard SG, Rewald B, Zadworny M (2015) Redefining fine roots improves understanding of below-ground contributions to terrestrial biosphere processes. *New Phytol* 207: 505-518. doi: 10.1111/nph.13363.

570 McCree KJ (1974) Equations for the rate of dark respiration of white clove and grain sorghum, as
571 functions of dry weight, photosynthetic rate, and temperature. *Crop Sci* 14: 509-514.

572 Mizoguchi Y, Ohtani Y, Takanashi S, Iwata H, Yasuda Y, Nakai Y (2012) Seasonal and interannual
573 variation in net ecosystem production of an evergreen needleleaf forest in Japan. *J Forest*
574 *Res-Jpn* 17: 283-295. doi: 10.1007/s10310-011-0307-0.

575 Moyano F, Atkin OK, Bahn M, Bruhn D, Burton AJ, Heinemeyer A, Kutsch W, Wieser G (2009)
576 Respiration from roots and the mycorrhizosphere. In: W Kutsch, M Bahn, A Heinemeyer
577 (eds) *Soil Carbon Dynamics*. Cambridge University Press, Cambridge, UK.

578 Noguchi K, Sakata T, Mizoguchi T, Takahashi M (2005) Estimating the production and mortality
579 of fine roots in a Japanese cedar (*Cryptomeria japonica* D. Don) plantation using a
580 minirhizotron technique. *J Forest Res-Jpn* 10: 435-441. doi: 10.1007/s10310-005-0163-x.

581 Pan Y, Birdsey RA, Fang J, Houghton R, Kauppi PE, Kurz WA, Phillips OL, Shvidenko A, Lewis
582 SL, Canadell JG, Ciais P, Jackson RB, Pacala SW, McGuire AD, Piao S, Rautiainen A, Sitch
583 S, Hayes D (2011) A large and persistent carbon sink in the world's forests. *Science* 333:
584 988-993. doi: 10.1126/science.1201609.

585 Penning de Vries FWT (1974) Substrate utilization and respiration in relation to growth and
586 maintenance in higher plants. *Netherlands Journal of Agricultural Science* 22: 40-44.

587 Radvile L, McCormack ML, Post E, Eissenstat DM (2016) Root phenology in a changing climate. *J*
588 *Exp Bot* 67: 3617-3628. doi: 10.1093/jxb/erw062.

589 Ravindranath NH, Ostwald M (2008) *Carbon Inventory Methods: Handbook for Greenhouse Gas*
590 *Inventory, Carbon Mitigation and Roundwood Production Projects*. Springer.

591 Reichstein M, Janssens IA (2009) Semi-empirical modeling of the response of soil respiration to
592 environmental factors in laboratory and field conditions. In: W Kutsch, M Bahn, A
593 Heinemeyer (eds) *Soil Carbon Dynamics*. Cambridge University Press, Cambridge, UK.

594 Richter DD, Markewitz D, Trumbore SE, Wells CG (1999) Rapid accumulation and turnover of soil
595 carbon in a re-establishing forest. *Nature* 400: 56-58.

596 Sakai Y, Takahashi M, Tanaka N (2007) Root biomass and distribution of a *Picea*—*Abies* stand and
597 a *Larix*—*Betula* stand in pumiceous Entisols in Japan. *J Forest Res-Jpn* 12: 120-125. doi:
598 10.1007/s10310-006-0270-3.

599 Sano T, Hirano T, Liang N, Hirata R, Fujinuma Y (2010) Carbon dioxide exchange of a larch forest
600 after a typhoon disturbance. *Forest Ecol Manag* 260: 2214-2223. doi:
601 10.1016/j.foreco.2010.09.026.

602 Silver WL, Miya RK (2001) Global patterns in root decomposition: comparisons of climate and litter
603 quality effects. *Oecologia* 129: 407-419. doi: 10.1007/s004420100740.

604 Steinaker DF, Wilson SD, Peltzer DA (2010) Asynchronicity in root and shoot phenology in grasses
605 and woody plants. *Global Change Biol* 16: 2241-2251. doi: 10.1111/j.1365-2486.2009.02065.x.

606 Subke J-A, Inglis I, Francesca Cotrufo M (2006) Trends and methodological impacts in soil CO₂
607 efflux partitioning: A metaanalytical review. *Global Change Biol* 12: 921-943. doi:

10.1111/j.1365-2486.2006.01117.x.

- Sun LF, Teramoto M, Liang N, Yazaki T, Hirano T (2017) Comparison of litter-bag and chamber methods for measuring CO₂ emissions from leaf litter decomposition in a temperate forest. *Journal of Agricultural Meteorology* 73: 59-67. doi: 10.2480/agrmet.D-16-00012.
- Sweetlove LJ, Williams TC, Cheung CY, Ratcliffe RG (2013) Modelling metabolic CO₂ evolution--a fresh perspective on respiration. *Plant Cell Environ* 36: 1631-1640. doi: 10.1111/pce.12105.
- Teramoto M, Liang N, Zeng J, Saigusa N, Takahashi Y (2017) Long-term chamber measurements reveal strong impacts of soil temperature on seasonal and inter-annual variation in understory CO₂ fluxes in a Japanese larch (*Larix kaempferi* Sarg.) forest. *Agr Forest Meteorol* 247: 194-206. doi: 10.1016/j.agrformet.2017.07.024.
- Thongo M'Bou A, Saint-André L, de Grandcourt A, Nouvellon Y, Jourdan C, Mialoundama F, Epron D (2010) Growth and maintenance respiration of roots of clonal Eucalyptus cuttings: scaling to stand-level. *Plant Soil* 332: 41-53. doi: 10.1007/s11104-009-0272-y.
- Thornley JHM (1970) Respiration, growth and maintenance in plants. *Nature* 227: 304-305.
- Thornton PE, Rosenbloom NA (2005) Ecosystem model spin-up: Estimating steady state conditions in a coupled terrestrial carbon and nitrogen cycle model. *Ecol Model* 189: 25-48. doi: 10.1016/j.ecolmodel.2005.04.008.
- Tierney G, Fahey TJ, Groffman PM, Hardy JP, Fitzhugh RD, Driscoll CT, Yavitt JB (2003) Environmental control of fine root dynamics in a northern hardwood forest. *Global Change Biol* 9: 670-679.
- Vogt KA, Vogt DJ, Bloomfield J (1998) Analysis of some direct and indirect methods for estimating root biomass and production of forests at an ecosystem level. *Plant Soil* 200: 71-89.
- Wang W, Zu Y, Wang H, Hirano T, Takagi K, Sasa K, Koike T (2005) Effect of collar insertion on soil respiration in a larch forest measured with a LI-6400 soil CO₂ flux system. *J Forest Res-Jpn* 10: 57-60. doi: 10.1007/s10310-004-0102-2.
- Warren JM, Hanson PJ, Iversen CM, Kumar J, Walker AP, Wullschlegel SD (2015) Root structural and functional dynamics in terrestrial biosphere models--evaluation and recommendations. *New Phytol* 205: 59-78. doi: 10.1111/nph.13034.
- Wieder RK, Lang GE (1982) A critique of the analytical methods used in examining decomposition data obtained from litter bags. *Ecology* 63: 1636-1642.
- Yazaki T, Hirano T, Sano T (2016) Biomass Accumulation and Net Primary Production during the Early Stage of Secondary Succession after a Severe Forest Disturbance in Northern Japan. *Forests* 7. doi: 10.3390/f7110287.
- Yuan ZY, Chen H (2010) Fine Root Biomass, Production, Turnover Rates, and Nutrient Contents in Boreal Forest Ecosystems in Relation to Species, Climate, Fertility, and Stand Age: Literature Review and Meta-Analyses. *Crit Rev Plant Sci* 29: 204-221. doi: 10.1080/07352689.2010.483579.

Table 1 Soil temperature (T_s) at a depth of 5 cm and soil water content (SWC) of 5-cm-thick surface soil in chamber collars with four different treatments (CC, SC, LC and TC). T_s and SWC were measured simultaneously with soil CO₂ efflux once or twice a month between May 2015 and November 2015. Mean $\pm$ 1 SD of all measurements is shown ($n = 12$). Different letters in the same row denote significant difference among the treatments and between the sites ($P < 0.05$) according to Tukey's HSD after ANOVA ($P < 0.01$).

Treatment	Site	T_s (°C)	SWC (m ³ m ⁻³)
CC	DF	14.7 $\pm$ 0.9a	0.25 $\pm$ 0.05a
	EF	12.4 $\pm$ 0.4b	0.28 $\pm$ 0.04a
SC	DF	14.7 $\pm$ 0.9a	0.27 $\pm$ 0.07a
	EF	12.5 $\pm$ 0.4b	0.26 $\pm$ 0.07a
LC	DF	14.6 $\pm$ 0.8a	0.29 $\pm$ 0.08a
	EF	12.5 $\pm$ 0.4b	0.25 $\pm$ 0.06a
TC	DF	14.4 $\pm$ 0.8a	0.43 $\pm$ 0.09b
	EF	12.3 $\pm$ 0.3b	0.35 $\pm$ 0.06b
ANOVA (P value)	Site	<0.0001	0.25
	Treatment	0.85	0.0004
	Interaction	0.98	0.35

Table 2 Biomass of tree fine (< 2 mm), medium (< 10 mm) and coarse (> 10 mm) roots sampled in November 2014 from 30-cm-thick surface soil. Mean $\pm$ 1 SD ($n = 5$) is shown. The decomposition constant (k) determined by curve fitting to Eq. 5 (R^2) and annual decomposition (R_d) of tree dead roots in trenched collars (TC) were also shown. Numbers in parentheses denote the percentage against Total.

Site	Root class	Initial biomass (g DM m ⁻²)	k (yr ⁻¹)	r^2	RD (g C m ⁻² yr ⁻¹)
DF	Fine	262 $\pm$ 128 (20)	0.42	0.98	45
	Medium	278 $\pm$ 152 (22)	0.25	0.91	31
	Coarse	750 $\pm$ 636 (58)	0.16	0.97	55
	Total	1290 (100)			131
EF	Fine	362 $\pm$ 104 (33)	0.29	0.98	46
	Medium	454 $\pm$ 194 (42)	0.14	0.91	30

Coarse	276 ± 384 (25)	0.10	0.91	13
Total	1092 (100)			89

Table 3 Annual sums (g C m⁻² year⁻¹) of soil respiration (R_s), leaf litter decomposition (R_l), root respiration (R_r), belowground heterotrophic respiration (R_h), soil CO₂ efflux on trenched collars (R_{TC}), and decomposition of tree dead roots in trenched collars (R_d) from November 2014 through November 2015 (mean ± 1 SD, n = 5). Numbers in parentheses denote the percentage against R_s .

Site	R_s	R_l	R_r	R_h	R_{TC}	R_d
DF	1255 ± 401	211 ± 224	610 ± 447	434 ± 257	776 ± 83	131
	(100)	(17)	(49)	(35)	(62)	(10)
EF	1139 ± 386	112 ± 118	393 ± 356	634 ± 248	835 ± 172	89
	(100)	(10)	(35)	(56)	(73)	(8)

Table 4 Summary of tree fine root dynamics in the root mat and A horizon. Annual production (P), biomass change (ΔB), mortality (M), mean biomass (B) and turnover rates between November 2014 and November 2015 are shown as mean ± 1 SD (n = 5). Numbers in parentheses denote herbaceous fine root biomass.

Site	Root distribution	P (g DM m ⁻² year ⁻¹)	ΔB (g DM m ⁻² year ⁻¹)	M (g DM m ⁻² year ⁻¹)	Mean B (g DM m ⁻²)	Turnover rate (year ⁻¹)
DF	Root mat	178 ± 24	-25 ± 43	204 ± 66	190 ± 58 (64 ± 14)	0.94 ± 0.42
	A horizon	190 ± 22	39 ± 73	151 ± 63	150 ± 44 (46 ± 14)	1.27 ± 0.44
	Total	368 ± 42	14 ± 67	355 ± 73	340 ± 58 (110 ± 20)	1.08 ± 0.36
EF	Root mat	158 ± 42	-58 ± 42	216 ± 49	258 ± 32	0.61 ± 0.17
	A horizon	194 ± 46	10 ± 71	185 ± 87	258 ± 40	0.75 ± 0.18
	Total	352 ± 62	-48 ± 96	400 ± 126	516 ± 38	0.68 ± 0.12

Table 5 Fitted parameters of the model (Eq. 8) (± 1 standard error).

	DF	EF
c (g C g DM ⁻¹)	0.32 ± 0.33	0.24 ± 0.20
d (g C g DM ⁻¹ day ⁻¹)	0.00041 ± 0.0019	0.00036 ± 0.0010

$f(^{\circ}\text{C}^{-1})$	0.12 ± 0.21	0.094 ± 0.12
$h \text{ (g C m}^{-2} \text{ day}^{-1}\text{)}$	0.94 ± 0.66	0.35 ± 0.68
Adjusted R^2	0.87	0.93
P value	0.087	0.065

Table 6 Annual sums (g C m⁻² year⁻¹) of root respiration (R_r) in Table 3, tree fine root growth respiration (R_g), maintenance respiration (R_m), residual (h), and total ($R_g + R_m + h$) from November 2014 through November 2015. These values were calculated using Eq. 8 with parameters in Table 5. Numbers in parentheses denote the percentage against the total. Uncertainties (± 1 SD) were propagated from those of parameters and measurements.

	DF	EF
R_r	610	393
R_g	121 (19)	86 (22)
R_m	166 (26)	182 (46)
h	351 (55)	128 (32)
Total	638 (100)	396 (100)

Fig. 1 Layout of collars for control (CC), sampling (SC), litter removal (LC), and trenching (TC) treatments.

Fig. 2 Seasonal variations in daily mean a) soil temperature (T_s) at a depth of 5 cm ($n = 2$) and b) soil water content (SWC) at a depth of 3 cm ($n = 4$) from September 2014 through December 2015. SWC was measured at a nearby re-growing forest site.

Fig. 3 Seasonal variations in soil CO_2 efflux measured on SC and TC collars in a) DF and b) EF sites. Mean ± 1 SD are shown ($n = 5$).

Fig. 4 Relationship between soil CO₂ efflux and soil temperature at a depth of 5 cm for SC and TC collars in a) DF and b) EF sites. Mean $\pm$ 1 SD are shown ($n = 5$). An exponential equation (Eq. 3) is fitted significantly ($P < 0.001$).

Fig. 5 Temperature-normalized soil CO₂ efflux (R_b) against soil water content (SWC) of 5-cm-thick surface soil for SC and TC collars in a) DF (at 14°C) and b) EF (at 12 °C) sites. Mean $\pm$ 1 SD are shown ($n = 5$).

Fig. 6 Seasonal variations in a) fine root biomass in DF site, b) tree fine root biomass (*B*) in DF site, c) *B* in EF site, d) tree fine root production (*P*) in DF site and e) *P* in EF site from November 2014 through November 2015. Mean $\pm$ 1 SD are shown ($n = 5$) in a) – c).

Fig. 7 Relationship between tree fine root production (P) and soil temperature at a depth of 5 cm. Mean $\pm$ 1 SD are shown ($n = 5$). A line is fitted significantly ($P < 0.05$).

Fig. 8 Relationship between root respiration (R_r) and tree fine root production (P). A line is fitted significantly ($p < 0.05$).