



Title	The adjoint group of a Coxeter quandle
Author(s)	Akita, Toshiyuki
Citation	Kyoto journal of mathematics, 60(4), 1245-1260 https://doi.org/10.1215/21562261-2019-0061
Issue Date	2020-12
Doc URL	http://hdl.handle.net/2115/80388
Type	article (author version)
File Information	Kyoto J. Math.60-4_1245-1260.pdf



[Instructions for use](#)

THE ADJOINT GROUP OF A COXETER QUANDLE

TOSHIYUKI AKITA

ABSTRACT. We give explicit descriptions of the adjoint group $\text{Ad}(Q_W)$ of the Coxeter quandle Q_W associated with an arbitrary Coxeter group W . The adjoint group $\text{Ad}(Q_W)$ turns out to be an intermediate group between W and the corresponding Artin group A_W , and fits into a central extension of W by a finitely generated free abelian group. We construct 2-cocycles of W corresponding to the central extension. In addition, we prove that the commutator subgroup of the adjoint group $\text{Ad}(Q_W)$ is isomorphic to the commutator subgroup of W . Finally, the root system Φ_W associated with a Coxeter group W turns out to be a rack. We prove that the adjoint group $\text{Ad}(\Phi_W)$ of Φ_W is isomorphic to the adjoint group of Q_W .

1. INTRODUCTION

A nonempty set Q equipped with a binary operation $Q \times Q \rightarrow Q$, $(x, y) \mapsto x * y$ is called a *quandle* if it satisfies the following three conditions:

- (1) $x * x = x$ ($x \in Q$),
- (2) $(x * y) * z = (x * z) * (y * z)$ ($x, y, z \in Q$),
- (3) For all $y \in Q$, the map $Q \rightarrow Q$ defined by $x \mapsto x * y$ is bijective.

If X satisfies (2) and (3) but not necessarily (1), then X is called a *rack*. Quandles and racks have been studied in low dimensional topology as well as in Hopf algebras (see Nosaka [19] and Andruskiewitsch-Graña [3] for instance). To any quandle or rack Q one can associate a group $\text{Ad}(Q)$ called the *adjoint group* of Q (also called the associated group or the enveloping group in the literature). It is defined by the presentation

$$\text{Ad}(Q) := \langle e_x \ (x \in Q) \mid e_y^{-1} e_x e_y = e_{x*y} \ (x, y \in Q) \rangle.$$

The assignment $Q \mapsto \text{Ad}(Q)$ is a functor from the category of quandles or racks to the category of groups. Although adjoint groups play important roles in the study of quandles and racks, not much is known about the structure of them, partly because the definition of $\text{Ad}(Q)$ by a possibly infinite presentation is difficult to work with in explicit calculations. We refer Eisermann [12] and Nosaka [18, 19] for generality of adjoint groups, and [9, 12, 18, 19] for descriptions of adjoint groups of certain classes of quandles.

In this paper, we will study the adjoint group of a Coxeter quandle. Let (W, S) be a Coxeter system, a pair of a Coxeter group W and the set S of Coxeter generators

Key words and phrases. quandle, rack, Coxeter group, root system, Artin group, pure Artin group, group cohomology.

of W . Following Nosaka [18], we define the *Coxeter quandle* Q_W associated with (W, S) to be the set of all reflections of W :

$$Q_W := \bigcup_{w \in W} w^{-1}Sw.$$

The quandle operation is given by the conjugation $x * y := y^{-1}xy = yxy$. The symmetric group \mathfrak{S}_n of n letters is a Coxeter group (of type A_{n-1}), and the associated Coxeter quandle $Q_{\mathfrak{S}_n}$ is nothing but the set of all transpositions. In their paper [2], Andruskiewitsch-Fantino-García-Vendramin obtained remarkable results concerning with $\text{Ad}(Q_{\mathfrak{S}_n})$. Namely, they proved that $\text{Ad}(Q_{\mathfrak{S}_n})$ is an intermediate group between \mathfrak{S}_n and the braid group B_n of n strands, in the sense that the canonical projection $B_n \twoheadrightarrow \mathfrak{S}_n$ splits into a sequence of surjections $B_n \twoheadrightarrow \text{Ad}(Q_{\mathfrak{S}_n}) \twoheadrightarrow \mathfrak{S}_n$, and that $\text{Ad}(Q_{\mathfrak{S}_n})$ fits into a central extension of the form

$$(1.1) \quad 0 \rightarrow \mathbb{Z} \rightarrow \text{Ad}(Q_{\mathfrak{S}_n}) \rightarrow \mathfrak{S}_n \rightarrow 1.$$

See [2, Proposition 3.2] and the proofs therein. Furthermore, the central extension (1.1) turns out to be the unique nontrivial central extension of \mathfrak{S}_n by \mathbb{Z} (see Corollary 4.6).

The primary purpose of this paper is to generalize those results to arbitrary Coxeter quandles. We will show that $\text{Ad}(Q_W)$ is an intermediate group between W and the Artin group A_W associated with W (Proposition 3.3), and that $\text{Ad}(Q_W)$ fits into a central extension of the form

$$(1.2) \quad 0 \rightarrow \mathbb{Z}^{c(W)} \rightarrow \text{Ad}(Q_W) \xrightarrow{\phi} W \rightarrow 1,$$

where $c(W)$ is the number of conjugacy classes of elements in Q_W (Theorem 3.1). As a byproduct of Theorem 3.1, we will determine the rational cohomology ring of $\text{Ad}(Q_W)$ (Corollary 3.6).

In case $c(W) = 1$, the central extension (1.2) turns out to be the unique nontrivial central extension of W by \mathbb{Z} (Corollary 4.6). As is known, the central extension (1.2) corresponds to the cohomology class $u_\phi \in H^2(W, \mathbb{Z}^{c(W)})$. We will construct 2-cocycles of W representing u_ϕ (Proposition 4.3 and Theorem 4.8).

Alternatively, Eisermann [12] claimed that $\text{Ad}(Q_{\mathfrak{S}_n})$ is isomorphic to the semidirect product $A_n \rtimes \mathbb{Z}$ where A_n is the alternating group on n letters, but he did not write down the proof. We will generalize his result to Coxeter quandles (Theorem 5.1 and Corollary 5.2).

In the final section, we deal with root systems. To each Coxeter system (W, S) , one can associate the root system $\Phi_W \subset V$ by using the geometric representation $W \rightarrow GL(V)$, where V is a real vector space with the basis $\{\alpha_s \mid s \in S\}$. The root system Φ_W turns out to be a rack with the rack operation $\alpha * \beta := t_\beta(\alpha)$ where $t_\beta \in GL(V)$ is the reflection along β . We close this paper by proving $\text{Ad}(\Phi_W) \cong \text{Ad}(Q_W)$ (Theorem 6.3).

Notation. Let G be a group, $g, h \in G$ elements of G and $m \geq 2$ a natural number. Define

$$(gh)_m := \underbrace{ghghg \cdots}_m \in G.$$

For example, $(gh)_2 = gh$, $(gh)_3 = ghg$, $(gh)_4 = ghgh$. Let $G_{\text{Ab}} := G/[G, G]$ be the abelianization of G , $\text{Ab}_G : G \rightarrow G_{\text{Ab}}$ the natural projection, and write $[g] := g[G, G] \in G_{\text{Ab}}$ for $g \in G$. For a quandle Q and elements $x_k \in Q$ ($k = 1, 2, 3, \dots$), we denote

$$x_1 * x_2 * x_3 * \cdots * x_\ell := (\cdots ((x_1 * x_2) * x_3) * \cdots) * x_\ell,$$

for simplicity.

2. COXETER GROUPS AND COXETER QUANDLES

2.1. Coxeter groups. Let S be a finite set and $m : S \times S \rightarrow \mathbb{N} \cup \{\infty\}$ a map satisfying the following conditions:

- (1) $m(s, s) = 1$ for all $s \in S$;
- (2) $2 \leq m(s, t) = m(t, s) \leq \infty$ for all distinct $s, t \in S$.

The map m is represented by the *Coxeter graph* Γ whose vertex set is S and whose edges are the unordered pairs $\{s, t\} \subset S$ such that $m(s, t) \geq 3$. The edges with $m(s, t) \geq 4$ are labeled by the number $m(s, t)$. The *Coxeter system* associated with Γ is the pair (W, S) , where W is the group generated by $s \in S$ and the fundamental relations $(st)^{m(s,t)} = 1$ ($s, t \in S, m(s, t) < \infty$):

$$(2.1) \quad W := \langle s \in S \mid (st)^{m(s,t)} = 1 (s, t \in S, m(s, t) < \infty) \rangle.$$

The group W is called the *Coxeter group* (of type Γ), and elements of S are called *Coxeter generators* of W (also called simple reflections in the literature). Note that the order of the product st is precisely $m(s, t)$. In particular, every Coxeter generator $s \in S$ has order 2. It is easy to check that the defining relations in (2.1) are equivalent to the following relations

$$(2.2) \quad s^2 = 1 (s \in S), (st)_{m(s,t)} = (ts)_{m(s,t)} (s, t \in S, s \neq t, m(s, t) < \infty).$$

Finally, the *odd subgraph* Γ_{odd} is a subgraph of Γ whose vertex set is S and whose edges are the unordered pairs $\{s, t\} \subset S$ such that $m(s, t)$ is an odd integer. We refer Björner-Brenti [4], Bourbaki [5], Davis [10], and Humphreys [15] for further details of Coxeter groups.

2.2. Conjugacy classes of reflections. Let

$$Q_W := \bigcup_{w \in W} w^{-1} S w$$

be the set of reflections in W as in the introduction (the underlying set of the Coxeter quandle). Let \mathcal{O}_W be the set of conjugacy classes of elements of Q_W under W , and \mathcal{R}_W a complete set of representatives of conjugacy classes. We may choose such that $\mathcal{R}_W \subseteq S$. Let $c(W)$ be the cardinality of \mathcal{O}_W . The following two propositions are well-known and easy to prove (see Björner-Brenti [4, Chapter 1, Exercises 16–17] for instance).

Proposition 2.1. *The elements of \mathcal{O}_W are in one-to-one correspondence with the connected components of Γ_{odd} . Consequently, $c(W)$ equals to the number of connected components of Γ_{odd} .*

To be precise, the conjugacy class of $s \in S$ corresponds to the connected component of Γ_{odd} containing $s \in S$.

Proposition 2.2. W_{Ab} is the elementary abelian 2-group with a basis $\{[s] \in W_{\text{Ab}} \mid s \in \mathcal{R}_W\}$. In particular, $W_{\text{Ab}} \cong (\mathbb{Z}/2)^{c(W)}$.

2.3. Coxeter quandles. Now we turn our attention to Coxeter quandles. Let Q_W be the Coxeter quandle associated with (W, S) and

$$\text{Ad}(Q_W) := \langle e_x (x \in Q_W) \mid e_y^{-1} e_x e_y = e_{x*y} (x, y \in Q_W) \rangle$$

the adjoint group of Q_W . Observe that

$$(2.3) \quad e_y^{-1} e_x e_y = e_{x*y} = e_{y^{-1}xy}$$

and

$$(2.4) \quad e_y e_x e_y^{-1} = e_{x*y},$$

where (2.4) follows from $e_y^{-1} e_{x*y} e_y = e_{x*y*y} = e_x$.

Proposition 2.3. $\text{Ad}(Q_W)$ is generated by $e_s (s \in S)$.

Proof. Given $x \in Q_W$, we can express x as $x = (s_1 s_2 \cdots s_\ell)^{-1} s_0 (s_1 s_2 \cdots s_\ell)$ for some $s_i \in S$ ($0 \leq i \leq \ell$) by the definition of Q_W . Applying (2.3), we have

$$e_x = e_{(s_1 s_2 \cdots s_\ell)^{-1} s_0 (s_1 s_2 \cdots s_\ell)} = e_{s_0 * s_1 * s_2 * \cdots * s_\ell} = (e_{s_1} e_{s_2} \cdots e_{s_\ell})^{-1} e_{s_0} (e_{s_1} e_{s_2} \cdots e_{s_\ell}),$$

proving the proposition. \square

Proposition 2.4. $\text{Ad}(Q_W)_{\text{Ab}}$ is the free abelian group with a basis $\{[e_s] \mid s \in \mathcal{R}_W\}$. In particular, $\text{Ad}(Q_W)_{\text{Ab}} \cong \mathbb{Z}^{c(W)}$.

Proof. By the definition of $\text{Ad}(Q_W)$, the abelianization $\text{Ad}(Q_W)_{\text{Ab}}$ is generated by $[e_x] (x \in Q_W)$ subject to the relations $[e_x] = [e_{y^{-1}xy}]$, $[e_x][e_y] = [e_y][e_x] (x, y \in Q_W)$. Consequently, $[e_x] = [e_y] \in \text{Ad}(Q_W)_{\text{Ab}}$ if and only if $x, y \in Q_W$ are conjugate in W . We conclude that $\text{Ad}(Q_W)_{\text{Ab}}$ is the free abelian group with a basis $\{[e_s] \mid s \in \mathcal{R}_W\}$. \square

Let $\phi : \text{Ad}(Q_W) \rightarrow W$ be a surjective homomorphism defined by $e_x \mapsto x$, which is well-defined by virtue of (2.3), and let $C_W := \ker \phi$ be its kernel.

Lemma 2.5. C_W is a central subgroup of $\text{Ad}(Q_W)$.

Proof. Given $g \in C_W$, it suffices to prove $g^{-1} e_x g = e_x$ for all $x \in Q_W$. To do so, set $g = e_{y_1}^{\varepsilon_1} e_{y_2}^{\varepsilon_2} \cdots e_{y_\ell}^{\varepsilon_\ell} (\varepsilon_i \in \{\pm 1\})$. Applying (2.3) and (2.4), we have

$$g^{-1} e_x g = (e_{y_1}^{\varepsilon_1} e_{y_2}^{\varepsilon_2} \cdots e_{y_\ell}^{\varepsilon_\ell})^{-1} e_x (e_{y_1}^{\varepsilon_1} e_{y_2}^{\varepsilon_2} \cdots e_{y_\ell}^{\varepsilon_\ell}) = e_{x*y_1*y_2*\cdots*y_\ell}.$$

Now $x*y_1*y_2*\cdots*y_\ell = (y_1 y_2 \cdots y_\ell)^{-1} x (y_1 y_2 \cdots y_\ell)$, and the proposition follows from $\phi(g) = y_1^{\varepsilon_1} y_2^{\varepsilon_2} \cdots y_\ell^{\varepsilon_\ell} = y_1 y_2 \cdots y_\ell = 1$. \square

Lemma 2.6. If $x, y \in Q_W$ are conjugate in W , then $e_x^2 = e_y^2 \in C_W$.

Proof. It is obvious that $e_x^2 \in C_W$ for all $x \in Q_W$. Since C_W is a central subgroup, $e_x^2 = e_y^{-1} e_x^2 e_y = (e_y^{-1} e_x e_y)^2 = e_{y^{-1}xy}^2$ holds for all $x, y \in Q_W$, which implies the lemma. \square

3. ARTIN GROUPS AND THE PROOF OF THE MAIN RESULT

Now we state the main result of this paper:

Theorem 3.1. *The central subgroup C_W is the free abelian group with a basis $\{e_s^2 \mid s \in \mathcal{R}_W\}$. In particular, $C_W \cong \mathbb{Z}^{c(W)}$.*

As a consequence of Theorem 3.1, $\text{Ad}(Q_W)$ fits into a central extension of the form $0 \rightarrow \mathbb{Z}^{c(W)} \rightarrow \text{Ad}(Q_W) \rightarrow W \rightarrow 1$ as stated in the introduction. We begin with the determination of the rank of C_W .

Proposition 3.2. $\text{rank}(C_W) = c(W)$.

Proof. The central extension $1 \rightarrow C_W \rightarrow \text{Ad}(Q_W) \xrightarrow{\phi} W \rightarrow 1$ yields the following exact sequence for the rational homology of groups:

$$H_2(W, \mathbb{Q}) \rightarrow H_1(C_W, \mathbb{Q})_W \rightarrow H_1(\text{Ad}(Q_W), \mathbb{Q}) \rightarrow H_1(W, \mathbb{Q})$$

(see Brown [8, Corollary VII.6.4]). Here the co-invariants $H_1(C_W, \mathbb{Q})_W$ coincides with $H_1(C_W, \mathbb{Q})$ because C_W is a central subgroup of $\text{Ad}(Q_W)$. It is known that the rational (co)homology of a Coxeter group is trivial (see Akita [1, Proposition 5.2] or Davis [10, Theorem 15.1.1]). As a result, we have an isomorphism $H_1(C_W, \mathbb{Q}) \cong H_1(\text{Ad}(Q_W), \mathbb{Q})$ and hence we have

$$\begin{aligned} \text{rank}(C_W) &= \dim_{\mathbb{Q}}(C_W \otimes \mathbb{Q}) = \dim_{\mathbb{Q}} H_1(C_W, \mathbb{Q}) \\ &= \dim_{\mathbb{Q}} H_1(\text{Ad}(Q_W), \mathbb{Q}) = \dim_{\mathbb{Q}}(\text{Ad}(Q_W)_{\text{Ab}} \otimes \mathbb{Q}) = c(W) \end{aligned}$$

as desired. \square

To proceed further, we need the notions of Artin groups and pure Artin groups. Given a Coxeter system (W, S) , the *Artin group* A_W associated with (W, S) is the group defined by the presentation

$$A_W := \langle a_s \ (s \in S) \mid (a_s a_t)_{m(s,t)} = (a_t a_s)_{m(s,t)} \ (s, t \in S, s \neq t, m(s,t) < \infty) \rangle.$$

In view of (2.2), there is an obvious surjective homomorphism $\pi : A_W \rightarrow W$ defined by $a_s \mapsto s$ ($s \in S$). The *pure Artin group* P_W associated with (W, S) is defined to be the kernel of π so that there is an extension

$$1 \rightarrow P_W \hookrightarrow A_W \xrightarrow{\pi} W \rightarrow 1.$$

In case W is the symmetric group on n letters, A_W is the braid group of n strands, and P_W is the pure braid group of n strands. Artin groups were introduced by Brieskorn-Saito [7]. Little is known about the structure of general Artin groups. Among others, the following questions are still open.

- (1) Are Artin groups torsion free?
- (2) What is the center of Artin groups?
- (3) Do Artin groups have solvable word problem?
- (4) Are there finite $K(\pi, 1)$ -complexes for Artin groups?

See survey articles by Paris [20–22] for further details of Artin groups.

Proposition 3.3. *The assignment $a_s \mapsto e_s$ ($s \in S$) yields a well-defined surjective homomorphism $\psi : A_W \rightarrow \text{Ad}(Q_W)$.*

Proof. As for the well-definedness, it suffices to show $(e_s e_t)_{m(s,t)} = (e_t e_s)_{m(s,t)}$ for all distinct $s, t \in S$ with $m(s,t) < \infty$. Applying the relation $e_x e_y = e_y e_{x*y}$ ($x, y \in Q_W$) repeatedly as in

$$(e_t e_s)_{m(s,t)} = e_t e_s e_t e_s \cdots = e_s e_{t*s} e_t e_s \cdots = e_s e_t e_{t*s*t} e_s \cdots = \cdots,$$

we obtain

$$(e_t e_s)_{m(s,t)} = (e_s e_t)_{m(s,t)-1} e_x,$$

where

$$\begin{aligned} x &= \overbrace{t * s * t * s * \cdots}^{m(s,t)} = (st)_{m(s,t)-1}^{-1} t (st)_{m(s,t)-1} \\ &= (st)_{m(s,t)-1}^{-1} (ts)_{m(s,t)} \\ &= (st)_{m(s,t)-1}^{-1} (st)_{m(s,t)}. \end{aligned}$$

In the last equality we used the relation $(st)_{m(s,t)} = (ts)_{m(s,t)}$. It follows that x is the last letter in $(st)_{m(s,t)}$, i.e. $x = s$ or $x = t$ according as $m(s,t)$ is odd or $m(s,t)$ is even. We conclude that $(e_s e_t)_{m(s,t)-1} e_x = (e_s e_t)_{m(s,t)}$ as desired. Finally, the surjectivity follows from Proposition 2.3. \square

As a result, the adjoint group $\text{Ad}(Q_W)$ is an intermediate group between a Coxeter group W and the corresponding Artin group A_W , in the sense that the canonical surjection $\pi : A_W \twoheadrightarrow W$, $a_s \mapsto s$ ($s \in S$) splits into a sequence of surjections

$$A_W \xrightarrow{\psi} \text{Ad}(Q_W) \xrightarrow{\phi} W, \quad a_s \mapsto e_s \mapsto s \quad (s \in S).$$

Proposition 3.4. P_W is the normal closure of $\{a_s^2 \mid s \in S\}$ in A_W . In other words, P_W is generated by $g^{-1} a_s^2 g$ ($s \in S, g \in A_W$).

Proof. Given a Coxeter system (W, S) , let $F(S)$ be the free group on S and put

$$R := \{(st)_{m(s,t)} (ts)_{m(s,t)}^{-1} \mid s, t \in S, s \neq t, m(s,t) < \infty\}, \quad Q := \{s^2 \mid s \in S\}.$$

Let $N(R), N(Q), N(R \cup Q)$ be the normal closure of $R, Q, R \cup Q$ in $F(S)$, respectively. The third isomorphism theorem yields a short exact sequence of groups

$$1 \rightarrow \frac{N(R)N(Q)}{N(R)} \rightarrow \frac{F(S)}{N(R)} \xrightarrow{p} \frac{F(S)}{N(R)N(Q)} \rightarrow 1.$$

Observe that the left term $N(R)N(Q)/N(R)$ is nothing but the normal closure of Q in $F(S)/N(R)$. Now $F(S)/N(R)N(Q) = F(S)/N(R \cup Q) = W$ by the definition of W and $F(S)/N(R)$ is identified with A_W via $s \mapsto a_s$ ($s \in S$). Under this identification, the map p is the canonical surjection $\pi : A_W \twoheadrightarrow W$ and hence the left term $N(R)N(Q)/N(R)$ coincides with P_W . The proposition follows. \square

Remark 3.5. Digne-Gomi [11, Corollary 6] obtained a presentation of P_W by using Reidemeister-Schreier method. Their presentation is infinite whenever W is infinite. Although Proposition 3.4 may be read off from their presentation, we wrote down the proof because our proof is much simpler than the arguments in [11].

Proof of Theorem 3.1. Consider the commutative diagram

$$\begin{array}{ccccccc}
 1 & \longrightarrow & P_W & \hookrightarrow & A_W & \xrightarrow{\pi} & W \longrightarrow 1 \\
 & & \downarrow \psi|_{P_W} & & \downarrow \psi & & \parallel \\
 1 & \longrightarrow & C_W & \hookrightarrow & \text{Ad}(Q_W) & \xrightarrow{\phi} & W \longrightarrow 1
 \end{array}$$

whose rows are exact. Since $\psi : A_W \rightarrow \text{Ad}(Q_W)$ is surjective, one can check that its restriction $\psi|_{P_W} : P_W \rightarrow C_W$ is also surjective. As P_W is generated by $g^{-1}a_s^2g$ ($s \in S, g \in A_W$) by Proposition 3.4, C_W is generated by

$$\psi(g^{-1}a_s^2g) = \psi(g)^{-1}e_s^2\psi(g) = e_s^2 \quad (s \in S, g \in A_W),$$

where the last equality follows from the fact that C_W is central by Lemma 2.5. Combining with Lemma 2.6, we see that C_W is generated by $\{e_s^2 \mid s \in \mathcal{R}_W\}$. Now $\text{rank}(C_W) = c(W)$ by Proposition 3.2 and $|\mathcal{R}_W| = c(W)$ by the definition of \mathcal{R}_W , C_W must be a free abelian group of rank $c(W)$ and $\{e_s^2 \mid s \in \mathcal{R}_W\}$ must be a basis of C_W . \square

As an immediate cosequence of Theorem 3.1, we can determine the rational cohomology ring of $\text{Ad}(Q_W)$:

Corollary 3.6. *For any Coxeter system (W, S) , the inclusion $C_W \hookrightarrow \text{Ad}(Q_W)$ induces an isomorphism*

$$H^*(\text{Ad}(Q_W), \mathbb{Q}) \xrightarrow{\cong} H^*(C_W, \mathbb{Q}) \cong \wedge_{\mathbb{Q}}(u_1, u_2, \dots, u_{c(W)})$$

with $\deg u_k = 1$ ($1 \leq k \leq c(W)$).

Proof. In the Lyndon-Hochschild-Serre spectral sequence

$$E_2^{pq} = H^p(W, H^q(C_W, \mathbb{Q})) \Rightarrow H^{p+q}(\text{Ad}(Q_W), \mathbb{Q})$$

associated with the central extension $1 \rightarrow C_W \rightarrow \text{Ad}(Q_W) \rightarrow W \rightarrow 1$, one has

$$E_2^{pq} \cong H^p(W, \mathbb{Q}) \otimes_{\mathbb{Q}} H^q(C_W, \mathbb{Q}) \cong \begin{cases} H^q(C_W, \mathbb{Q}) & p = 0 \\ 0 & p \neq 0 \end{cases}$$

because $H^p(W, \mathbb{Q}) = 0$ for $p \neq 0$ (see the proof of Proposition 3.2). The first isomorphism follows immediately. The second isomorphism follows from the fact $C_W \cong \mathbb{Z}^{c(W)}$. \square

Remark 3.7. For any right-angled Coxeter group W , Kishimoto [16, Theorem 5.3] determined the mod 2 cohomology ring of $\text{Ad}(Q_W)$.

4. CONSTRUCTION OF 2-COCYCLES

Throughout this section, we assume that the reader is familiar with group cohomology and Coxeter groups. The central extension

$$(4.1) \quad 1 \rightarrow C_W \rightarrow \text{Ad}(Q_W) \xrightarrow{\phi} W \rightarrow 1$$

corresponds to the cohomology class $u_\phi \in H^2(W, C_W)$ (see Brown [8, §IV.3] for precise). In this section, we will construct 2-cocycles representing u_ϕ . Before doing so, we claim $u_\phi \neq 0$. The claim is equivalent to the following lemma.

Lemma 4.1. *The central extension $1 \rightarrow C_W \rightarrow \text{Ad}(Q_W) \xrightarrow{\phi} W \rightarrow 1$ is nontrivial.*

Proof. If the central extension is trivial, then $\text{Ad}(Q_W) \cong C_W \times W$. But this is not the case because $\text{Ad}(Q_W)_{\text{Ab}} \cong \mathbb{Z}^{c(W)}$ by Proposition 2.4 while $(C_W \times W)_{\text{Ab}} \cong C_W \times W_{\text{Ab}} \cong \mathbb{Z}^{c(W)} \times (\mathbb{Z}/2)^{c(W)}$ by Proposition 2.2. \square

Now we invoke the celebrated Matsumoto's theorem:

Theorem 4.2 (Matsumoto [17]). *Let (W, S) be a Coxeter system, M a monoid and $f : S \rightarrow M$ a map such that $(f(s)f(t))_{m(s,t)} = (f(t)f(s))_{m(s,t)}$ for all $s, t \in S$ with $s \neq t$, $m(s,t) < \infty$. Then there exists a unique map $F : W \rightarrow M$ such that $F(w) = f(s_1) \cdots f(s_k)$ whenever $w = s_1 \cdots s_k$ ($s_i \in S$) is a reduced expression.*

The proof also can be found in [5, Chapitre IV, §1, Proposition 5] and [14, Theorem 1.2.2]. Define a map $f : S \rightarrow \text{Ad}(Q_W)$ by $s \mapsto e_s$, then f satisfies the assumption of Theorem 4.2 as in the proof of Proposition 3.3, and hence there exists a unique map $F : W \rightarrow \text{Ad}(Q_W)$ such that $F(w) = f(s_1) \cdots f(s_k) = e_{s_1} \cdots e_{s_k}$ whenever $w = s_1 \cdots s_k$ ($s_i \in S$) is a reduced expression. It is clear that $F : W \rightarrow \text{Ad}(Q_W)$ is a set-theoretical section of $\phi : \text{Ad}(Q_W) \rightarrow W$. Define $c : W \times W \rightarrow C_W$ by

$$c(w_1, w_2) = F(w_1)F(w_2)F(w_1w_2)^{-1}.$$

The standard argument in group cohomology (see Brown [8, §IV.3]) implies the following result:

Proposition 4.3. *c is a normalized 2-cocycle and $[c] = u_\phi \in H^2(W, C_W)$.*

Remark 4.4. In case W is the symmetric group of n letters \mathfrak{S}_n , Proposition 4.3 was stated in [2, Remark 3.3].

Now we deal with the case $c(W) = 1$ more precisely. If $c(W) = 1$ then the odd subgraph Γ_{odd} of W is connected and hence W must be irreducible. All finite irreducible Coxeter groups of type other than B_n ($n \geq 2$), F_4 , $I_2(m)$ (m even) satisfy $c(W) = 1$. Among affine irreducible Coxeter groups, those of type \tilde{A}_n ($n \geq 2$), \tilde{D}_n ($n \geq 4$), \tilde{E}_6 , \tilde{E}_7 and \tilde{E}_8 fulfill $c(W) = 1$. For simplifying the notation, we will identify C_W with \mathbb{Z} by $e_s^2 \mapsto 1$ and denote our central extension by

$$0 \rightarrow \mathbb{Z} \rightarrow \text{Ad}(Q_W) \xrightarrow{\phi} W \rightarrow 1.$$

Proposition 4.5. *If $c(W) = 1$ then $H^2(W, \mathbb{Z}) \cong \mathbb{Z}/2$.*

Proof. A short exact sequence $0 \rightarrow \mathbb{Z} \hookrightarrow \mathbb{C} \rightarrow \mathbb{C}^\times \rightarrow 1$ of abelian groups, where $\mathbb{C} \rightarrow \mathbb{C}^\times$ is defined by $z \mapsto \exp(2\pi\sqrt{-1}z)$, induces the exact sequence

$$H^1(W, \mathbb{C}) \rightarrow H^1(W, \mathbb{C}^\times) \xrightarrow{\delta} H^2(W, \mathbb{Z}) \rightarrow H^2(W, \mathbb{C})$$

(see Brown [8, Proposition III.6.1]). It is known that $H^k(W, \mathbb{C}) = 0$ for $k > 0$ (see the proof of Proposition 3.2), which implies that the connecting homomorphism $\delta : H^1(W, \mathbb{C}^\times) \rightarrow H^2(W, \mathbb{Z})$ is an isomorphism. We claim that $H^1(W, \mathbb{C}^\times) = \text{Hom}(W, \mathbb{C}^\times) \cong \mathbb{Z}/2$. Indeed, W is generated by S consisting of elements of order 2, and all elements of S are mutually conjugate by the assumption $c(W) = 1$. Thus $\text{Hom}(W, \mathbb{C}^\times)$ consists of the trivial homomorphism and the homomorphism ρ defined by $\rho(s) = -1$ ($s \in S$). \square

Combining Lemma 4.1 and Proposition 4.5, we obtain the following corollary.

Corollary 4.6. *If $c(W) = 1$ then $0 \rightarrow \mathbb{Z} \rightarrow \text{Ad}(Q_W) \xrightarrow{\phi} W \rightarrow 1$ is the unique non-trivial central extension of W by \mathbb{Z} .*

In general, given a homomorphism of groups $f : G \rightarrow \mathbb{C}^\times$, the cohomology class $\delta f \in H^2(G, \mathbb{Z})$, where $\delta : \text{Hom}(G, \mathbb{C}^\times) \rightarrow H^2(G, \mathbb{Z})$ is the connecting homomorphism as above, can be described as follows. For each $g \in G$, choose a branch of $\log(f(g))$. We argue such that $\log(f(1)) = 0$. Define $\tau_f : G \times G \rightarrow \mathbb{Z}$ by

$$(4.2) \quad \tau_f(g, h) = \frac{1}{2\pi\sqrt{-1}} \{ \log(f(g)) + \log(f(h)) - \log(f(gh)) \} \in \mathbb{Z}.$$

By a diagram chase, one can prove the following:

Proposition 4.7. *τ_f is a normalized 2-cocycle and $[\tau_f] = \delta f \in H^2(G, \mathbb{Z})$.*

Assuming $c(W) = 1$, let $\rho : W \rightarrow \mathbb{C}^\times$ be the homomorphism defined by $\rho(s) = -1$ ($s \in S$) as in the proof of Proposition 4.5. Note that $\rho(w) = (-1)^{\ell(w)}$ ($w \in W$) where $\ell(w)$ is the *length* of w (see Humphreys [15, §5.2]). For each $w \in W$, choose a branch of $\log(\rho(w))$ as

$$\log(\rho(w)) = \begin{cases} 0 & \text{if } \ell(w) \text{ is even,} \\ \pi\sqrt{-1} & \text{if } \ell(w) \text{ is odd.} \end{cases}$$

Applying (4.2), the corresponding map $\tau_\rho : W \times W \rightarrow \mathbb{Z}$ is given by

$$(4.3) \quad \tau_\rho(w_1, w_2) = \begin{cases} 1 & \text{if } \ell(w_1) \text{ and } \ell(w_2) \text{ are odd,} \\ 0 & \text{otherwise.} \end{cases}$$

Combining Lemma 4.1, Corollary 4.6, and Proposition 4.7, we obtain the following theorem:

Theorem 4.8. *If $c(W) = 1$ then $\tau_\rho : W \times W \rightarrow \mathbb{Z}$ defined by (4.3) is a normalized 2-cocycle and $[\tau_\rho] = u_\phi \in H^2(W, \mathbb{Z})$.*

5. COMMUTATOR SUBGROUPS OF ADJOINT GROUPS

As was stated in the introduction, Eisermann [12, Example 1.18] claimed that $\text{Ad}(Q_{\varepsilon_n})$ is isomorphic to the semidirect product $A_n \rtimes \mathbb{Z}$ where A_n is the alternating group on n letters. We will generalize his result to Coxeter quandles by showing the following theorem:

Theorem 5.1. $\phi : \text{Ad}(Q_W) \rightarrow W$ induces an isomorphism

$$\phi : [\text{Ad}(Q_W), \text{Ad}(Q_W)] \xrightarrow{\cong} [W, W].$$

Proof. Consider the following commutative diagram with exact rows and columns.

$$\begin{array}{ccccccc}
 & & 1 & & 1 & & \\
 & & \downarrow & & \downarrow & & \\
 & & C_W & \xrightarrow[\cong]{\text{Ab}_{\text{Ad}(Q_W)}} & \ker \phi_{\text{Ab}} & & \\
 & & \downarrow & & \downarrow & & \\
 1 & \longrightarrow & [\text{Ad}(Q_W), \text{Ad}(Q_W)] & \hookrightarrow & \text{Ad}(Q_W) & \xrightarrow{\text{Ab}_{\text{Ad}(Q_W)}} & \text{Ad}(Q_W)_{\text{Ab}} \longrightarrow 1 \\
 & & \downarrow \phi & & \downarrow \phi & & \downarrow \phi_{\text{Ab}} \\
 1 & \longrightarrow & [W, W] & \hookrightarrow & W & \xrightarrow{\text{Ab}_W} & W_{\text{Ab}} \longrightarrow 1 \\
 & & & & \downarrow & & \downarrow \\
 & & & & 1 & & 1
 \end{array}$$

From Proposition 2.2 and Proposition 2.4, we see that $\ker \phi_{\text{Ab}}$ is the free abelian group with a basis $\{[e_s^2] \mid s \in \mathcal{R}_W\}$, which implies that $\text{Ab}_{\text{Ad}(Q_W)} : C_W \rightarrow \ker \phi_{\text{Ab}}$ is an isomorphism because it assigns $[e_s^2]$ to e_s^2 ($s \in \mathcal{R}_W$). Since ϕ is surjective, it is obvious that $\phi([\text{Ad}(Q_W), \text{Ad}(Q_W)]) = [W, W]$. We will show that $\phi : [\text{Ad}(Q_W), \text{Ad}(Q_W)] \rightarrow [W, W]$ is injective. To do so, let $g \in [\text{Ad}(Q_W), \text{Ad}(Q_W)]$ be an element with $g \in \ker \phi = C_W$. Then $[g] := \text{Ab}_{\text{Ad}(Q_W)}(g) = 1$ by the exactness of the middle row. But it implies $g = 1$ because $\text{Ab}_{\text{Ad}(Q_W)} : C_W \rightarrow \ker \phi_{\text{Ab}}$ is an isomorphism. \square

Corollary 5.2. *There is a group extension of the form*

$$1 \rightarrow [W, W] \rightarrow \text{Ad}(Q_W) \xrightarrow{\text{Ab}_{\text{Ad}(Q_W)}} \mathbb{Z}^{c(W)} \rightarrow 1.$$

If $c(W) = 1$ then the extension splits and $\text{Ad}(Q_W) \cong [W, W] \rtimes \mathbb{Z}$.

Since $c(\mathfrak{S}_n) = 1$ and $[\mathfrak{S}_n, \mathfrak{S}_n] = A_n$, we recover the claim by Eisermann mentioned above.

Remark 5.3. By using Theorem 5.1, Kishimoto [16, Theorem 2.5] proved that the following commutative square which appeared in the proof of Theorem 5.1 is actually a pull-back.

$$\begin{array}{ccc}
 \text{Ad}(Q_W) & \xrightarrow{\text{Ab}_{\text{Ad}(Q_W)}} & \text{Ad}(Q_W)_{\text{Ab}} \\
 \downarrow \phi & & \downarrow \phi_{\text{Ab}} \\
 W & \xrightarrow{\text{Ab}_W} & W_{\text{Ab}}
 \end{array}$$

6. ROOT SYSTEMS

As was pointed out by Andruskiewitsch-Graña [3, §1.3.5], Brieskorn [6, p. 58, Example 9], and Fenn-Rourke [13, Example 10], a root system turns out to be a

rack. In this final section, we turn our attention to root systems. Let us recall the definition and properties of the root system associated with a Coxeter system (W, S) . See Björner-Brenti [4, Chapter 4] or Humphreys [15, Part II, Chapter 5] for precise. Let V be the real vector space with a basis $\{\alpha_s \mid s \in S\}$ in one to one correspondence with S . Define a symmetric bilinear form $B(-, -)$ on V by

$$B(\alpha_s, \alpha_t) := -\cos \frac{\pi}{m(s, t)}.$$

Here we understand $B(\alpha_s, \alpha_t) = -1$ in case $m(s, t) = \infty$. For each $s \in S$, define a reflection $\sigma_s \in GL(V)$ by

$$\sigma_s(\lambda) = \lambda - 2B(\alpha_s, \lambda)\alpha_s.$$

The *geometric representation* of W (or the canonical representation in the literature) is the unique homomorphism $\sigma : W \rightarrow GL(V)$ satisfying $s \mapsto \sigma_s$ ($s \in S$). The geometric representation is faithful and preserves the form $B(-, -)$. For simplicity, we may write $w(\lambda)$ in place of $\sigma(w)(\lambda)$ ($w \in W, \lambda \in V$). The *root system* Φ_W associated with (W, S) is defined by

$$\Phi_W := \{w(\alpha_s) \mid s \in S, w \in W\}.$$

For each root $\alpha \in \Phi_W$, define a reflection $t_\alpha \in GL(V)$ by $t_\alpha(\lambda) := \lambda - 2B(\alpha, \lambda)\alpha$. Note that if $\alpha = \alpha_s$ ($s \in S$) then $t_\alpha = \sigma_s$.

Lemma 6.1. *The root system Φ_W turns out to be a rack with the rack operation $\alpha * \beta = t_\beta(\alpha)$ ($\alpha, \beta \in \Phi_W$).*

Proof. It suffices to prove $(\alpha * \beta) * \gamma = (\alpha * \gamma) * (\beta * \gamma)$ for $\alpha, \beta, \gamma \in \Phi_W$. One has

$$(\alpha * \beta) * \gamma = t_\gamma(\alpha - 2B(\beta, \alpha)\beta) = t_\gamma(\alpha) - 2B(\beta, \alpha)t_\gamma(\beta)$$

while

$$(\alpha * \gamma) * (\beta * \gamma) = t_\gamma(\alpha) * t_\gamma(\beta) = t_\gamma(\alpha) - 2B(t_\gamma(\beta), t_\gamma(\alpha))t_\gamma(\beta),$$

and the assertion follows from $B(t_\gamma(\beta), t_\gamma(\alpha)) = B(\beta, \alpha)$. Note that Φ_W is not a quandle because $\alpha * \alpha = t_\alpha(\alpha) = -\alpha$. \square

Now let Q_W be the Coxeter quandle associated with a Coxeter system (W, S) as before, and define a map $p : \Phi_W \rightarrow Q_W$ by $\alpha \mapsto t_\alpha$. Here we identify elements of $\sigma(W)$ with those of W , which is possible since σ is injective. Note that t_α belongs to Q_W because $t_\alpha = wsw^{-1}$ holds for $\alpha = w(\alpha_s)$, and that the map p is surjective and two-to-one (see Humphreys [15, p. 116] or Björner-Brenti [4, p. 104]). Indeed, p maps $\pm\alpha \in \Phi_W$ to the same element $t_\alpha = t_{-\alpha}$.

Lemma 6.2. *The map $p : \Phi_W \rightarrow Q_W$ is a morphism of racks.*

Proof. In general, $wt_\alpha w^{-1} = t_{w(\alpha)}$ holds for $\alpha \in \Phi_W$ and $w \in W$ (see Björner-Brenti [4, p. 104]). Now for $\alpha, \beta \in \Phi_W$, set $\gamma := t_\beta(\alpha)$ and we have

$$p(\alpha * \beta) = p(\gamma) = t_\gamma = t_\beta t_\alpha t_\beta^{-1} = t_\beta^{-1} t_\alpha t_\beta = t_\alpha * t_\beta = p(\alpha) * p(\beta)$$

as desired. \square

We close this paper by proving the following result:

Theorem 6.3. *The morphism $p: \Phi_W \rightarrow Q_W$ of racks induces an isomorphism $\text{Ad}(\Phi_W) \cong \text{Ad}(Q_W)$.*

Proof. Recall that $\text{Ad}(\Phi_W)$ is defined by the presentation

$$\text{Ad}(\Phi_W) = \langle e_\alpha \ (\alpha \in \Phi_W) \mid e_{\alpha*\beta} = e_\beta^{-1} e_\alpha e_\beta \rangle.$$

For each $\alpha \in \Phi_W$, we have $e_{\alpha*\alpha} = e_{-\alpha}$ while $e_\alpha^{-1} e_\alpha e_\alpha = e_\alpha$, which proves that $e_{-\alpha} = e_\alpha$. On the other hand, as $p: \Phi_W \rightarrow Q_W$ is surjective, $\text{Ad}(Q_W)$ may be defined as

$$\text{Ad}(Q_W) = \langle e_{p(\alpha)} \ (\alpha \in \Phi_W) \mid e_{p(\alpha)*p(\beta)} = e_{p(\beta)}^{-1} e_{p(\alpha)} e_{p(\beta)} \rangle.$$

Since $p^{-1}(\alpha) = \{\pm\alpha\}$ and $p(\alpha*\beta) = p(\alpha)*p(\beta)$ by Lemma 6.2, the assignment $e_\alpha \mapsto e_{p(\alpha)}$ ($\alpha \in \Phi_W$) induces an isomorphism $\text{Ad}(\Phi_W) \rightarrow \text{Ad}(Q_W)$. \square

Remark 6.4. For a (crystallographic) root system Φ and its Weyl group $W(\Phi)$, one can prove that $\text{Ad}(\Phi) \cong \text{Ad}(Q_{W(\Phi)})$ in a similar way.

Acknowledgement. The author thanks Daisuke Kishimoto and Takefumi Nosaka for valuable comments and discussions with the author. He also thanks Ye Liu for informing the paper Digne-Gomi [11] and careful reading of drafts of this paper. This work was partially supported by JSPS KAKENHI Grant Number 26400077, and by the Research Institute for Mathematical Sciences, a Joint Usage/Research Center located in Kyoto University.

REFERENCES

- [1] Toshiyuki Akita, *Euler characteristics of Coxeter groups, PL-triangulations of closed manifolds, and cohomology of subgroups of Artin groups*, J. London Math. Soc. (2) **61** (2000), no. 3, 721–736, DOI 10.1112/S0024610700008693. MR1766100 (2001f:20080)
- [2] N. Andruskiewitsch, F. Fantino, G. A. García, and L. Vendramin, *On Nichols algebras associated to simple racks*, Groups, algebras and applications, Contemp. Math., vol. 537, Amer. Math. Soc., Providence, RI, 2011, pp. 31–56, DOI 10.1090/conm/537/10565. MR2799090 (2012g:16065)
- [3] Nicolás Andruskiewitsch and Matías Graña, *From racks to pointed Hopf algebras*, Adv. Math. **178** (2003), no. 2, 177–243, DOI 10.1016/S0001-8708(02)00071-3. MR1994219
- [4] Anders Björner and Francesco Brenti, *Combinatorics of Coxeter groups*, Graduate Texts in Mathematics, vol. 231, Springer, New York, 2005. MR2133266
- [5] Nicolas Bourbaki, *Éléments de mathématique*, Masson, Paris, 1981 (French). Groupes et algèbres de Lie. Chapitres 4, 5 et 6. [Lie groups and Lie algebras. Chapters 4, 5 and 6]. MR647314 (83g:17001)
- [6] E. Brieskorn, *Automorphic sets and braids and singularities*, Braids (Santa Cruz, CA, 1986), Contemp. Math., vol. 78, Amer. Math. Soc., Providence, RI, 1988, pp. 45–115. MR975077
- [7] Egbert Brieskorn and Kyoji Saito, *Artin-Gruppen und Coxeter-Gruppen*, Invent. Math. **17** (1972), 245–271, DOI 10.1007/BF01406235 (German). MR0323910
- [8] Kenneth S. Brown, *Cohomology of groups*, Graduate Texts in Mathematics, vol. 87, Springer-Verlag, New York, 1982. MR672956 (83k:20002)
- [9] F.J.-B.J. Clauwens, *The adjoint group of an Alexander quandle* (2011), available at <http://arxiv.org/abs/1011.1587>.
- [10] Michael W. Davis, *The geometry and topology of Coxeter groups*, London Mathematical Society Monographs Series, vol. 32, Princeton University Press, Princeton, NJ, 2008. MR2360474 (2008k:20091)

- [11] F. Digne and Y. Gomi, *Presentation of pure braid groups*, J. Knot Theory Ramifications **10** (2001), no. 4, 609–623, DOI 10.1142/S0218216501001037. MR1831679
- [12] Michael Eisermann, *Quandle coverings and their Galois correspondence*, Fund. Math. **225** (2014), no. 1, 103–168, DOI 10.4064/fm225-1-7. MR3205568
- [13] Roger Fenn and Colin Rourke, *Racks and links in codimension two*, J. Knot Theory Ramifications **1** (1992), no. 4, 343–406. MR1194995
- [14] Meinolf Geck and Götz Pfeiffer, *Characters of finite Coxeter groups and Iwahori-Hecke algebras*, London Mathematical Society Monographs. New Series, vol. 21, The Clarendon Press, Oxford University Press, New York, 2000. MR1778802
- [15] James E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics, vol. 29, Cambridge University Press, Cambridge, 1990. MR1066460 (92h:20002)
- [16] Daisuke Kishimoto, *Right-angled Coxeter quandles and polyhedral products* (2017), available at <http://arxiv.org/abs/1706.06209>.
- [17] Hideya Matsumoto, *Générateurs et relations des groupes de Weyl généralisés*, C. R. Acad. Sci. Paris **258** (1964), 3419–3422 (French). MR0183818
- [18] Takefumi Nosaka, *Central extensions of groups and adjoint groups of quandles*, Geometry and analysis of discrete groups and hyperbolic spaces, RIMS Kôkyûroku Bessatsu, B66, Res. Inst. Math. Sci. (RIMS), Kyoto, 2017, pp. 167–184.
- [19] ———, *Quandles and topological pairs*, SpringerBriefs in Mathematics, Springer, Singapore, 2017. Symmetry, knots, and cohomology. MR3729413
- [20] Luis Paris, *Braid groups and Artin groups*, Handbook of Teichmüller theory. Vol. II, IRMA Lect. Math. Theor. Phys., vol. 13, Eur. Math. Soc., Zürich, 2009, pp. 389–451, DOI 10.4171/055-1/12. MR2497781
- [21] ———, *$K(\pi, 1)$ conjecture for Artin groups*, Ann. Fac. Sci. Toulouse Math. (6) **23** (2014), no. 2, 361–415, DOI 10.5802/afst.1411 (English, with English and French summaries). MR3205598
- [22] ———, *Lectures on Artin groups and the $K(\pi, 1)$ conjecture*, Groups of exceptional type, Coxeter groups and related geometries, Springer Proc. Math. Stat., vol. 82, Springer, New Delhi, 2014, pp. 239–257, DOI 10.1007/978-81-322-1814-2_13. MR3207280

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, HOKKAIDO UNIVERSITY, SAPPORO, 060-0810 JAPAN

E-mail address: akita@math.sci.hokudai.ac.jp