Title	Paradoxapseudes shimojiensis sp. nov. (Crustacea: Tanaidacea: Apseudidae) from a submarine limestone cave in Japan, with notes on its chelipedal morphology and sexual system
Author(s)	Kakui, Keiichi; Fujita, Yoshihisa
Citation	Marine Biology Research, 16(3), 195-207 https://doi.org/10.1080/17451000.2020.1720249
Issue Date	2020-03-20
Doc URL	http://hdl.handle.net/2115/80409
Rights	This is an Accepted Manuscript of an article published by Taylor & Francis in Marine Biology Research on 12/02/2020, available online: https://www.tandfonline.com/doi/full/10.1080/17451000.2020.1720249 .
Type	article (author version)
File Information	MAR BIOL RES16_195-207.pdf

[Instructions for use](#)

This is an Accepted Manuscript of an article published by Taylor & Francis
in *Marine Biology Research* on 12 Feb. 2020, available online:
<http://www.tandfonline.com/10.1080/17451000.2020.1720249>.

***Paradoxapseudes shimojiensis* sp. nov. (Crustacea: Tanaidacea: Apseudidae) from a submarine limestone cave in Japan, with notes on its chelipedal morphology and sexual system**

Keiichi Kakui^a and Yoshihisa Fujita^b

^a*Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan;* ^b*Okinawa Prefectural University of Arts, Naha 903-8602, Okinawa, Japan*

Corresponding author. Keiichi Kakui: kakui@eis.hokudai.ac.jp; Faculty of Science, Hokkaido University, N10 W8, Kita-ku, Sapporo 060-0810, Japan

Email address for 2nd author: fujitayo@okigei.ac.jp

Acknowledgments

We thank Masaru Mizuyama, Katrine Worsaae, Peter Rask Møller, Yuji Ise, Hiroki Ichi, Hirotaka Ichi, and Go Tomitani for assistance in scuba sampling in the submarine cave; and Matthew H. Dick for reviewing the manuscript and editing our English.

Funding

This study was supported by the Japan Society for the Promotion of Science (JSPS) under KAKENHI grants JP16K18597 and JP19K06800 to KK, and JP16K07490 and JP16H06309 to YF.

ABSTRACT

We describe *Paradoxapseudes shimojiensis* sp. nov. from a submarine limestone cave at Shimoji-jima Island, Ryukyu Islands, southwestern Japan. This species resembles five species (*P. basibidens*, *P. bassoprofundo*, *P. bermudeus*, *P. edgari*, and *P. heroeae*, among 17 congeners) that bear a long pleotelson, but differs from them in having i) a naked antennal article 1, ii) the maxillipedal basis with one inner distal plumose seta, iii) the chelipedal basis with one dorsodistal and one ventro-subproximal simple setae, iv) the pereopod-1 basis with four dorsal simple setae longer than the width of the pereopod-1 basis, but without ventrodistal spiniform setae, v) the pereopod-1 merus with three mid-inner ventral simple setae, and vi) the pleopodal protopod with two inner plumose setae. We determined partial nucleotide sequences for the cytochrome *c* oxidase subunit I (COI) and 18S rRNA (18S) genes in *P. shimojiensis* for future use in DNA barcoding and phylogeny. *Paradoxapseudes shimojiensis* has serial ridges on the inner surfaces of the left and right chelipedal bases that quite resemble the stridulatory organs in harvestmen (Opiliones); by analogy, we speculate that these ridges may be stridulatory sound-producing organs. Two specimens had both a fully-developed marsupium and genital cone, suggesting that *P. shimojiensis* is simultaneously hermaphroditic.

KEYWORDS: Malacostraca; Peracarida; stygophilic; hermaphrodite; acoustic; benthos

Introduction

Since the pioneering work by Băcescu (1980), 17 species of tanaidaceans in 15 genera have been reported from submarine caves (Kakui and Fujita 2018; Kakui et al. 2019), although most of these species are not restricted to caves (García-Herrero et al. 2019). For species that have been reported only from caves, it remains unclear whether they are obligate cave dwellers, both because they lack obvious external adaptations to cave life and because there are insufficient data from the benthic areas outside the caves (Kakui and Fujita 2018). As submarine caves are difficult to access and investigate, and minute animals like tanaidaceans can easily be overlooked, research on submarine-cave tanaidaceans, including their taxonomy, is still in its infancy.

The apseudid genus *Paradoxapseudes* Guțu, 1991, consisting of 17 species (Tzeng and Hsueh 2014; Bird 2015), is the only tanaidacean genus with multiple submarine-cave dwellers, including *P. bermudeus* (Băcescu, 1980), *P. intermedius* (Hansen, 1895), and *P. mortoni* (Bamber, 1997) (Guțu and Iliffe 1985; Bamber 2000; Gerovasileiou et al. 2016). Only one congener, *P. littoralis* (Shiino, 1952), has been reported from Japanese waters, but not as a cave dweller (Shiino 1952). In 2018, our faunal survey of a submarine limestone cave at Shimoji-jima Island, Miyako Island Group, southwestern Japan, yielded one *Paradoxapseudes* species, which turned out to be undescribed. Here we describe this species as new, provide partial sequences for the cytochrome *c* oxidase subunit I (COI) and 18S rRNA (18S) genes, and discuss the possible function of unusual serial ridges on the chelipeds, as well as indications that the species is simultaneously hermaphroditic.

Materials and methods

All tanaidaceans were collected by SCUBA diving on 26 October 2018 in the

submarine limestone cave Akuma-no-Yakata (“Devil’s Palace” in Japanese), located on a reef slope at Shimoji-jima Island, Miyako Island Group, southwestern Japan (24°49'22.51"N, 125°08'07.84"E), with the entrance at a depth of about 35 m; see Osawa and Fujita (2019) for detailed information on the cave. Mud deposited in holes or fissures in the cave wall was collected with a commercially made aquatic suction sampler (yabby pump), from the “middle zone” of the cave (50–80 m from the entrance, ca. 20 m depth, no light, ca. 28‰ salinity; rocky or rubble-covered substrate) (Osawa and Fujita, 2019). Tanaidaceans were sorted from the mud sample and fixed and preserved in 99% ethanol. The specimens studied were deposited in the University Museum Fужukan, University of the Ryukyus (RUMF), Okinawa, Japan, and in the Invertebrate Collection of the Hokkaido University Museum (ICHUM), Sapporo. The methods used for dissection, preparation of slides, light microscopy, scanning electron microscopy (SEM), and drawing were as described by Kakui and Angsupanich (2012).

Orientation and terminology here follow Larsen (2003), except that the term “plumose sensory seta” (PSS; Bird 2011) is used instead of “broom seta” and the term “protopod” is used instead of “basal article” for pleopods and uropods. Body length (BL) was measured from the base of the antennules to the tip of the pleotelson, and body width at the widest portion of the cephalothorax (CW, cephalothorax width). Pleotelson length (PL) was measured between the anterior and posterior edges of the pleotelson, and pleotelson width (PW) at the widest portion of the segment (in many specimens, the width across the lateral processes on the pleotelson). Appendages were measured in the holotype. Measurements were made axially from digital images by using ImageJ (Rasband 2019): dorsally on the body, antennules, antennae, and uropods; laterally on the chelipeds, pereopods, and pleopods. The length and width of congeners

were measured from original illustrations if these measurements were not provided in descriptions.

Total DNA was extracted from the left cheliped of the holotype by using a NucleoSpin Tissue XS Kit (TaKaRa Bio, Japan). Table I lists the primers used for DNA amplification and sequencing. PCR amplification conditions for COI with TaKaRa Ex Taq DNA polymerase (TaKaRa Bio) were 94°C for 1 min; 35 cycles of 98°C for 10 s, 62°C for 30 s, and 72°C for 50 s; and 72°C for 2 min. Conditions for 18S amplification with KOD FX Neo polymerase (Toyobo, Japan) were 94°C for 2 min; 45 cycles of 98°C for 10 s, 65°C for 30 s, and 68°C for 90 s; and 68°C for 3 min. Nucleotide sequences were determined by direct sequencing with a BigDye Terminator Kit ver. 3.1 and a 3730 Genetic Analyzer (Life Technologies, California, USA), and segments were concatenated by using MEGA7 (Kumar et al. 2016). All sequences we determined were deposited in the International Nucleotide Sequence Database (INSD) through the DNA Data Bank of Japan (DDBJ).

The determined COI sequence (658 nt, encoding 219 amino acids) was aligned with that of *P. intermedius* obtained from the INSD (accession number HM016216; Drumm 2010; originally identified as *Apseudes* cf. *bermudeus*) by using CLUSTAL W (Thompson et al. 1994); the aligned sequences were trimmed in MEGA7 to the shortest length between them (402 nt). Our 18S sequence (2176 nt) was aligned with those of *P. bermudeus* and *P. littoralis* obtained from INSD (accession numbers GQ175865 and AB618175; Wilson 2009; Kakui et al. 2011) by using MAFFT ver. 7 (Katoh and Standley 2013) with the Q-INS-i strategy (Katoh and Toh 2008); the aligned sequences were trimmed to the shortest length among them (2036 nt). Kimura (1980) 2-parameter (K2P) distances among the aligned sequences in each dataset were calculated with MEGA7.

126

127 **Systematics**128 **Family Apseudidae Leach, 1814**129 **Genus *Paradoxapseudes* Guțu, 1991**

130 *Paradoxapseudes* Guțu, 1991: 349. Type species: *Paradoxapseudes cubensis* Guțu,
 131 1991.

132

133 ***Amended diagnosis modified after Guțu (2008)***

134 Apseudinae with small sized, dorsoventrally flattened body. Carapace longer than wide;
 135 lateral margins without spiniform processes; posterior lateral expansions slight or absent
 136 (= branchial regions not evident in dorsal view); rostrum prominent, acuted anteriorly;
 137 ocular lobes well defined, generally with pigmented visual elements; epistome single,
 138 thin, spiniform. Pereonites 3–6 similar and approximately equal; anterior half with at
 139 most one pair of lateral processes. Pleon with five short pleonites having some lateral
 140 circumplumose setae. Antennular article 1 with or without inner proximal fine
 141 denticulation; inner flagellum with two or three articles (accidentally, with only one).
 142 Antennal articles 4 and 5 long and thin; squama with 3–6 setae. Mandibles with
 143 triarticulate palp; article 1 with 3–6 simple setae. Labium palp ovate, slightly narrower
 144 distally, with three distal slender spiniform setae. Maxillule with biarticulate palp.
 145 Maxilliped generally with inner distal long plumose seta on basis and palp-article 1;
 146 palp-article 2 with spiniform or simple seta on outer distal corner. Epignath cup-shaped;
 147 terminal seta well developed, slender, setulate in distal half. Cheliped with exopod.
 148 Pereopod 1 fossorial, with exopod; coxa with small anterior rounded process bearing 2–
 149 5 setae; basis with 3–5 dorsal plumose setae (apparently simple in many cases);

propodus with 3–4 ventral spiniform setae. Propodus of pereopods 5 and 6 with inner ventral row of blade-like spiniform setae. Pleopods biramous, in five pairs. Uropodal exopod with five articles; endopod with at most 24 articles. Oostegites in five pairs.

Remarks

Since Guțu (2008) provided the latest diagnosis of *Paradoxapseudes*, five species were added to the genus (Bird 2015), requiring some minor amendments to the diagnosis (e.g., the condition/number of setae on the pereopod-1 coxa and the maxillipedal basis), which are reflected in the above diagnosis. And, as Guțu and Iliffe (1989) suggested the similarity between *Paradoxapseudes bermudeus* and *Falsapseudes bowmani* (Guțu and Iliffe, 1989), we added two characters to distinguish them: “epistome single, thin, spiniform” and “anterior half [of pereonites 3–6] with at most one pair of lateral processes”. The characters “lateral margins [of carapace] without spiniform processes; posterior lateral expansions [on carapace] slight or absent (= branchial regions not evident in dorsal view)” were also added.

***Paradoxapseudes shimojiensis* sp. nov.**

[New Japanese name: *Shimoji-apuseudesu*]

(Figures 1–5)

Diagnosis

Pleotelson long (PL/PW 2.57–2.61), with two pairs of slight lateral processes. Antennal article 1 naked. Maxillipedal basis with inner distal plumose seta. Chelipedal basis with inner serial ridges and one dorsodistal and one ventro-subproximal simple setae.

Chelipedal merus with ventrodistal spiniform seta. Pereopod-1 basis with four dorsal

simple setae, each longer than width of pereopod-1 basis; ventrodistal region without spiniform setae. Pereopod-1 merus with three mid-inner ventral simple setae. Pleopodal protopod with two inner plumose setae.

Etymology

The specific name is an adjective referring to the type locality.

Material examined

Holotype: ovigerous simultaneous hermaphrodite, RUMF-ZC-6081 (BL 2.27 mm, CW 0.35 mm), dissected, 8 slides and 1 vial; INSD accession number LC510354 (COI) and LC510355 (18S); Akuma-no-Yakata (Devil's Palace), Shimoji-jima Island, Miyako Island Group, southwestern Japan, northwestern Pacific Ocean (24°49'22.51"N, 125°08'07.84"E), 20 m depth, mud, 26.x.2018, collected by Y. Fujita.

Paratype: simultaneous hermaphrodite with empty, fully-developed marsupium, ICHUM6029 (CW 0.34 mm), dissected, 4 slides, 1 SEM stub, and 1 vial; same collection information as for holotype.

Description of simultaneous hermaphrodite, based primarily on holotype, with observation of epignath from paratype

Body (Figures 1, 2A, a1, B) dorsoventrally flattened, 6.73 times as long as CW, translucent, whitish when alive (Figure 1A, B); body wall not heavily calcified. Cephalothorax 0.17 times as long as BL, 1.16 times as long as wide, with pair of mid-lateral simple setae; rostrum hastate; eyes white when alive (Figure 1A, B); large, acute keel (hyposphenium) present posterior to maxillipeds (Figure 1B). Pereonites 1–6 with length ratio 1.00:1.14:1.63:1.85:1.77:1.46; pereonites 1 and 2 wider than long; pereonites 3–6 almost as long as wide; all pereonites with 1–3 pairs of simple setae;

pereonites 3–6 with pair of slight lateral processes in anterior half; pereonite 6 with genital cone (Figures 1C, 2B). Pleon 0.31 times as long as BL. Pleonites narrower than pereonite 6; all wider than long, similar in shape, with triangular epimeron bearing several plumose setae. Pleotelson 2.57 times as long as wide, slightly narrower than pleonite 5, with two pairs of slight lateral processes and several pairs of simple setae.

Antennule (Figure 2C, D, d1) with seven-articulate outer flagellum and triarticulate inner flagellum. Combined length of peduncle and outer flagellum 2.06 times as long as cephalothorax. Peduncular articles 1–4 with length ratio of 1.00:0.30:0.27:0.11; article 1 with one mid-outer, one outer distal, and four inner simple setae, eight outer PSS, and numerous inner fine setae; article 2 with three outer distal, one inner subproximal, and three inner distal simple setae, and four distal PSS; article 3 with one outer distal and three inner distal simple setae; article 4 with two PSS at insertion of inner flagellum. Outer flagellum 0.63 times as long as peduncle; articles 1 and 3 naked; articles 2, 4, 5, 6 with three, three, one, two simple setae, respectively; articles 4 and 6 each with aesthetasc; article 7 with two middle and four distal simple setae, and distal PSS. Inner flagellum 0.46 times as long as outer flagellum; articles 1 and 3 with two and three simple setae, respectively; articles 1–3 each with distal PSS.

Antenna (Figure 2E) with ten articles, 0.58 times as long as antennule; articles 1–10 with length ratio 1.00:2.35:0.52:1.74:1.62:1.01:0.96:1.21:0.69:0.85. Article 1 with inner distal triangular process, naked. Article 2 with one outer subproximal, one outer distal, and one mid-inner simple setae, and two inner distal spines. Article 3 with inner distal simple seta and two (one small and one large) inner distal spines. Article 4 with two inner PSS. Article 5 with two distal simple setae and six distal PSS. Article 6 with outer distal simple seta. Articles 7–9 each with one outer distal and one inner distal simple

225 setae; article 8 with distal PSS. Article 10 with five distal simple setae. Squama 0.13
 226 times as long as antenna, narrow, with five simple setae.

227 Epistome thin, spiniform (Figures 1B, 2B). Labrum (Figure 2F) bipartite, with
 228 lateral fine setae and distal serrations. Mandibles (Figure 2G–J) with serrations on body
 229 and well-developed molar process. Left mandibular incisor (Figure 2G) with four teeth;
 230 setiferous lobe with several complex setae; lacinia mobilis with four teeth. Right
 231 mandibular incisor (Figure 2H) with five teeth; setiferous lobe with trifurcate seta and
 232 several complex setae. Palp (Figure 2J, j1, j2) with three articles; articles 1–3 with
 233 length ratio 1.00:2.61:1.43; article 1 with four ventral simple setae; article 2 with mid-
 234 ventral simple seta and seven ventral blade-like spiniform setae (Figure 2j1); article 3
 235 with three simple setae, four serrate setae, and pectinate seta (Figure 2j2) in distal
 236 region. Labium (Figure 2K) with lobe bearing inner and outer setation. Palp setulate,
 237 with three distal spiniform setae. Maxillule (Figure 3A–C, c1) with biarticulate palp
 238 bearing four biserrate-tipped setae (Figure 3c1) at end. Outer endite (Figure 3A) with
 239 twelve distal spiniform setae, two subdistal setae, and outer and inner setation. Inner
 240 endite (Figure 3B) with distal simple seta, two distal plumose setae, distal trifurcate seta,
 241 outer clumps of fine setae, and outer process. Maxilla (Figure 3D, d1, d2) with outer
 242 and inner serrations on body. Inner lobe of fixed endite with serrate spiniform seta and
 243 17 basally swollen setae. Outer lobe of fixed endite (Figure 3D, d1, 2) with six simple
 244 setae (one broken), five upswept spiniform setae (one pinnate), three trifurcate
 245 spiniform setae, branched seta (Figure 3d2), and serrate spiniform seta; outer margin
 246 serrate. Inner lobe of movable endite with five setae (one pinnate). Outer lobe of
 247 movable endite with eight simple setae and two basally setulate setae. Maxilliped
 248 (Figure 3E, F, f1) with naked coxa. Basis with (right) or without (left) two outer distal
 249 spiniform processes; inner distal region with plumose seta. Endite with outer region

serrate and setulate; inner region with two (right) or one (left) coupling hooks, five basally swollen setae (two plumose); distal region with ventral simple seta, ventral pinnate spiniform seta, and eight complex spiniform setae. Palp article 1 with outer distal simple seta and inner distal plumose seta; article 2 with one outer distal, one inner ventrodistal, and 17 inner simple setae, and two inner ventral plumose setae; article 3 with nine inner simple setae; article 4 with one outer ventrodistal and two outer distal simple setae, and four inner distal spiniform setae. Epignath (Figure 3G) cup-shaped, with setulate terminal seta.

Cheliped (Figures 4A, B, b1, 5) with triarticulate exopod bearing four plumose setae at tip. Basis 2.06 times as long as wide, with one dorsodistal, one ventro-subproximal, one mid-ventral, and two ventrodistal simple setae, and inner serial ridges (Figure 5). Merus with one dorsodistal, one outer distal, and two ventro-subdistal simple setae, and ventrodistal spiniform seta. Carpus 1.37 times as long as basis, 2.50 times as long as wide, with six dorsal, two outer ventral, and six ventral simple setae. Propodus with three outer ventral, five ventral, and two inner ventral simple setae, and triangular claw; dorsal region of palm with three simple setae; three inner simple setae and inner short plumose seta at insertion of dactylus; cutting surface with three proximal and six outer distal simple setae, proximal small triangular process, subproximal large triangular process bearing three denticles, and several tiny spiniform setae. Dactylus (Figure 4b1) longer than fixed finger, with three inner subdistal simple setae, four ventral small triangular processes, and several ventral spiniform setae; unguis pointed.

Pereopods 1–6 cylindrical, with length ratio 1.00:0.93:0.90:0.83:0.91:1.00. Pereopod 1 (Figure 4C, c1) 0.35 times as long as BL, with length ratio of basis, ischium, merus, carpus, propodus, and dactylus–unguis 1.00:0.11:0.45:0.47:0.35:0.38. Coxa with anterior process bearing three simple setae; inner region with several plumose setae.

275 Basis cylindrical, narrow, 4.57 times as long as wide, with four dorsal simple setae each
 276 longer than width of pereopod-1 basis, one ventro-subproximal, one ventro-subdistal,
 277 and three ventrodistal simple setae, and two subproximal PSS. Ischium with three
 278 ventrodistal simple setae. Merus with three dorsodistal, three mid-inner ventral, two
 279 ventral, and two ventrodistal simple setae, and one dorsodistal and one ventrodistal
 280 spiniform setae. Carpus with seven dorsal, three long ventral, and three short ventral
 281 simple setae, and one dorsodistal and two ventral spiniform setae. Propodus with two
 282 dorsal, one mid-outer, and four ventral simple setae, two dorsal and three ventral
 283 spiniform setae, mid-dorsal PSS, and ventrodistal pectinate spiniform seta (Figure 4c1).
 284 Dactylus with one dorsal and two ventrodistal simple setae, and ventral serration;
 285 unguis half of dactylus length, naked. Exopod triarticulate; article 3 with five plumose
 286 setae. Pereopod 2 (Figure 4D) with length ratio of articles from basis to dactylus–unguis
 287 1.00:0.14:0.25:0.43:0.46:0.56. Coxa with simple seta. Basis cylindrical, narrow, 4.69
 288 times as long as wide, with one inner subproximal, one ventro-subproximal, one ventro-
 289 subdistal, and three ventrodistal simple setae, and three dorsal PSS. Ischium with two
 290 ventrodistal simple setae. Merus with two dorsodistal, one outer distal, and one ventral
 291 simple setae, and ventrodistal spiniform seta. Carpus with four dorsodistal and two
 292 ventral simple setae, and one mid-outer ventral, one outer distal, and one ventrodistal
 293 spiniform setae. Propodus with two dorsal and two ventral (one tiny) simple setae, two
 294 dorsal, one mid-outer, and two ventral spiniform setae, and mid-dorsal PSS. Dactylus
 295 with one dorsal and two ventrodistal simple setae; unguis slightly shorter than dactylus,
 296 naked. Pereopod 3 (Figure 4E, e1) with length ratio of articles from basis to dactylus–
 297 unguis 1.00:0.15:0.20:0.36:0.43:0.52. Similar to pereopod 2, except: basis without
 298 ventro-subproximal simple seta; basis with two ventrodistal simple setae; merus with
 299 dorsodistal simple seta; carpus with three dorsodistal simple setae; carpus without mid-

300 ventral simple seta; carpus with two outer distal spiniform setae; propodus with three
 301 dorsal spiniform setae; propodus without mid-outer spiniform seta. Pereopod 4 (Figure
 302 4F, f1) with length ratio of articles from basis to dactylus–unguis
 303 1.00:0.12:0.26:0.40:0.40:0.42. Coxa with simple seta. Basis cylindrical, narrow, 3.52
 304 times as long as wide, with one outer-subproximal, one inner-subproximal, and two
 305 ventrodistal simple setae, and three dorsal and one ventral PSS. Ischium with two
 306 ventrodistal simple setae. Merus with one dorsodistal and two ventrodistal simple setae,
 307 and two ventrodistal spiniform setae. Carpus with two outer distal and two inner distal
 308 simple setae, and two ventral, two outer distal, and two inner distal spiniform setae.
 309 Propodus with dorsodistal long simple seta, mid-dorsal PSS, dorso-subdistal row of five
 310 pectinate setae, and dorsodistal row of six pectinate setae. Dactylus with one dorsal and
 311 one ventrodistal simple setae; unguis as long as dactylus, naked. Pereopod 5 (Figure 4G,
 312 g1) with length ratio of articles from basis to dactylus–unguis
 313 1.00:0.12:0.32:0.44:0.41:0.55. Coxa, basis, ischium, merus, and dactylus–unguis similar
 314 to those of pereopod 4, except: basis without inner subproximal simple seta; basis with
 315 mid-inner ventral simple seta; ischium with three ventrodistal simple setae. Carpus with
 316 three dorsodistal and one ventro-subdistal simple setae, and three ventral spiniform
 317 setae. Propodus with two dorsal simple setae, one dorsodistal, one outer subdistal, and
 318 two ventral spiniform setae, mid-dorsal PSS, and inner ventral row of nine blade-like
 319 spiniform setae. Pereopod 6 (Figure 4H, h1–4) with length ratio of articles from basis to
 320 dactylus–unguis 1.00:0.12:0.42:0.56:0.48:0.57. Coxa with simple seta and plumose seta.
 321 Basis with one ventro-subproximal, one ventro-subdistal, and two ventrodistal simple
 322 setae, and four dorsal plumose setae; proximal region with four PSS. Ischium with three
 323 ventrodistal simple setae. Merus with two ventrodistal simple setae and three dorsal
 324 plumose setae. Carpus with two outer distal and two ventral simple setae, ventrodistal

spiniiform seta, and three dorsal plumose setae. Propodus with mid-dorsal simple seta, two outer subdistal, one ventro-subproximal, and one inner subdistal spiniiform setae, mid-dorsal PSS, two dorsodistal serrate setae (Figure 4h1), one outer subdistal and four inner dorsodistal pectinate setae (Figure 4h2), and inner ventral row of eleven blade-like spiniiform setae (Figure 4h3). Dactylus–unguis similar to pereopod 5.

Pleopods (Figure 4I, i1) five pairs, all similar. Protopod biarticulate, narrow, with two inner plumose setae. Exopod uniarticulate, 1.27 times as long as protopod, with eight plumose setae. Endopod uniarticulate, 1.23 times as long as exopod, with six plumose setae and mid-inner distally-hooked plumose seta (Figure 4i1).

Uropod (Figures 1B, 4J) with protopod bearing one inner distal and four outer simple setae. Exopod with article 1 bearing simple seta; other articles lost before fixation. Endopod nearly twice as long as pleotelson (Figure 1B); article 1 naked; other articles lost before fixation.

Variation and stability

In addition to the holotype (RUMF-ZC-6081), one paratype (ICHUM6029) was dissected, and its antennule, antenna, maxilliped, cheliped, pereopods 1–6, and pleopod 1 were observed (as with the holotype, most uropodal articles were lost before fixation). The numbers of setae, aesthetascs, and processes on these appendages were identical between the two specimens, with the following exceptions (ranges in parentheses; the state in the holotype in bold). Antennule: inner distal simple setae (**3**–4) and distal PSS (**4**–5) on article 2. Antenna: outer simple setae on article 6 (**1**–2); PSS on article 8 (0–**1**). Maxilliped: outer distal spiniiform processes on the basis (**0** or **2**); distal complex spiniiform setae on the endite (**8**–9); inner simple setae on palp articles 2 and 3 (**17**–**18**, **8**–**9**, respectively). Cheliped: ventral simple setae on the carpus (**5**–**6**); proximal simple

setae on the cutting surface of the fixed finger (3–4); dorsal simple setae on the palm (3–4). Pereopod 1: dorsal simple setae on the carpus (6–7); inner ventral simple setae proximal to the most proximal spiniform seta on the carpus (1–2). Pereopod 2: ventrodistal simple setae on the basis (3–4); ventral simple setae on the merus (1–2); dorsodistal simple setae (4–5) and mid-outer ventral spiniform setae (0–1) on the carpus. Pereopod 3: ventroproximal PSS on the basis (0–1); ventral simple setae on the carpus (0–1). Pereopod 4: ventro-subdistal simple setae on the basis (0–1); ventral spiniform setae on the propodus (0–1). Pereopod 5: outer distal simple setae (0–1) and ventro-subdistal simple setae (0–1) on the carpus. Pereopod 6: dorsal plumose setae on the basis (4–5); dorsal plumose setae (2–3) and inner distal simple setae (0–1) on the carpus; inner ventral blade-like spiniform setae (11 or 13) and inner dorsodistal pectinate setae (4–5) on the propodus.

Inner serial ridges on the chelipedal bases, a fully-developed marsupium, and a genital cone were observed on both specimens. The PL/PW ratio of the holotype and paratype were 2.57 and 2.61, respectively.

Genetic information

The partial COI sequence (658 nt, encoding 219 amino acids) and the nearly complete 18S sequence (2176 nt) were determined from the holotype specimen; their INSD accession numbers are LC510354 and LC510355, respectively. For COI (402 nt in the aligned dataset), the K2P distance between *Paradoxapseudes shimojiensis* and *P. intermedius* was 34.3%. For 18S (2036 nt in the aligned dataset), K2P distances between *P. shimojiensis* and two congeners (*P. bermudeus* and *P. littoralis*) were 5.7% and 5.5%, respectively.

372 **Remarks**

373 *Paradoxapseudes shimojiensis* sp. nov. is the 18th species described in this genus and
 374 the fourth *Paradoxapseudes* species found in submarine caves. Five congeners share a
 375 long pleotelson, defined here as one with PL/PW greater than 1.7, with *P. shimojiensis*:
 376 *P. basibidens* (Guțu, 2007) from Tanzania; *P. bassoprofundo* Bamber and Błażewicz-
 377 Paszkowycz, 2013 from Australia; *P. bermudeus* from Bermuda (submarine cave); *P.*
 378 *edgari* Guțu, 2008 from the Cook Islands; and *P. heroae* Sieg, 1985 from Argentina.

379 *Paradoxapseudes shimojiensis* can be distinguished from these five species by
 380 the combinations of six character states presented in Table II. In addition to the features
 381 in the table, *P. shimojiensis* differs from *P. basibidens* in having more-setose
 382 maxillipedal palp articles 2 and 3 (eleven and five simple setae on the two articles in *P.*
 383 *basibidens*), the chelipedal merus with one ventrodistal spiniform seta (absent in *P.*
 384 *basibidens*), the pereopod-1 carpus with six or seven dorsal simple setae (four in *P.*
 385 *basibidens*), the bases of pereopods 4 and 5 bearing ventrodistal simple setae (lacking in
 386 *P. basibidens*), and the ischia of pereopods 1–6 without dorsal setae (present in *P.*
 387 *basibidens*) (Guțu 2007); from *P. bassoprofundo* in having the chelipedal basis with a
 388 single ventro-subproximal simple seta (two in *P. bassoprofundo*), the chelipedal merus
 389 with one ventrodistal spiniform seta (absent in *P. bassoprofundo*), the chelipedal carpus
 390 with several dorsal simple setae (absent in *P. bassoprofundo*), and the ischia of
 391 pereopods 4–6 without dorsal setae (present in *P. bassoprofundo*) (Bamber and
 392 Błażewicz-Paszkowycz 2013); from *P. bermudeus* in having the ventro-subproximal
 393 seta on the pereopod-1 basis simple (plumose in *P. bermudeus*), the chelipedal merus
 394 with one ventrodistal spiniform seta (absent in *P. bermudeus*), and the ischia of
 395 pereopods 5 and 6 without dorsal setae (present in *P. bermudeus*) (Băcescu 1980; Guțu
 396 2008); from *P. edgari* in the pleotelson having two pairs of slight lateral processes

(lateral margins of pleotelson straight in *P. edgari*), the chelipedal basis with one dorsodistal simple seta (two in *P. edgari*), the chelipedal merus with one ventrodistal spiniform seta (absent in *P. edgari*), and the ventro-subproximal seta on the pereopod-1 basis simple (plumose in *P. edgari*) (Guțu 2008); and from *P. heroae* in having the chelipedal basis without dorsal processes (present in *P. heroae*), the inner distal seta on article 1 of the maxillipedal palp plumose (simple in *P. heroae*), the pereopod-1 basis with one ventro-subproximal simple seta (absent in *P. heroae*), the pereopod-1 carpus with six or seven dorsal simple setae (11 in *P. heroae*), and the ischia of pereopods 4 and 5 without dorsal setae (present in *P. heroae*) (Sieg 1986).

Paradoxapseudes shimojiensis differs from another Japanese species, *P. littoralis*, in having a long pleotelson ($PL/PW = 1.2$ in *P. littoralis*), antennal article 1 naked (with one inner seta in *P. littoralis*), the chelipedal merus with one ventrodistal spiniform seta (absent in *P. littoralis*), the pereopod-6 basis with four or five dorsal plumose setae (eight in *P. littoralis*), and the dorsal setae on the pereopod-6 carpus plumose (simple in *P. littoralis*) (Shiino 1952).

Discussion

Paradoxapseudes shimojiensis sp. nov. is the 18th tanaidacean species reported from submarine caves worldwide, and the second from Japan—the other being the pseudozeuxid *Haimormus shimojiensis* Kakui and Fujita, 2018, originally described from the same cave as *P. shimojiensis*. Submarine caves are extreme environments under dark, food scarce, and hypoxic conditions; thus morphological, behavioral, and physiological adaptations to improve food finding capability, to resist starvation, and to reduce energy demand via reduced metabolism may be required for life in caves (Iliffe and Bishop 2007). Although cave endemic species are known to exist (e.g., Okanishi

and Fujita 2018), as caves open to the surrounding marine environment, animals thriving in caves can also be found in suitable habitats outside of caves. In addition, organisms which can survive in caves for only a short time can accidentally invade caves (Iliffe and Bishop 2007). Two of our *P. shimojiensis* individuals were collected from the completely dark, middle zone of the cave, and both had a fully-developed marsupium, implying this species may be thriving in the caves (= stygophilic taxon). But, since we lack faunal data from the surrounding benthic environment outside the cave, it remains unclear whether *P. shimojiensis* is restricted to submarine caves. Furthermore, while cave-dwelling animals often lose eye pigments or eyes (Iliffe and Bishop 2007), the lack of eye pigments (Figure 1A, B) in *P. shimojiensis* is not necessarily an adaptation to cave dwelling. Species in *Paradoxapseudes*, including *P. shimojiensis*, have a fossorial-type pereopod 1 (Guțu and Sieg 1999), indicating that they are burrowers (or tube-dwellers; cf. Kakui and Hiruta 2017). Non-cave-dwelling congeners thus also likely live in dim or dark microhabitats, and the loss of eye pigments in *P. shimojiensis* could be related either to cave dwelling, or to a burrowing or tube-dwelling mode of life.

We found inner serial ridges on the left and right chelipeds of *P. shimojiensis* (Figure 5), which is the first case known in *Paradoxapseudes*. The ridges form an arc on the inner surface of the basis, and those on the left and right chelipeds are opposed to each other; each ridge is orthogonal to the swing path of the basis. This face-to-face relationship of serial ridges or hatching on chelipeds has been found in some other tanaidacean genera (*Apseudomorpha* Miller, 1940, *Nesotanaïs* Shiino, 1968, and *Tanaopsis* Sars, 1899), but the location of the ridges in these taxa is different from that in *P. shimojiensis*: the ridges occur on the carpi in *Apseudomorpha* (Menzies 1953); on the propodi and dactyli in *Nesotanaïs* (Kakui et al. 2010); and on the dactyli in

Tanaopsis (Kakui and Shimada 2017). The location of the serial ridges in *P. shimojiensis* is quite similar to that of the ridges in *Histicostoma harvestmen* (Nemastomatidae, Opiliones; Juberthie 1957), where they form an arc on the inner surface of the basal segment of the chelicerae and are considered to be a stridulatory organ. This similarity suggests that the paired serial ridges in *P. shimojiensis* may also be a stridulatory organ and function in producing sound. As possible stridulatory organs have also been found in tanaidaceans not dwelling in submarine caves (e.g., Kakui et al. 2010), and no other submarine cave species bear these ridges, the serial ridges may not be an adaptation to cave dwelling.

The two specimens of *P. shimojiensis* we observed bear external female and male reproductive features in the form of a fully-developed marsupium and genital cone (Figure 1). This indicates that *P. shimojiensis* is simultaneously hermaphroditic. *Paradoxapseudes* contains three possibly hermaphroditic species (*P. bermudeus*, *P. intermedius*, and *P. shimojiensis*; Băcescu 1980; Guțu 2006; this study), which are three of the four congeners reported from submarine caves. Simultaneous or sequential hermaphroditism has so far been confirmed or inferred for about 30 tanaidacean species (Rumbold et al. 2015; Kakui et al. 2019), or about 2% of the tanaidacean species currently known worldwide (ca. 1400 species; Anderson 2017). Among taxa reported from submarine caves, however, six of 18 species (ca. 33%) have been confirmed or suggested to be hermaphrodites: *Apseudopsis latreilli* (Milne-Edwards, 1828), the three *Paradoxapseudes* species mentioned above, *Falsapseudes bowmani* (Guțu and Iliffe, 1989), and *Chondrochelia savignyi* (Krøyer, 1842) (Masunari 1983; Esquete et al. 2012; Kakui et al. 2019). As tanaidaceans are benthic animals and lack a planktonic larval stage (obligate dispersal phase) in their life cycle, they have fewer opportunities to colonize submarine caves than organisms with planktonic larvae. In the case of two

random individuals reaching the inside of a cave, gonochoristic species require both a male and a female for propagation; in comparison, protandrous or protogynous species have a twofold chance of propagation, and simultaneous hermaphrodites have a threefold chance. For simultaneous hermaphrodites that can self-fertilize, such as one species of *Apseudes* (Kakui and Hiruta 2013), only one individual is needed to establish a population. These advantages of hermaphrodites in colonization may be reflected in the larger proportion of hermaphroditic species in submarine caves.

Acknowledgments

We thank Masaru Mizuyama, Katrine Worsaae, Peter Rask Møller, Yuji Ise, Hiroki Ichi, Hirotaka Ichi, and Go Tomitani for assistance in scuba sampling in the submarine cave; and Matthew H. Dick for reviewing the manuscript and editing our English. This study was supported by the Japan Society for the Promotion of Science (JSPS) under KAKENHI grants JP16K18597 and JP19K06800 to KK, and JP16K07490 and JP16H06309 to YF.

References

- Anderson G. 2017. Tanaidacea—Thirty Years of Scholarship, Version 2.5. [accessed 2019 August 15]. <https://aquila.usm.edu/tan aids30/4/>.
- Băcescu M. 1980. *Apseudes bermudeus* n. sp. from caves around Bermude Islands. Acta Adriatica. 21:401–407.

- 493 Bamber RN. 2000. Additions to the apseudomorph tanaidaceans (Crustacea: Peracarida)
 494 of Hong Kong. In: Morton B, editor. The marine flora and fauna of Hong Kong
 495 and southern China. Vol. V. Hong Kong: Hong Kong University Press; p. 37–51.
- 496 Bamber RN, Błażewicz-Paszkowycz M. 2013. Another inordinate fondness[†]: diversity
 497 of the tanaidacean fauna of Australia, with description of three new taxa. Journal of
 498 Natural History. 47:1767–1789.
- 499 Bird GJ. 2011. Paratanaoidean tanaidaceans (Crustacea: Peracarida) from littoral and
 500 shallow sublittoral habitats in New Zealand, with descriptions of three new genera
 501 and seven new species. Zootaxa. 2891:1–62.
- 502 Bird GJ. 2015. Tanaidacea (Crustacea: Peracarida) of the Kermadec Biodiscovery
 503 Expedition 2011, with a new sub-family of Paratanaidae: Metatanainae. Bulletin of
 504 the Auckland Museum. 20:369–404.
- 505 Drumm DT. 2010. Phylogenetic relationships of Tanaidacea (Eumalacostraca:
 506 Peracarida) inferred from three molecular loci. Journal of Crustacean Biology.
 507 30:692–698.
- 508 Esquete P, Bamber RN, Moreira J, Troncoso JS. 2012. Redescription and postmarsupial
 509 development of *Apseudopsis latreillii* (Crustacea: Tanaidacea). Journal of the
 510 Marine Biological Association of the United Kingdom. 92:1023–1041.
- 511 Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. 1994. DNA primers for
 512 amplification of mitochondrial cytochrome *c* oxidase subunit I from diverse
 513 metazoan invertebrates. Molecular Marine Biology and Biotechnology. 3:294–299.
- 514 García-Herrero Á, Sánchez N, García-Gómez G, Pardos F, Martinez A. 2019. Two new
 515 stygophilic tanaidomorphs (Peracarida, Tanaidacea) from Canary Islands and
 516 southeastern Iberian Peninsula. Marine Biodiversity. 49:107–130.

- 517 Gerovasileiou V, Chintiroglou CC, Konstantinou D, Voltsiadou E. 2016. Sponges as
518 “living hotels” in Mediterranean marine caves. *Scientia Marina*. 80:279–289.
- 519 Guțu M. 1991. The description of *Paradoxapseudes cubensis*, a new genus and a new
520 species of Tanapseudidae (Crustacea, Tanaidacea). *Travaux du Muséum d’Histoire*
521 *Naturelle “Grigore Antipa”*. 31:349–354.
- 522 Guțu M. 2006. New apseudomorph taxa (Crustacea, Tanaidacea) of the world ocean.
523 Bucharest: Curtea Veche.
- 524 Guțu M. 2007. Contribution to the knowledge of the Indo-West-Pacific Apseudomorpha
525 (Crustacea: Tanaidacea). *Travaux du Muséum National d’Histoire Naturelle*
526 *“Grigore Antipa”*. 50:47–86.
- 527 Guțu M. 2008. New data on the genus *Paradoxapseudes* Guțu, 1991, including the
528 description of a new species. The synonymysation of *Gollumudes* Bamber, 2000
529 with *Paradoxapseudes* and the description of a new apseudid genus (Crustacea:
530 Tanaidacea). *Travaux du Muséum National d’Histoire Naturelle “Grigore Antipa”*.
531 51:17–42.
- 532 Guțu M, Iliffe TM. 1985. The redescription of *Apseudes* (?) *propinquus* Richardson,
533 1902 (Crustacea, Tanaidacea) from Bermuda caves. *Travaux du Muséum*
534 *d’Histoire Naturelle “Grigore Antipa”*. 27:55–62.
- 535 Guțu M, Iliffe TM. 1989. Description of two new species of Tanaidacea (Crustacea)
536 from the marine water caves of the Palau Islands (Pacific Ocean). *Travaux du*
537 *Muséum National d’Histoire Naturelle “Grigore Antipa”*. 30:169–180.
- 538 Guțu M, Sieg J. 1999. *Ordre des Tanaïdacés (Tanaidacea Hansen, 1895) [Order*
539 *Tanaidacea (Tanaidacea Hansen, 1895)]*. *Mémoires de l’Institut Océanographique*,
540 Monaco. 19:353–389.

- 541 Iliffe TM, Bishop RE. 2007. Adaptations to life in marine caves. In: Safran P, editor.
 542 Fisheries and aquaculture. Vol. 5. Oxford: Eolss Publishers; p. 183–205.
- 543 Juberthie C. 1957. Présence d’organes de stridulation chez deux Nemastomatidae
 544 (Opilions) [Presence of stridulation organs in two Nemastomatidae (Opilions)].
 545 Bulletin du Muséum National d’Histoire Naturelle. 2nd Series. 29:210–212.
- 546 Kakui K, Angsupanich S. 2012. *Birdotanais songkhlaensis*, a new genus and species of
 547 Nototanaidae (Crustacea: Tanaidacea) from Thailand. Raffles Bulletin of Zoology.
 548 60:421–432.
- 549 Kakui K, Fujita Y. 2018. *Haimormus shimojiensis*, a new genus and species of
 550 Pseudozeuxidae (Crustacea: Tanaidacea) from a submarine limestone cave in
 551 Northwestern Pacific. PeerJ. 6:e4720.
- 552 Kakui K, Hiruta C. 2013. Selfing in a malacostracan crustacean: why a tanaidacean but
 553 not decapods. Naturwissenschaften. 100:891–894.
- 554 Kakui K, Hiruta C. 2017. Tube construction by a tanaidacean crustacean using a novel
 555 mucus secretion system involving the anal opening. Zoological Letters. 3:20.
- 556 Kakui K, Hiruta C, Uyeno D. 2019. Sexual system in the tanaidacean *Falsapseudes*
 557 *bowmani* (Crustacea: Malacostraca: Peracarida). Invertebrate Biology. 138:e12257.
- 558 Kakui K, Kajihara H, Mawatari SF. 2010. A new species of *Nesotanais* Shiino, 1968
 559 (Crustacea, Tanaidacea) from Japan, with a key to species and a note on male
 560 chelipeds. ZooKeys. 33:1–17.
- 561 Kakui K, Katoh T, Hiruta SF, Kobayashi N, Kajihara H. 2011. Molecular systematics of
 562 Tanaidacea (Crustacea: Peracarida) based on 18S sequence data, with an
 563 amendment of suborder/superfamily-level classification. Zoological Science.
 564 28:749–757.

- 565 Kakui K, Shimada D. 2017. A new species of *Tanaopsis* (Crustacea: Tanaidacea) from
566 Japan, with remarks on the functions of serial ridges and grooves on the
567 appendages. *Zootaxa*. 4282:324–336.
- 568 Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
569 improvements in performance and usability. *Molecular Biology and Evolution*.
570 30:772–780.
- 571 Katoh K, Toh H. 2008. Recent developments in the MAFFT multiple sequence
572 alignment program. *Briefings in Bioinformatics*. 9:286–298.
- 573 Kimura M. 1980. A simple method for estimating evolutionary rates of base
574 substitutions through comparative studies of nucleotide sequences. *Journal of*
575 *Molecular Evolution*. 16:111–120.
- 576 Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular evolutionary genetics
577 analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*.
578 33:1870–1874.
- 579 Larsen K. 2003. Proposed new standardized anatomical terminology for the Tanaidacea
580 (Peracarida). *Journal of Crustacean Biology*. 23:644–661.
- 581 Masunari S. 1983. Postmarsupial development and population dynamics of *Leptochelia*
582 *savignyi* (Krøyer, 1842) (Tanaidacea). *Crustaceana*. 44:151–162.
- 583 Menzies RJ. 1953. The apseudid chelifera of the eastern tropical and north temperate
584 Pacific Ocean. *Bulletin of the Museum of Comparative Zoology at Harvard*
585 *College*. 107:443–496.
- 586 Nakayama T, Watanabe S, Mitsui K, Uchida H, Inoue I. 1996. The phylogenetic
587 relationship between the Chlamydomonadales and Chlorococcales inferred from
588 18SrDNA sequence data. *Phycological Research*. 44:47–55.

- 589 Okanishi M, Fujita Y. 2018. First finding of anchialine and submarine cave dwelling
590 brittle stars from the Pacific Ocean, with descriptions of new species of *Ophiolepis*
591 and *Ophiozonella* (Echinodermata: Ophiuroidea: Amphilepidida). Zootaxa.
592 4377:1–20.
- 593 Osawa M, Fujita Y. 2019. Submarine cave hermit crabs (Crustacea: Decapoda:
594 Anomura: Paguroidea) from three islands of the Ryukyu Islands, southwestern
595 Japan. Zootaxa. 4560:463–482.
- 596 Rasband WS. 2019. ImageJ [software]. [accessed 2019 August 15].
597 <http://imagej.nih.gov/ij>.
- 598 Rumbold CE, Obenat SM, Leonardi MS, Spivak ED. 2015. Intersex in the
599 gonochoristic crustacean *Tanais dulongii* (Audouin, 1826) (Peracarida:
600 Tanaidacea: Tanaididae): a comparison of external reproductive characteristics.
601 Journal of Natural History. 49:775–788.
- 602 Shiino SM. 1952. A new genus and two new species of the order Tanaidacea found at
603 Seto. Publications of the Seto Marine Biological Laboratory. 2:53–68.
- 604 Sieg J. 1986. Crustacea Tanaidacea of the Antarctic and the Subantarctic 1. On material
605 collected at Tierra del Fuego, Isla de los Estados, and the west coast of the
606 Antarctic Peninsula. Washington D.C.: American Geophysical Union.
- 607 Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: Improving the sensitivity
608 of progressive multiple sequence alignment through sequence weighting, position-
609 specific gap penalties and weight matrix choice. Nucleic Acids Research. 22:4673–
610 4680.
- 611 Tzeng Y-W, Hsueh P-W. 2014. New species and records of Apseudomorpha
612 (Crustacea: Tanaidacea) from Taiwan. Zootaxa. 3869:313–337.

- 613 Wilson GDF. 2009. The phylogenetic position of the Isopoda in the Peracarida
614 (Crustacea: Malacostraca). *Arthropod Systematics & Phylogeny*. 67:159–198.
615

616 Table I. List of PCR and cycle sequencing (CS) primers used in this study

Marker	Primer	Sequence	Reaction	Source
COI	LCO1490	GGTCAACAAATCATAAAGATATTGG	PCR & CS	Folmer et al. (1994)
	HCO2198	TAAACTTCAGGGTGACCAAAAAATCA	PCR & CS	Folmer et al. (1994)
18S	SR-1	TACCTGGTTGATCCTGCCAG	PCR	Nakayama et al. (1996)
	18S-b3F	CCTGAGAAACGGCTACCACAT	CS	Kakui and Shimada (2017)
	18S-b4R	TCCAACTACGAGCTTTTAAACC	CS	Kakui et al. (2011)
	18S-b8R	TCTAAGGGCATCACAGACCTG	CS	Kakui et al. (2011)
	18S-b8F	GGTCTGTGATGCCCTTAGATG	CS	Kakui et al. (2011)
	SR-12	CCTTCCGCAGGTTACCTAC	PCR & CS	Nakayama et al. (1996)

617

618

619 Table II. Comparison of selected characters for six *Paradoxapseudes* species having a long pleotelson (PL/PW greater than 1.7). *,
 620 hermaphroditic (others, gonochoristic). ^aLong, as long as or longer than width of pereopod-1 basis; short, shorter than width of pereopod-1 basis.
 621 –, no data. Abbreviations: pl.seta, plumose seta; s.seta, simple seta; sp.seta, spiniform seta

	Antenna	Maxilliped	Pereopod 1			Pleopod
	Article 1	Basis	Basis		Merus	Protopod
	Inner s.seta	Inner distal seta	Dorsal seta ^a	Ventrodistal sp.seta	Mid-inner ventral s.seta	Inner pl.seta
<i>P. shimojiensis</i> sp. nov.*	0	1 pl.seta	4 long s.setae	0	3	2
<i>P. basibidens</i>	0	1 pl.seta	5 long pl.setae	1	0	2
<i>P. bassoprofundo</i>	1	0	1 long and 3 short s.setae	0	3	1
<i>P. bermudeus</i> *	0	–	4 long s.setae	0	3	0
<i>P. edgari</i>	0	1 pl.seta	4 long s.setae	1	4	2
<i>P. heroae</i>	1	2 s.setae	3 short s.setae	0	4	2

Figure 1. *Paradoxapseudes shimojiensis* sp. nov.: A, B, holotype in living state, left dorsal (A) and left (B) views; C, paratype, fixed, in glycerine, showing genital cone with two male gonopores (arrow) on pereonite 6, and empty fully-developed marsupium; right pereopods detached, right ventral view. Abbreviations: *p6*, pereonite 6; *pl1*, pleonite 1.

Figure 2. *Paradoxapseudes shimojiensis* sp. nov., holotype: A, body, dorsal view; a1, pleotelson, dorsal view; B, body, left view; C, right antennule, articles 1 and 2, ventral view; D, right antennule, articles 3 and 4, and outer and inner flagella, ventral view; d1, outer flagellum, article 7, ventral view (most setae omitted); E, right antenna, ventral view; F, labrum; G, H, mandible, left and right distal portions, respectively; I, left mandibular molar; J, right mandibular palp; j1, blade-like spiniform seta; j2, pectinate seta; K, labium. Arrows, left slight lateral processes.

Figure 3. *Paradoxapseudes shimojiensis* sp. nov. (A–F, holotype; G, paratype): A, B, outer and inner endites of left maxillule, respectively; C, right maxillular palp; c1, distal portion of biserrate-tipped seta; D, left maxilla (most setae on outer lobe of fixed endite omitted), anterior view; d1, same, outer lobe of fixed endite, anterior view; d2, branched seta; E, left maxillipedal endite; F, right maxilliped (setae on endite omitted); f1, same, palp article-4; G, right epignath.

Figure 4. *Paradoxapseudes shimojiensis* sp. nov., holotype: A, left cheliped, outer view (inner serial ridges omitted); B, right chela, inner view (dactylus and most setae omitted); b1, same, dactylus, inner view; C–H, right pereopods 1–6, outer view; c1, pectinate spiniform seta; e1, dactylus (part) and unguis of right pereopod 3, outer view;

647 f1, distal portion of right pereopod-4 propodus, outer view; g1, blade-like spiniform
648 seta; h1, serrate seta; h2, pectinate seta; h3, blade-like spiniform seta; h4, distal portion
649 of right pereopod-6 propodus, inner view; I, right pleopod 1 (most setal ornamentation
650 omitted); i1, distally-hooked plumose seta; J, right uropod (most articles of endopod and
651 exopod lost). Arrow, broken portion on pereopod-1 coxa; arrowhead, artificially bent
652 point on pereopod-4 dactylus.

653

654 **Figure 5.** *Paradoxapseudes shimojiensis* sp. nov., paratype, SEM image: A, right
655 cheliped, inner view, with rectangle indicating position of enlargement in B; B, inner
656 serial ridges on basis.

A

B

A, B: 0.5 mm

C

A **B**