

HOKKAIDO UNIVERSITY

Title	A theory of Miyawaki liftings : the Hilbert-Siegel case
Author(s)	Atobe, Hiraku
Citation	Mathematische annalen, 376(3-4), 1467-1535 https://doi.org/10.1007/s00208-019-01946-w
Issue Date	2020-04
Doc URL	http://hdl.handle.net/2115/80775
Rights	This is a post-peer-review, pre-copyedit version of an article published in Mathematische Annalen. The final authenticated version is available online at: https://doi.org/10.1007/s00208-019-01946-w
Туре	article (author version)
File Information	Math. Ann. 376-3-4_1467-1535.pdf



A THEORY OF MIYAWAKI LIFTINGS: THE HILBERT-SIEGEL CASE

HIRAKU ATOBE

ABSTRACT. The Miyawaki liftings are defined by the pullbacks of Ikeda liftings. Recently, Ikeda and Yamana extended the theory of Ikeda liftings to Hilbert–Siegel modular forms. In this paper, using their results, we establish a theory of Miyawaki liftings, both locally and globally. In the local theory, we describe the Miyawaki liftings for almost tempered unitary representations explicitly. In the global theory, we discuss the non-vanishing of the Miyawaki liftings using seesaw identities and the global Gan–Gross–Prasad conjecture. As an application of local Miyawaki liftings, we prove a new case of the local Gan–Gross–Prasad conjecture.

CONTENTS

1. Introduc	tion	1
2. Ikeda lif	2. Ikeda liftings and their Fourier–Jacobi coefficients	
3. Local Miyawaki liftings		17
4. Global M	fiyawaki liftings	31
5. A relation	n between Miyawaki liftings and the Gan–Gross–Prasad conjecture	37
Appendix A	Jacquet modules of representations of metaplectic groups	40
Appendix B.	Local Langlands correspondence and Arthur's multiplicity formula	42
Appendix C	Gan–Gross–Prasad conjectures	49
References		52

1. INTRODUCTION

In 1992, Miyawaki [39] predicted the existence of certain Siegel modular forms. Let $S_k(\text{Sp}_n(\mathbb{Z}))$ be the space of Siegel cusp forms of degree n, weight k, and level one.

Conjecture 1.1 (Miyawaki [39]). For normalized Hecke eigenforms $f \in S_{2k-4}(SL_2(\mathbb{Z}))$ and $g \in S_k(SL_2(\mathbb{Z}))$, there should exist a Hecke eigenform $F_{f,g} \in S_k(Sp_3(\mathbb{Z}))$ whose standard L-function is given by

$$L(s, F_{f,g}, st) = L(s, g, Ad)L(s + k - 2, f)L(s + k - 3, f).$$

In 2006, to approach Miyawaki's conjecture, Ikeda [28] constructed certain liftings, which are now called the Miyawaki liftings, as follows: For positive even k, a normalized Hecke eigenform $f \in S_{2k-2(n+r)}(\mathrm{SL}_2(\mathbb{Z}))$ gives the Ikeda lift $F^{(2n+2r)} \in S_k(\mathrm{Sp}_{2n+2r}(\mathbb{Z}))$ (defined up to a constant). For the (classical) Ikeda lifting, see [27]. For a Hecke eigenform $g \in S_k(\mathrm{Sp}_r(\mathbb{Z}))$, Ikeda [28] defined the Miyawaki lift $\mathcal{M}^{(2n+r)}(g, F^{(2n+2r)})$ by the integral

$$\mathcal{M}^{(2n+r)}\left(g,F^{(2n+2r)}\right)\left(Z_{2n+r}\right)$$

²⁰¹⁰ Mathematics Subject Classification. Primary 11F70; Secondary 11F46.

Key words and phrases. Automorphic representations; Miyawaki liftings; Gan-Gross-Prasad conjecture.

$$= \int_{\operatorname{Sp}_r(\mathbb{Z})\backslash\mathfrak{H}_r} F^{(2n+2r)}\left(\begin{pmatrix} Z_{2n+r} & 0\\ 0 & Z_r \end{pmatrix}\right) \overline{g^c(Z_r)} (\det \operatorname{Im} Z_r)^{k-r-1} dZ_r,$$

where \mathfrak{H}_r is the Siegel upper half space of genus r and we set $g^c(Z_r) = \overline{g(-\overline{Z_r})}$. It is easy to see that $\mathcal{M}^{(2n+r)}(g, F^{(2n+2r)}) \in S_k(\operatorname{Sp}_{2n+r}(\mathbb{Z}))$. Ikeda proved the following:

Theorem 1.2 (Ikeda [28, Theorem 1.1]). If $\mathcal{M}^{(2n+r)}(g, F^{(2n+2r)})$ is not identically zero, then it is a Hecke eigenform with standard L-function

$$L\left(s, \mathcal{M}^{(2n+r)}\left(g, F^{(2n+2r)}\right), \mathrm{st}\right) = L(s, g, \mathrm{st}) \prod_{i=1}^{2n} L(s+k-r-i, f)$$

Therefore Miyawaki's conjecture (Conjecture 1.1) was reduced to the non-vanishing of $\mathcal{M}^{(3)}(g, F^{(4)})$. Ikeda gave a conjectural formula for the Petersson norm of $\mathcal{M}^{(2n+r)}(g, F^{(2n+2r)})$, and proposed the following conjecture.

- **Conjecture 1.3** (Ikeda [28, Conjecture 5.1]). (1) When n = 0, the Miyawaki lift $\mathcal{M}^{(r)}(g, F^{(2r)})$ is nonzero if and only if the central value of the tensor product L-function $L(s, \operatorname{st}(g) \boxtimes f)$ is nonzero.
 - (2) When n > 0, the Miyawaki lift $\mathcal{M}^{(2n+r)}(g, F^{(2n+2r)})$ is always nonzero.

Ichino [25] and Xue [57] proved Conjecture 1.3 (1) for the case where r = 1 independently. Garrett-Heim [17] established a Hecke duality of Ikeda liftings and gave a preliminary answer to Conjecture 1.3 (1).

Nowadays Miyawaki's conjecture (Conjecture 1.1) follows from Arthur's multiplicity formula ([3, Theorem 1.5.2]), which was established in 2013. However, this formula tells us only the existence of modular forms (or automorphic representations). The non-vanishing of Miyawaki liftings (Conjecture 1.3) is still open and interesting. The integral representations of Miyawaki liftings would imply several properties, which do not follow from Arthur's multiplicity formula.

Recently, Ikeda–Yamana [29] extended the theory of Ikeda liftings to Hilbert–Siegel modular forms using representation theory. The purpose of this paper is to establish a theory of Miyawaki liftings using the extended Ikeda liftings. We will define the Miyawaki liftings more generally, and give their several properties. Moreover, we will approach the non-vanishing problem of Miyawaki liftings using the Gan–Gross–Prasad conjecture.

To describe our results, let F be a totally real number field, and \mathbb{A} be the ring of adeles of F. The ring of finite adeles of F is denoted by \mathbb{A}_{fin} . For a place v of F, we write $v < \infty$ (resp. $v \mid \infty$) if v is a finite place (resp. if v is an infinite place). Fix a non-trivial unitary character ψ of \mathbb{A}/F such that for $v \mid \infty$, the local component ψ_v is of the form $\psi_v(x_v) = \exp(2\pi a_v \sqrt{-1}x_v)$ for $x_v \in F_v \cong \mathbb{R}$ with fixed $a_v > 0$. Let $\tau = \otimes'_v \tau_v$ be an irreducible cuspidal automorphic representation of PGL₂(\mathbb{A}) satisfying the following conditions:

- (A1) For $v < \infty$, τ_v is a principal series $\mu_v \times \mu_v^{-1} \coloneqq \operatorname{Ind}_{P_{(1,1)}(F_v)}^{\operatorname{GL}_2(F_v)}(\mu_v \boxtimes \mu_v^{-1})$, where $P_{(1,1)}$ is the Borel subgroup of GL₂ consisting of upper triangular matrices. (Note that $\mu_v \times \mu_v^{-1}$ has the trivial central character so that $\operatorname{PGL}_2(F_v)$ acts on it.)
- (A2) For $v \mid \infty, \tau_v$ is a discrete series representation with lowest weight $\pm 2k_v$, where $k_v > 0$.
- (A3) The root number $\varepsilon(1/2,\tau)$ is equal to 1.

Let Sp_n be the symplectic group of rank n. For each place v, we denote the metaplectic double cover of $\operatorname{Sp}_n(F_v)$ by $\widetilde{\operatorname{Sp}}_n(F_v)$. Identify $\widetilde{\operatorname{Sp}}_n(F_v) = \operatorname{Sp}_n(F_v) \times \{\pm 1\}$ as sets. Let P_n be the Siegel parabolic subgroup of Sp_n , and $\widetilde{P}_n(F_v)$ be the inverse image of $P_n(F)$ in $\widetilde{\operatorname{Sp}}_n(F_v)$. When $v < \infty$, we set

$$I_{\psi_v}^{(n)}(\tau_v) = \operatorname{Ind}_{\widetilde{P}_n(F_v)}^{\widetilde{\operatorname{Sp}}_n(F_v)}(\mu_v^{(n)})$$

to be a degenerate principal series. Here,

$$\mu_v^{(n)}\left(\begin{pmatrix} A & 0\\ 0 & tA^{-1} \end{pmatrix}, \zeta\right) = \zeta^n \left(\frac{\alpha_{\psi_v}(1)}{\alpha_{\psi_v}(\det A)}\right)^n \mu_v(\det A)$$

for $A \in \operatorname{GL}_n(F_v)$ and $\zeta \in \{\pm 1\}$, where $\alpha_{\psi_v}(a)$ is the Weil constant (see §2.2). It is known that $I_{\psi_v}^{(n)}(\tau_v)$ is irreducible ([35, 50]). Set $k + (n/2) = (k_v + (n/2))_v \in \prod_{v \mid \infty} \mathbb{Z}$. Let $\widetilde{\operatorname{Sp}}_n(\mathbb{A})$ be the metaplectic double cover of the adelic group $\operatorname{Sp}_n(\mathbb{A})$. We denote the set of holomorphic cusp forms of weight k + (n/2) by $\mathcal{S}_{k+(n/2)}\left(\operatorname{Sp}_n(F) \setminus \widetilde{\operatorname{Sp}}_n(\mathbb{A})\right)$. For the definition, see §2.5. Ikeda– Yamana [29] showed that the irreducible representation $I_{\psi}^{(n)}(\tau) = \otimes_{v < \infty}' I_{\psi_v}^{(n)}(\tau_v)$ of $\widetilde{\operatorname{Sp}}_n(\mathbb{A}_{\mathrm{fin}})$ appears in $\mathcal{S}_{k+(n/2)}\left(\operatorname{Sp}_n(F) \setminus \widetilde{\operatorname{Sp}}_n(\mathbb{A})\right)$ with multiplicity one. We denote the unique subrepresentation of $\mathcal{S}_{k+(n/2)}\left(\operatorname{Sp}_n(F) \setminus \widetilde{\operatorname{Sp}}_n(\mathbb{A})\right)$ which is isomorphic to $I_{\psi}^{(n)}(\tau)$ by $\operatorname{Ik}_{\psi}^{(n)}(\tau)$, and call it the Ikeda lift of τ .

Now let n,r be non-negative integers. Then we have an embedding $\iota\colon \mathrm{Sp}_n\times\mathrm{Sp}_r\to\mathrm{Sp}_{n+r}$ given by

$$\iota\left(\begin{pmatrix} A_1 & B_1 \\ C_1 & D_1 \end{pmatrix}, \begin{pmatrix} A_2 & B_2 \\ C_2 & D_2 \end{pmatrix}\right) = \begin{pmatrix} A_1 & 0 & B_1 & 0 \\ 0 & A_2 & 0 & B_2 \\ \hline C_1 & 0 & D_1 & 0 \\ 0 & C_2 & 0 & D_2 \end{pmatrix}$$

For an admissible (not necessarily irreducible) representation π of $\widetilde{\operatorname{Sp}}_r(\mathbb{A}_{\operatorname{fin}})$ occurring in the space $\mathcal{S}_{k+(n+r)/2}\left(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A})\right)$, we define the (global) Miyawaki lift $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ of π by the representation of $\widetilde{\operatorname{Sp}}_n(\mathbb{A}_{\operatorname{fin}})$ generated by the integrals

$$\mathcal{M}^{(n)}((g_n,\zeta_n);\varphi,\mathcal{F}) = \int_{\mathrm{Sp}_r(F)\backslash \mathrm{Sp}_r(\mathbb{A})} \mathcal{F}(\iota(g_n,g_r),\zeta_n\zeta_r)\overline{\varphi(g_r,\zeta_r)} dg_r$$

for $\varphi \in \pi$, $\mathcal{F} \in \mathrm{Ik}_{\psi}^{(n+r)}(\tau)$ and $(g_n, \zeta_n) \in \widetilde{\mathrm{Sp}}_n(\mathbb{A})$. This is a subrepresentation of $\mathcal{S}_{k+(n+r)/2}(\mathrm{Sp}_n(F) \setminus \widetilde{\mathrm{Sp}}_n(\mathbb{A}))$. We summarize several properties of Miyawaki liftings.

Theorem 1.4. Let π be an irreducible representation of $\widetilde{\operatorname{Sp}}_r(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+(n+r)/2}(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A}))$, and $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ be its Miyawaki lift. Suppose that $\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \neq 0$ and $n \geq r$.

(1) If π has an A-parameter Ψ , then $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ has an A-parameter

$$\Psi \boxplus \tau \chi_{-1}^{[(n+r)/2]}[n-r],$$

where χ_{-1} is the quadratic character of $\mathbb{A}^{\times}/F^{\times}$ corresponding to $F(\sqrt{-1})/F$. For the notation of A-parameters, see §4.2, or Appendix B.4 for a more detailed explanation.

(2) Suppose that π has a tempered A-parameter. Then $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ is irreducible, and $\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \cong \otimes_{v<\infty}' \mathcal{M}_{\psi_v,\tau_v}^{(n)}(\pi_v)$. Here, $\mathcal{M}_{\psi_v,\tau_v}^{(n)}(\pi_v)$ is the local Miyawaki lift of π_v described below. (3) We have

$$\pi \subset \mathcal{M}_{\psi,\tau}^{(r)}\left(\mathcal{M}_{\psi,\tau}^{(n)}(\pi)\right).$$

If π has a tempered A-parameter and if $r \leq n \leq r+1$ or n > 2r, then the inclusion is in fact an equality.

The statements in Theorem 1.4 are proven in Proposition 4.3, Proposition 4.4, and Theorem 4.5. Remark that Arthur's multiplicity formula ([3, Theorem 1.5.2]) should imply the existence of an irreducible subrepresentation of $S_{k+(n+r)/2}(\operatorname{Sp}_r(\mathcal{A}))$ satisfying Theorem 1.4 (1) at least when n + r is even.

The definition of Miyawaki liftings is similar to the one of theta liftings. As Miyawaki liftings are given by the pullbacks of Ikeda liftings, theta liftings are defined by the pullbacks of theta functions. One of the most important properties of theta liftings is the seesaw identities, which have several applications. For instance, by using seesaw identities, Ichino [25] computed the pullbacks of Saito–Kurokawa liftings, which is regarded as a special case of Ikeda's conjectural formula for the Petersson norm of a certain Miyawaki lift. For another example, Xue [56] reduced the refined version of the Gan–Gross–Prasad conjecture for the symplectic-metaplectic case to the one for the special orthogonal case.

Miyawaki liftings also satisfy certain seesaw identities. To state these identities, we introduce the Fourier–Jacobi periods. Fix a totally positive element $\xi \in F^{\times}$ and set $\psi_{\xi}(x) = \psi(\xi x)$ for $x \in \mathbb{A}$. Let n' = n or n' = n - 1. For $\varphi \in S_{l/2}\left(\operatorname{Sp}_n(F) \setminus \widetilde{\operatorname{Sp}}_n(\mathbb{A})\right), \ \varphi' \in S_{l'/2}\left(\operatorname{Sp}_{n'}(F) \setminus \widetilde{\operatorname{Sp}}_{n'}(\mathbb{A})\right)$, and $\phi \in \mathcal{S}(\mathbb{A}^{n'})$, we define the Fourier–Jacobi period $\mathcal{P}_{n,n',\psi_{\xi}}\left(\varphi, \overline{\varphi'}, \phi\right)$ by the integral

$$\begin{cases} \int_{\operatorname{Sp}_{n}(F)\setminus\operatorname{Sp}_{n}(\mathbb{A})} \varphi(g,\zeta)\overline{\varphi'(g,\zeta)}\overline{\Theta}_{\psi_{\xi}}^{\phi}(g,\zeta)}dg, & \text{if } n'=n, \\ \int_{V_{n-1}(F)\setminus V_{n-1}(\mathbb{A})} \int_{\operatorname{Sp}_{n-1}(F)\setminus\operatorname{Sp}_{n-1}(\mathbb{A})} \varphi(v(g,\zeta))\overline{\varphi'(g,\zeta)}\overline{\Theta}_{\psi_{\xi}}^{\phi}(v(g,\zeta))}dgdv, & \text{if } n'=n-1\end{cases}$$

Here, $\Theta_{\psi_{\xi}}^{\phi}$ is the theta function associated to $\phi \in \mathcal{S}(\mathbb{A}^{n'})$, which is a genuine automorphic form on a Jacobi group $\widetilde{J}_{n'}(\mathbb{A}) = \widetilde{\operatorname{Sp}}_{n'}(\mathbb{A}) \ltimes V_{n'}(\mathbb{A})$, where $V_{n'} \cong F^{2n'} \oplus F$ is a Heisenberg group. Let $\mathcal{S}(\mathbb{A}^{n'})_{\xi}$ be the subspace of $\mathcal{S}(\mathbb{A}^{n'})$ consisting of lowest weight vectors of the Weil representation of $\widetilde{J}_{n'}(\mathbb{A})$ with respect to ψ_{ξ} .

Proposition 1.5 (Seesaw identity (Proposition 4.7)). Let π and π' be irreducible representations of $\widetilde{\operatorname{Sp}}_r(\mathbb{A}_{\operatorname{fin}})$ and $\widetilde{\operatorname{Sp}}_{n-1}(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+(n+r)/2}\left(\operatorname{Sp}_r(F)\backslash\widetilde{\operatorname{Sp}}_r(\mathbb{A})\right)$ and $\mathcal{S}_{k+(n-1+r)/2}\left(\operatorname{Sp}_{n-1}(F)\backslash\widetilde{\operatorname{Sp}}_{n-1}(\mathbb{A})\right)$, respectively.

(1) If there exist $\mathcal{M}^{(n)}(\varphi, \mathcal{F}_1) \in \mathcal{M}^{(n)}_{\psi, \tau}(\pi)$ with $\varphi \in \pi, \varphi' \in \pi'$, and $\phi_1 \in \mathcal{S}(\mathbb{A}^{n-1})_{\xi}$ such that

$$\mathcal{P}_{n,n-1,\psi_{\xi}}\left(\mathcal{M}^{(n)}(\varphi,\mathcal{F}_{1}),\overline{\varphi'},\phi_{1}\right)\neq 0,$$

then there exist $\mathcal{M}^{(r)}(\varphi', \mathcal{F}'_2) \in \mathcal{M}^{(r)}_{\psi, \tau \chi_{\xi}}(\pi')$ and $\phi_2 \in \mathcal{S}(\mathbb{A}^r)_{\xi}$ such that

$$\mathcal{P}_{r,r,\psi_{\xi}}\left(\varphi,\overline{\mathcal{M}^{(r)}(\varphi',\mathcal{F}_{2}')},\phi_{2}\right)\neq0.$$

(2) Assume that $n + r \ge 2$. If there exist $\varphi \in \pi$, $\mathcal{M}^{(r)}(\varphi', \mathcal{F}'_2) \in \mathcal{M}^{(r)}_{\psi, \tau \chi_{\xi}}(\pi')$ with $\varphi' \in \pi'$, and $\phi_2 \in \mathcal{S}(\mathbb{A}^r)_{\xi}$ such that

$$\mathcal{P}_{r,r,\psi_{\xi}}\left(\varphi,\overline{\mathcal{M}^{(r)}(\varphi',\mathcal{F}'_{2})},\phi_{2}\right)\neq0,$$

then there exist $\mathcal{M}^{(n)}(\varphi, \mathcal{F}_1) \in \mathcal{M}^{(n)}_{\psi, \tau}(\pi)$ and $\phi_1 \in \mathcal{S}(\mathbb{A}^{n-1})_{\xi}$ such that

$$\mathcal{P}_{n,n-1,\psi_{\xi}}\left(\mathcal{M}^{(n)}(\varphi,\mathcal{F}_{1}),\overline{\varphi'},\phi_{1}\right)\neq 0.$$

We shall write these properties as the following seesaw diagram:



Remark that the proof of Proposition 1.5 uses the non-vanishing of Fourier–Jacobi coefficients of Ikeda liftings (Proposition 2.7 (3)). In particular, the seesaw identities would not follow from Arthur's multiplicity formula.

Finally, following Ikeda's conjecture (Conjecture 1.3), we formulate a conjecture on the nonvanishing of Miyawaki liftings.

Conjecture 1.6 (Conjecture 5.1). Let π be an irreducible representation of $\widetilde{\text{Sp}}_r(\mathbb{A}_{\text{fin}})$ occurring in $\mathcal{S}_{k+(n+r)/2}\left(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A})\right).$

- (1) When n = r, the Miyawaki lift M^(r)_{ψ,τ}(π) is nonzero if and only if the central value of the Rankin–Selberg L-function L(s, π × τχ^r₋₁) is nonzero.
 (2) When n > n = the L(t) = the transformed (n)
- (2) When n > r, the Miyawaki lift $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ is always nonzero.

Using the seesaw identities (Proposition 1.5), one can prove the easiest case.

Proposition 1.7 (Corollary 5.6). Suppose that n = r = 1. Let π be an irreducible representation of $\operatorname{SL}_2(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+1}(\operatorname{SL}_2(F)\backslash \operatorname{SL}_2(\mathbb{A}))$. Then $\mathcal{M}^{(1)}_{\psi,\tau}(\pi) \neq 0$ if and only if $L(1/2, \pi \times \tau \chi_{-1}) \neq 0$ 0.

Outline of the proof. The Miyawaki lift $\mathcal{M}_{\psi,\tau}^{(1)}(\pi)$ is nonzero if and only if there exists a vector in $\mathcal{M}^{(1)}_{\psi,\tau}(\pi)$ whose ξ -th Fourier coefficient is nonzero for some totally positive element $\xi \in F^{\times}$. This condition means that $\mathcal{P}_{1,0,\psi_{\xi}}$ is nonzero on $\mathcal{M}_{\psi,\tau}^{(1)}(\pi) \times \overline{\eta} \times \mathcal{S}(\mathbb{A}^0)_{\xi}$, where η is the nontrivial character of $\widetilde{\mathrm{Sp}}_0(\mathbb{A}) = \{\pm 1\}$. By the seesaw identity (Proposition 1.5), it is equivalent that $\mathcal{P}_{1,1,\psi_{\mathcal{E}}}$ is nonzero on $\pi \times \overline{\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(1)}(\eta)} \times \mathcal{S}(\mathbb{A}^{1})_{\xi}$. Note that $\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(1)}(\eta)$ is the Ikeda lift $I_{\psi}^{(1)}(\tau\chi_{\xi})$, i.e., the Shimura correspondence of $\tau \chi_{\xi}$. By the global Gan–Gross–Prasad conjecture for $SL_2 \times \widetilde{SL}_2$ proven by Gan– Gurevich [12], Qiu [44], and Xue [56], the non-vanishing of the Fourier–Jacobi period $\mathcal{P}_{1,1,\psi_{\mathcal{E}}}$ on $\pi \times I_{\psi}^{(1)}(\tau \chi_{\xi}) \times \mathcal{S}(\mathbb{A}^1)_{\xi}$ is equivalent to the non-vanishing of the central value of $L(s, \pi \times \tau \chi_{-1})$. \Box

By a similar argument, we can relate Conjecture 1.6 to the Gan–Gross–Prasad conjecture (see Conjecture 5.2 below). The following theorem summarizes Theorems 5.5 and 5.7.

- Theorem 1.8. (1) Assume the Gan–Gross–Prasad conjecture (Conjecture 5.2) and Hypothesis 5.3 (A) below. Then Conjecture 1.6 for n = r, r + 1 holds when π has a tempered Aparameter.
 - (2) Assume Hypothesis 5.3 (B) or (C) below. Then Conjecture 1.6 for n > r is reduced to the case where n - r = 1.

Hypothesis 5.3 is an expectation of the existence of certain (tempered) automorphic representations. This was also considered in the theory of twisted automorphic descents introduced by Jiang–Zhang [32, Conjecture 2.3]. For more comments, see Remark 5.4 below.

There is a local analogue of Miyawaki liftings. Now let F be a non-archimedean local field of characteristic zero, and ψ be a non-trivial additive character of F. Let χ_{-1} be the quadratic character of F^{\times} corresponding to $F(\sqrt{-1})/F$. A local analogue of the Ikeda lifting is the degenerate principal series $I_{\psi}^{(n+r)}(\tau) = \operatorname{Ind}_{\widetilde{P}_{n+r}(F)}^{\widetilde{\operatorname{Sp}}_{n+r}(F)}(\mu^{(n+r)})$, where $\tau = \mu \times \mu^{-1}$ is a principal series of $\operatorname{GL}_2(F)$ with μ being a unitary character of F^{\times} . Recall that there is an embedding $\iota: \operatorname{Sp}_n \times \operatorname{Sp}_r \hookrightarrow \operatorname{Sp}_{n+r}$. For an irreducible representation π of $\widetilde{\operatorname{Sp}}_r(F)$, on which the kernel $\{\pm 1\}$ of the covering map $\widetilde{\operatorname{Sp}}_r(F) \twoheadrightarrow \operatorname{Sp}_r(F)$ acts by $(\pm 1)^{n+r}$, the maximal π -isotypic quotient of $I_{\psi}^{(n+r)}(\tau)$ is of the form

$$\mathcal{M}_{\psi,\tau}^{(n)}(\pi)\boxtimes\pi$$

for some smooth representation $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ of $\widetilde{\mathrm{Sp}}_n(F)$, on which the kernel $\{\pm 1\}$ of the covering map $\widetilde{\operatorname{Sp}}_n(F) \twoheadrightarrow \operatorname{Sp}_n(F)$ acts by $(\pm 1)^{n+r}$. We call $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ the local Miyawaki lift of π . The following is the main local theorem.

Theorem 1.9 (Theorem 3.1). Let μ be a unitary character of F^{\times} , and π be an irreducible representation of $\widetilde{\operatorname{Sp}}_r(F)$ on which $\{\pm 1\}$ acts by $(\pm 1)^{n+r}$. Suppose that $n \geq r$.

- (1) The local Miyawaki lift $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ is nonzero and of finite length.
- (2) If π is almost tempered (see §3.2 below) and unitary, then
 - $\mathcal{M}^{(n)}_{\psi,\tau}(\pi)$ is irreducible;
 - $\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \cong (\mu' \circ \det_{n-r}) \rtimes \pi \coloneqq \operatorname{Ind}_{\widetilde{P}_{n-r}(F)}^{\widetilde{\operatorname{Sp}}_n(F)}((\mu' \circ \det_{n-r}) \boxtimes \pi) \text{ with } \mu' = \mu \chi_{-1}^{[(n+r)/2]};$
 - $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ is isomorphic to the unique irreducible quotient of the induced representation

$$\begin{cases} \tau'|\cdot|^{\frac{n-r-1}{2}} \times \tau'|\cdot|^{\frac{n-r-3}{2}} \times \cdots \times \tau'|\cdot|^{\frac{1}{2}} \rtimes \pi & \text{if } n+r \equiv 0 \mod 2, \\ \tau'|\cdot|^{\frac{n-r-1}{2}} \times \tau'|\cdot|^{\frac{n-r-3}{2}} \times \cdots \times \tau'|\cdot|^{1} \times \mu' \rtimes \pi & \text{if } n+r \equiv 1 \mod 2, \end{cases}$$

where $\tau' = \tau \otimes \chi_{-1}^{[(n+r)/2]} = \mu' \times \mu'^{-1}$.

For the notation of parabolic inductions, see $\S3.2$ below.

(3) For any irreducible almost tempered unitary representations π_1 and π_2 , we have

$$\mathcal{M}_{\psi,\tau}^{(n)}(\pi_1) \cong \mathcal{M}_{\psi,\tau}^{(n)}(\pi_2) \implies \pi_1 \cong \pi_2.$$

- (4) Assume one of the following:
 - π is almost tempered and unitary, and $r \leq n \leq r+1$ or n > 2r;

• π is discrete series so that $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ is irreducible. Set

$$\pi' = \mathcal{M}_{\psi,\tau}^{(r)} \left(\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \right).$$

Then all irreducible subquotients of π' are isomorphic to π , and the maximal semisimple quotient of π' is irreducible.

(5) Suppose that μ is unramified, and set $\alpha = (\mu \chi_{-1}^{[(n+r)/2]})(\varpi)$, where ϖ is a uniformizer of F. If π is an irreducible unramified representation of $Sp_r(F)$ with the Satake parameter

 $\{\beta_1^{\pm 1}, \ldots, \beta_r^{\pm 1}\}$, then $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ has a unique irreducible unramified quotient. Its Satake parameter is equal to

$$\{\beta_1^{\pm 1}, \dots, \beta_r^{\pm 1}\} \cup \left\{\alpha^{\pm 1}q^{\frac{n-r-1}{2}}, \alpha^{\pm 1}q^{\frac{n-r-3}{2}}, \dots, \alpha^{\pm 1}q^{-\frac{n-r-1}{2}}\right\}$$

as multisets, where q is the cardinality of the residue field of F.

There is also a local analogue of seesaw identities (Proposition 1.5). For $\xi \in F^{\times}$, we denote the Weil representations of the Jacobi group $\widetilde{J}_{n-1}(F) = \widetilde{\operatorname{Sp}}_{n-1}(F) \ltimes V_{n-1}(F)$ and the metaplectic group $\widetilde{\operatorname{Sp}}_r(F)$ with respect to ψ_{ξ} by $\omega_{\psi_{\xi}}^{(n-1)}$ and $\omega_{\psi_{\xi}}^{(r)}$, respectively.

Proposition 1.10 (Seesaw identity (Proposition 3.10)). Let π and π' be irreducible representations of $\widetilde{\operatorname{Sp}}_r(F)$ and $\widetilde{\operatorname{Sp}}_{n-1}(F)$, on which $\{\pm 1\}$ acts by $(\pm 1)^{r+n}$ and $(\pm 1)^{r+n-1}$, respectively. Then

$$\operatorname{Hom}_{\widetilde{J}_{n-1}(F)}\left(\left.\mathcal{M}_{\psi,\tau}^{(n)}(\pi)\right|_{\widetilde{J}_{n-1}(F)},\pi'\otimes\omega_{\psi_{\xi}}^{(n-1)}\right)\cong\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_{r}(F)}\left(\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi')\otimes\omega_{\psi_{\xi}}^{(r)},\pi\right).$$

As an application, the seesaw identity gives a quite new example of the local Gan–Gross–Prasad conjecture in a non-generic case (Theorem C.5).

This paper is organized as follows. In §2, we recall the theory of Ikeda liftings extended by Ikeda–Yamana [29]. The local and global theories of Miyawaki liftings are explained in §3 and §4, respectively. In §5, we discuss Conjecture 1.6 and its relation with the Gan–Gross–Prasad conjecture. In Appendices A, B and C, we recall results on Jacquet modules of representations of metaplectic groups, the local and global Langlands program, and the Gan–Gross–Prasad conjecture, respectively.

Acknowledgments. The author is grateful to Shunsuke Yamana, Atsushi Ichino, and Tamotsu Ikeda for their helpful comments. Thanks are also due to the referees for helpful comments. This work was supported by the Foundation for Research Fellowships of Japan Society for the Promotion of Science for Young Scientists (PD) Grant 29-193.

2. Ikeda liftings and their Fourier–Jacobi coefficients

In this section, we recall the theory of Ikeda liftings along with [29].

2.1. Metaplectic group and its representations. Let F be a totally real number field. The symplectic group Sp_n is an algebraic group defined over F given by

$$\operatorname{Sp}_{n}(F) = \left\{ g \in \operatorname{GL}_{2n}(F) \mid {}^{t}g \begin{pmatrix} 0 & -\mathbf{1}_{n} \\ \mathbf{1}_{n} & 0 \end{pmatrix} g = \begin{pmatrix} 0 & -\mathbf{1}_{n} \\ \mathbf{1}_{n} & 0 \end{pmatrix} \right\}.$$

The set of symmetric matrices of size n with coefficients in F is denoted by $\operatorname{Sym}_n(F)$. For $A \in \operatorname{GL}_n(F)$ and $B \in \operatorname{Sym}_n(F)$, set

$$\mathbf{m}(A) = \begin{pmatrix} A & 0\\ 0 & {}^{t}A^{-1} \end{pmatrix}, \quad \mathbf{n}(B) = \begin{pmatrix} \mathbf{1}_{n} & B\\ 0 & \mathbf{1}_{n} \end{pmatrix} \in \operatorname{Sp}_{n}(F).$$

For each k = 1, ..., n, we define a standard maximal parabolic subgroup $P_k(F)$ of $\text{Sp}_n(F)$ by

$$P_{k}(F) = \left\{ \begin{pmatrix} a & * & * & * \\ 0 & A & * & B \\ \hline 0 & 0 & ^{t}a^{-1} & 0 \\ 0 & C & * & D \end{pmatrix} \middle| a \in \operatorname{GL}_{k}(F), \ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \operatorname{Sp}_{n-k}(F) \right\}.$$

Let $P_k(F) = M_k(F)N_k(F)$ be the standard Levi decomposition, so that the Levi $M_k(F)$ is isomorphic to $\operatorname{GL}_k(F) \times \operatorname{Sp}_{n-k}(F)$. In particular, $P_n(F) = M_n(F)N_n(F)$ is the Siegel parabolic subgroup with $M_n(F) = \{\mathbf{m}(A) \mid A \in \operatorname{GL}_n(F)\}$ and $N_n(F) = \{\mathbf{n}(B) \mid B \in \operatorname{Sym}_n(F)\}$. Let $B_n = \bigcap_{k=1}^n P_k$ be a Borel subgroup of Sp_n . A parabolic subgroup P of Sp_n is called standard if P contains B_n .

For each place v of F, we denote by $\operatorname{Sp}_n(F_v)$ the metaplectic group, i.e., the topological double cover of the symplectic group $\operatorname{Sp}_n(F_v)$. As sets, we identify $\widetilde{\operatorname{Sp}}_n(F_v)$ with $\operatorname{Sp}_n(F_v) \times \{\pm 1\}$. Then the group law of $\widetilde{\operatorname{Sp}}_n(F_v)$ is given by

$$(g_1,\zeta_1)(g_2,\zeta_2) = (g_1g_2,c_v(g_1,g_2)\zeta_1\zeta_2)$$

for $g_1, g_2 \in \text{Sp}_n(F_v)$ and $\zeta_1, \zeta_2 \in \{\pm 1\}$, where $c_v(g_1, g_2)$ is Rao's 2-cocycle of $\text{Sp}_n(F_v)$ with values in $\{\pm 1\}$. The double cover $\widetilde{\text{Sp}}_n(F_v) \to \text{Sp}_n(F_v)$ splits over the subgroup $N_n(F_v)$ by $\mathbf{n}(B) \mapsto (\mathbf{n}(B), 1)$. If F_v is a non-archimedean local field whose residue characteristic is not 2, there is a unique splitting

$$\operatorname{Sp}_n(\mathfrak{o}_v) \to \widetilde{\operatorname{Sp}}_n(F_v), \ g \mapsto (g, s(g)).$$

Here, we denote by \mathfrak{o}_v the ring of integers of F_v . We identify $N_n(F_v)$ and $\operatorname{Sp}_n(\mathfrak{o}_v)$ with the images of these splittings. If H is a subgroup of $\operatorname{Sp}_n(F)$, the inverse image of H in $\widetilde{\operatorname{Sp}}_n(F)$ is denoted by \widetilde{H} .

Next, we define the global metaplectic group. We denote the adele ring of F by \mathbb{A} . Let \mathfrak{S} be a finite set of places of F, which contains all places above 2 and ∞ . Put

$$\operatorname{Sp}_n(\mathbb{A})_{\mathfrak{S}} = \prod_{v \in \mathfrak{S}} \operatorname{Sp}_n(F_v) \times \prod_{v \notin \mathfrak{S}} \operatorname{Sp}(\mathfrak{o}_v)$$

Then the double cover $\widetilde{\operatorname{Sp}}_n(\mathbb{A})_{\mathfrak{S}} \to \operatorname{Sp}_n(\mathbb{A})_{\mathfrak{S}}$ is defined by the 2-cocycle $\prod_{v \in \mathfrak{S}} c_v(g_{1,v}, g_{2,v})$. For $\mathfrak{S}_1 \subset \mathfrak{S}_2$, there exists an embedding $\widetilde{\operatorname{Sp}}_n(\mathbb{A})_{\mathfrak{S}_1} \hookrightarrow \widetilde{\operatorname{Sp}}_n(\mathbb{A})_{\mathfrak{S}_2}$ given by

$$((g_v)_v,\zeta)\mapsto \left((g_v)_v,\zeta\prod_{v\in\mathfrak{S}_2\backslash\mathfrak{S}_1}s_v(g_v)
ight).$$

The global metaplectic group $\widetilde{\mathrm{Sp}}_n(\mathbb{A})$ is defined by the inductive limit

$$\widetilde{\operatorname{Sp}}_n(\mathbb{A}) = \varinjlim_{\mathfrak{S}} \widetilde{\operatorname{Sp}}_n(\mathbb{A})_{\mathfrak{S}},$$

where \mathfrak{S} runs over all finite sets of places of F containing all places above 2 and ∞ . The covering $\widetilde{\operatorname{Sp}}_n(\mathbb{A}) \to \operatorname{Sp}_n(\mathbb{A})$ splits over $\operatorname{Sp}_n(F)$ uniquely. We identify $\operatorname{Sp}_n(F)$ with the image of the splitting.

2.2. Weil representations of Jacobi groups. We recall the Weil representation on a Jacobi group in the local setting. Let F be a local field of characteristic zero. Fix a non-trivial unitary character ψ of F. For $\xi \in F^{\times}$, we define a new non-trivial unitary character ψ_{ξ} by

$$\psi_{\xi}(x) = \psi(\xi x)$$

for $x \in F$. Let \langle , \rangle be the quadratic Hilbert symbol. For $x, \xi \in F^{\times}$, we set $\chi_{\xi}(x) = \langle x, \xi \rangle$. For each Schwartz function $f \in \mathcal{S}(F)$, the Fourier transform \hat{f} (with respect to ψ_{ξ}) is defined by

$$\hat{f}(x) = \int_{F} f(y)\psi_{\xi}(xy)dy,$$

where dy is the self-dual Haar measure on F with respect to ψ_{ξ} . For $a \in F^{\times}$, there exists an 8-th root of unity $\alpha_{\psi_{\xi}}(a)$ such that

$$\int_{F} f(x)\psi_{\xi}(ax^{2})dx = \alpha_{\psi_{\xi}}(a)|2a|^{-\frac{1}{2}}\int_{F} \hat{f}(x)\psi_{\xi}(-\frac{x^{2}}{4a})dx$$

for any $f \in \mathcal{S}(F)$. The constant $\alpha_{\psi_{\xi}}(a)$ is called the Weil constant. It satisfies that $\alpha_{\psi_{\xi}}(ab^2) = \alpha_{\psi_{\xi}}(a)$ and

$$\frac{\alpha_{\psi_{\xi}}(a)\alpha_{\psi_{\xi}}(b)}{\alpha_{\psi_{\xi}}(1)\alpha_{\psi_{\xi}}(ab)} = \langle a, b \rangle$$

for $a, b \in F^{\times}$. In particular,

$$\left(\frac{\alpha_{\psi_{\xi}}(1)}{\alpha_{\psi_{\xi}}(a)}\right)^2 = \chi_{-1}(a),$$

where $\chi_{-1} = \langle \cdot, -1 \rangle$ is the quadratic character associated to $F(\sqrt{-1})/F$. Put

$$\mathbf{v}(x,y,z) = \begin{pmatrix} 1 & x & z & y \\ 0 & \mathbf{1}_{n-1} & {}^t y & 0 \\ \hline 0 & 0 & 1 & 0 \\ 0 & 0 & -{}^t x & \mathbf{1}_{n-1} \end{pmatrix} \in \operatorname{Sp}_n(F),$$

where $x, y \in F^{n-1}$ are row vectors and $z \in F$. We set

$$V(F) = V_{n-1}(F) = \{ \mathbf{v}(x, y, z) \mid x, y \in F^{n-1}, z \in F \}$$

$$X(F) = X_{n-1}(F) = \{ \mathbf{v}(x, 0, 0) \mid x \in F^{n-1} \},$$

$$Y(F) = Y_{n-1}(F) = \{ \mathbf{v}(0, y, 0) \mid y \in F^{n-1} \},$$

$$Z(F) = Z_{n-1}(F) = \{ \mathbf{v}(0, 0, z) \mid z \in F \}.$$

Note that V(F) is a Heisenberg group. We regard $\widetilde{\operatorname{Sp}}_{n-1}(F)$ as a subgroup of $\widetilde{\operatorname{Sp}}_n(F)$ by the embedding

$$\left(\begin{pmatrix} A & B \\ C & D \end{pmatrix}, \zeta \right) \mapsto \left(\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & A & 0 & B \\ \hline 0 & 0 & 1 & 0 \\ 0 & C & 0 & D \end{pmatrix}, \zeta \right).$$

By the Stone–von Neumann theorem, there is a unique irreducible admissible representation $\omega_{\psi_{\xi}}$ of V(F) on which the center Z(F) acts by ψ_{ξ} . This representation $\omega_{\psi_{\xi}}$ extends to the Weil representation of the group $\widetilde{J}_{n-1}(F) = V(F) \rtimes \widetilde{\mathrm{Sp}}_{n-1}(F)$. We call $J_{n-1}(F) = V(F) \rtimes \mathrm{Sp}_{n-1}(F)$ a Jacobi group. The representation $\omega_{\psi_{\xi}}$ is realized on the Schwartz space $\mathcal{S}(X(F))$ explicitly as follows:

$$\begin{split} \omega_{\psi_{\xi}}(\mathbf{v}(x,y,z))\phi(t) &= \psi_{\xi}(z+2t\cdot{}^{t}y+x\cdot{}^{t}y)\phi(t+x),\\ \omega_{\psi_{\xi}}(\mathbf{m}(A),\zeta)\phi(t) &= \zeta \frac{\alpha_{\psi_{\xi}}(1)}{\alpha_{\psi_{\xi}}(\det A)} |\det A|^{\frac{1}{2}}\phi(tA),\\ \omega_{\psi_{\xi}}(\mathbf{n}(B),\zeta)\phi(t) &= \zeta\psi_{\xi}(t\cdot B\cdot{}^{t}t)\phi(t),\\ \omega_{\psi_{\xi}}\left(\begin{pmatrix} 0 & -\mathbf{1}_{n-1}\\ \mathbf{1}_{n-1} & 0 \end{pmatrix},\zeta\right)\phi(t) &= \zeta\alpha_{\psi_{\xi}}(1)^{-n+1} |2|^{\frac{n-1}{2}} \int_{X(F)} \phi(u)\overline{\psi_{\xi}(2t\cdot{}^{t}u)}du \end{split}$$

for $\zeta \in \{\pm 1\}$, $\mathbf{v}(x, y, z) \in V(F)$, $A \in \operatorname{GL}_{n-1}(F)$, $B \in \operatorname{Sym}_{n-1}(F)$, and $\phi \in \mathcal{S}(X(F))$. Here, $du = \prod_i du_i$ is the Haar measure on X(F) with du_i being the self-dual Haar measure on F with respect to ψ_{ξ} . The Weil representation $\omega_{\psi_{\xi}}$ is unitary with respect to the inner product

$$(\phi_1, \phi_2) = \int_{X(F)} \phi_1(t) \overline{\phi_2(t)} dt$$

for $\phi_1, \phi_2 \in \mathcal{S}(X(F))$.

2.3. Non-archimedean case. In this subsection, we assume that F is non-archimedean. For a smooth representation Π of $\widetilde{J}_{n-1}(F)$, we put

$$\mathrm{FJ}_{\psi_{\xi}}(\Pi) = (\Pi \otimes \overline{\omega_{\psi_{\xi}}})_{V(F)}.$$

Here, $(\cdot)_{V(F)}$ means the maximal quotient on which V(F) acts trivially. We call $\mathrm{FJ}_{\psi_{\xi}}(\Pi)$ the Fourier–Jacobi module of Π with index ψ_{ξ} . We regard $\mathrm{FJ}_{\psi_{\xi}}(\Pi)$ as a representation of $\widetilde{\mathrm{Sp}}_{n-1}(F)$. Since V(F) is unipotent, the correspondence $\Pi \mapsto \mathrm{FJ}_{\psi_{\xi}}(\Pi)$ is an exact functor from the category of smooth representations of $\widetilde{\mathrm{Sp}}_{n-1}(F)$.

Conversely, for a smooth representation π of $\widetilde{\mathrm{Sp}}_{n-1}(F)$, one can consider the tensor product

 $\pi\otimes\omega_{\psi_{\mathcal{E}}}$

which is a smooth representation of $\widetilde{J}_{n-1}(F)$. The correspondence $\pi \mapsto \pi \otimes \omega_{\psi_{\xi}}$ is an exact functor from the category of smooth representations of $\widetilde{\operatorname{Sp}}_{n-1}(F)$ to the category of smooth representations of $\widetilde{J}_{n-1}(F)$.

The following proposition seems to be well-known, but we give a proof for the convenience of the readers. (For an archimedean analogue, see [49].)

Proposition 2.1. Suppose that F is non-archimedean. The map

 $\pi \mapsto \pi \otimes \omega_{\psi_{\xi}}$

gives a 1-1 correspondence between irreducible smooth representations of $\widetilde{\operatorname{Sp}}_{n-1}(F)$ and irreducible smooth representations of $\widetilde{J}_{n-1}(F)$ on which Z(F) acts by ψ_{ξ} . The inverse mapping is given by the Fourier-Jacobi module $\Pi \mapsto \operatorname{FJ}_{\psi_{\xi}}(\Pi)$.

Proof. Since any smooth representation of V(F) on which Z(F) acts by ψ_{ξ} is a direct sum of copies of the Weil representation $\omega_{\psi_{\xi}}$, for any (nonzero) smooth representation Π of \widetilde{J}_{n-1} , we can write $\Pi \cong \mathcal{V} \otimes \mathcal{S}(X(F))$ as a representation of V(F) for some (nonzero) vector space \mathcal{V} on which V(F)acts trivially. For $g \in \widetilde{\mathrm{Sp}}_{n-1}(F)$, the operator $\Pi(g) \circ (\mathbf{1}_{\mathcal{V}} \otimes \omega_{\psi_{\xi}}(g)^{-1})$ commutes with the action of V(F). By Schur's lemma, we have

$$\Pi(g) \circ (\mathbf{1}_{\mathcal{V}} \otimes \omega_{\psi_{\xi}}(g)^{-1}) = \pi(g) \otimes \mathbf{1}_{\mathcal{S}(X(F))}$$

for some $\pi(g) \in \operatorname{Aut}(\mathcal{V})$. Then π gives a group homomorphism $\pi \colon \widetilde{\operatorname{Sp}}_{n-1}(F) \to \operatorname{Aut}(\mathcal{V})$. It is smooth since Π and $\omega_{\psi_{\xi}}$ are smooth. We conclude that $\Pi \cong \pi \otimes \omega_{\psi_{\xi}}$ as representations of $\widetilde{J}_{n-1}(F)$. Note that $\operatorname{FJ}_{\psi_{\xi}}(\pi \otimes \omega_{\psi_{\xi}}) \cong \pi$ as representations of $\widetilde{\operatorname{Sp}}_{n-1}(F)$. In particular, $\operatorname{FJ}_{\psi_{\xi}}(\Pi) \neq 0$ for any nonzero smooth representation of $\widetilde{J}_{n-1}(F)$. Since $- \otimes \omega_{\psi_{\xi}}$ and $\operatorname{FJ}_{\psi_{\xi}}$ are exact functors, we see that π is irreducible as a representation of $\widetilde{\operatorname{Sp}}_{n-1}(F)$ if and only if $\pi \otimes \omega_{\psi_{\xi}}$ is irreducible as a representation of $\widetilde{J}_{n-1}(F)$. For an irreducible smooth representation Π of $\widetilde{\text{Sp}}_n(F)$, we denote by $\Pi_{\psi_{\xi}}$ the maximal quotient of Π on which Z(F) acts by ψ_{ξ} . Then $\Pi_{\psi_{\xi}}$ is a smooth representation of $\widetilde{J}_{n-1}(F)$. By a similar argument to the proof of Proposition 2.1, we have

$$\Pi_{\psi_{\xi}} \cong \mathrm{FJ}_{\psi_{\xi}}(\Pi | \widetilde{J}_{n-1}(F)) \otimes \omega_{\psi_{\xi}}$$

Let μ be a unitary character of F^{\times} . For each integer n, we define a character $\mu^{(n)}$ of $\widetilde{M}_n(F)$ by

$$\mu^{(n)}((\mathbf{m}(A),\zeta)) = \zeta^n \left(\frac{\alpha_{\psi}(1)}{\alpha_{\psi}(\det A)}\right)^n \mu(\det A).$$

It is also regarded as a character of $\widetilde{P}_n(F)$. If

$$\tau = \mu \times \mu^{-1} = \operatorname{Ind}_{P_{(1,1)}(F)}^{\operatorname{GL}_2(F)}(\mu \otimes \mu^{-1})$$

is an irreducible parabolic induction of $GL_2(F)$, where $P_{(1,1)}$ is the Borel subgroup of GL_2 consisting of upper triangular matrices, then the degenerate principal series

$$I_{\psi}^{(n)}(\tau) = I_{\psi}(\mu^{(n)}) = \operatorname{Ind}_{\widetilde{P}_{n}(F)}^{\widetilde{\operatorname{Sp}}_{n}(F)}(\mu^{(n)})$$

is irreducible by [35, 50]. See also [29, Propositions 3.1, 5.1].

In general, the Fourier–Jacobi module $\operatorname{FJ}_{\psi_{\xi}}(\Pi|J_{n-1}(F))$ is rarely irreducible since the restriction of Π to $\widetilde{J}_{n-1}(F)$ is often reducible. However, $\operatorname{FJ}_{\psi_{\xi}}$ sends degenerate principal series of $\widetilde{\operatorname{Sp}}_n(F)$ to ones of $\widetilde{\operatorname{Sp}}_{n-1}(F)$.

Proposition 2.2. For any $\xi \in F^{\times}$, we have an isomorphism

$$\mathrm{FJ}_{\psi_{\xi}}(I_{\psi}^{(n)}(\tau)) \cong I_{\psi}^{(n-1)}(\tau\chi_{\xi}),$$

where $\tau \chi_{\xi} = \tau \otimes \chi_{\xi}$. In particular, we have

$$\left(I_{\psi}^{(n)}(\tau)\right)_{\psi_{\xi}} \cong I_{\psi}^{(n-1)}(\tau\chi_{\xi}) \otimes \omega_{\psi_{\xi}}$$

as $\widetilde{J}_{n-1}(F)$ -modules.

Proof. See [21, Theorem 3.1] and [55].

2.4. Archimedean case. In this subsection, we consider the case where $F = \mathbb{R}$. We assume that the non-trivial unitary character ψ of \mathbb{R} is of the form

$$\psi(x) = \exp(2\pi a \sqrt{-1}x)$$

for $x \in \mathbb{R}$ with a > 0. Then the Weil constant $\alpha_{\psi}(t)$ is given by

$$\alpha_{\psi}(t) = \begin{cases} \exp(\pi\sqrt{-1}/4) & \text{if } t > 0, \\ \exp(-\pi\sqrt{-1}/4) & \text{if } t < 0. \end{cases}$$

For $\xi \in \mathbb{R}^{\times}$ with $\xi > 0$, we define $\phi_{\xi}^{0} \in \mathcal{S}(X(\mathbb{R}))$ by

$$\phi_{\xi}^{0}(x) = \exp(-2\pi a\xi(x_{1}^{2} + \dots + x_{n-1}^{2}))$$

for $x = (x_1, \dots, x_{n-1}) \in X(\mathbb{R})$. Let $K_{\infty} = \left\{ \begin{pmatrix} \alpha & \beta \\ -\beta & \alpha \end{pmatrix} \middle| \alpha, \beta \in \operatorname{Mat}_n(\mathbb{R}), \ {}^t\alpha\beta = {}^t\beta\alpha, \ {}^t\alpha\alpha + {}^t\beta\beta = \mathbf{1}_n \right\}$

be the usual maximal compact subgroup of $\operatorname{Sp}_n(\mathbb{R})$. For $u = \begin{pmatrix} \alpha & \beta \\ -\beta & \alpha \end{pmatrix} \in K_{\infty}$, we write $\det(\alpha + \sqrt{-1\beta}) = e^{\sqrt{-1}\theta}$ with $-\pi < \theta \leq \pi$, and we set

$$\det^{1/2}(u,\zeta) = \zeta e^{\sqrt{-1}\theta/2}.$$

Then $\omega_{\psi_{\xi}}(u,\zeta)\phi_{\xi}^{0} = \det^{1/2}(u,\zeta) \cdot \phi_{\xi}^{0}$ for any $(u,\zeta) \in \widetilde{K}_{\infty} \cap \widetilde{\mathrm{Sp}}_{n-1}(\mathbb{R})$. In particular, $\det^{1/2}$ is a genuine character of \widetilde{K}_{∞} . For an integer l, we set $\det^{l/2}(u,\zeta) = (\det^{1/2}(u,\zeta))^{l}$. We denote the irreducible lowest weight representation of $\widetilde{\mathrm{Sp}}_{n}(\mathbb{R})$ with lowest \widetilde{K}_{∞} -type $\det^{l/2}$ by $\mathcal{D}_{l/2}^{(n)}$.

Let

$$\mathfrak{H}_n = \{ Z \in \operatorname{Mat}_n(\mathbb{C}) \mid {}^t Z = Z, \ \operatorname{Im}(Z) > 0 \}$$

be the Siegel upper half space of genus n. Here, for a symmetric matrix B, we write B > 0 if B is positive definite. Then $\text{Sp}_n(\mathbb{R})$ acts on \mathfrak{H}_n by

$$g(Z) = (AZ + B)(CZ + D)^{-1}, \quad g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \operatorname{Sp}_n(\mathbb{R}).$$

Note that the stabilizer of $\mathbf{i} = \sqrt{-1} \cdot \mathbf{1}_n \in \mathfrak{H}_n$ in $\operatorname{Sp}_n(\mathbb{R})$ is equal to K_{∞} . We set $j(g, Z) = \det(CZ + D)$. Then there exists a unique automorphy factor $\tilde{j}((g,\zeta), Z)$ of $\widetilde{\operatorname{Sp}}_n(\mathbb{R})$ such that $\tilde{j}((g,\zeta), Z)^2 = j(g, Z)$ for any $(g,\zeta) \in \widetilde{\operatorname{Sp}}_n(\mathbb{R})$ and $Z \in \mathfrak{H}_n$. The following lemma might be well-known, but we give a proof for the convenience of the readers.

Lemma 2.3. For $A \in GL_n(\mathbb{R})$ with det A > 0, $B \in Sym_n(\mathbb{R})$, and $(u, \zeta) \in K_\infty$, we have

$$\widetilde{j}(\mathbf{n}(B)(\mathbf{m}(A),1)(u,\zeta),\mathbf{i}) = (\det A)^{-\frac{1}{2}} \det^{-1/2}(u,\zeta)$$

Proof. Since $\tilde{j}((g,\zeta), Z)$ is an automorphy factor of $\widetilde{\operatorname{Sp}}_n(\mathbb{R})$, for any $(g,\zeta) \in \widetilde{\operatorname{Sp}}_n(\mathbb{R})$, the function

$$\mathfrak{H}_n \to \mathbb{C}, \ Z \mapsto \widetilde{j}((g,\zeta),Z)$$

is holomorphic.

First, we claim that $\tilde{j}(\mathbf{n}(B), Z) = 1$ for any $B \in \text{Sym}_n(\mathbb{R})$ and $Z \in \mathfrak{H}_n$. Note that $\tilde{j}(\mathbf{n}(B), Z)^2 = j(\mathbf{n}(B), Z) = 1$. Since \mathfrak{H}_n is connected, $\tilde{j}(\mathbf{n}(B), Z)$ is independent of Z. In particular, $B \mapsto \tilde{j}(\mathbf{n}(B), Z)$ gives a group homomorphism $\text{Sym}_n(\mathbb{R}) \to \{\pm 1\}$. Since $\text{Sym}_n(\mathbb{R})$ is divisible, it must be the trivial character. Hence $\tilde{j}(\mathbf{n}(B), Z) = 1$.

Next, we claim that $\tilde{j}((\mathbf{m}(A), 1), Z) = (\det A)^{-1/2}$ for $A \in \operatorname{GL}_2(\mathbb{R})$ with $\det A > 0$ and $Z \in \mathfrak{H}_n$. Since \mathfrak{H}_n is connected, the function $\mathfrak{H}_n \ni Z \mapsto (\det A)^{1/2} \tilde{j}((\mathbf{m}(A), 1), Z) \in \{\pm 1\}$ is a constant for any $A \in \operatorname{GL}_2(\mathbb{R})$ with $\det A > 0$. In particular, the map $A \mapsto (\det A)^{1/2} \tilde{j}((\mathbf{m}(A), 1), Z) \in \{\pm 1\}$ gives a group homomorphism. Since $(e^{(1/2)X})^2 = e^X$ for $X \in \mathfrak{gl}_n(\mathbb{R})$, we have $(\det A)^{1/2} \tilde{j}((\mathbf{m}(A), 1), Z) = 1$ when $A = e^X$ for some $X \in \mathfrak{gl}_n(\mathbb{R})$. Since any $A \in \operatorname{GL}_n(\mathbb{R})$ with $\det A > 0$ can be written as a product $A = e^{X_1} \cdots e^{X_k}$ for some $X_1, \ldots, X_k \in \mathfrak{gl}_n(\mathbb{R})$, we have $(\det A)^{1/2} \tilde{j}((\mathbf{m}(A), 1), Z) = 1$ for any $A \in \operatorname{GL}_2(\mathbb{R})$ with $\det A > 0$.

Similarly, $\tilde{j}((\mathbf{1}_{2n}, -1), Z) \in \{\pm 1\}$ is independent of $Z \in \mathfrak{H}_n$. Choose $A \in \mathcal{O}(n)$ such that $A^2 = \mathbf{1}_n$ and det A = -1. Then $(\mathbf{m}(A), 1)^2 = (\mathbf{1}_{2n}, -1)$ so that $\tilde{j}((\mathbf{1}_{2n}, -1), \mathbf{i}) = \tilde{j}((\mathbf{m}(A), 1), \mathbf{i})^2 = j(\mathbf{m}(A), \mathbf{i}) = \det(A)^{-1} = -1$. Hence $\tilde{j}((\mathbf{1}_{2n}, -1), Z) = -1$ for any $Z \in \mathfrak{H}_n$.

Finally, we see that $\det^{1/2}(u,\zeta)\tilde{j}((u,\zeta),\mathbf{i}) \in \{\pm 1\}$ gives a group homomorphism $\widetilde{K}_{\infty} \to \{\pm 1\}$, which factors through K_{∞} . Since K_{∞} is divisible, it must be the trivial character. Hence $\tilde{j}((u,\zeta),\mathbf{i}) = \det^{-1/2}(u,\zeta)$. This completes the proof. Let $\operatorname{Sym}_n^+(\mathbb{R}) = \{B \in \operatorname{Sym}_n(\mathbb{R}) \mid B > 0\}$. Fix an integer k > 0. For $B \in \operatorname{Sym}_n^+(\mathbb{R})$, we define a function W_B^0 on $\widetilde{\operatorname{Sp}}_n(\mathbb{R})$ by

$$W_B^0(g) = (\det B)^{(2k+n)/4} \psi(\operatorname{Tr}(Bg(\mathbf{i}))) \tilde{j}(g, \mathbf{i})^{-(2k+n)} = \psi(\operatorname{Tr}(Bz)) \det(B[A])^{(2k+n)/4} \exp(-2\pi a \cdot \operatorname{Tr}(B[A])) \det^{k+(n/2)}(\tilde{u})$$

for $g = \mathbf{n}(z)(\mathbf{m}(A), 1)\widetilde{u} \in \widetilde{\mathrm{Sp}}_n(\mathbb{R})$ with $z \in \mathrm{Sym}_n(\mathbb{R})$, $A \in \mathrm{GL}_n(\mathbb{R})$, $\det A > 0$, and $\widetilde{u} \in \widetilde{K}_{\infty}$. Here, we put $B[A] = {}^tABA$.

Proposition 2.4 ([29, Lemma 7.6]). For $\xi \in \mathbb{R}^{\times}$ with $\xi > 0$ and $B' \in \text{Sym}_{n-1}^{+}(\mathbb{R})$, we put

$$B = \begin{pmatrix} \xi & 0\\ 0 & B' \end{pmatrix} \in \operatorname{Sym}_n^+(\mathbb{R})$$

Then we have

$$\int_{X(\mathbb{R})} W_B^0(\mathbf{v}(x,0,0)g') \overline{\omega_{\psi_{\xi}}(g')\phi_{\xi}^0(x)} dx = |\det B|^{1/4} e^{-2\pi a\xi} W_{B'}^0(g')$$

for $g' \in \widetilde{\mathrm{Sp}}_{n-1}(\mathbb{R})$.

The Lie algebras $\mathfrak{sp}_n(\mathbb{R})$ and \mathfrak{k} of $\mathrm{Sp}_n(\mathbb{R})$ and K_∞ are given by

$$\mathfrak{sp}_{n}(\mathbb{R}) = \left\{ X \in \operatorname{Mat}_{2n}(\mathbb{R}) \middle| {}^{t}X \begin{pmatrix} 0 & -\mathbf{1}_{n} \\ \mathbf{1}_{n} & 0 \end{pmatrix} + {}^{t}X \begin{pmatrix} 0 & -\mathbf{1}_{n} \\ \mathbf{1}_{n} & 0 \end{pmatrix} X = 0 \right\}$$
$$= \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \operatorname{Mat}_{2n}(\mathbb{R}) \middle| B = {}^{t}B, \ C = {}^{t}C, \ A = -{}^{t}D \right\},$$
$$\mathfrak{k} = \left\{ \begin{pmatrix} A & B \\ -B & A \end{pmatrix} \in \operatorname{Mat}_{2n}(\mathbb{R}) \middle| {}^{t}A = -A, \ {}^{t}B = B \right\},$$

respectively. Note that \mathfrak{k} is the 1-eigenspace of the Cartan involution $\theta X = -^t X$ on $\mathfrak{sp}_n(\mathbb{R})$. The (-1)-eigenspace is given by

$$\mathfrak{p} = \left\{ \begin{pmatrix} A & B \\ B & -A \end{pmatrix} \in \operatorname{Mat}_{2n}(\mathbb{R}) \mid A = {}^{t}A, B = {}^{t}B \right\}.$$

Hence $\mathfrak{sp}_n(\mathbb{R}) = \mathfrak{k} \oplus \mathfrak{p}$. The homeomorphism

$$\rho: \operatorname{Sp}_n(\mathbb{R})/K_\infty \to \mathfrak{H}_n, \ g \mapsto g(\mathbf{i})$$

induces an isomorphism

$$d\rho\colon \mathfrak{sp}_n(\mathbb{R})/\mathfrak{k}=\mathfrak{p}\xrightarrow{\sim} T_{\mathbf{i}}\mathfrak{H}_n\cong \mathrm{Sym}_n(\mathbb{C})$$

where $T_{\mathbf{i}}\mathfrak{H}_n$ is the tangent space at \mathbf{i} on \mathfrak{H}_n . This map is given by

$$d\rho\left(\begin{pmatrix} A & B\\ B & -A \end{pmatrix}\right) = 2(B + \sqrt{-1}A)$$

Then the complex structure (i.e., multiplication with $\sqrt{-1}$) on $\operatorname{Sym}_n(\mathbb{C})$ gives a map

$$J: \mathfrak{p} \to \mathfrak{p}, \ \begin{pmatrix} A & B \\ B & -A \end{pmatrix} \mapsto \begin{pmatrix} B & -A \\ -A & -B \end{pmatrix}.$$

Let $\mathfrak{sp}_n(\mathbb{C})$, $\mathfrak{k}_{\mathbb{C}}$, and $\mathfrak{p}_{\mathbb{C}}$ be the complexifications of $\mathfrak{sp}_n(\mathbb{R})$, \mathfrak{k} , and \mathfrak{p} , respectively. We denote the $(\pm \sqrt{-1})$ -eigenspace of J on $\mathfrak{p}_{\mathbb{C}}$ by $\mathfrak{p}_{\mathbb{C}}^{\pm}$. Then

$$\mathfrak{p}_{\mathbb{C}}^{\pm} = \left\{ \begin{pmatrix} A & B \\ B & -A \end{pmatrix} \otimes 1 \pm \begin{pmatrix} -B & A \\ A & B \end{pmatrix} \otimes \sqrt{-1} \mid A, B \in \operatorname{Sym}_{n}(\mathbb{R}) \right\}$$

$$= \left\{ \begin{pmatrix} A & \pm \sqrt{-1}A \\ \pm \sqrt{-1}A & -A \end{pmatrix} \in \operatorname{Mat}_{2n}(\mathbb{C}) \ \middle| \ A \in \operatorname{Sym}_{n}(\mathbb{C}) \right\}.$$

The elements of $\mathfrak{p}^+_{\mathbb{C}}$ and $\mathfrak{p}^-_{\mathbb{C}}$ correspond to the linear combinations of differential operators

$$\left\{\frac{d}{dz_j}\right\}_j$$
 and $\left\{\frac{d}{d\overline{z_j}}\right\}_j$,

respectively, where $z_j = x_j + \sqrt{-1}y_j$ are coordinates on \mathfrak{H}_n at the point **i**.

The element $X \in \mathfrak{p}$ acts on a smooth function W on $\mathrm{Sp}_n(\mathbb{R})$ by

$$X \cdot W(g) = \left. \frac{d}{dt} \right|_{t=0} W(g(e^{tX}, 1))$$

for $g \in \widetilde{\operatorname{Sp}}_n(\mathbb{R})$. This action is extended to $\mathfrak{p}_{\mathbb{C}}$ linearly. It is easy to check that $\mathfrak{p}_{\mathbb{C}}^-$ acts on W_B^0 by zero for any $B \in \operatorname{Sym}_n^+(\mathbb{R})$ (cf. [4, Lemma 7]). Hence W_B^0 is a lowest weight vector in $\mathcal{D}_{k+(n/2)}^{(n)}$. Similarly, $\mathfrak{p}_{\mathbb{C}}^- \cap \mathfrak{sp}_{n-1}(\mathbb{C})$ acts on ϕ_{ξ}^0 by zero. This fact can be proven by using the Fock model of $\omega_{\psi_{\xi}}$.

2.5. Global case. Now we let F be a totally real number field, and ψ be a non-trivial unitary character of \mathbb{A}/F . We assume that for each infinite place v of F, there exists $a_v \in F_v \cong \mathbb{R}$ with $a_v > 0$ such that $\psi(x_v) = \exp(2\pi a_v \sqrt{-1}x_v)$ for $x_v \in F_v$. For a place v of F, we define a maximal compact subgroup K_v of $\operatorname{Sp}_n(F_v)$ by

$$K_v = \begin{cases} \operatorname{Sp}_n(\mathfrak{o}_v) & \text{if } v \text{ is non-archimedean,} \\ K_\infty & \text{if } v \text{ is real.} \end{cases}$$

Here, we denote by \mathfrak{o}_v the ring of integers of F_v when v is non-archimedean. If v is archimedean, we define $(\mathfrak{p}_v)^-_{\mathbb{C}} \subset \operatorname{Lie}(\operatorname{Sp}_n(F_v)) \otimes_{\mathbb{R}} \mathbb{C}$ as in the previous subsection.

Recall that a function $\varphi \colon \mathrm{Sp}_n(F) \backslash \mathrm{Sp}_n(\mathbb{A}) \to \mathbb{C}$ is a cusp form if

- φ is smooth and of moderate growth;
- φ is right *K*-finite, where $K = \prod_{v} K_{v}$;
- φ is \mathfrak{z} -finite, where \mathfrak{z} is the center of the universal enveloping algebra of $\operatorname{Lie}(\operatorname{Sp}_n(F \otimes_{\mathbb{O}} \mathbb{R})) \otimes_{\mathbb{R}} \mathbb{C}$;
- there exists $\delta \in \{0,1\}$ such that $\varphi((g,\zeta)) = \zeta^{\delta} \varphi((g,1))$ for any $g \in \operatorname{Sp}_n(\mathbb{A})$;
- For any proper F-parabolic subgroup P of Sp_n , the constant term along P

$$\int_{N(F)\setminus N(\mathbb{A})}\varphi(ug)du$$

is zero for any $g \in \widetilde{\mathrm{Sp}}_n(\mathbb{A})$, where N is the unipotent radical of P.

We say that φ is genuine if $\delta = 1$. Let $l = (l_v)_v \in \prod_{v \mid \infty} \mathbb{Z}$. We say that a cusp form φ is holomorphic of weight l/2 if

- $X_v \cdot \varphi = 0$ for any $X_v \in (\mathfrak{p}_v)_{\mathbb{C}}^-$;
- $\varphi(g\widetilde{u}_v) = \det^{l_v/2}(\widetilde{u}_v)\varphi(g)$ for $g \in \widetilde{\mathrm{Sp}}_n(\mathbb{A})$ and $\widetilde{u}_v \in \widetilde{K}_v$

for any infinite place v. We denote the space of holomorphic cusp forms of weight l/2 by

$$\mathcal{S}_{l/2}\left(\operatorname{Sp}_n(F)\backslash \widetilde{\operatorname{Sp}}_n(\mathbb{A})\right).$$

The group $\widetilde{\operatorname{Sp}}_n(\mathbb{A}_{\operatorname{fin}})$ acts on $\mathcal{S}_{l/2}\left(\operatorname{Sp}_n(F)\backslash \widetilde{\operatorname{Sp}}_n(\mathbb{A})\right)$ by the right translation.

Let $\tau = \otimes'_v \tau_v$ be an irreducible unitary cuspidal automorphic representation of $\operatorname{GL}_2(\mathbb{A})$. Assume that

- (A1) for any finite place v, the local factor τ_v is an irreducible principal representation $\mu_v \times \mu_v^{-1}$;
- (A2) for any infinite place v, the local factor τ_v is a discrete series representation with lowest weight $\pm 2k_v$ where $k_v > 0$;
- (A3) the root number

$$\varepsilon(\tau) = \varepsilon\left(\frac{1}{2}, \tau, \psi\right) = \prod_{v < \infty} \mu_v(-1) \cdot \prod_{v \mid \infty} (-1)^{k_v}$$

is equal to 1.

For $v < \infty$, we notice that $|\mu_v(\varpi_v)|_v = 1$ by the Ramanujan conjecture proven in the case of Hilbert modular forms by Blasius [8], where ϖ_v is a uniformizer of F_v . For a more general result on the Ramanujan conjecture, see [10]. We refer to $\pm 2k = (\pm 2k_v)_v \in \prod_{v \mid \infty} \mathbb{Z}$ as the weight of $\otimes_{v \mid \infty} \tau_v$.

Put $k + (n/2) = (k_v + n/2)_v \in \prod_{v \mid \infty} \mathbb{Z}$ for each integer n > 0. For $v < \infty$, we let $I_{\psi_v}^{(n)}(\tau_v) = \operatorname{Ind}_{\widetilde{P}_n(F_v)}^{\widetilde{\operatorname{Sp}}_n(F_v)}(\mu_v^{(n)})$ be the degenerate principal series defined in §2.3. Set $I_{\psi}^{(n)}(\tau) = \otimes_{v < \infty}' I_{\psi_v}^{(n)}(\tau_v)$, which is an irreducible representation of $\widetilde{\operatorname{Sp}}_n(\mathbb{A}_{\operatorname{fin}})$. The following is a part of the main theorem in [29].

Theorem 2.5 ([29, Theorems 1.1, 1.2]). Let $\tau = \bigotimes_{v}' \tau_{v}$ be an irreducible unitary cuspidal automorphic representation of $\operatorname{GL}_{2}(\mathbb{A})$ satisfying the conditions (A1), (A2) and (A3). Then the representation $I_{\psi}^{(n)}(\tau)$ occurs in $\mathcal{S}_{k+(n/2)}\left(\operatorname{Sp}_{n}(F)\backslash\widetilde{\operatorname{Sp}}_{n}(\mathbb{A})\right)$ with multiplicity one.

We denote the unique subrepresentation of $\mathcal{S}_{k+(n/2)}\left(\operatorname{Sp}_n(F)\setminus\widetilde{\operatorname{Sp}}_n(\mathbb{A})\right)$ which is isomorphic to $I_{\psi}^{(n)}(\tau)$ by $\operatorname{Ik}_{\psi}^{(n)}(\tau)$, and call it the Ikeda lift of τ .

Let $\xi \in F^{\times}$ be a totally positive element. For $\phi \in \mathcal{S}(X(\mathbb{A}))$, the theta function $\Theta_{\psi_{\xi}}^{\phi}(vg')$ is defined by

$$\Theta_{\psi_{\xi}}^{\phi}(\mathbf{v}(x,y,z)g') = \sum_{t \in X(F)} \psi_{\xi}(z+2t \cdot {}^{t}y+x \cdot {}^{t}y)\omega_{\psi_{\xi}}(g')\phi(t+x)$$

for $\mathbf{v}(x, y, z) \in V(\mathbb{A})$ and $g' \in \widetilde{\mathrm{Sp}}_{n-1}(\mathbb{A})$. It is a genuine automorphic form on $\widetilde{J}_{n-1}(\mathbb{A})$. One can easily check that

$$\int_{Z(\mathbb{A})V(F)\setminus V(\mathbb{A})} \Theta_{\psi_{\xi}}^{\phi_{1}}(vg') \overline{\Theta_{\psi_{\xi}}^{\phi_{2}}(vg')} dv = (\phi_{1}, \phi_{2}) = \int_{X(\mathbb{A})} \phi_{1}(x) \overline{\phi_{2}(x)} dx$$

for $\phi_1, \phi_2 \in \mathcal{S}(X(\mathbb{A}))$ and $g' \in \operatorname{Sp}_{n-1}(\mathbb{A})$.

We denote by $\mathcal{S}(X(\mathbb{A}))_{\xi}$ the subspace of $\mathcal{S}(X(\mathbb{A}))$ spanned by $\phi = \bigotimes_{v} \phi_{v}$ such that $\phi_{v} = \phi_{\xi_{v}}^{0}$ for each infinite place v. For $\varphi \in \mathcal{S}_{l/2}\left(\operatorname{Sp}_{n}(F) \setminus \widetilde{\operatorname{Sp}}_{n}(\mathbb{A})\right)$ and $\phi \in \mathcal{S}(X(\mathbb{A}))_{\xi}$, the Fourier–Jacobi coefficient associated to (φ, ϕ) is defined by

$$\mathrm{FJ}_{\psi_{\xi}}^{\phi}(g';\varphi) = \int_{V(F) \setminus V(\mathbb{A})} \varphi(vg') \overline{\Theta_{\psi_{\xi}}^{\phi}(vg')} dv.$$

This is a cusp form on $\text{Sp}_{n-1}(\mathbb{A})$. (See also [19, Lemma 2.3] and [18, Theorem 8].) Moreover, by Proposition 2.4, we conclude that

$$\mathrm{FJ}_{\psi_{\xi}}^{\phi}(\varphi) \in \mathcal{S}_{(l-1)/2}\left(\mathrm{Sp}_{n-1}(F) \setminus \widetilde{\mathrm{Sp}}_{n-1}(\mathbb{A})\right)$$

for any $\varphi \in \mathcal{S}_{l/2}\left(\operatorname{Sp}_n(F) \setminus \widetilde{\operatorname{Sp}}_n(\mathbb{A})\right)$ and $\phi \in \mathcal{S}(X(\mathbb{A}))_{\xi}$.

Proposition 2.6. For $\mathcal{F} \in \text{Ik}_{\psi}^{(n)}(\tau)$ and $\phi \in \mathcal{S}(X(\mathbb{A}))_{\xi}$, we have $\text{FJ}_{\psi_{\xi}}^{\phi}(\mathcal{F}) \in \text{Ik}_{\psi}^{(n-1)}(\tau\chi_{\xi})$.

Proof. We may assume that $\mathrm{FJ}_{\psi_{\xi}}^{\phi}(\mathcal{F}) \neq 0$. Let Π be the representation of $\widetilde{\mathrm{Sp}}_{n-1}(\mathbb{A})$ generated by $\mathrm{FJ}_{\psi_{\xi}}^{\phi}(\mathcal{F})$ for $\mathcal{F} \in \mathrm{Ik}_{\psi}^{(n)}(\tau)$ and $\phi \in \mathcal{S}(X(\mathbb{A}))_{\xi}$. Since Π is cuspidal, it is a direct sum of irreducible representations. Choose an irreducible direct summand π of Π and a projection $\Pi \twoheadrightarrow \pi$. Then the map

$$\operatorname{Ik}_{\psi}^{(n)}(\tau) \otimes \overline{\omega_{\psi_{\xi}}} \xrightarrow{\operatorname{FJ}_{\psi_{\xi}}} \Pi \twoheadrightarrow \pi$$

is a (nonzero) $V(\mathbb{A}_{\text{fin}})$ -invariant map so that it factors through $\operatorname{FJ}_{\psi_{\xi}}(\operatorname{Ik}_{\psi}^{(n)}(\tau))$. Since $\operatorname{FJ}_{\psi_{\xi}}(\operatorname{Ik}_{\psi}^{(n)}(\tau))$ is nonzero, we have $\operatorname{FJ}_{\psi_{\xi}}(\operatorname{Ik}_{\psi}^{(n)}(\tau)) = \operatorname{Ik}_{\psi}^{(n-1)}(\tau\chi_{\xi})$ by Proposition 2.2. Since it is irreducible, we have $\pi \cong \operatorname{Ik}_{\psi}^{(n-1)}(\tau\chi_{\xi})$. Hence Π is isomorphic to a direct sum of some copies of $\operatorname{Ik}_{\psi}^{(n-1)}(\tau\chi_{\xi})$. However the Ikeda lift appears in $\mathcal{S}_{k+(n-1)/2}\left(\operatorname{Sp}_{n-1}(F)\setminus\widetilde{\operatorname{Sp}}_{n-1}(\mathbb{A})\right)$ with multiplicity one, we see that Π is irreducible and is equal to $\operatorname{Ik}_{\psi}^{(n-1)}(\tau\chi_{\xi})$. \Box

For $\varphi \in \mathcal{S}_{l/2}\left(\operatorname{Sp}_n(F) \setminus \widetilde{\operatorname{Sp}}_n(\mathbb{A})\right)$, we define the ξ -th Fourier–Jacobi coefficient $\varphi_{\psi_{\xi}}$ by

$$\varphi_{\psi_{\xi}}(vg') = \int_{Z(F)\backslash Z(\mathbb{A})} \varphi(zvg') \overline{\psi_{\xi}(z)} dz$$

for $v \in V(\mathbb{A})$ and $g' \in \widetilde{\mathrm{Sp}}_{n-1}(\mathbb{A})$. If Π be a subspace of $\mathcal{S}_{l/2}\left(\mathrm{Sp}_n(F) \setminus \widetilde{\mathrm{Sp}}_n(\mathbb{A})\right)$, we put $\Pi_{\psi_{\mathcal{E}}} = \left\{\varphi_{\psi_{\mathcal{E}}} \mid \varphi \in \Pi\right\}$.

This is the maximal quotient of Π on which $Z(\mathbb{A}_{fin})$ acts by ψ_{ξ} .

Proposition 2.7. (1) For $\mathcal{F} \in \text{Ik}_{\psi}^{(n)}(\tau)$, there exist $\mathcal{F}'_1, \ldots, \mathcal{F}'_r \in \text{Ik}_{\psi}^{(n-1)}(\tau\chi_{\xi})$ and $\phi_1, \ldots, \phi_r \in \mathcal{S}(X(\mathbb{A}))_{\xi}$ such that

$$\mathcal{F}_{\psi_{\xi}}(vg') = \sum_{i=1}^{r} \mathcal{F}'_{i}(g') \Theta_{\psi_{\xi}}^{\phi_{i}}(vg')$$

for $v \in V(\mathbb{A})$ and $g' \in \widetilde{\mathrm{Sp}}_{n-1}(\mathbb{A})$.

(2) Suppose that $\left(\operatorname{Ik}_{\psi}^{(n)}(\tau)\right)_{\psi_{\xi}}$ is nonzero. Then for $\mathcal{F}' \in \operatorname{Ik}_{\psi}^{(n-1)}(\tau\chi_{\xi})$ and $\phi \in \mathcal{S}(X(\mathbb{A}))_{\xi}$, there exists $\mathcal{F} \in \operatorname{Ik}_{\psi}^{(n)}(\tau)$ such that

$$\mathcal{F}_{\psi_{\xi}}(vg') = \mathcal{F}'(g')\Theta^{\phi}_{\psi_{\xi}}(vg')$$

for $v \in V(\mathbb{A})$ and $g' \in \widetilde{\operatorname{Sp}}_{n-1}(\mathbb{A})$. (3) If $n \ge 2$, then $\left(\operatorname{Ik}_{\psi}^{(n)}(\tau)\right)_{\psi_{\xi}} \neq 0$ for any totally positive $\xi \in F^{\times}$.

Proof. By [26, Proposition 1.3], for $\mathcal{F} \in \text{Ik}_{\psi}^{(n)}(\tau)$, there exist $\phi_1, \ldots, \phi_r \in \mathcal{S}(X(\mathbb{A}))_{\xi}$ with $(\phi_i, \phi_j) = \delta_{i,j}$ such that

$$\mathcal{F}_{\psi_{\xi}}(vg') = \sum_{i=1}^{r} \mathrm{FJ}_{\psi_{\xi}}^{\phi_{i}}(\mathcal{F})(g') \cdot \Theta_{\psi_{\xi}}^{\phi_{i}}(vg')$$

for $v \in V(\mathbb{A})$ and $g' \in \widetilde{\operatorname{Sp}}_{n-1}(\mathbb{A})$. Since $\operatorname{FJ}_{\psi_{\xi}}^{\phi_i}(\mathcal{F})(g') \in \operatorname{Ik}_{\psi}^{(n-1)}(\tau\chi_{\xi})$ by Proposition 2.6, we obtain (1).

Therefore, there is a $\widetilde{J}_{n-1}(\mathbb{A}_{\mathrm{fin}})\text{-equivariant inclusion}$

$$\left(\mathrm{Ik}_{\psi}^{(n)}(\tau)\right)_{\psi_{\xi}} \subset \mathrm{Ik}_{\psi}^{(n-1)}(\tau\chi_{\xi}) \otimes \omega_{\psi_{\xi}}.$$

If $\left(\operatorname{Ik}_{\psi}^{(n)}(\tau)\right)_{\psi_{\xi}}$ is nonzero, this inclusion must be equal since $\operatorname{Ik}_{\psi}^{(n-1)}(\tau\chi_{\xi})\otimes\omega_{\psi_{\xi}}$ is irreducible. Hence we obtain (2).

Let $\operatorname{Sym}_n^+(F)$ be the subset of $\operatorname{Sym}_n(F)$ consisting of symmetric matrices whose images in $\operatorname{Sym}_n(F_v)$ are positive definite for each infinite place v. By [29, Lemmas 8.5, 5.4], there exists $\mathcal{F} \in \operatorname{Ik}_{\psi}^{(n)}(\tau)$ such that for $B \in \operatorname{Sym}_n^+(F)$, the *B*-th Fourier coefficient

$$W_B(g) = \int_{\operatorname{Sym}_n(F) \setminus \operatorname{Sym}_n(\mathbb{A})} \mathcal{F}(\mathbf{n}(X)g) \overline{\psi(\operatorname{Tr}(BX))} dX$$

is not identically zero if and only if $L(1/2, \tau \otimes \chi_{\det(B)}) \neq 0$. When $B = \operatorname{diag}(\xi, \xi', 1, \ldots, 1)$ with $\xi' \in F^{\times}$ totally positive, the *B*-th Fourier coefficient W_B is given by

$$W_B(g) = \int_{Z(\mathbb{A})\operatorname{Sym}_n(F)\backslash\operatorname{Sym}_n(\mathbb{A})} \mathcal{F}_{\psi_{\xi}}(\mathbf{n}(X)g)\overline{\psi(\operatorname{Tr}(BX))} dX.$$

Hence the ξ -th Fourier–Jacobi coefficient $\mathcal{F}_{\psi_{\xi}}$ is not identically zero if $W_B \neq 0$. Since there exists a totally positive element $\xi' \in F^{\times}$ such that $L(1/2, \tau \otimes \chi_{\xi\xi'}) \neq 0$ by [53, Théorème 4], we obtain (3).

3. Local Miyawaki liftings

In this section, we define the local Miyawaki lifting, and prove basic properties.

3.1. **Definition.** Let F be a non-archimedean local field of characteristic zero, and ψ be a non-trivial additive character of F. For two non-negative integers n and r, we define an embedding $\iota: \operatorname{Sp}_n \times \operatorname{Sp}_r \hookrightarrow \operatorname{Sp}_{n+r}$ by

$$\iota\left(\begin{pmatrix}A_1 & B_1\\C_1 & D_1\end{pmatrix}, \begin{pmatrix}A_2 & B_2\\C_2 & D_2\end{pmatrix}\right) = \begin{pmatrix}A_1 & 0 & B_1 & 0\\0 & A_2 & 0 & B_2\\\hline C_1 & 0 & D_1 & 0\\0 & C_2 & 0 & D_2\end{pmatrix},$$

and we identify $\operatorname{Sp}_n \times \operatorname{Sp}_r$ with the image. For $\tau = \mu \times \mu^{-1}$ with μ being a unitary character of F^{\times} , we consider the local Ikeda lift $I_{\psi}^{(n+r)}(\tau) = \operatorname{Ind}_{\widetilde{P}_{n+r}(F)}^{\widetilde{\operatorname{Sp}}_{n+r}(F)}(\mu^{(n+r)})$, which is a degenerate principal series of $\widetilde{\operatorname{Sp}}_{n+r}(F)$. For an irreducible representation π of $\widetilde{\operatorname{Sp}}_r(F)$, on which the kernel $\{\pm 1\}$ of the covering map $\widetilde{\operatorname{Sp}}_r(F) \twoheadrightarrow \operatorname{Sp}_r(F)$ acts by $(\pm 1)^{n+r}$, the maximal π -isotypic quotient of $I_{\psi}^{(n+r)}(\tau)$ is of the form

$$\mathcal{M}^{(n)}_{\psi,\tau}(\pi) \boxtimes \pi$$

for some smooth representation $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ of $\widetilde{\operatorname{Sp}}_n(F)$, on which the kernel $\{\pm 1\}$ of the covering map $\widetilde{\operatorname{Sp}}_n(F) \twoheadrightarrow \operatorname{Sp}_n(F)$ acts by $(\pm 1)^{n+r}$. We call $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ the local Miyawaki lift of π .

3.2. **Preliminary.** In this subsection, we recall some basic terminologies of representations of $\widetilde{\operatorname{Sp}}_r(F)$.

A parabolic induction

$$\tau_1|\cdot|^{s_1}\times\cdots\times\tau_l|\cdot|^{s_l}\rtimes\pi_0\coloneqq\operatorname{Ind}_{P(F)}^{\operatorname{Sp}_r(F)}(\tau_1|\cdot|^{s_1}\otimes\cdots\otimes\tau_l|\cdot|^{s_l}\otimes\pi_0),$$

is standard if

- P = MN is a standard parabolic subgroup of Sp_r with $M \cong \operatorname{GL}_{k_1} \times \cdots \times \operatorname{GL}_{k_l} \times \operatorname{Sp}_{r_0}$;
- τ_i (resp. π_0) is an irreducible tempered representation of $\operatorname{GL}_{k_i}(F)$ (resp. $\operatorname{Sp}_{r_0}(F)$);
- s_1, \ldots, s_l are real numbers such that $s_1 > \cdots > s_l > 0$.

We understand that each irreducible tempered representation of $\operatorname{Sp}_r(F)$ is a standard module (the case where l = 0 and $r_0 = r$). The Langlands classification asserts that any irreducible smooth representation π of $\operatorname{Sp}_r(F)$ is a unique irreducible quotient of a standard module $\tau_1 |\cdot|^{s_1} \times \cdots \times \tau_l |\cdot|^{s_l} \rtimes \pi_0$, which is called the Langlands quotient and is denoted by $J(\tau_1 |\cdot|^{s_1}, \ldots, \tau_l |\cdot|^{s_l}, \pi_0)$. For any irreducible representation π , the datum $(P, \{\tau_i |\cdot|^{s_i}\}, \pi_0)$ is determined uniquely up to an isomorphism.

When π is a genuine irreducible representation of $\widetilde{\mathrm{Sp}}_r$, one should replace the parabolic induction with

$$\operatorname{Ind}_{\widetilde{P}(F)}^{\widetilde{\operatorname{Sp}}_r(F)}(\tau_{1,\psi}|\cdot|^{s_1}\otimes\cdots\otimes\tau_{l,\psi}|\cdot|^{s_l}\otimes\pi_0),$$

where

$$\tau_{i,\psi}(a,\zeta) = \zeta \frac{\alpha_{\psi}(1)}{\alpha_{\psi}(\det a)} \tau_i(a)$$

is a genuine irreducible representation of the double cover $\widetilde{\operatorname{GL}}_{k_i}(F)$ of $\operatorname{GL}_{k_i}(F)$ (which is identified with $\operatorname{GL}_{k_i}(F) \times \{\pm 1\}$ as sets), and π_0 is a genuine irreducible representation of $\widetilde{\operatorname{Sp}}_{r_0}(F)$. For more precision, see [15]. We identify τ_{ψ} with τ itself, and we use the same notation $\tau_1 |\cdot|^{s_1} \times \cdots \times \tau_l |\cdot|^{s_l} \rtimes \pi_0$ and $J(\tau_1 |\cdot|^{s_1}, \ldots, \tau_l |\cdot|^{s_l}, \pi_0)$ as in the non-genuine case.

We say that an irreducible representation π of $\operatorname{Sp}_r(F)$ is almost tempered if $\pi = J(\tau_1|\cdot|^{s_1},\ldots,\tau_l|\cdot|^{s_l},\pi_0)$ with $0 < s_l < \cdots < s_1 < 1/2$. We understand that irreducible tempered representations are almost tempered (the case where l = 0). Then the standard module $\tau_1|\cdot|^{s_1} \times \cdots \times \tau_l|\cdot|^{s_l} \rtimes \pi_0$ is irreducible by Corollary B.3, so that π is equal to this standard module.

When we consider unramified representations, we always assume that the residue characteristic of F is greater than 2. Then the covering map $\widetilde{\operatorname{Sp}}_r(F) \to \operatorname{Sp}_r(F)$ splits over the maximal compact subgroup $\operatorname{Sp}_r(\mathfrak{o})$, where \mathfrak{o} is the ring of integers of F. Recall that an irreducible smooth representation π of $\widetilde{\operatorname{Sp}}_r(F)$ is unramified if π has a nonzero $\operatorname{Sp}_r(\mathfrak{o})$ -fixed vector. If π is unramified, then there exist unramified characters χ_1, \ldots, χ_r such that π is a unique unramified constituent of the induced representation $\operatorname{Ind}_{\widetilde{B}_r(F)}^{\widetilde{\operatorname{Sp}}_r(F)}(\chi_1 \otimes \cdots \otimes \chi_r)$. We call the multiset $\{\chi_1(\varpi)^{\pm 1}, \ldots, \chi_r(\varpi)^{\pm 1}\}$ the Satake parameter of π , where ϖ is a uniformizer of F. We write $q = |\varpi|^{-1}$ for the cardinality of the residue field of F.

Recall that we can associate an irreducible representation π of $\widetilde{\operatorname{Sp}}_r(F)$ to an *L*-parameter ϕ , which is a self-dual representation of the Weil–Deligne group $WD_F = W_F \times \operatorname{SL}_2(\mathbb{C})$ of *F*. This is symplectic (resp. orthogonal) if π is genuine (resp. not genuine). More precisely, see Appendix B. We say that an *L*-parameter ϕ is of good parity if ϕ is a sum of irreducible self-dual representations of the same type as ϕ . The unique irreducible algebraic representation of $\operatorname{SL}_2(\mathbb{C})$ of dimension *d* is denoted by S_d . 3.3. Main local theorem. Recall that $\tau = \mu \times \mu^{-1}$ with μ being a unitary character of F^{\times} . Let χ_{-1} be the quadratic character associated to the extension $F(\sqrt{-1})/F$. For a real number a, we denote by [a] the maximal integer which is not greater than a. The determinant character of $GL_t(F)$ is denoted by det_t .

The following is the local main theorem.

Theorem 3.1. Let μ be a unitary character of F^{\times} , and π be an irreducible representation of $\widetilde{\operatorname{Sp}}_r(F)$ on which $\{\pm 1\}$ acts by $(\pm 1)^{n+r}$. Suppose that $n \ge r$.

- The local Miyawaki lift M⁽ⁿ⁾_{ψ,τ}(π) is nonzero and of finite length.
 If π is almost tempered and unitary, then
 - - $\mathcal{M}_{\psi.\tau}^{(n)}(\pi)$ is irreducible;
 - $\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \cong (\mu' \circ \det_{n-r}) \rtimes \pi \text{ with } \mu' = \mu \chi_{-1}^{[(n+r)/2]};$
 - $\mathcal{M}^{(n)}_{\psi_{\tau}}(\pi)$ is isomorphic to the unique irreducible quotient of the induced representation

$$\begin{cases} \tau'|\cdot|^{\frac{n-r-1}{2}} \times \tau'|\cdot|^{\frac{n-r-3}{2}} \times \cdots \times \tau'|\cdot|^{\frac{1}{2}} \rtimes \pi & \text{if } n+r \equiv 0 \mod 2, \\ \tau'|\cdot|^{\frac{n-r-1}{2}} \times \tau'|\cdot|^{\frac{n-r-3}{2}} \times \cdots \times \tau'|\cdot|^{1} \times \mu' \rtimes \pi & \text{if } n+r \equiv 1 \mod 2, \end{cases}$$

where $\tau' = \tau \otimes \chi_{-1}^{[(n+r)/2]} = \mu' \times \mu'^{-1}$.

(3) For any irreducible almost tempered unitary representations π_1 and π_2 , we have

$$\mathcal{M}_{\psi,\tau}^{(n)}(\pi_1) \cong \mathcal{M}_{\psi,\tau}^{(n)}(\pi_2) \implies \pi_1 \cong \pi_2.$$

(4) Suppose that π is almost tempered and unitary so that $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ is irreducible by (2). Assume further that one of the following conditions holds:

(a) The L-parameter ϕ does not contain $\mu^{\pm 1}S_d$ for any $d \ge n - r$ with $d \equiv n - r \mod 2$; (b) n = r or n = r + 1.

Set

$$\pi' = \mathcal{M}_{\psi,\tau}^{(r)} \left(\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \right).$$

Then all irreducible subquotients of π' are isomorphic to π , and the maximal semisimple quotient of π' is irreducible.

(5) Suppose that μ is unramified, and set $\alpha = (\mu \chi_{-1}^{[(n+r)/2]})(\varpi)$. If π is an irreducible unramified representation of $\widetilde{\operatorname{Sp}}_r(F)$ with the Satake parameter $\{\beta_1^{\pm 1}, \ldots, \beta_r^{\pm 1}\}$, then $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ has a unique irreducible unramified quotient. Its Satake parameter is equal to

$$\left\{\beta_1^{\pm 1}, \dots, \beta_r^{\pm 1}\right\} \cup \left\{\alpha^{\pm 1} q^{\frac{n-r-1}{2}}, \alpha^{\pm 1} q^{\frac{n-r-3}{2}}, \dots, \alpha^{\pm 1} q^{-\frac{n-r-1}{2}}\right\}$$

as multisets.

As in Remark 3.9 (1) below, the condition (a) of Theorem 3.1 (4) holds when n > 2r or π is discrete series.

The assertion (2) gives the Langlands data for $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ explicitly. In particular, it deduces (3). The proof of (5) for the non-genuine case (i.e., the case where n + r is even) is [28, Proposition 3.1]. The genuine case is proven similarly.

We prove Theorem 3.1 (1), (2) and (4) in §3.5, §3.6 and §3.7, respectively.

3.4. Miyawaki liftings and degenerate induced representations. In this subsection, we show that $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ is of finite length, and that for almost tempered π , there is a surjection

$$(\mu' \circ \det_{n-r}) \rtimes \pi \twoheadrightarrow \mathcal{M}^{(n)}_{\psi,\tau}(\pi).$$

We need the following lemma (see [36] and [16, Lemma 2.2]).

Lemma 3.2. Suppose that $n \ge r$. Set $\mu' = \mu \chi_{-1}^{[(n+r)/2]}$. The local Ikeda lift (the degenerate principal series) $I_{\psi}^{(n+r)}(\tau) = \operatorname{Ind}_{\widetilde{P}_{n+r}(F)}^{\widetilde{\operatorname{Sp}}_{n+r}(F)}(\mu^{(n+r)})$ has an $\widetilde{\operatorname{Sp}}_{n}(F) \times \widetilde{\operatorname{Sp}}_{r}(F)$ -equivariant filtration

$$0 \subset I_0 \subset I_1 \subset \cdots \subset I_r = I_{\psi}^{(n+r)}(\tau)$$

with successive quotients

$$R_t = I_t / I_{t-1} = \operatorname{Ind}_{\widetilde{P}_{t+(n-r)}(F) \times \widetilde{P}_t(F)}^{\widetilde{Sp}_n(F) \times \widetilde{Sp}_r(F)} \left(\left(\mu' |\det_{t+(n-r)}|^{\frac{t}{2}} \boxtimes \mu' |\det_t|^{\frac{t+(n-r)}{2}} \right) \otimes C_c^{\infty} \left(\widetilde{Sp}_{r-t}(F) \right) \right).$$

Here the induction is normalized, and $\widetilde{\operatorname{Sp}}_{r-t}(F) \times \widetilde{\operatorname{Sp}}_{r-t}(F)$ acts on $C_c^{\infty}\left(\widetilde{\operatorname{Sp}}_{r-t}(F)\right)$ by

$$((g_1,g_2)\varphi)(x) = \varphi(g_1^{-1} \cdot x \cdot \alpha g_2 \alpha^{-1}) \quad with \quad \alpha = \begin{pmatrix} \mathbf{1}_{r-t} & 0\\ 0 & -\mathbf{1}_{r-t} \end{pmatrix}.$$

In particular,

$$R_0 = \operatorname{Ind}_{\widetilde{P}_{n-r}(F) \times \widetilde{\operatorname{Sp}}_r(F)}^{\widetilde{\operatorname{Sp}}_n(F) \times \widetilde{\operatorname{Sp}}_r(F)} \left(\mu' \circ \det_{n-r} \otimes C_c^{\infty} \left(\widetilde{\operatorname{Sp}}_r(F) \right) \right).$$

Using this lemma, we have the following.

Proposition 3.3. Let π and π' be smooth representations of $\widetilde{\operatorname{Sp}}_r(F)$ and $\widetilde{\operatorname{Sp}}_n(F)$, respectively. Set $\mu' = \mu \chi_{-1}^{[(n+r)/2]}$. Suppose that π is irreducible, almost tempered and that

$$\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_n(F)\times\widetilde{\operatorname{Sp}}_r(F)}\left(I_{\psi}^{(n+r)}(\tau),\pi'\boxtimes\pi\right)\neq 0.$$

Then we have

$$\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_n(F)}\left((\mu'\circ \det_{n-r})\rtimes \pi,\pi'\right)\neq 0.$$

Moreover, if there is an $\widetilde{\operatorname{Sp}}_n(F) \times \widetilde{\operatorname{Sp}}_r(F)$ -equivariant surjection

$$I_{\psi}^{(n+r)}(\tau) \twoheadrightarrow \pi' \boxtimes \pi,$$

then there is an $\widetilde{\operatorname{Sp}}_n(F)$ -equivariant surjection

$$(\mu' \circ \det_{n-r}) \rtimes \pi \twoheadrightarrow \pi'.$$

Proof. By Lemma 3.2, for some $0 \leq t \leq r$, we must have $\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_n(F) \times \widetilde{\operatorname{Sp}}_r(F)}(R_t, \pi' \boxtimes \pi) \neq 0$. By Bernstein's Frobenius reciprocity (see e.g., [9]), this Hom-space is isomorphic to the space of $(\widetilde{M}_{t+(n-r)}(F) \times \widetilde{M}_t(F))$ -equivariant maps

$$\left(\mu'|\det_{t+(n-r)}|^{\frac{t}{2}}\boxtimes\mu'|\det_t|^{\frac{t+(n-r)}{2}}\right)\otimes C_c^{\infty}\left(\widetilde{\operatorname{Sp}}_{r-t}(F)\right)\to R_{\overline{P_{t+(n-r)}(F)}}(\pi')\boxtimes R_{\overline{P_t(F)}}(\pi),$$

where $R_{\overline{P_t(F)}}(\pi)$ is the normalized Jacquet module of π with respect to (the double cover of) the opposite parabolic subgroup $\overline{P_t(F)}$ to $P_t(F)$. First, we assume that t > 0. By taking the contragredient, we have

$$\operatorname{Hom}_{\widetilde{\operatorname{GL}}_t(F)}\left(R_{P_t(F)}(\pi^{\vee}), \mu'^{-1} |\operatorname{det}_t|^{\frac{-t-(n-r)}{2}}\right) \neq 0.$$

Since π^{\vee} is almost tempered, we must have -t - (n - r) > -1. This contradicts $n \ge r$ and t > 0. Hence we must have t = 0, and

$$\operatorname{Hom}_{\widetilde{M}_{n-r}(F)\times\widetilde{\operatorname{Sp}}_{r}(F)}\left(\mu'\circ\operatorname{det}_{n-r}\otimes C_{c}^{\infty}\left(\widetilde{\operatorname{Sp}}_{r}(F)\right), R_{\overline{P_{n-r}(F)}}(\pi')\boxtimes\pi\right)\neq 0.$$

Taking care of the action of $\widetilde{\operatorname{Sp}}_r(F) \times \widetilde{\operatorname{Sp}}_r(F)$ on $C_c^{\infty}\left(\widetilde{\operatorname{Sp}}_r(F)\right)$, we see that this Hom-space is isomorphic to

$$\operatorname{Hom}_{\widetilde{M}_{n-r}(F)}\left(\mu'\circ\operatorname{det}_{n-r}\otimes\pi,R_{\overline{P_{n-r}(F)}}(\pi')\right).$$

Using Bernstein's Frobenius reciprocity, we obtain the first assertion.

Suppose that there is an $\operatorname{Sp}_n(F) \times \operatorname{Sp}_r(F)$ -equivariant surjection $I_r \twoheadrightarrow \pi' \boxtimes \pi$. Then the image of I_0 is of the form $\pi'_0 \boxtimes \pi$ for some $\widetilde{\operatorname{Sp}}_n(F)$ -subrepresentation π'_0 of π' , and it induces a surjection $I_r/I_0 \twoheadrightarrow (\pi'/\pi'_0) \boxtimes \pi$. The above argument implies $(\pi'/\pi'_0) = 0$ so that the restriction gives an $\widetilde{\operatorname{Sp}}_n(F) \times \widetilde{\operatorname{Sp}}_r(F)$ -equivariant surjection $I_0 \twoheadrightarrow \pi' \boxtimes \pi$. This induces an $\widetilde{\operatorname{Sp}}_n(F)$ -equivariant surjection $(\mu' \circ \det_{n-r}) \rtimes \pi \twoheadrightarrow \pi'$.

Let π and π' be smooth representations of $\widetilde{\operatorname{Sp}}_r(F)$ and $\widetilde{\operatorname{Sp}}_n(F)$, respectively. Suppose that π is irreducible, and that there exists a surjection

$$I_{\psi}^{(n+r)}(\tau) \twoheadrightarrow \pi' \boxtimes \pi.$$

Then Lemma 3.2 gives a filtration

$$0 \subset \pi'_0 \subset \pi'_1 \subset \cdots \subset \pi'_r = \pi'.$$

More precisely, the restriction of the surjection to $I_t \subset I_{\psi}^{(n+r)}(\tau)$ defines a subrepresentation π'_t of π' so that the image of I_t is $\pi'_t \boxtimes \pi$. By a similar argument to Proposition 3.3, one can show that the successive quotients π'_t/π'_{t-1} are of finite length. Therefore π' must also be of finite length.

3.5. Non-vanishing of local Miyawaki liftings. Next, we show Theorem 3.1 (1). We have seen that $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ is of finite length. We show that $\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \neq 0$. In fact, we will prove that

$$\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_n(F)\times\widetilde{\operatorname{Sp}}_r(F)}\left(I_{\psi}^{(n+r)}(\tau),\left((\mu'\circ \det_{n-r})\rtimes\pi\right)\boxtimes\pi\right)\neq 0$$

for any irreducible representation π of $\widetilde{\mathrm{Sp}}_r(F)$.

Using

$$\operatorname{Sp}_{2r}(F) \hookrightarrow \operatorname{Sp}_{n+r}(F), \ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \mapsto \begin{pmatrix} \mathbf{1}_{n-r} & & \\ & A & B \\ \hline & & \\ & & C & D \end{pmatrix}$$

we regard $\widetilde{\operatorname{Sp}}_{2r}(F)$ as a subgroup of $\widetilde{\operatorname{Sp}}_{n+r}(F)$. Then for $\Phi \in I_{\psi}^{(n+r)}(\tau)$, we have

$$\Phi|_{\widetilde{\operatorname{Sp}}_{2r}(F)} \in \operatorname{Ind}_{\widetilde{P}_{2r}(F)}^{\widetilde{\operatorname{Sp}}_{2r}(F)} \left(\mu \chi_{-1}^{[\frac{n+r}{2}]} |\det_{2r}|^{\frac{n-r}{2}} \right).$$

By the theory of the doubling method [38], [43], a doubling zeta integral gives a nonzero element

$$\mathcal{Z}_{r,r} \in \operatorname{Hom}_{\widetilde{\operatorname{Sp}}_{r}(F) \times \widetilde{\operatorname{Sp}}_{r}(F)} \left(\operatorname{Ind}_{\widetilde{P}_{2r}(F)}^{\widetilde{\operatorname{Sp}}_{2r}(F)} \left(\mu \chi_{-1}^{\left[\frac{n+r}{2}\right]} |\det_{2r}|^{\frac{n-r}{2}} \right) \otimes \left(\pi \boxtimes \pi \right), \mathbb{C} \right).$$

Note that the embedding $\iota: \operatorname{Sp}_r(F) \times \operatorname{Sp}_r(F) \hookrightarrow \operatorname{Sp}_{2r}(F)$ is not the usual one in this theory (see [38]). For $\Phi \in I_{\psi}^{(n+r)}(\tau), f \in (\mu^{-1}\chi_{-1}^{[-(n+r)/2]} \circ \operatorname{det}_{n-r}) \rtimes \pi$, and $v \in \pi$, we consider the integral

$$\mathcal{Z}_{r,n}(\Phi, f, v) = \int_{P_{n-r}(F) \setminus \operatorname{Sp}_n(F)} \mathcal{Z}_{r,r}\left((\iota(g, 1), \zeta) \Phi|_{\widetilde{\operatorname{Sp}}_{2r}(F)} \otimes f(g, \zeta) \otimes v \right) dg$$

Proposition 3.4. The integral $\mathcal{Z}_{r,n}(\Phi, f, v)$ is well-defined and gives a nonzero element in

$$\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_{n}(F)\times\widetilde{\operatorname{Sp}}_{r}(F)}\left(I_{\psi}^{(n+r)}(\tau)\otimes\left(\left(\left(\mu^{-1}\chi_{-1}^{[-(n+r)/2]}\circ\operatorname{det}_{n-r}\right)\rtimes\pi\right)\boxtimes\pi\right),\mathbb{C}\right).$$

Proof. Since

$$h \cdot \iota(u, 1) \cdot h^{-1} = \begin{pmatrix} \mathbf{1}_{n-r} & * & * & * \\ 0 & \mathbf{1}_{2r} & * & 0 \\ \hline 0 & 0 & \mathbf{1}_{n-r} & 0 \\ 0 & 0 & * & \mathbf{1}_{2r} \end{pmatrix}$$

for $u \in N_{n-r}(F) \subset \operatorname{Sp}_n(F)$ and $h \in \widetilde{\operatorname{Sp}}_{2r}(F)$, we have $\Phi(h \cdot (\iota(ug, 1), \zeta)) = \Phi(h \cdot (\iota(g, 1), \zeta))$. Similarly, we have

$$\begin{split} \Phi\left(h\cdot \left(\iota \left(\left(\begin{array}{c|c|c} a & 0 & 0 & 0\\ 0 & \mathbf{1}_r & 0 & 0\\ \hline 0 & 0 & ta^{-1} & 0\\ 0 & 0 & 0 & \mathbf{1}_r\end{array}\right)g, 1\right), \zeta\right)\right) \otimes f\left(\left(\begin{array}{c|c|c} a & 0 & 0 & 0\\ \hline 0 & \mathbf{1}_r & 0 & 0\\ \hline 0 & 0 & ta^{-1} & 0\\ \hline 0 & 0 & 0 & \mathbf{1}_r\end{array}\right)g, \zeta\right) \\ &= |\det a|^{\frac{n+r+1}{2}} \Phi(h\cdot(\iota(g,1),\zeta)) \otimes f(g,\zeta) \end{split}$$

for $a \in \operatorname{GL}_{n-r}(F)$. Hence we have

$$\begin{aligned} \mathcal{Z}_{r,r}\left((\iota(pg,1),\zeta)\Phi|_{\widetilde{\operatorname{Sp}}_{2r}(F)}\otimes f(pg,\zeta)\otimes v\right)\\ &=\delta_{P_{n-r}}(p)\cdot\mathcal{Z}_{r,r}\left((\iota(g,1),\zeta)\Phi|_{\widetilde{\operatorname{Sp}}_{2r}(F)}\otimes f(g,\zeta)\otimes v\right)\end{aligned}$$

for $p \in P_{n-r}(F) \subset \operatorname{Sp}_n(F)$, where $\delta_{P_{n-r}}$ is the modulus character of $P_{n-r}(F)$. This shows that $\mathcal{Z}_{r,n}(\Phi, f, v)$ is well-defined.

It is easy to see that

$$\mathcal{Z}_{r,n} \in \operatorname{Hom}_{\widetilde{\operatorname{Sp}}_{n}(F) \times \widetilde{\operatorname{Sp}}_{r}(F)} \left(I_{\psi}^{(n+r)}(\tau) \otimes \left(\left(\left(\mu^{-1} \chi_{-1}^{[-(n+r)/2]} \circ \det_{n-r} \right) \rtimes \pi \right) \otimes \pi \right), \mathbb{C} \right).$$

Since

$$I_{\psi}^{(n+r)}(\tau) \to \operatorname{Ind}_{\widetilde{P}_{2r}(F)}^{\widetilde{\operatorname{Sp}}_{2r}(F)} \left(\mu \chi_{-1}^{[\frac{n+r}{2}]} |\det_{2r}|^{\frac{n-r}{2}} \right), \ \Phi \mapsto \Phi|_{\widetilde{\operatorname{Sp}}_{2r}(F)}$$

is surjective, we see that $\mathcal{Z}_{r,n} \neq 0$.

Applying Proposition 3.4 for π^{\vee} , we conclude that

$$\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_n(F)\times\widetilde{\operatorname{Sp}}_r(F)}(I_{\psi}^{(n+r)}(\tau),((\mu'\circ\det_{n-r})\rtimes\pi)\boxtimes\pi)\neq 0$$

since $((\mu' \circ \det_{n-r}) \rtimes \pi)^{\vee} \cong (\mu^{-1}\chi_{-1}^{[-(n+r)/2]} \circ \det_{n-r}) \rtimes \pi^{\vee}.$

3.6. Irreducibility of degenerate induced representations. To show Theorem 3.1 (2), we need to prove the irreducibility of the induced representation $(\mu \circ \det_{n-r}) \rtimes \pi$, where π is an irreducible almost tempered unitary representation of $\widetilde{\text{Sp}}_r(F)$, and μ is a unitary character of F^{\times} . When π is supercuspidal and $\delta = 0$, the irreducibility of $(\mu \circ \det_{n-r}) \rtimes \pi$ was proven by Tadić [52, Theorem 9.1], and its Langlands data was given by Jantzen [31]. We imitate their proofs.

We first show the irreducibility of other induced representations. Let π be an irreducible representation of $\widetilde{\text{Sp}}_r(F)$. We set $\delta = 1$ if π is genuine, and $\delta = 0$ otherwise. For a smooth representation Π of $\widetilde{\text{Sp}}_n(F)$, we write s.s.(Π) for the semisimplification of Π .

Proposition 3.5. Let μ be a unitary character of F^{\times} , and π be an irreducible tempered representation of $\widetilde{\operatorname{Sp}}_r(F)$. Suppose that the L-parameter ϕ for π is of good parity (see §3.2). Then for $l \geq k-\delta$ with $l \equiv k - \delta \mod 2$, the induced representation

$$\mu^{-1} |\det_k|^{\frac{l}{2}} \rtimes \pi$$

is irreducible.

Proof. We prove the proposition by induction on k. When k = 1 and l > 0, this is Corollary B.4. When k = 1 and l = 0 so that $\delta = 1$, the irreducibility of $\mu^{-1} \rtimes \pi$ follows from Theorem B.1 (6).

Suppose that k > 1 and that $\mu^{-1} |\det_k|^{\frac{l}{2}} \rtimes \pi$ is reducible. Let $\sigma_1, \ldots, \sigma_t$ be irreducible representations of $\widetilde{\operatorname{Sp}}_{k+r}(F)$ such that $\operatorname{s.s.}(\mu^{-1} |\det_k|^{\frac{l}{2}} \rtimes \pi) = \sigma_1 \oplus \cdots \oplus \sigma_t$, and that σ_1 is a submodule and σ_t is a quotient of $\mu^{-1} |\det_k|^{\frac{l}{2}} \rtimes \pi$:

$$\sigma_1 \hookrightarrow \mu^{-1} |\det_k|^{\frac{l}{2}} \rtimes \pi \twoheadrightarrow \sigma_t.$$

Since the Langlands quotient appears in the standard module with multiplicity one as a subquotients, we have $\sigma_i \not\cong \sigma_t$ for $i \neq t$.

By Proposition A.4, we have

$$s.s.R_{P_1(F)}(\mu^{-1}|\det_k|^{\frac{l}{2}} \rtimes \pi) = \mu|\cdot|^{-\frac{l+k-1}{2}} \boxtimes \left(\mu^{-1}|\det_{k-1}|^{\frac{l-1}{2}} \rtimes \pi\right)$$
$$\oplus \mu^{-1}|\cdot|^{\frac{l-k+1}{2}} \boxtimes \left(\mu^{-1}|\det_{k-1}|^{\frac{l+1}{2}} \rtimes \pi\right)$$
$$\oplus s.s.\left(\bigoplus_{\lambda} \chi_{\lambda}|\cdot|^{\alpha_{\lambda}} \boxtimes \left(\mu^{-1}|\det_k|^{\frac{l}{2}} \rtimes \pi_{\lambda}\right)\right),$$

where s.s. $R_{P_1(F)}(\pi) = \bigoplus_{\lambda} \chi_{\lambda} |\cdot|^{\alpha_{\lambda}} \boxtimes \pi_{\lambda}$ with χ_{λ} unitary and $\alpha_{\lambda} \in \mathbb{R}$. By the induction hypothesis, we see that $\mu^{-1} |\det_{k-1}|^{(l\pm 1)/2} \rtimes \pi$ is irreducible. By Corollary B.5, we have $2\alpha_{\lambda} \in \mathbb{Z}$ and $2\alpha_{\lambda} \equiv l - k \mod 2$. In particular, the first two summands appear in s.s. $R_{P_1(F)}(\mu^{-1} |\det_k|^{\frac{l}{2}} \rtimes \pi)$ with multiplicity one.

Since
$$\sigma_1 \hookrightarrow \mu^{-1} |\det_k|^{\frac{l}{2}} \rtimes \pi$$
 and $\sigma_t \hookrightarrow \mu |\det_k|^{-\frac{l}{2}} \rtimes \pi$, we have
 $\operatorname{s.s.} R_{P_1(F)}(\sigma_i) \supset \mu^{-1} |\cdot|^{\frac{l-k+1}{2}} \boxtimes \left(\mu^{-1} |\det_{k-1}|^{\frac{l+1}{2}} \rtimes \pi\right) \iff i = 1$
 $\operatorname{s.s.} R_{P_1(F)}(\sigma_i) \supset \mu |\cdot|^{-\frac{l+k-1}{2}} \boxtimes \left(\mu^{-1} |\det_{k-1}|^{\frac{l-1}{2}} \rtimes \pi\right) \iff i = t.$

On the other hand, by Proposition A.4, we have

s.s.
$$R_{P_k(F)}(\mu^{-1}|\det_k|^{\frac{l}{2}} \rtimes \pi) \supset$$
 s.s. $\left(\mu|\det_{k-1}|^{-\frac{l+1}{2}} \times \mu^{-1}| \cdot |^{\frac{l-k+1}{2}}\right) \boxtimes \pi.$

Note that $\mu |\det_{k-1}|^{-\frac{l+1}{2}} = \langle \mu; -(l+k-1)/2, \dots, (-l+k-3)/2 \rangle$ and $\mu^{-1}|\cdot|^{\frac{l-k+1}{2}} = \langle \mu^{-1}; (l-k+1)/2 \rangle$ in the notation in Appendix A.1. Unless $\delta = 1$, l = k - 1 and $\mu^{-1} = \mu$, the representation

 $\mu |\det_{k-1}|^{-\frac{l+1}{2}} \times \mu^{-1}| \cdot |^{\frac{l-k+1}{2}}$ is irreducible by Theorem A.2. When it is irreducible, there exists *i* such that

s.s.
$$R_{P_k(F)}(\sigma_i) \supset \left(\mu |\det_{k-1}|^{-\frac{l+1}{2}} \times \mu^{-1}| \cdot |^{\frac{l-k+1}{2}}\right) \boxtimes \pi$$

Considering the Jacquet module with respect to $P_k \cap P_1$, we see that

s.s.
$$R_{P_1(F)}(\sigma_i) \supset \mu^{-1}| \cdot |^{\frac{l-k+1}{2}} \boxtimes \left(\mu^{-1} |\det_{k-1}|^{\frac{l+1}{2}} \rtimes \pi \right) \oplus \mu| \cdot |^{-\frac{l+k-1}{2}} \boxtimes \left(\mu^{-1} |\det_{k-1}|^{\frac{l-1}{2}} \rtimes \pi \right).$$

This implies that i = 1 and i = t. We obtain a contradiction.

Similarly, when $k \geq 3$, we have

s.s.
$$R_{P_k(F)}(\mu^{-1}|\det_k|^{\frac{l}{2}} \rtimes \pi) \supset \left(\mu|\det_{k-2}|^{-\frac{l+2}{2}} \times \mu^{-1}|\det_2|^{\frac{l-k+2}{2}}\right) \boxtimes \pi.$$

The right hand side is irreducible by Theorem A.2 since $\mu |\det_{k-2}|^{-\frac{l+2}{2}} = \langle \mu; -(l+k-1)/2, \dots, (-l+k-5)/2 \rangle$ and $\mu^{-1} |\det_2|^{\frac{l-k+2}{2}} = \langle \mu^{-1}; (l-k+1)/2, (l-k+3)/2 \rangle$. By the same argument, we obtain a contradiction.

Therefore, the proposition is reduced to the case where $\delta = 1$, k = l + 1 = 2 and $\mu^{-1} = \mu$. We treat this case in the following lemma.

Lemma 3.6. Let π be an irreducible genuine tempered representation of $\widetilde{\operatorname{Sp}}_r(F)$ so that $\delta = 1$. Suppose that μ is a quadratic character, and that the L-parameter ϕ for π is of good parity. Then the induced representation

$$\mu |\det_2|^{\frac{1}{2}} \rtimes \pi$$

is irreducible.

Proof. First, we show that $(\mu \circ \det_3) \rtimes \pi$ is irreducible in this case. By Theorem A.2 and Corollary B.4, we have

$$\begin{aligned} \tau|\cdot|\times\mu\rtimes\pi\cong\mu|\cdot|\times\mu|\cdot|\times\mu\rtimes\pi\\ \twoheadrightarrow\mu|\cdot|\times\mu|\mathrm{det}_2|^{\frac{1}{2}}\rtimes\pi\\ \cong\mu|\mathrm{det}_2|^{\frac{1}{2}}\times\mu|\cdot|\rtimes\pi\\ \cong\mu|\mathrm{det}_2|^{\frac{1}{2}}\times\mu|\cdot|^{-1}\rtimes\pi\\ \twoheadrightarrow(\mu\circ\mathrm{det}_3)\rtimes\pi. \end{aligned}$$

Since π is unitary, $(\mu \circ \det_3) \rtimes \pi$ is semisimple. Therefore we deduce that $(\mu \circ \det_3) \rtimes \pi \cong J(\tau | \cdot |, \mu \rtimes \pi)$.

Now we start to prove the lemma. Suppose for the sake of contradiction that $\mu |\det_2|^{\frac{1}{2}} \rtimes \pi$ is reducible. Take an irreducible submodule of σ , and an irreducible quotient σ' of $\mu |\det_2|^{\frac{1}{2}} \rtimes \pi$:

$$\sigma \hookrightarrow \mu |\det_2|^{\frac{1}{2}} \rtimes \pi \twoheadrightarrow \sigma'.$$

Then $\sigma \neq \sigma'$. Moreover, we have

$$\mu|\cdot|^1 \rtimes \sigma \hookrightarrow \mu|\cdot|^1 \times \mu|\det_2|^{\frac{1}{2}} \rtimes \pi \twoheadrightarrow \mu|\cdot|^1 \rtimes \sigma'.$$

Since $\tau |\cdot| \times \mu \rtimes \pi \twoheadrightarrow \mu |\cdot|^1 \times \mu |\det_2|^{\frac{1}{2}} \rtimes \pi$, we have $\mu |\cdot|^1 \rtimes \sigma' \twoheadrightarrow (\mu \circ \det_3) \rtimes \pi$. Moreover, since any standard module has its Langlands quotient with multiplicity one as a subquotient, we have

$$(\mu \circ \det_3) \rtimes \pi \not\subset \text{s.s.}(\mu | \cdot |^1 \rtimes \sigma).$$

Using proposition A.4, we compute the Jacquet modules $\mathrm{s.s.}R_{P_1(F)}(\mu|\cdot|^1\times\mu|\mathrm{det}_2|^{\frac{1}{2}}\rtimes\pi)$ and s.s. $R_{P_1(F)}((\mu \circ \det_3) \rtimes \pi)$. Then the sums of all irreducible representations of the form $\mu | \cdot |^{-1} \boxtimes \Sigma$ which appear in s.s. $R_{P_1(F)}(\mu|\cdot|^1 \times \mu |\det_2|^{\frac{1}{2}} \rtimes \pi)$ and s.s. $R_{P_1(F)}((\mu \circ \det_3) \rtimes \pi)$ are isomorphic to

s.s.
$$\left(\mu |\cdot|^{-1} \boxtimes \mu |\det_2|^{\frac{1}{2}} \rtimes \pi\right) \oplus$$
 s.s. $\left(\mu |\cdot|^{-1} \boxtimes (\mu |\cdot|^1 \times \mu) \rtimes \pi\right)$

and

s.s.
$$\left(\mu |\cdot|^{-1} \boxtimes \mu |\det_2|^{-\frac{1}{2}} \rtimes \pi\right) \oplus$$
 s.s. $\left(\mu |\cdot|^{-1} \boxtimes \mu |\det_2|^{\frac{1}{2}} \rtimes \pi\right)$,

respectively. Hence the difference is

$$\mu|\cdot|^{-1}\boxtimes\mu|\cdot|^{\frac{1}{2}}\operatorname{St}_{2}\rtimes\pi,$$

where $\mu |\cdot|^{\frac{1}{2}} St_2 = \langle \mu; 1, 0 \rangle$ is a Steinberg representation. By Proposition B.2, we see that $\mu |\cdot|^{\frac{1}{2}} St_2 \rtimes \pi$ is irreducible. Since

s.s. $R_{P_1(F)}(\mu|\cdot|^1 \rtimes \sigma) \supset \mu|\cdot|^{-1} \boxtimes \sigma$,

we deduce that $\sigma = \mu |\cdot|^{\frac{1}{2}} \operatorname{St}_2 \rtimes \pi$. However, we have

$$\mathrm{s.s.}R_{P_1(F)}(\mu|\cdot|^{\frac{1}{2}}\mathrm{St}_2\rtimes\pi)\supset\mu|\cdot|^{1}\boxtimes\mu\rtimes\pi\not\subset\mathrm{s.s.}R_{P_1(F)}(\mu|\mathrm{det}_2|^{\frac{1}{2}}\rtimes\pi).$$

Therefore, $\mu |\cdot|^{\frac{1}{2}} \operatorname{St}_2 \rtimes \pi \not\subset \mu |\det_2|^{\frac{1}{2}} \rtimes \pi$, which is a contradiction.

Now we show the following.

Theorem 3.7. Let π be an irreducible almost tempered representation of $\widetilde{\operatorname{Sp}}_r(F)$, and μ be a unitary character of F^{\times} . Set $\tau = \mu \times \mu^{-1}$. Then there exists a surjection

$$\begin{cases} \tau |\cdot|^{k-\frac{1}{2}} \times \tau |\cdot|^{k-\frac{3}{2}} \times \dots \times \tau |\cdot|^{\frac{1}{2}} \rtimes \pi \twoheadrightarrow (\mu \circ \det_{2k}) \rtimes \pi & \text{if } \delta = 0, \\ \tau |\cdot|^{k-1} \times \tau |\cdot|^{k-2} \times \dots \times \tau |\cdot|^{1} \times \mu \rtimes \pi \twoheadrightarrow (\mu \circ \det_{2k-1}) \rtimes \pi & \text{if } \delta = 1. \end{cases}$$

In particular, if π is unitary, then $(\mu \circ \det_{2k-\delta}) \rtimes \pi$ is irreducible.

Proof. By Theorem B.1, we have

$$\pi \cong \tau_1 \times \cdots \times \tau_l \rtimes \pi_0,$$

where

- τ_i = | · |^{s_i}τ'_i with τ'_i being an irreducible discrete series representation of GL_{k_i}(F);
 1/2 > s₁ ≥ s₂ ≥ · · · ≥ s_l ≥ 0;
- when $s_i = 0$, the irreducible representation ϕ_i of WD_F corresponding to τ_i is not orthogonal if $\delta = 0$, and is not symplectic if $\delta = 1$;
- π_0 is an irreducible representation of $\operatorname{Sp}_{r_0}(F)$ whose L-parameter is of good parity.

The segments corresponding to $\mu^{-1} |\det_k|^{\frac{k-\delta}{2}}$ and $\mu |\det_k|^{-\frac{k-\delta}{2}}$ are $[(1-\delta)/2, k-(1+\delta)/2]$ and $\left[-k + (1+\delta)/2, -(1-\delta)/2\right]$, respectively, which contain $(1-\delta)/2$ or $-(1-\delta)/2$. Hence by Theorem A.2, we see that

$$\mu^{-1} |\det_k|^{\frac{k-\delta}{2}} \times \tau_i \cong \tau_i \times \mu^{-1} |\det_k|^{\frac{k-\delta}{2}},$$
$$\mu |\det_k|^{-\frac{k-\delta}{2}} \times \tau_i \cong \tau_i \times \mu |\det_k|^{-\frac{k-\delta}{2}}.$$

By Proposition 3.5, we have

$$\mu^{-1} |\det_k|^{\frac{k-\delta}{2}} \rtimes \pi_0 \cong \mu |\det_k|^{-\frac{k-\delta}{2}} \rtimes \pi_0.$$

Therefore,

$$\begin{aligned} \tau |\cdot|^{k-\frac{1+\delta}{2}} &\times \tau |\cdot|^{k-\frac{1+\delta}{2}-1} \times \dots \times \tau |\cdot|^{1+\delta} \times (\mu \circ \det_{\delta}) \rtimes \pi \\ & \twoheadrightarrow \mu |\det_{k-\delta}|^{\frac{k}{2}} \times \mu^{-1} |\det_{k}|^{\frac{k-\delta}{2}} \rtimes \pi \\ & \cong \mu |\det_{k-\delta}|^{\frac{k}{2}} \times \tau_{1} \times \dots \times \tau_{l} \times \mu^{-1} |\det_{k}|^{\frac{k-\delta}{2}} \rtimes \pi_{0} \\ & \cong \mu |\det_{k-\delta}|^{\frac{k}{2}} \times \tau_{1} \times \dots \times \tau_{l} \times \mu |\det_{k}|^{-\frac{k-\delta}{2}} \rtimes \pi_{0} \\ & \cong \mu |\det_{k-\delta}|^{\frac{k}{2}} \times \mu |\det_{k}|^{-\frac{k-\delta}{2}} \rtimes \pi \\ & \twoheadrightarrow (\mu \circ \det_{2k-\delta}) \rtimes \pi. \end{aligned}$$

If π is unitary, then $(\mu \circ \det_{2k-\delta}) \rtimes \pi$ is semisimple. Therefore, by the uniqueness of the Langlands quotient, we conclude that $(\mu \circ \det_{2k-\delta}) \rtimes \pi$ is irreducible. \Box

This theorem together with Proposition 3.3 implies Theorem 3.1 (2).

3.7. Going down case. Finally, we show Theorem 3.1 (4).

Let π be an irreducible almost tempered unitary representation of $\widetilde{\operatorname{Sp}}_r(F)$. Note that $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ is irreducible by Theorem 3.1 (2). Set

$$\pi' = \mathcal{M}_{\psi,\tau}^{(r)} \left(\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \right).$$

Since there is a surjection $\pi' \to \pi$, we have $\pi' \neq 0$. Moreover, by the definition and Theorem 3.1 (2), we obtain a surjection

$$I_{\psi}^{(n+r)}(\tau) \twoheadrightarrow \left((\mu' \circ \det_{n-r}) \rtimes \pi \right) \boxtimes \pi'.$$

By Lemma 3.2, there is $0 \le t \le r$ such that $\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_n(F) \times \widetilde{\operatorname{Sp}}_r(F)}(R_t, ((\mu' \circ \det_{n-r}) \rtimes \pi) \boxtimes \pi') \ne 0$. By Bernstein's Frobenius reciprocity, this Hom-space is isomorphic to the space of $(\widetilde{M}_{t+(n-r)}(F) \times \widetilde{M}_t(F))$ -equivariant maps

$$\begin{pmatrix} \mu' |\det_{t+(n-r)}|^{\frac{t}{2}} \boxtimes \mu' |\det_{t}|^{\frac{t+(n-r)}{2}} \end{pmatrix} \otimes C_{c}^{\infty}(\widetilde{\operatorname{Sp}}_{r-t}(F)) \to R_{\overline{P_{t+(n-r)}(F)}}((\mu' \circ \det_{n-r}) \rtimes \pi) \boxtimes R_{\overline{P_{t}(F)}}(\pi').$$

In particular, we have a nonzero $\widetilde{\operatorname{GL}}_{t+(n-r)}(F)$ -equivariant map

$$R_{P_{t+(n-r)}(F)}((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi^{\vee}) \to \mu'^{-1} |\det_{t+(n-r)}|^{-\frac{t}{2}}.$$

By Proposition A.4, s.s. $R_{P_{t+(n-r)}(F)}((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi^{\vee})$ is the sum of

s.s.
$$\left(\mu' |\det_{n-r-a}|^{-\frac{a}{2}} \times \mu'^{-1} |\det_b|^{-\frac{n-r-b}{2}} \times \tau_{\lambda}\right) \boxtimes \left(\mu'^{-1} |\det_{a-b}|^{-\frac{n-r-a-b}{2}} \rtimes \pi_{\lambda}\right),$$

where (a, b) runs over the pairs of integers such that $0 \le b \le a \le n - r$, and $\tau_{\lambda} \boxtimes \pi_{\lambda}$ runs over all irreducible subquotients of $R_{P_{a-b+t}(F)}(\pi^{\vee})$. Hence we have

s.s.
$$\left(\mu' |\det_{n-r-a}|^{-\frac{a}{2}} \times \mu'^{-1} |\det_{b}|^{-\frac{n-r-b}{2}} \times \tau_{\lambda}\right) \supset \mu'^{-1} |\det_{t+(n-r)}|^{-\frac{t}{2}}$$

for some (a, b) and λ . Note that the segments corresponding to $\mu' |\det_{n-r-a}|^{-\frac{a}{2}}, \ \mu'^{-1} |\det_b|^{-\frac{n-r-b}{2}}$ and $\mu'^{-1} |\det_{t+(n-r)}|^{-\frac{t}{2}}$ are $[-(n-r-1)/2, (n-r-1)/2 - a], \ [-(n-r-1)/2, -(n-r-1)/2 + b - 1]$ and [-(n-r-1)/2 - t, (n-r-1)/2], respectively. If $t \geq 1$, then s.s. $R_{P_1(F)}(\pi^{\vee})$ must contain a

nonzero representation on which $\widetilde{\operatorname{GL}}_1(F)$ acts by $\mu'^{-1}| \cdot |^{-(n-r-1)/2-t}$. This contradicts that π is almost tempered, by Casselman's criterion (see e.g., [34, Lemma 2.4]). Hence we must have t = 0.

Therefore, we obtain a surjection

$$\operatorname{Ind}_{\widetilde{P}_{n-r}(F)\times\widetilde{\operatorname{Sp}}_{r}(F)}^{\widetilde{\operatorname{Sp}}_{r}(F)}\left((\mu'\circ \det_{n-r})\otimes C_{c}^{\infty}(\widetilde{\operatorname{Sp}}_{r}(F))\right)\twoheadrightarrow\left((\mu'\circ \det_{n-r})\rtimes\pi\right)\boxtimes\pi'.$$

Now Theorem 3.1 (4) follows from the following lemma.

Lemma 3.8. Let π be an irreducible almost tempered unitary representation of $\widetilde{\operatorname{Sp}}_r(F)$. Then the maximal $((\mu' \circ \det_{n-r}) \rtimes \pi)$ -isotypic quotient of

$$\operatorname{Ind}_{\widetilde{P}_{n-r}(F)\times\widetilde{\operatorname{Sp}}_{r}(F)}^{\widetilde{\operatorname{Sp}}_{r}(F)\times\widetilde{\operatorname{Sp}}_{r}(F)}\left(\left(\mu'\circ \det_{n-r}\right)\otimes C_{c}^{\infty}\left(\widetilde{\operatorname{Sp}}_{r}(F)\right)\right)$$

is isomorphic to $((\mu' \circ \det_{n-r}) \rtimes \pi) \boxtimes \Pi$, where Π is determined so that the subrepresentation of $R_{P_{n-r}(F)}((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi)$ on which $\widetilde{\operatorname{GL}}_{n-r}(F)$ acts by $\mu'^{-1} \circ \det_{n-r}$ is isomorphic to $(\mu'^{-1} \circ \det_{n-r}) \boxtimes \Pi$. Moreover, if π satisfies one of the conditions in Theorem 3.1 (4), then s.s.(Π) = π^{α} for some integer $\alpha \geq 1$, and that

$$\dim \operatorname{Hom}_{\widetilde{\operatorname{Sp}}}(F)(\Pi, \pi) = 1.$$

Proof. The linear dual Π^* of Π , which is not necessarily smooth, is given by

$$\Pi^* \cong \operatorname{Hom}_{\widetilde{\operatorname{Sp}}_n(F)} \left(\operatorname{Ind}_{\widetilde{P}_{n-r}(F) \times \widetilde{\operatorname{Sp}}_r(F)}^{\widetilde{\operatorname{Sp}}_r(F)} \left((\mu' \circ \det_{n-r}) \otimes C_c^{\infty}(\widetilde{\operatorname{Sp}}_r(F)) \right), (\mu' \circ \det_{n-r}) \rtimes \pi \right).$$

As representations of $Sp_n(F)$, we have

$$\operatorname{Ind}_{\widetilde{P}_{n-r}(F)\times\widetilde{\operatorname{Sp}}_{r}(F)}^{\widetilde{\operatorname{Sp}}_{r}(F)}\left((\mu'\circ\det_{n-r})\otimes C_{c}^{\infty}(\widetilde{\operatorname{Sp}}_{r}(F))\right)\cong\operatorname{ind}_{\widetilde{P}_{n-r}^{\circ}(F)}^{\widetilde{\operatorname{Sp}}_{n}(F)}(\chi),$$

where

- $P_{n-r}^{\circ}(F) \subset P_{n-r}(F)$ is generated by $\operatorname{GL}_{n-r}(F)$ and $N_{n-r}(F)$, so that $P_{n-r}(F)/P_{n-r}^{\circ}(F) \cong \operatorname{Sp}_{r}(F)$;
- $\operatorname{ind}_{\widetilde{P}_{n-r}^{\circ}(F)}^{\widetilde{\operatorname{Sp}}_n(F)}$ is the (unnormalized) compact induction functor;
- the character $\chi \colon \widetilde{P}_{n-r}^{\circ}(F) \to \mathbb{C}^{\times}$ is given by

$$\chi\left(\begin{pmatrix} a & * & * & * \\ 0 & \mathbf{1}_r & * & 0 \\ \hline 0 & 0 & ta^{-1} & 0 \\ 0 & 0 & * & \mathbf{1}_r \end{pmatrix}, \zeta\right) = \zeta^{n+r} \left(\frac{\alpha_{\psi}(1)}{\alpha_{\psi}(\det a)}\right)^{n+r} \mu(\det a) |\det a|^{\frac{n+r+1}{2}}$$

Moreover, $\widetilde{\operatorname{Sp}}_r(F)$ acts on $\operatorname{ind}_{\widetilde{P}_{n-r}(F)}^{\widetilde{\operatorname{Sp}}_n(F)}(\chi)$ by

$$(g_r \cdot \varphi)(g_n) = \varphi((g_r^{-1})^{\alpha}g_n)$$

for $g_r \in \widetilde{\operatorname{Sp}}_r(F)$, $g_n \in \widetilde{\operatorname{Sp}}_n(F)$ and $\varphi \in \operatorname{ind}_{\widetilde{P}_{n-r}^{\circ}(F)}^{\widetilde{\operatorname{Sp}}_n(F)}(\chi)$. Here, we set

$$g_r^{\alpha} = \begin{pmatrix} \mathbf{1}_r & 0\\ 0 & -\mathbf{1}_r \end{pmatrix} g_r \begin{pmatrix} \mathbf{1}_r & 0\\ 0 & -\mathbf{1}_r \end{pmatrix}^{-1},$$

and we identify $\widetilde{\operatorname{Sp}}_r(F)$ with the image of the composition $\widetilde{\operatorname{Sp}}_r(F) \hookrightarrow \widetilde{M}_{n-r}(F) \hookrightarrow \widetilde{\operatorname{Sp}}_n(F)$. By the Frobenius reciprocity, we have

$$\Pi^* \cong \operatorname{Hom}_{\widetilde{\operatorname{Sp}}_n(F)} \left((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi^{\vee}, \operatorname{Ind}_{\widetilde{P}_{n-r}^{\circ}(F)}^{\widetilde{\operatorname{Sp}}_n(F)} (\chi^{-1} \delta_{P_{n-r}}) \right)$$
$$\cong \operatorname{Hom}_{\widetilde{P}_{n-r}^{\circ}(F)} \left((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi^{\vee} |_{\widetilde{P}_{n-r}^{\circ}(F)}, \chi^{-1} \delta_{P_{n-r}} \right)$$
$$\cong \operatorname{Hom}_{\widetilde{\operatorname{GL}}_{n-r}(F)} \left(R_{P_{n-r}(F)} \left((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi^{\vee} \right) |_{\widetilde{\operatorname{GL}}_{n-r}(F)}, \mu'^{-1} \circ \det_{n-r} \right).$$

Moreover, $\widetilde{\operatorname{Sp}}_r(F)$ acts on this Hom-space by

$$(g_r \cdot \Phi)(f) = \Phi((g_r^{-1})^{\alpha} \cdot f)$$

for Φ in this Hom-space, and for $f \in R_{P_{n-r}(F)}((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi^{\vee})$. Since the representation $[g_r \mapsto \pi^{\vee}(g_r^{\alpha})]$ is isomorphic to π , by the definition of the Jacquet module, we see that the action $f \mapsto g_r^{\alpha} \cdot f$ is isomorphic to $R_{P_{n-r}(F)}((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi)$. Hence

$$\Pi^* \cong \operatorname{Hom}_{\widetilde{\operatorname{GL}}_{n-r}(F)} \left(R_{P_{n-r}(F)} \left((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi \right) |_{\widetilde{\operatorname{GL}}_{n-r}(F)}, \mu'^{-1} \circ \det_{n-r} \right).$$

as representations of $\operatorname{Sp}_r(F)$. This means that Π is determined so that the subrepresentation of $R_{P_{n-r}(F)}((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi)$ on which $\widetilde{\operatorname{GL}}_{n-r}(F)$ acts by $\mu'^{-1} \circ \det_{n-r}$ is isomorphic to $(\mu'^{-1} \circ \det_{n-r}) \boxtimes \Pi$ since $R_{P_{n-r}(F)}((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi)$ is admissible.

Note that s.s. $R_{P_{n-r}(F)}((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi)$ is the sum of

$$I_{(a,b),\lambda} = \text{s.s.} \left(\mu' |\det_{n-r-a}|^{-\frac{a}{2}} \times \mu'^{-1} |\det_b|^{-\frac{n-r-b}{2}} \times \tau_\lambda \right) \boxtimes \left(\mu'^{-1} |\det_{a-b}|^{-\frac{n-r-a-b}{2}} \rtimes \pi_\lambda \right),$$

where (a, b) runs over the pairs of integers such that $0 \le b \le a \le n - r$, and $\tau_{\lambda} \boxtimes \pi_{\lambda}$ runs over all irreducible subquotients of $R_{P_{a-b}(F)}(\pi)$. We claim that when π satisfies (a) or (b) in Theorem 3.1 (4), if

$$I_{(a,b),\lambda}|_{\widetilde{\operatorname{GL}}_{n-r}(F)} \supset \mu'^{-1} \circ \det_{n-r},$$

then $I_{(a,b),\lambda} \cong \mu'^{-1} \circ \det_{n-r} \boxtimes \pi$.

Note that the segments corresponding to $\mu' |\det_{n-r-a}|^{-\frac{a}{2}}, \mu'^{-1} |\det_b|^{-\frac{n-r-b}{2}}$ and $\mu'^{-1} \circ \det_{n-r}$ are [-(n-r-1)/2, (n-r-1)/2-a], [-(n-r-1)/2, -(n-r-1)/2+b-1] and [-(n-r-1)/2, (n-r-1)/2], respectively. When the *L*-parameter ϕ does not contain $\mu^{\pm 1}S_d$ for any $d \ge n-r$ with $d \equiv n-r \mod 2$, by computing the Jacquet module with respect to the Borel subgroup of $\widetilde{\operatorname{GL}}_{n-r}(F)$, we see that if

s.s.
$$\left(\mu' |\det_{n-r-a}|^{-\frac{a}{2}} \times \mu'^{-1} |\det_b|^{-\frac{n-r-b}{2}} \times \tau_\lambda\right) \supset \mu'^{-1} \circ \det_{n-r},$$

then (a,b) = (0,0) or (a,b) = (n-r,n-r). In these cases, we have $I_{(a,b),\lambda} = \mu'^{\pm 1} \circ \det_{n-r} \boxtimes \pi$, as desired.

When n = r, there is nothing to prove. Now we assume that n = r + 1. We take the maximal integer a such that

$$\pi \cong \underbrace{\mu^{\prime-1} \times \cdots \times \mu^{\prime-1}}_{a-1} \rtimes \pi_0$$

for some irreducible representation π_0 . Then by induction on a, one can show that the subrepresentation of s.s. $R_{P_1(F)}(\mu'^{-1} \rtimes \pi)$ on which $\widetilde{\operatorname{GL}}_1(F)$ acts by $\mu'^{\pm 1}$ is isomorphic to

$$\left((\mu'\boxtimes\pi)\oplus(\mu'^{-1}\boxtimes\pi)\right)^{\oplus a}.$$

This proves the claim when n = r + 1.

When π satisfies one of the conditions in Theorem 3.1 (4), by the claim, we see that s.s.(Π) is of the form $\pi^{\oplus \alpha}$ for some integer $\alpha \geq 1$. Moreover, we have

$$\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_{r}(F)}(\Pi,\pi) \cong (\Pi^{*} \otimes \pi)^{\operatorname{Sp}_{r}(F)}$$
$$\cong \operatorname{Hom}_{\widetilde{\operatorname{GL}}_{n-r}(F)}\left(R_{P_{n-r}(F)}\left((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi\right)|_{\widetilde{\operatorname{GL}}_{n-r}(F)}, (\mu'^{-1} \circ \det_{n-r}) \boxtimes \pi\right)^{\widetilde{\operatorname{Sp}}_{r}(F)}$$
$$\cong \operatorname{End}_{\widetilde{\operatorname{Sp}}_{r}(F)}((\mu'^{-1} \circ \det_{n-r}) \rtimes \pi),$$

which is one dimensional by Theorem 3.1 (2).

Remark 3.9. (1) If the L-parameter ϕ contains one of μS_d or $\mu^{-1}S_d$ for some $d \ge n-r$ with $d \equiv n-r \mod 2$, then $\phi \supset \mu S_d \oplus \mu^{-1}S_d$. By comparing the dimensions, we have

$$2r + 1 - \delta \ge 2d \ge 2(n - r).$$

Hence we must have $n \leq 2r$. In other words, when n > 2r, the condition (a) in Theorem 3.1 (4) always holds.

(2) In general, Π in Lemma 3.8 might have irreducible subquotients other than π . For example, consider the case where r = 2, n = 4 and $\pi = \mu'^{-1} \operatorname{St}_2 \rtimes \mathbf{1}_{\operatorname{Sp}_0(F)}$. Then one can see that

s.s.(
$$\Pi$$
) $\supset (\mu'^{-1} \circ \det_2) \rtimes \mathbf{1}_{\mathrm{Sp}_0(F)} \not\cong \pi.$

3.8. Seesaw identities. In the theory of theta liftings, seesaw identities are useful tools. The following proposition is an analogy for Miyawaki liftings. Recall that $J_{n-1}(F) = \operatorname{Sp}_{n-1}(F) \ltimes V_{n-1}(F)$ is a Jacobi subgroup of $\operatorname{Sp}_n(F)$. The center of the Heisenberg group $V_{n-1}(F)$ is denoted by $Z_{n-1}(F)$. For $\xi \in F^{\times}$, we denote the Weil representations of $\widetilde{J}_{n-1}(F)$ and $\widetilde{\operatorname{Sp}}_r(F)$ with respect to ψ_{ξ} by $\omega_{\psi_{\xi}}^{(n-1)}$ and $\omega_{\psi_{\xi}}^{(r)}$, respectively.

Proposition 3.10 (Seesaw identity). Let π and π' be irreducible representations of $\widetilde{\operatorname{Sp}}_r(F)$ and $\widetilde{\operatorname{Sp}}_{n-1}(F)$, on which $\{\pm 1\}$ acts by $(\pm 1)^{r+n}$ and $(\pm 1)^{r+n-1}$, respectively. Then

$$\operatorname{Hom}_{\widetilde{J}_{n-1}(F)}\left(\left.\mathcal{M}_{\psi,\tau}^{(n)}(\pi)\right|_{\widetilde{J}_{n-1}(F)},\pi'\otimes\omega_{\psi_{\xi}}^{(n-1)}\right)\cong\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_{r}(F)}\left(\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi')\otimes\omega_{\psi_{\xi}}^{(r)},\pi\right).$$

We shall write this property as the following seesaw diagram:



Proof. By proposition 2.2, there exists a $J_{n+r-1}(F)$ -surjection

$$I_{\psi}^{(n+r)}(\tau) \twoheadrightarrow I_{\psi}^{(n+r-1)}(\tau\chi_{\xi}) \otimes \omega_{\psi_{\xi}}^{(n+r-1)}.$$

Note that as $\widetilde{J}_{n-1}(F) \times \widetilde{\operatorname{Sp}}_r(F)$ -modules, we have $\omega_{\psi_{\xi}}^{(n+r-1)} \cong \omega_{\psi_{\xi}}^{(n-1)} \boxtimes \omega_{\psi_{\xi}}^{(r)}$. Composing a surjective $\widetilde{\operatorname{Sp}}_{n-1}(F) \times \widetilde{\operatorname{Sp}}_r(F)$ -map

$$I_{\psi}^{(n+r-1)}(\tau\chi_{\xi}) \twoheadrightarrow \pi' \boxtimes \mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi')$$

we obtain a nonzero $\widetilde{J}_{n-1}(F) \times \widetilde{\operatorname{Sp}}_r(F)$ -map

$$I_{\psi}^{(n+r)}(\tau) \twoheadrightarrow \left(\pi' \otimes \omega_{\psi_{\xi}}^{(n-1)}\right) \boxtimes \left(\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi') \otimes \omega_{\psi_{\xi}}^{(r)}\right)$$

If $\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_r(F)}\left(\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi')\otimes\omega_{\psi_{\xi}}^{(r)},\pi\right)\neq 0$, then we obtain a nonzero $\widetilde{J}_{n-1}(F)\times\widetilde{\operatorname{Sp}}_r(F)$ -map $I_{\psi}^{(n+r)}(\tau) \twoheadrightarrow \left(\pi' \otimes \omega_{\psi_{\mathcal{E}}}^{(n-1)}\right) \boxtimes \pi.$

By the definition of local Miyawaki liftings, it implies a nonzero $\widetilde{J}_{n-1}(F)$ -map $\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \to \pi' \otimes \omega_{\psi_{\mathcal{E}}}^{(n-1)}$ so that

$$\operatorname{Hom}_{\widetilde{J}_{n-1}(F)}\left(\left.\mathcal{M}_{\psi,\tau}^{(n)}(\pi)\right|_{\widetilde{J}_{n-1}(F)},\pi'\otimes\omega_{\psi_{\xi}}^{(n-1)}\right)\neq0$$

Conversely, if $\operatorname{Hom}_{\widetilde{J}_{n-1}(F)}\left(\left.\mathcal{M}_{\psi,\tau}^{(n)}(\pi)\right|_{\widetilde{J}_{n-1}(F)}, \pi'\otimes\omega_{\psi_{\xi}}^{(n-1)}\right)\neq 0$, then we have a nonzero $\widetilde{J}_{n-1}(F)\times \mathbb{C}_{\mathcal{J}_{n-1}(F)}$

 $\widetilde{\operatorname{Sp}}_r(F)$ -map

$$I_{\psi}^{(n+r)}(\tau) \twoheadrightarrow \mathcal{M}_{\psi,\tau}^{(n)}(\pi) \boxtimes \pi \to (\pi' \otimes \omega_{\psi_{\xi}}^{(n-1)}) \boxtimes \pi.$$

It factors through

$$\left(I_{\psi}^{(n+r)}(\tau)\right)_{\psi_{\xi}}$$

which is the maximal quotient of $I_{\psi}^{(n+r)}(\tau)$ on which $Z_{n+r-1}(F)$ acts by ψ_{ξ} . By the proof of Proposition 2.1 together with Proposition 2.2, we see that

$$\left(I_{\psi}^{(n+r)}(\tau)\right)_{\psi_{\xi}} \cong I_{\psi}^{(n+r-1)}(\tau\chi_{\xi}) \otimes \omega_{\psi_{\xi}}^{(n+r-1)}$$

as $\widetilde{J}_{n+r-1}(F)$ -modules. Hence we have a nonzero $\widetilde{J}_{n-1}(F) \times \widetilde{\operatorname{Sp}}_r(F)$ -map

$$I_{\psi}^{(n+r-1)}(\tau\chi_{\xi}) \otimes \omega_{\psi_{\xi}}^{(n+r-1)} \to \left(\pi' \otimes \omega_{\psi_{\xi}}^{(n-1)}\right) \boxtimes \pi.$$

This implies that

$$\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_{r}(F)}\left(\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi')\otimes\omega_{\psi_{\xi}}^{(r)},\pi\right)\neq 0.$$

Hence we obtain two homomorphisms

$$\operatorname{Hom}_{\widetilde{J}_{n-1}(F)}\left(\left.\mathcal{M}_{\psi,\tau}^{(n)}(\pi)\right|_{\widetilde{J}_{n-1}(F)},\pi'\otimes\omega_{\psi_{\xi}}^{(n-1)}\right)\rightleftharpoons\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_{r}(F)}\left(\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi')\otimes\omega_{\psi_{\xi}}^{(r)},\pi\right).$$

These are obtained by the following diagram of $\widetilde{J}_{n-1}(F) \times \widetilde{\operatorname{Sp}}_r(F)$ -homomorphisms

$$I_{\psi}^{(n+r)}(\tau) \longrightarrow \mathcal{M}_{\psi,\tau}^{(n)}(\pi) \boxtimes \pi$$

$$\downarrow \varphi_{1} \otimes \mathrm{id}$$

$$\left(\pi' \otimes \omega_{\psi_{\xi}}^{(n-1)}\right) \boxtimes \pi$$

$$\uparrow^{\mathrm{id} \otimes \varphi_{2}}$$

$$I_{\psi}^{(n+r-1)}(\tau\chi_{\xi}) \otimes \omega_{\psi_{\xi}}^{(n+r-1)} \longrightarrow \left(\pi' \otimes \omega_{\psi_{\xi}}^{(n-1)}\right) \boxtimes \left(\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi') \otimes \omega_{\psi_{\xi}}^{(r)}\right)$$

$$= \int_{\mathcal{M}_{\psi}} \left(\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(n)}(\pi)\right|_{\varphi}, \quad \pi' \otimes \omega_{\psi_{\xi}}^{(n-1)}\right) \text{ and } \varphi_{2} \in \mathrm{Hom}_{\widetilde{\Theta_{\psi}}}(\pi) \left(\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi') \otimes \omega_{\psi_{\xi}}^{(r)}, \pi\right)$$

for $\varphi_1 \in \operatorname{Hom}_{\widetilde{J}_{n-1}(F)}\left(\left.\mathcal{M}_{\psi,\tau}^{(n)}(\pi)\right|_{\widetilde{J}_{n-1}(F)}, \pi' \otimes \omega_{\psi_{\xi}}^{(n-1)}\right)$ and $\varphi_2 \in \operatorname{Hom}_{\widetilde{\operatorname{Sp}}_r(F)}\left(\mathcal{M}_{\psi,\tau\chi_{\xi}}^{\vee,\tau}(\pi') \otimes \operatorname{Clearly}, \text{ the two maps } \varphi_1 \mapsto \varphi_2 \text{ and } \varphi_2 \mapsto \varphi_1 \text{ are inverse to each other.}\right)$ ψ_{ξ} *"*.).

4. GLOBAL MIYAWAKI LIFTINGS

Now we let F be a totally real number field, and ψ be a non-trivial unitary character of \mathbb{A}/F . We assume that for each infinite place v of F, there exists $a_v \in F_v \cong \mathbb{R}$ with $a_v > 0$ such that $\psi(x_v) = \exp(2\pi a_v \sqrt{-1}x_v)$ for $x_v \in F_v$.

In this section, we define the global Miyawaki liftings as the pullbacks of Ikeda liftings, and establish basic properties. One may regard the global Miyawaki liftings as an analogue of the global theta liftings, which are defined by the pullbacks of theta functions.

4.1. **Definition.** Let $\tau = \otimes_v' \tau_v$ be an irreducible unitary cuspidal automorphic representation of $\operatorname{GL}_2(\mathbb{A})$ satisfying the conditions (A1), (A2) and (A3) in §2.5. We denote the weight of $\otimes_{v|\infty} \tau_v$ by $\pm 2k = (\pm 2k_v)_v$ with $k_v > 0$. Then we have the Ikeda lift $\operatorname{Ik}_{\psi}^{(n+r)}(\tau)$, which is an irreducible admissible representation of $\widetilde{\operatorname{Sp}}_{n+r}(\mathbb{A}_{\operatorname{fin}})$ occurring in the space $\mathcal{S}_{k+(n+r)/2}\left(\operatorname{Sp}_{n+r}(F)\setminus\widetilde{\operatorname{Sp}}_{n+r}(\mathbb{A})\right)$ of holomorphic cusp forms on $\widetilde{\operatorname{Sp}}_{n+r}(\mathbb{A})$ of weight k + (n+r)/2. For $\mathcal{F} \in \operatorname{Ik}_{\psi}^{(n+r)}(\tau)$ and $\varphi \in \mathcal{S}_{k+(n+r)/2}\left(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A})\right)$, consider the integral

$$\mathcal{M}^{(n)}((g_n,\zeta_n);\varphi,\mathcal{F}) = \int_{\mathrm{Sp}_r(F)\backslash \mathrm{Sp}_r(\mathbb{A})} \mathcal{F}(\iota(g_n,g_r),\zeta_n\zeta_r)\overline{\varphi(g_r,\zeta_r)} dg_r$$

for $(g_n, \zeta_n) \in \widetilde{\operatorname{Sp}}_n(\mathbb{A})$, where dg_r is the Tamagawa measure on $\operatorname{Sp}_r(F) \setminus \operatorname{Sp}_r(\mathbb{A})$. Note that this integral does not depend on the choice of $\zeta_r \in \{\pm 1\}$, and that $\mathcal{M}^{(n)}(\varphi, \mathcal{F})$ is genuine if and only if n + r is odd.

Lemma 4.1. We have

$$\mathcal{M}^{(n)}(\varphi, \mathcal{F}) \in \mathcal{S}_{k+(n+r)/2}\left(\operatorname{Sp}_n(F) \setminus \widetilde{\operatorname{Sp}}_n(\mathbb{A})\right).$$

Proof. The non-trivial part is the cuspidality. Note that $\mathcal{M}^{(n)}(\varphi, \mathcal{F})$ has a Fourier expansion of the form

$$\mathcal{M}^{(n)}((g_n,\zeta_n);\varphi,\mathcal{F}) = \sum_{B\in \operatorname{Sym}_n^+(F)} W_B(g_n,\zeta_n),$$

where

$$W_B(g_n,\zeta_n) = \int_{\mathrm{Sym}_n(F)\backslash \mathrm{Sym}_n(\mathbb{A})} \mathcal{M}^{(n)}(\mathbf{n}(X)(g_n,\zeta_n);\varphi,\mathcal{F})\overline{\psi(\mathrm{Tr}(BX))} dX$$

Since

$$\mathbf{v}(0,y,z)\mapsto\psi\left(\operatorname{Tr}\left(B\begin{pmatrix}z&y\\ty&0\end{pmatrix}\right)\right)$$

is a non-trivial character of $Y(\mathbb{A})Z(\mathbb{A})$ for any $B \in \text{Sym}_n^+(F)$, we have

$$\int_{Y(F)Z(F)\setminus Y(\mathbb{A})Z(\mathbb{A})} \mathcal{M}^{(n)}(\mathbf{v}(0,y,z)(g_n,\zeta_n);\varphi,\mathcal{F}) dy dz = 0.$$

Since YZ is a normal subgroup of N_k , which is the unipotent radical of the standard maximal parabolic subgroup P_k , the constant term of $\mathcal{M}^{(n)}(\varphi, \mathcal{F})$ along P_k must be zero. Hence $\mathcal{M}^{(n)}(\varphi, \mathcal{F})$ is cuspidal.

Let π be an admissible (not necessarily irreducible) representation of $\widetilde{\operatorname{Sp}}_r(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+(n+r)/2}\left(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A})\right)$. The (global) Miyawaki lift $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ of π is defined by the representation of $\widetilde{\operatorname{Sp}}_n(\mathbb{A}_{\operatorname{fin}})$ generated by

$$\left\{ \mathcal{M}^{(n)}(\varphi, \mathcal{F}) \in \mathcal{S}_{k+(n+r)/2}(\mathrm{Sp}_n(F) \setminus \widetilde{\mathrm{Sp}}_n(\mathbb{A})) \mid \mathcal{F} \in \mathrm{Ik}_{\psi}^{(n+r)}(\tau), \ \varphi \in \pi \right\}.$$

4.2. A-parameters. Arthur's multiplicity formula established by Arthur [3] and Gan–Ichino [14] describes the discrete spectrum of automorphic forms on $\operatorname{Sp}_n(F) \setminus \widetilde{\operatorname{Sp}}_n(\mathbb{A})$ in terms of global A-parameters. In this subsection, we review Arthur's multiplicity formula for holomorphic cusp forms. For more precision, see also Appendix B.4.

A discrete global A-parameter for $\operatorname{Sp}_n(F)$ (resp. $\widetilde{\operatorname{Sp}}_n(F)$) is a symbol

$$\Psi = \tau_1[d_1] \boxplus \cdots \boxplus \tau_t[d_t],$$

where τ_i is an irreducible cuspidal unitary automorphic representation of $\operatorname{GL}_{m_i}(\mathbb{A})$, and d_i is a positive integer satisfying several conditions, e.g., $\sum_{i=1}^t m_i d_i$ is equal to 2n + 1 (resp. 2n). The precise definition is given in Appendix B.4. Two A-parameters $\Psi = \bigoplus_{i=1}^t \tau_i[d_i]$ and $\Psi' = \bigoplus_{i=1}^{t'} \tau_i'[d_i']$ are said to be equivalent if t = t' and there exists a permutation $\sigma \in \mathfrak{S}_t$ such that $d'_i = d_{\sigma(i)}$ and $\tau'_i \cong \tau_{\sigma(i)}$ for each *i*. We denote the set of equivalence classes of discrete global A-parameters for $\operatorname{Sp}_n(F)$ (resp. $\widetilde{\operatorname{Sp}}_n(F)$) by $\Psi_2(\operatorname{Sp}_n(F))$ (resp. $\Psi_2(\widetilde{\operatorname{Sp}}_n(F))$). We call an A-parameter $\Psi = \bigoplus_{i=1}^t \tau_i[d_i]$ tempered if $d_i = 1$ for any *i*. In this case, we write $\Psi = \bigoplus_{i=1}^t \tau_i$ for simplicity.

We state Arthur's multiplicity formula ([3, Theorem 1.5.2] and [14, Theorems 1.1, 1.3]) for holomorphic cusp forms.

Theorem 4.2 (Arthur's multiplicity formula). Let $l = (l_v) \in \prod_{v \mid \infty} \mathbb{Z}$ with $l_v > 0$ and $l_v \equiv l_{v'} \mod 2$ for any $v, v' \mid \infty$.

(1) For $\Psi \in \Psi_2(\operatorname{Sp}_n(F))$ if l_v is even, and for $\Psi \in \Psi_2(\widetilde{\operatorname{Sp}}_n(F))$ if l_v is odd, there exists an $\widetilde{\operatorname{Sp}}_n(\mathbb{A}_{\operatorname{fin}})$ -stable subspace $\mathcal{S}_{l/2,\Psi}$ of $\mathcal{S}_{l/2}\left(\operatorname{Sp}_n(F)\setminus\widetilde{\operatorname{Sp}}_n(\mathbb{A})\right)$ (possibly zero) such that

$$\mathcal{S}_{l/2}\left(\mathrm{Sp}_n(F)\backslash\widetilde{\mathrm{Sp}}_n(\mathbb{A})\right) = \bigoplus_{\Psi} \mathcal{S}_{l/2,\Psi},$$

where Ψ runs over $\Psi_2(\operatorname{Sp}_n(F))$ if l_v is even, and over $\Psi_2(\widetilde{\operatorname{Sp}}_n(F))$ if l_v is odd.

(2) Suppose that $\pi = \otimes_{v < \infty}' \pi_v$ is an irreducible subrepresentation of $S_{l/2,\Psi}$ with $\Psi = \boxplus_{i=1}^t \tau_i[d_i]$. Then for almost all $v < \infty$, the local factors π_v of π and $\tau_{i,v}$ of τ_i are unramified for any *i*. Moreover, if we denote the Satake parameter for $\tau_{i,v}$ by $\{c_{i,v,1}, \ldots, c_{i,v,m_i}\}$, then the Satake parameter of π_v is equal to

$$\bigcup_{i=1}^{t}\bigcup_{j=1}^{m_{i}}\left\{c_{i,v,j}q^{-\frac{d_{i}-1}{2}},c_{i,v,j}q^{-\frac{d_{i}-3}{2}},\ldots c_{i,v,j}q^{\frac{d_{i}-1}{2}}\right\}$$

as multisets.

(3) If Ψ is a tempered A-parameter, then $S_{l/2,\Psi}$ is multiplicity-free as a representation of $\widetilde{Sp}_n(\mathbb{A}_{fin})$.

If an admissible representation π of $\widetilde{\mathrm{Sp}}_n(\mathbb{A}_{\mathrm{fin}})$ is contained in $\mathcal{S}_{l/2,\Psi}$, we say that π has an A-parameter Ψ , and Ψ is the A-parameter for π .

4.3. **Basic properties.** We establish basic properties of Miyawaki liftings. Let $\pi = \bigotimes_{v < \infty}' \pi_v$ be an irreducible admissible representation of $\widetilde{\operatorname{Sp}}_r(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+(n+r)/2}\left(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A})\right)$.

First we compute the A-parameter for Miyawaki liftings.

Proposition 4.3. Suppose that $\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \neq 0$ with $n \geq r$. If π has an A-parameter Ψ , then $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ has an A-parameter

$$\Psi \boxplus \tau \chi_{-1}^{[(n+r)/2]}[n-r].$$

Here, when n = r, we omit $\tau \chi_{-1}^{[(n+r)/2]}[n-r]$ so that $\mathcal{M}_{\psi,\tau}^{(r)}(\pi)$ has the same A-parameter as π .

Proof. This follows from the computation of Satake parameters for $\mathcal{M}_{\psi_v,\tau_v}^{(n)}(\pi_v)$ (Theorem 3.1 (5)).

Miyawaki liftings have the following duality property.

Proposition 4.4. Suppose that $\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \neq 0$. Then

$$\pi \subset \mathcal{M}_{\psi,\tau}^{(r)}\left(\mathcal{M}_{\psi,\tau}^{(n)}(\pi)\right).$$

Proof. For $\varphi_1, \varphi_2 \in \mathcal{S}_{k+(n+r)/2}\left(\operatorname{Sp}_r(F) \setminus \widetilde{\operatorname{Sp}}_r(\mathbb{A})\right)$, we define the Petersson inner product by

$$\langle \varphi_1, \varphi_2 \rangle = \int_{\operatorname{Sp}_r(F) \setminus \operatorname{Sp}_r(\mathbb{A})} \varphi_1(g, \zeta) \overline{\varphi_2(g, \zeta)} dg.$$

If $\mathcal{M}^{(n)}(\varphi, \mathcal{F}) \neq 0$ for $\varphi \in \pi$ and $\mathcal{F} \in \mathrm{Ik}_{\psi}^{(n+r)}(\tau)$, then

$$\left\langle \mathcal{M}^{(r)} \left(\mathcal{M}^{(n)}(\varphi, \mathcal{F}), \mathcal{F} \right), \varphi \right\rangle$$

= $\int_{\operatorname{Sp}_r(F) \setminus \operatorname{Sp}_r(\mathbb{A})} \left(\int_{\operatorname{Sp}_n(F) \setminus \operatorname{Sp}_n(\mathbb{A})} \mathcal{F}(\iota(g_n, g_r), \zeta_n \zeta_r) \overline{\mathcal{M}^{(n)}((g_n, \zeta_n)\varphi, \mathcal{F})} dg_n \right) \overline{\varphi(g_r, \zeta_r)} dg_r$
= $\left\langle \mathcal{M}^{(n)}(\varphi, \mathcal{F}), \mathcal{M}^{(n)}(\varphi, \mathcal{F}) \right\rangle \neq 0.$

Hence \langle,\rangle is nonzero on $\mathcal{M}_{\psi,\tau}^{(r)}\left(\mathcal{M}_{\psi,\tau}^{(n)}(\pi)\right) \times \pi$. This shows that $\pi \subset \mathcal{M}_{\psi,\tau}^{(r)}\left(\mathcal{M}_{\psi,\tau}^{(n)}(\pi)\right)$.

The local irreducibility (Theorem 3.1) implies the global irreducibility.

Theorem 4.5. Suppose that π has a tempered A-parameter, and $\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \neq 0$ with $n \geq r$. Then:

- (1) $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ is irreducible, and $\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \cong \otimes_{v<\infty}' \mathcal{M}_{\psi_v,\tau_v}^{(n)}(\pi_v).$
- (2) When $n \leq r+1$ or n > 2r, we have

$$\mathcal{M}_{\psi,\tau}^{(r)}\left(\mathcal{M}_{\psi,\tau}^{(n)}(\pi)\right) = \pi.$$

Proof. Since π has a tempered A-parameter, the local factor π_v is almost tempered by Lemma B.8. By Theorem 3.1 (2), the local Miyawaki lift $\mathcal{M}_{\psi_v,\tau_v}^{(n)}(\pi_v)$ is irreducible. Hence the π -isotypic quotient of $\mathrm{Ik}_{\psi}^{(n+r)}(\tau)$ is of the form $\pi' \boxtimes \pi$ for some irreducible representation π' of $\widetilde{\mathrm{Sp}}_n(\mathbb{A}_{\mathrm{fin}})$. The $\widetilde{\mathrm{Sp}}_r(\mathbb{A}_{\mathrm{fin}})$ -invariant surjection

$$\mathrm{Ik}_{\psi}^{(n+r)}(\tau) \otimes \overline{\pi} \twoheadrightarrow \mathcal{M}_{\psi,\tau}^{(n)}(\pi), \ \mathcal{F} \otimes \overline{\varphi} \mapsto \mathcal{M}^{(n)}(\varphi,\mathcal{F})$$

factors through a surjective $\widetilde{\mathrm{Sp}}_n(\mathbb{A}_{\mathrm{fin}})$ -homomorphism $\pi' \twoheadrightarrow \mathcal{M}_{\psi,\tau}^{(n)}(\pi)$. Since π' is irreducible, this map must be injective if $\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \neq 0$. Hence $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ is irreducible. Moreover, the above $\widetilde{\mathrm{Sp}}_r(\mathbb{A}_{\mathrm{fin}})$ -invariant surjection implies that the local component at $v < \infty$ of $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ is isomorphic to $\mathcal{M}_{\psi,\tau,\tau_v}^{(n)}(\pi_v)$. Hence we obtain (1).

By Proposition 4.4, we know that

$$\mathcal{M}_{\psi,\tau}^{(r)}\left(\mathcal{M}_{\psi,\tau}^{(n)}(\pi)\right)\supset\pi.$$

By a similar argument to (1), when $r \le n \le r+1$ or n > 2r, Theorem 3.1 (4) implies that the left hand side is irreducible. Hence the inclusion is an equality, and we obtain (2).

Corollary 4.6. Suppose that π has a tempered A-parameter Ψ .

- (1) If n = r and $\mathcal{M}_{\psi,\tau}^{(r)}(\pi) \neq 0$, then $\mathcal{M}_{\psi,\tau}^{(r)}(\pi) = \pi$ as a subrepresentation of $\mathcal{S}_{k+r}(\mathrm{Sp}_r(F) \setminus \mathrm{Sp}_r(\mathbb{A}))$.
- (2) If n = r+1 and $\mathcal{M}_{\psi,\tau}^{(r+1)}(\pi) \neq 0$, then $\mathcal{M}_{\psi,\tau}^{(r+1)}(\pi)$ is a unique irreducible subrepresentation of $\mathcal{S}_{k+r+1/2}\left(\operatorname{Sp}_{r+1}(F)\setminus\widetilde{\operatorname{Sp}}_{r+1}(\mathbb{A})\right)$ whose local component at $v < \infty$ is isomorphic to $\mu'_v \rtimes \pi_v$ with $\mu'_v = \mu_v \chi_{-1}^r$.

Proof. By Theorem 3.1 (3), the local Miyawaki lift $\mathcal{M}_{\psi_v,\tau_v}^{(n)}(\pi_v)$ is isomorphic to π_v if n = r, and to $\mu'_v \rtimes \pi_v$ if n = r + 1. By Proposition 4.3, when n = r (resp. n = r + 1), we see that $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ has a tempered A-parameter $\Psi' = \Psi$ (resp. $\Psi' = \Psi \boxplus \tau \chi_{-1}^r$). Since $\mathcal{S}_{k+(n+r)/2,\Psi'}$ is multiplicity-free by Theorem 4.2 (3), we have $\mathcal{M}_{\psi,\tau}^{(r)}(\pi) = \pi$ if n = r, and $\mathcal{M}_{\psi,\tau}^{(r+1)}(\pi)$ satisfies the desired uniqueness property if n = r + 1.

Let n' = n or n' = n - 1. Fix $l, l' \in \prod_{v \mid \infty} \mathbb{Z}$ such that $l_v \equiv l_{v'} \neq l'_v \equiv l'_{v'} \mod 2$ for any $v, v' \mid \infty$. For $\varphi \in \mathcal{S}_{l/2}\left(\operatorname{Sp}_n(F) \setminus \widetilde{\operatorname{Sp}}_n(\mathbb{A})\right), \ \varphi' \in \mathcal{S}_{l'/2}\left(\operatorname{Sp}_{n'}(F) \setminus \widetilde{\operatorname{Sp}}_{n'}(\mathbb{A})\right)$, and $\phi \in \mathcal{S}(X_{n'}(\mathbb{A}))$, we define a Fourier–Jacobi period $\mathcal{P}_{n,n',\psi_{\xi}}(\varphi, \overline{\varphi'}, \phi)$ by the integral

$$\begin{cases} \int_{\operatorname{Sp}_{n}(F)\backslash\operatorname{Sp}_{n}(\mathbb{A})} \varphi(g,\zeta) \overline{\varphi'(g,\zeta)} \overline{\Theta}_{\psi_{\xi}}^{\phi}(g,\zeta) dg, & \text{if } n' = n, \\ \int_{V_{n-1}(F)\backslash V_{n-1}(\mathbb{A})} \int_{\operatorname{Sp}_{n-1}(F)\backslash\operatorname{Sp}_{n-1}(\mathbb{A})} \varphi(v(g,\zeta)) \overline{\varphi'(g,\zeta)} \overline{\Theta}_{\psi_{\xi}}^{\phi}(v(g,\zeta)) dg dv, & \text{if } n' = n-1 \end{cases}$$

Fix a totally positive element $\xi \in F^{\times}$. Recall that $\mathcal{S}(X(\mathbb{A}))_{\xi}$ is the subspace of $\mathcal{S}(X(\mathbb{A}))$ spanned by $\phi = \bigotimes_{v} \phi_{v}$ such that $\phi_{v} = \phi_{\xi_{v}}^{0}$ for each infinite place v. The local seesaw identity (Proposition 3.10) is a local analogue of the following result.

Proposition 4.7 (Seesaw identity). Let π and π' be irreducible representations of $\widetilde{\operatorname{Sp}}_r(\mathbb{A}_{\operatorname{fin}})$ and $\widetilde{\operatorname{Sp}}_{n-1}(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+(n+r)/2}(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A}))$ and $\mathcal{S}_{k+(n-1+r)/2}(\operatorname{Sp}_{n-1}(F)\setminus\widetilde{\operatorname{Sp}}_{n-1}(\mathbb{A}))$, respectively.

(1) If there exist $\mathcal{M}^{(n)}(\varphi, \mathcal{F}_1) \in \mathcal{M}^{(n)}_{\psi, \tau}(\pi)$ with $\varphi \in \pi$, $\varphi' \in \pi'$, and $\phi_1 \in \mathcal{S}(X_{n-1}(\mathbb{A}))_{\xi}$ such that $\mathcal{P}_{n,n-1,\psi_{\xi}}\left(\mathcal{M}^{(n)}(\varphi, \mathcal{F}_1), \overline{\varphi'}, \phi_1\right) \neq 0,$

then there exist $\mathcal{M}^{(r)}(\varphi', \mathcal{F}'_2) \in \mathcal{M}^{(r)}_{\psi, \tau \chi_{\xi}}(\pi')$ and $\phi_2 \in \mathcal{S}(X_r(\mathbb{A}))_{\xi}$ such that

$$\mathcal{P}_{r,r,\psi_{\xi}}\left(\varphi,\overline{\mathcal{M}^{(r)}(\varphi',\mathcal{F}'_{2})},\phi_{2}\right)\neq 0.$$

(2) Assume that $n + r \geq 2$. If there exist $\varphi \in \pi$, $\mathcal{M}^{(r)}(\varphi', \mathcal{F}'_2) \in \mathcal{M}^{(r)}_{\psi, \tau\chi_{\xi}}(\pi')$ with $\varphi' \in \pi'$, and $\phi_2 \in \mathcal{S}(X_r(\mathbb{A}))_{\xi}$ such that

$$\mathcal{P}_{r,r,\psi_{\xi}}\left(\varphi,\overline{\mathcal{M}^{(r)}(\varphi',\mathcal{F}_{2}')},\phi_{2}\right)\neq0,$$

then there exist $\mathcal{M}^{(n)}(\varphi, \mathcal{F}_1) \in \mathcal{M}^{(n)}_{\psi, \tau}(\pi)$ and $\phi_1 \in \mathcal{S}(X_{n-1}(\mathbb{A}))_{\xi}$ such that

$$\mathcal{P}_{n,n-1,\psi_{\xi}}\left(\mathcal{M}^{(n)}(\varphi,\mathcal{F}_{1}),\overline{\varphi'},\phi_{1}\right)\neq 0.$$

Proof. Suppose that there exist $\mathcal{M}^{(n)}(\varphi, \mathcal{F}) \in \mathcal{M}^{(n)}_{\psi, \tau}(\pi)$ with $\varphi \in \pi, \varphi' \in \pi'$, and $\phi \in \mathcal{S}(X_{n-1}(\mathbb{A}))_{\xi}$ such that

$$\mathcal{P}_{n,n-1,\psi_{\xi}}\left(\mathcal{M}^{(n)}(\varphi,\mathcal{F}),\overline{\varphi'},\phi\right)\neq 0.$$

It is equal to

$$\begin{split} &\int_{V_{n-1}(F)\setminus V_{n-1}(\mathbb{A})} \int_{\mathrm{Sp}_{n-1}(F)\setminus \mathrm{Sp}_{n-1}(\mathbb{A})} \\ &\times \left(\int_{\mathrm{Sp}_r(F)\setminus \mathrm{Sp}_r(\mathbb{A})} \mathcal{F}(\iota(v,1)\cdot\iota(g_{n-1},g_r),\zeta_n\zeta_r) \overline{\varphi(g_r,\zeta_n)} dg_r \right) \\ &\times \overline{\varphi'(g_{n-1},\zeta_n)} \overline{\Theta}_{\psi_{\xi}}^{\phi}(v(g_{n-1},\zeta_n)) dg_{n-1} dv. \end{split}$$

First, if we compute the integral on $Z_{n-1}(F) \setminus Z_{n-1}(\mathbb{A})$, then $\mathcal{F}_{\psi_{\xi}}$ appears. By Proposition 2.7, there exist $\mathcal{F}'_1, \ldots, \mathcal{F}'_r \in \mathrm{Ik}_{\psi}^{(n+r-1)}(\tau \otimes \chi_{\xi})$ and $\phi_1, \ldots, \phi_r \in \mathcal{S}(X_{n+r-1}(\mathbb{A}))_{\xi}$ such that

$$\mathcal{F}_{\psi_{\xi}}(\iota(v,1)\cdot\iota(g_{n-1},g_r),\zeta_n\zeta_r) = \sum_{i=1}^r \mathcal{F}'_i(\iota(g_{n-1},g_r),\zeta_n\zeta_r)\Theta^{\phi_i}_{\psi_{\xi}}(\iota(v,1)\cdot\iota(g_{n-1},g_r),\zeta_n\zeta_r).$$

Note that $X_{n+r-1}(\mathbb{A}) \cong X_{n-1}(\mathbb{A}) \oplus X_r(\mathbb{A})$. We may assume that $\phi_i = \phi_i^{(n-1)} \otimes \phi_i^{(r)}$ with $\phi_i^{(n-1)} \in \mathcal{S}(X_{n-1}(\mathbb{A}))_{\xi}$ and $\phi_i^{(r)} \in \mathcal{S}(X_r(\mathbb{A}))_{\xi}$. Then we have

$$\Theta_{\psi_{\xi}}^{\phi_{i}}(\iota(v,1)\cdot\iota(g_{n-1},g_{r}),\zeta_{n}\zeta_{r}) = \Theta_{\psi_{\xi}}^{\phi_{i}^{(n-1)}}(v(g_{n-1},\zeta_{n}))\Theta_{\psi_{\xi}}^{\phi_{i}^{(r)}}(g_{r},\zeta_{r}).$$

Hence there exists i such that

$$\int_{\operatorname{Sp}_{n-1}(F)\setminus\operatorname{Sp}_{n-1}(\mathbb{A})} \int_{\operatorname{Sp}_{r}(F)\setminus\operatorname{Sp}_{r}(\mathbb{A})} \mathcal{F}'_{i}(\iota(g_{n-1},g_{r}),\zeta_{n}\zeta_{r})\overline{\varphi(g_{r},\zeta_{r})\varphi'(g_{n-1},\zeta_{n})} \Theta_{\psi_{\xi}}^{\phi_{i}^{(r)}}(g_{r},\zeta_{r}) \times \left(\int_{Z_{n-1}(\mathbb{A})V_{n-1}(F)\setminus V_{n-1}(\mathbb{A})} \Theta_{\psi_{\xi}}^{\phi_{i}^{(n-1)}}(v(g_{n-1},\zeta_{n}))\overline{\Theta_{\psi_{\xi}}^{\phi}(v(g_{n-1},\zeta_{n}))} dv\right) dg_{r}dg_{n-1} \neq 0.$$

The integral on $Z_{n-1}(\mathbb{A})V_{n-1}(F)\setminus V_{n-1}(\mathbb{A})$ is equal to the inner product $(\phi_i^{(n-1)}, \phi)$, which does not depend on (g_{n-1}, ζ_n) . The other integral is equal to the complex conjugate of

 $\mathcal{P}_{r,r,\psi_{\xi}}\left(\varphi,\overline{\mathcal{M}^{(r)}(\varphi',\mathcal{F}'_{i})},\phi_{i}^{(r)}\right).$

Hence we obtain (1).

Next suppose that there exist $\varphi \in \pi$, $\mathcal{M}^{(r)}(\varphi', \mathcal{F}') \in \mathcal{M}^{(r)}_{\psi, \tau \chi_{\xi}}(\pi')$ with $\varphi' \in \pi'$, and $\phi_2 \in \mathcal{S}(X_r(\mathbb{A}))_{\xi}$ such that

$$\mathcal{P}_{r,r,\psi_{\xi}}\left(\varphi,\overline{\mathcal{M}^{(r)}(\varphi',\mathcal{F}')},\phi\right)\neq 0.$$

Choose a nonzero vector $\phi' \in \mathcal{S}(X_{n-1}(\mathbb{A}))_{\xi}$. Then by the same calculation above, we have

$$\begin{split} &(\phi',\phi')\mathcal{P}_{r,r,\psi_{\xi}}\left(\varphi,\overline{\mathcal{M}^{(r)}(\varphi',\mathcal{F}')},\phi\right) \\ &= \int_{\mathrm{Sp}_{r}(F)\backslash\mathrm{Sp}_{r}(\mathbb{A})}\int_{\mathrm{Sp}_{n-1}(F)\backslash\mathrm{Sp}_{n-1}(\mathbb{A})}\mathcal{F}'(\iota(g_{n-1},g_{r}),\zeta_{n}\zeta_{r})\overline{\varphi'(g_{n-1},\zeta_{n})\varphi(g_{r},\zeta_{r})}\Theta_{\psi_{\xi}}^{\phi}(g_{r},\zeta_{r}) \\ &\times \left(\int_{Z_{n-1}(\mathbb{A})V_{n-1}(F)\backslash V_{n-1}(\mathbb{A})}\Theta_{\psi_{\xi}}^{\phi'}(v(g_{n-1},\zeta_{n}))\overline{\Theta_{\psi_{\xi}}^{\phi'}(v(g_{n-1},\zeta_{n}))}dv\right)dg_{n-1}dg_{r} \\ &= \int_{Z_{n-1}(\mathbb{A})V_{n-1}(F)\backslash V_{n-1}(\mathbb{A})}\int_{\mathrm{Sp}_{n-1}(F)\backslash\mathrm{Sp}_{n-1}(\mathbb{A})} \\ &\times \left(\int_{\mathrm{Sp}_{r}(F)\backslash\mathrm{Sp}_{r}(\mathbb{A})}\mathcal{F}'(\iota(g_{n-1},g_{r}),\zeta_{n}\zeta_{r})\Theta_{\psi_{\xi}}^{\phi'\otimes\phi}(\iota(v,1)\cdot\iota(g_{n-1},g_{r}),\zeta_{n}\zeta_{r})\overline{\varphi(g_{r},\zeta_{r})}dg_{r}\right) \\ &\times \overline{\varphi'(g_{n-1},\zeta_{n})}\overline{\Theta_{\psi_{\xi}}^{\phi'}(v(g_{n-1},\zeta_{n}))}dg_{n-1}dv. \end{split}$$

If $n + r \ge 2$, then by Proposition 2.7 (2), (3), there exists $\mathcal{F} \in \text{Ik}_{\psi}^{(n+r)}(\tau)$ such that

$$\mathcal{F}_{\psi_{\xi}}(\iota(v,1)\cdot\iota(g_{n-1},g_r),\zeta_n\zeta_r) = \mathcal{F}'(\iota(g_{n-1},g_r),\zeta_n\zeta_r)\Theta_{\psi_{\xi}}^{\phi'\otimes\phi}(\iota(v,1)\cdot\iota(g_{n-1},g_r),\zeta_n\zeta_r).$$

Then

$$0 \neq (\phi', \phi') \mathcal{P}_{r,r,\psi_{\xi}}(\varphi, \overline{\mathcal{M}^{(r)}(\varphi', \mathcal{F}')}, \phi) = \mathcal{P}_{n,n-1,\psi_{\xi}}(\mathcal{M}^{(n)}(\varphi, \mathcal{F}), \overline{\varphi'}, \phi').$$

Hence we obtain (2).

As an application of this seesaw identity, we have a criterion for the non-vanishing of the Miyawaki liftings for the equal rank case.

Proposition 4.8. Let π be an irreducible representation of $\text{Sp}_r(\mathbb{A}_{\text{fin}})$ occurring in $\mathcal{S}_{k+r}(\text{Sp}_r(F) \setminus \text{Sp}_r(\mathbb{A}))$ with $r \geq 1$. Assume that π has a tempered A-parameter, and that the ξ -th Fourier–Jacobi module $\text{FJ}_{\psi_{\varepsilon}}(\pi)$ is nonzero. Then the following are equivalent:

- (a) $\mathcal{M}_{\psi,\tau}^{(r)}(\pi)$ is nonzero.
- (b) For any irreducible representation π' of $\widetilde{\operatorname{Sp}}_{r-1}(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+r-1/2}\left(\operatorname{Sp}_{r-1}(F)\setminus\widetilde{\operatorname{Sp}}_{r-1}(\mathbb{A})\right)$, if $\mathcal{P}_{r,r-1,\psi_{\xi}}$ is not identically zero on $\pi \times \overline{\pi'} \times \mathcal{S}(X_{r-1}(\mathbb{A}))_{\xi}$, then $\mathcal{P}_{r,r,\psi_{\xi}}$ is not identically zero on $\pi \times \overline{\mathcal{M}}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi') \times \mathcal{S}(X_{r}(\mathbb{A}))_{\xi}$.
- (c) There exists an irreducible representation π' of $\widetilde{\operatorname{Sp}}_{r-1}(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+r-1/2}\left(\operatorname{Sp}_{r-1}(F)\backslash\widetilde{\operatorname{Sp}}_{r-1}(\mathbb{A})\right)$ such that $\mathcal{P}_{r,r,\psi_{\xi}}$ is not identically zero on $\pi \times \overline{\mathcal{M}}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi') \times \mathcal{S}(X_{r}(\mathbb{A}))_{\xi}$.

Proof. First, we show that (a) implies (b). Suppose that $\mathcal{M}_{\psi,\tau}^{(r)}(\pi)$ is nonzero. Since π has a tempered A-parameter, by Corollary 4.6 (1), we have $\mathcal{M}_{\psi,\tau}^{(r)}(\pi) = \pi$. Then by Proposition 4.7 (1), we obtain the condition (b).

Next, we show that (b) implies (c). Since $FJ_{\psi_{\xi}}(\pi) \neq 0$, the map

$$\pi \otimes \overline{\omega_{\psi_{\xi}}} \to \mathcal{S}_{k+r-1/2}(\mathrm{Sp}_{r-1}(F) \backslash \widetilde{\mathrm{Sp}}_{r-1}(\mathbb{A})),$$
$$\varphi \otimes \overline{\phi} \mapsto \int_{V_{r-1}(F) \backslash V_{r-1}(\mathbb{A})} \varphi(vg') \cdot \overline{\Theta_{\psi_{\xi}}^{\phi}(vg')} dv$$

is nonzero. If we take an irreducible subrepresentation π' in the image of this map, then $\mathcal{P}_{r,r-1,\psi_{\mathcal{E}}}$ is not identically zero on $\pi \times \overline{\pi'} \times \mathcal{S}(X_{r-1}(\mathbb{A}))_{\xi}$. Applying (b) to π' , we see that π' satisfies the condition of (c).

Finally, we show that (c) implies (a). If $\mathcal{P}_{r,r,\psi_{\xi}}$ is not identically zero on $\pi \times \overline{\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi')} \times$ $\mathcal{S}(X_r(\mathbb{A}))_{\xi}$, then $\mathcal{P}_{r,r-1,\psi_{\xi}}$ is not identically zero on $\mathcal{M}_{\psi,\tau}^{(r)}(\pi) \times \overline{\pi'} \times \mathcal{S}(X_{r-1}(\mathbb{A}))_{\xi}$ by Proposition 4.7 (2). In particular, $\mathcal{M}_{\psi,\tau}^{(r)}(\pi) \neq 0$. This completes the proof.

5. A relation between Miyawaki liftings and the Gan-Gross-Prasad conjecture

In this section, we formulate a conjecture on the non-vanishing of global Miyawaki liftings. In addition, we relate this conjecture to the Gan-Gross-Prasad conjecture for the symplectic-metaplectic case.

5.1. Conjectures. Fix τ and ψ as in the previous section, and a totally positive element $\xi \in$ F^{\times} . We denote the weight of $\otimes_{v \mid \infty} \tau_v$ by $\pm 2k = (\pm 2k_v)_v$ with $k_v > 0$. Let π be an irreducible representation of $\widetilde{\operatorname{Sp}}_r(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+(n+r)/2}\left(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A})\right)$, and $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ be the Miyawaki lift of π . The Rankin–Selberg *L*-function attached to $\pi \times \tau \chi_{-1}^r$ is denoted by $L(s, \pi \times \tau \chi_{-1}^r)$.

Conjecture 5.1 $((M)_{r,n})$. Suppose that $n \ge r$.

- (1) When n = r, the Miyawaki lift $\mathcal{M}_{\psi,\tau}^{(r)}(\pi)$ is nonzero if and only if $L(1/2, \pi \times \tau \chi_{-1}^r) \neq 0$. (2) When n > r, the Miyawaki lift $\mathcal{M}_{\psi,\tau}^{(n)}(\pi)$ is always nonzero.

Note that when r = 0, Conjecture 5.1 is trivial since $\mathcal{M}_{\psi,\tau}^{(n)}(\eta^n) = \mathrm{Ik}_{\psi}^{(n)}(\tau)$, where η is the unique non-trivial character of $\widetilde{\mathrm{Sp}}_0(\mathbb{A}_{\mathrm{fin}}) \cong \mathbb{Z}/2\mathbb{Z}$. When $n \geq r$ and $n \equiv r \mod 2$, it is an extension of a part of Ikeda's conjecture [28, Conjecture 5.1]. The simplest case of this conjecture (the case where $F = \mathbb{Q}, n = r = 1$, and both $\tau \chi_{-1}$ and π are unramified everywhere) was proven by Ichino [25] and Xue [57] independently. When $n \ge r$ and $n \not\equiv r \mod 2$, some examples were given by Hayashida [22, Lemma 9.1].

In the rest of this section, we explain a relation between Conjecture 5.1 and the Gan–Gross– Prasad conjecture.

5.2. Gan-Gross-Prasad conjectures. The Gan-Gross-Prasad conjecture (the GGP conjecture) [11, Conjecture 24.1] relates the non-vanishing of the Fourier–Jacobi periods to the non-vanishing of the central values of the Rankin–Selberg L-functions. In this subsection, we review the GGP conjecture for holomorphic cusp forms. For more details, we refer the readers to Appendix C.

Suppose that n' = n or n' = n - 1. Let π and π' be irreducible representations of $\widetilde{Sp}_n(\mathbb{A}_{fin})$ and $\widetilde{\mathrm{Sp}}_{n'}(\mathbb{A}_{\mathrm{fin}})$ occurring in $\mathcal{S}_{l/2}\left(\mathrm{Sp}_n(F)\backslash\widetilde{\mathrm{Sp}}_n(\mathbb{A})\right)$ and $\mathcal{S}_{(l-1)/2}\left(\mathrm{Sp}_{n'}(F)\backslash\widetilde{\mathrm{Sp}}_{n'}(\mathbb{A})\right)$, respectively. Assume that π and π' have tempered A-parameters $\Psi = \bigoplus_{i=1}^{t} \tau_i$ and $\Psi' = \bigoplus_{j=1}^{t'} \tau'_j$, respectively. We define the Rankin–Selberg L-function $L(s, \pi \times \pi' \times \chi_{(-1)^{l-1}\xi})$ by

$$L\left(s, \pi \times \pi' \times \chi_{(-1)^{l-1}\xi}\right) = \prod_{i=1}^{t} \prod_{j=1}^{t'} L\left(s, \tau_i \times \tau'_j \times \chi_{(-1)^{l-1}\xi}\right).$$

Note that τ'_j is self-dual and unitary, so that $\overline{\tau'_j} \cong \tau'_j$ for any j.

Since we consider holomorphic cusp forms only, we state the GGP conjecture as the following unusual way.

Conjecture 5.2 (Gan–Gross–Prasad conjecture). Let $\pi = \bigotimes_{v < \infty}' \pi_v$ and $\pi' = \bigotimes_{v < \infty}' \pi'_v$ be as above.

(1) If the Fourier–Jacobi period $\mathcal{P}_{n,n',\psi_{\xi}}$ is not identically zero on $\pi \times \overline{\pi'} \times \mathcal{S}(X_{n'}(\mathbb{A}))_{\xi}$, then the central value $L(1/2, \pi \times \pi' \times \chi_{(-1)^{l-1}\xi})$ is nonzero and the local Hom-space

$$\operatorname{Hom}_{\operatorname{Sp}_{\pi'}(F_v)}\left(\pi_v\otimes\overline{\pi'_v}\otimes\overline{\omega_{\psi_{\mathcal{F}}}},\mathbb{C}\right)$$

is nonzero for any $v < \infty$.

(2) When n' = n, the converse of (1) holds.

For the usual statements and the relation with Conjecture 5.2, see Appendix C below. When n' = n, Conjecture 5.2 (1) was proven by Yamana [58].

5.3. Relation between Conjectures 5.1 and 5.2. In this subsection, we show Conjecture 5.1 using the GGP conjecture and the following hypothesis.

Hypothesis 5.3. Let π be an irreducible representation of $\widetilde{\operatorname{Sp}}_r(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{l/2}(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A}))$.

- (A) Suppose that π has a tempered A-parameter and that $\operatorname{FJ}_{\psi_{\xi}}(\pi) \neq 0$. Then there exists an irreducible representation π' of $\widetilde{\operatorname{Sp}}_{r-1}(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{(l-1)/2}\left(\operatorname{Sp}_{r-1}(F)\setminus\widetilde{\operatorname{Sp}}_{r-1}(\mathbb{A})\right)$ with tempered A-parameter such that $\mathcal{P}_{r,r-1,\psi_{\xi}}$ is not identically zero on $\pi \times \overline{\pi'} \times \mathcal{S}(X_{r-1}(\mathbb{A}))_{\xi}$.
- (B) There exists an irreducible representation π' of $\widetilde{\operatorname{Sp}}_r(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{(l-1)/2}\left(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A})\right)$ such that $\mathcal{P}_{r,r,\psi_{\mathcal{E}}}$ is not identically zero on $\pi \times \overline{\pi'} \times \mathcal{S}(X_r(\mathbb{A}))_{\mathcal{E}}$.
- (C) There exists an irreducible representation π' of $\widetilde{\operatorname{Sp}}_{r+1}(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{(l+1)/2}\left(\operatorname{Sp}_{r+1}(F)\setminus\widetilde{\operatorname{Sp}}_{r+1}(\mathbb{A})\right)$ such that $\mathcal{P}_{r+1,r,\psi_{\xi}}$ is not identically zero on $\pi' \times \overline{\pi} \times \mathcal{S}(X_r(\mathbb{A}))_{\xi}$.
- Remark 5.4. (1) Hypothesis 5.3 (A) may be regarded as a global analogue of [7, Lemma C.6]. It is also assumed in the theory of twisted automorphic descents. See Jiang–Zhang [32, Conjecture 2.3] and the remark after this conjecture for a nature of this expectation and some example.
 - (2) The main difficulty of Hypothesis 5.3 (A) is the tempered-ness of π' . Since all representations occurring in the space $S_{(l-1)/2}\left(\operatorname{Sp}_1(F)\setminus\widetilde{\operatorname{Sp}}_1(\mathbb{A})\right)$ have tempered A-parameters by Example B.9, Hypothesis 5.3 (A) is true when $r \leq 2$.
 - (3) Hypothesis 5.3 (B) means that the L^2 -orthogonal complement of

$$\left\{\varphi' \cdot \Theta_{\xi}^{\phi} \mid \varphi' \in \mathcal{S}_{(l-1)/2}\left(\operatorname{Sp}_{r}(F) \setminus \widetilde{\operatorname{Sp}}_{r}(\mathbb{A})\right), \ \phi \in \mathcal{S}(X_{r}(\mathbb{A}))_{\xi}\right\}$$

in $\mathcal{S}_{l/2}\left(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A})\right)$ is equal to $\{0\}$. Theta liftings might have potential for Hypothesis 5.3 (B).

- (4) In the classical setting, Hypothesis 5.3 (C) means the surjectivity of the ξ-th Fourier–Jacobi expansion of Siegel modular forms of degree r + 1. For r = 1, see Aoki [2] and Ibukiyama– Poor–Yuen [24].
- (5) In any case, Hypothesis 5.3 seems quite difficult. For the proof, it might be necessary to consider averages of Fourier–Jacobi periods (cf. see a paper of Nelson–Venkatesh [42]).

Using the GGP conjecture (Conjecture 5.2) and Hypothesis 5.3, we will show Conjecture 5.1. First, we consider the (almost) equal rank case, i.e., the case where n = r or n = r + 1.

Theorem 5.5. Assume the GGP conjecture (Conjecture 5.2) and Hypothesis 5.3 (A). Then for irreducible representations with tempered A-parameters, we have

$$(M)_{r-1,r} \implies (M)_{r,r} \implies (M)_{r,r+1}.$$

In particular, $(M)_{r,r}$ and $(M)_{r,r+1}$ for irreducible representations with tempered A-parameters are true for any $r \ge 0$.

Proof. First, we show $(M)_{r-1,r} \implies (M)_{r,r}$. Let π be an irreducible representation of $\widetilde{\operatorname{Sp}}_r(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+r}(\operatorname{Sp}_r(F)\backslash\operatorname{Sp}_r(\mathbb{A}))$ with tempered A-parameter Ψ . We choose a totally positive element $\xi \in F^{\times}$ such that $\operatorname{FJ}_{\psi_{\xi}}(\pi) \neq 0$. By Hypothesis 5.3 (A), we can find an irreducible representation π' of $\widetilde{\operatorname{Sp}}_{r-1}(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+r-1/2}\left(\operatorname{Sp}_{r-1}(F)\backslash\widetilde{\operatorname{Sp}}_{r-1}(\mathbb{A})\right)$ with tempered A-parameter Ψ' such that $\mathcal{P}_{r,r-1,\psi_{\xi}}$ is not identically zero on $\pi \times \overline{\pi'} \times \mathcal{S}(X_{r-1}(\mathbb{A}))_{\xi}$.

Suppose that $\mathcal{M}_{\psi,\tau}^{(r)}(\pi) \neq 0$. Then by Proposition 4.8 (b), we see that $\mathcal{P}_{r,r,\psi_{\xi}}$ is not identically zero on $\pi \times \overline{\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi')} \times \mathcal{S}(X_r(\mathbb{A}))_{\xi}$. Since π' has a tempered A-parameter Ψ' , by Theorem 4.5 and Proposition 4.3, the Miyawaki lifting $\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi')$ is irreducible and has a tempered A-parameter $\Psi' \boxplus \tau\chi_{(-1)^{r-1}\xi}$. By the GGP conjecture (Conjecture 5.2 (1)), we conclude that

$$L(1/2, \pi \times \pi' \times \chi_{-\xi})L(1/2, \pi \times (\tau \chi_{(-1)^{r-1}\xi}) \times \chi_{-\xi}) \neq 0,$$

so that $L(1/2, \pi \times \tau \chi_{(-1)^r}) \neq 0$.

Conversely, suppose that $L(1/2, \pi \times \tau \chi_{(-1)^r}) \neq 0$. By $(M)_{r-1,r}$, we have $\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi') \neq 0$. Hence by Theorem 4.5 and Proposition 4.3, it is irreducible and has a tempered A-parameter $\Psi' \boxplus \tau \chi_{(-1)^{r-1}\xi}$. By the GGP conjecture (Conjecture 5.2 (1), (2) and Theorem C.4), we see that $\mathcal{P}_{r,r,\psi_{\xi}}$ is not identically zero on $\pi \times \overline{\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi')} \times \mathcal{S}(X_r(\mathbb{A}))_{\xi}$. Hence by Proposition 4.8, we conclude that $\mathcal{M}_{\psi,\tau}^{(r)}(\pi) \neq 0$. This completes the proof of $(M)_{r-1,r} \implies (M)_{r,r}$. Next, we show $(M)_{r,r} \implies (M)_{r,r+1}$. By Lemma B.10, there exists an irreducible rep-

Next, we show $(M)_{r,r} \implies (M)_{r,r+1}$. By Lemma B.10, there exists an irreducible representation $\Pi = \bigotimes_{v < \infty}^{r} \Pi_v$ of $\widetilde{\operatorname{Sp}}_{r+1}(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+r+1/2}\left(\operatorname{Sp}_{r+1}(F)\setminus\widetilde{\operatorname{Sp}}_{r+1}(\mathbb{A})\right)$ such that $\Pi_v = \mathcal{M}_{\psi_v,\tau_v}^{(r+1)}(\pi_v)$ for each $v < \infty$, and the *A*-parameter for Π is equal to $\Psi \boxtimes \tau \chi_{-1}^r$. We choose a totally positive element $\xi \in F^{\times}$ such that $\operatorname{FJ}_{\psi_{\xi}}(\Pi) \neq 0$. Using Hypothesis 5.3 (A), we take an irreducible representation π' of $\operatorname{Sp}_r(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+r}\left(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A})\right)$ with tempered *A*-parameter Ψ' such that $\mathcal{P}_{r+1,r,\psi_{\xi}}$ is not identically zero on $\Pi \times \overline{\pi'} \times \mathcal{S}(X_r(\mathbb{A}))_{\xi}$. By the GGP conjecture (Conjecture 5.2 (1)), we see that

$$L(1/2, \Pi \times \pi' \times \chi_{\xi}) = L(1/2, \pi \times \pi' \times \chi_{\xi}) L(1/2, \pi' \times \tau \chi_{(-1)^r \xi})$$

is nonzero. Since $L(1/2, \pi' \times \tau \chi_{(-1)^r \xi}) \neq 0$, by $(M)_{r,r}$, we see that $\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi')$ is nonzero. By Corollary 4.6 (1), it is equal to π' itself. Since $L(1/2, \pi \times \pi' \times \chi_{\xi}) \neq 0$, by the GGP conjecture (Conjecture 5.2 (2) and Theorem C.4), we see that $\mathcal{P}_{r,r,\psi_{\xi}}$ is not identically zero on

$$\pi \times \overline{\pi'} \times \mathcal{S}(X_r(\mathbb{A}))_{\xi} = \pi \times \overline{\mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\pi')} \times \mathcal{S}(X_r(\mathbb{A}))_{\xi}.$$

Then by the seesaw identity (Proposition 4.7 (2)), we see that $\mathcal{P}_{r+1,r,\psi_{\xi}}$ is not identically zero on $\mathcal{M}_{\psi,\tau}^{(r+1)}(\pi) \times \overline{\pi'} \times \mathcal{S}(X_r(\mathbb{A}))_{\xi}$. In particular we have $\mathcal{M}_{\psi,\tau}^{(r+1)}(\pi) \neq 0$. This completes the proof of $(M)_{r,r} \implies (M)_{r,r+1}$.

Corollary 5.6. The conjecture $(M)_{1,1}$ is true.

Proof. Note that any irreducible representation occurring in $S_{k+1}(\operatorname{Sp}_1(F) \setminus \operatorname{Sp}_1(\mathbb{A}))$ has a tempered A-parameter (Remark B.9). The conjecture $(M)_{0,1}$ and Hypothesis 5.3 (A) for r = 1 are trivial.

The GGP conjecture for n = n' = 1 is known (see e.g., [12, Proof of Theorem 7.1], [44, Proposition 4.1, Theorem 4.5] and [56]). Hence we have $(M)_{1,1}$.

Next, we consider the going-up case, i.e., the case where n > r + 1.

Theorem 5.7. Suppose that $n \ge r+2$.

(1) Assuming Hypothesis 5.3 (B), we have

$$(M)_{r,n-1} \implies (M)_{r,n}.$$

(2) Assuming Hypothesis 5.3 (C), we have

$$(M)_{r+1,n} \implies (M)_{r,n}.$$

In particular, assuming one of Hypothesis 5.3 (B) or (C), the conjecture $(M)_{r,n}$ for $n \ge r+1$ is reduced to $(M)_{r,r+1}$ or $(M)_{n-1,n}$.

Proof. We prove only the first assertion. The proof of the second one is similar.

Let π be an irreducible representation of $\widetilde{\operatorname{Sp}}_r(\mathbb{A}_{\operatorname{fin}})$ occurring in $\mathcal{S}_{k+(n+r)/2}\left(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A})\right)$. Using Hypothesis 5.3 (B), we choose an irreducible representation π' of $\widetilde{\operatorname{Sp}}_r(\mathbb{A}_{\operatorname{fin}})$ which appears in $\mathcal{S}_{k+(n+r-1)/2}\left(\operatorname{Sp}_r(F)\setminus\widetilde{\operatorname{Sp}}_r(\mathbb{A})\right)$ such that $\mathcal{P}_{r,r,\psi_{\xi}}$ is not identically zero on $\pi \times \overline{\pi'} \times \mathcal{S}(X_r(\mathbb{A}))_{\xi}$. Applying $(M)_{r,n-1}$ to π' , we have $\Pi' = \mathcal{M}_{\psi,\tau\chi_{\xi}}^{(n-1)}(\pi') \neq 0$. By Proposition 4.4, we have $\pi' \subset \mathcal{M}_{\psi,\tau\chi_{\xi}}^{(r)}(\Pi')$. Hence $\mathcal{P}_{r,r,\psi_{\xi}}$ is not identically zero on $\pi \times \overline{\mathcal{M}}_{\psi,\tau\chi_{\xi}}^{(r)}(\Pi') \times \mathcal{S}(X_r(\mathbb{A}))_{\xi}$. By the seesaw identity (Proposition 4.7 (2)), we see that $\mathcal{P}_{n,n-1,\psi_{\xi}}$ is not identically zero on $\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \times \overline{\Pi'} \times \mathcal{S}(X_{n-1}(\mathbb{A}))_{\xi}$. In particular, we have $\mathcal{M}_{\psi,\tau}^{(n)}(\pi) \neq 0$. \Box

Appendix A. Jacquet modules of representations of metaplectic groups

Let F be a non-archimedean local field of characteristic zero. In this appendix, we recall computations of Jacquet modules of induced representations of $\operatorname{GL}_k(F)$ or $\widetilde{\operatorname{Sp}}_n(F)$.

A.1. Induced representations of general linear groups. Let P(F) = M(F)N(F) be a parabolic subgroup of $\operatorname{GL}_k(F)$ containing the Borel subgroup consisting of upper triangular matrices. Then the Levi part M(F) is of the form $\operatorname{GL}_{k_1}(F) \times \cdots \times \operatorname{GL}_{k_l}(F)$ with $k_1 + \cdots + k_l = k$. For representations τ_1, \ldots, τ_l of $\operatorname{GL}_{k_1}(F), \ldots, \operatorname{GL}_{k_l}(F)$, respectively, we denote the normalized induced representation by

$$\tau_1 \times \cdots \times \tau_l = \operatorname{Ind}_{P(F)}^{\operatorname{GL}_k(F)}(\tau_1 \otimes \cdots \otimes \tau_l).$$

A segment is a symbol [x, y], where $x, y \in \mathbb{R}$ with $x - y \in \mathbb{Z}$. We identify [x, y] as the set $\{x, x-1, \ldots, y\}$ if $x \geq y$, and $\{x, x+1, \ldots, y\}$ if $x \leq y$. Let ρ be an irreducible unitary supercuspidal representation of $\operatorname{GL}_d(F)$. Then the normalized induced representation

$$\rho|\cdot|^x imes \dots imes
ho|\cdot|^y$$

has a unique irreducible subrepresentation, which is denoted by

$$\langle \rho; x, \ldots, y \rangle$$

If $x \ge y$, this is called a Steinberg representation, which is an essentially discrete series representation of $\operatorname{GL}_{d(|x-y|+1)}(F)$. If x < y, this is called a Speh representation. For example, if $\rho = \mu$ be a unitary character (i.e., d = 1) and x < y, then $\langle \mu; x, \ldots, y \rangle = \mu |\det_{y-x+1}|^{(x+y)/2}$ is a character of $\operatorname{GL}_{y-x+1}(F)$, where we denote by \det_k the determinant character of $\operatorname{GL}_k(F)$.

Definition A.1. Let [x, y] and [x', y'] be two segments.

- (1) When $(x-y)(x-y') \ge 0$, we say that [x, y] and [x', y'] are linked if $[x, y] \not\subset [x', y']$, $[x', y'] \not\subset [x, y]$ as sets, and $[x, y] \cup [x, y']$ is also a segment.
- (2) When (x y)(x' y') < 0, we say that [x, y] and [x', y'] are linked if [y, x] and [x', y'] are linked, and $x, y \notin [x', y']$ and $x', y' \notin [x, y]$.

The linked-ness gives an irreducibility criterion for induced representations.

Theorem A.2 (Zelevinsky [59, Theorems 4.2, 9.7], Mœglin–Waldspurger [41]). Let [x, y] and [x', y'] be segments, and let ρ and ρ' be irreducible unitary supercuspidal representations of $\operatorname{GL}_d(F)$ and $\operatorname{GL}_{d'}(F)$, respectively. Then the induced representation

$$\langle \rho; x, \dots, y \rangle \times \langle \rho'; x', \dots, y' \rangle$$

is irreducible unless [x, y] are [x', y'] are linked, and $\rho \cong \rho'$.

For a partition (k_1, k_2) of k, we denote by $R_{(k_1, k_2)}$ the normalized Jacquet functor of representations of $\operatorname{GL}_k(F)$ with respect to the standard maximal parabolic subgroup P(F) = M(F)N(F)with $M(F) \cong \operatorname{GL}_{k_1}(F) \times \operatorname{GL}_{k_2}(F)$. The Jacquet module of $\langle \rho; x, \ldots, y \rangle$ is computed by Zelevinsky.

Proposition A.3 ([59, Propositions 3.4, 9.5]). Let ρ be an irreducible unitary supercuspidal representation of $\operatorname{GL}_d(F)$. Suppose that $x \neq y$ and set k = d(|x-y|+1). Then $R_{(k_1,k_2)}(\langle \rho; x, \ldots, y \rangle) = 0$ unless $k_1 \equiv 0 \mod d$. If $k_1 = dm$ with $1 \leq m \leq |x-y|$, we have

$$R_{(k_1,k_2)}(\langle \rho; x,\ldots,y\rangle) = \langle \rho; x,\ldots,x-\epsilon(m-1)\rangle \otimes \langle \rho; x-\epsilon m,\ldots,y\rangle_{\mathbb{R}}$$

where $\epsilon \in \{\pm 1\}$ is defined so that $\epsilon(x - y) > 0$.

A.2. Representations of double covers of general linear groups. Recall that for $a \in F^{\times}$, the Weil constant $\alpha_{\psi}(a)$ is an eighth root of unity, and satisfies that

$$\frac{\alpha_{\psi}(a)\alpha_{\psi}(b)}{\alpha_{\psi}(1)\alpha_{\psi}(ab)} = \langle a, b \rangle$$

for $a, b \in F$, where the right hand side is the Hilbert symbol. In particular,

$$\left(\frac{\alpha_{\psi}(1)}{\alpha_{\psi}(a)}\right)^2 = \chi_{-1}(a)$$

for $a \in F^{\times}$, where χ_{-1} is the quadratic character associated to $F(\sqrt{-1})/F$.

A double cover of $GL_k(F)$ is given by

$$\widetilde{\operatorname{GL}}_k(F) = \operatorname{GL}_k(F) \times \{\pm 1\}$$

with group law

$$(g_1,\epsilon_1)\cdot(g_2,\epsilon_2)=(g_1g_2,\epsilon_1\epsilon_2\cdot\langle\det g_1,\det g_2\rangle)$$

Let $\operatorname{Irr}(\operatorname{GL}_k(F))$ (resp. $\operatorname{Irr}(\widetilde{\operatorname{GL}}_k(F))$) be the set of equivalence classes of irreducible representations of $\operatorname{GL}_k(F)$ (resp. the set of equivalence classes of irreducible genuine representations of $\widetilde{\operatorname{GL}}_k(F)$). For an irreducible representation τ of $\operatorname{GL}_k(F)$ and $(g, \epsilon) \in \widetilde{\operatorname{GL}}_k(F)$, we set

$$\tau_{\psi}(g,\epsilon) = \epsilon \frac{\alpha_{\psi}(1)}{\alpha_{\psi}(\det g)} \tau(g).$$

Then τ_{ψ} is irreducible and genuine, and the map $\tau \mapsto \tau_{\psi}$ gives a bijection $\operatorname{Irr}(\operatorname{GL}_k(F)) \to \operatorname{Irr}(\operatorname{GL}_k(F))$. For a more precise representation theory for $\widetilde{\operatorname{GL}}_k(F)$, see [20, §4.1].

A.3. Induced representations of symplectic and metaplectic groups. We denote by $R_{P_t(F)}$ the normalized Jacquet functor of representations of $\widetilde{\operatorname{Sp}}_r(F)$ with respect to the maximal parabolic subgroup $\widetilde{P}_t(F)$. For a smooth representation Π of $\widetilde{\operatorname{Sp}}_r(F)$, we write s.s.(Π) for the semisimplification of Π .

Proposition A.4. Let π be a representation of $\widetilde{\operatorname{Sp}}_r(F)$, μ be a unitary character of F^{\times} , and $\alpha \in \mathbb{C}$. Then s.s. $R_{P_t(F)}(\mu |\det_k|^{\alpha} \rtimes \pi)$ is isomorphic to the direct sum of

s.s.
$$\left(\mu^{-1} |\det_{k-a}|^{-\alpha - \frac{a}{2}} \times \mu |\det_{b}|^{\alpha - \frac{k-b}{2}} \times \tau_{\lambda}\right) \boxtimes \left(\mu |\det_{a-b}|^{\alpha - \frac{k-a-b}{2}} \rtimes \pi_{\lambda}\right)$$

where (a, b) runs over the pairs of integers such that $0 \le b \le a \le k$ and $a - b \ge k - t$, and $\tau_{\lambda} \boxtimes \pi_{\lambda}$ runs over all irreducible subquotients of $R_{P_{a-b-k+t}(F)}(\pi)$ (with multiplicity).

Proof. This follows from Tadić's formula [51], [20, Proposition 4,5].

APPENDIX B. LOCAL LANGLANDS CORRESPONDENCE AND ARTHUR'S MULTIPLICITY FORMULA

In this appendix, we summarize the local Langlands correspondence and Arthur's multiplicity formula.

B.1. Local Langlands correspondence. We recall the local Langlands correspondence (LLC) for $\operatorname{Sp}_n(F)$ and $\operatorname{Sp}_n(F)$. Let F be a local field of characteristic zero. Fix a non-trivial unitary character ψ of F. We denote by W_F and WD_F the Weil group and the Weil–Deligne group of F, respectively, i.e.,

$$WD_F = \begin{cases} W_F \times \operatorname{SL}_2(\mathbb{C}) & \text{if } F \text{ is non-archimedean,} \\ W_F & \text{if } F \text{ is archimedean.} \end{cases}$$

A representation of WD_F is a homomorphism $\phi: WD_F \to \operatorname{GL}_N(\mathbb{C})$ such that

- $\phi(\text{Frob})$ is semi-simple if F is non-archimedean;
- $\phi|W_F$ is smooth if F is non-archimedean, and ϕ is continuous if F is archimedean;
- $\phi|\operatorname{SL}_2(\mathbb{C})$ is algebraic.

Here, Frob $\in W_F$ is a (geometric) Frobenius element if F is non-archimedean. When F is a nonarchimedean local field of residue characteristic p > 2, we call a representation ϕ of WD_F unramified if ϕ is trivial on $I_F \times SL_2(\mathbb{C})$, where I_F is the inertia subgroup of W_F . Set

$$\Phi(\operatorname{Sp}_n(F)) = \{\phi \colon WD_F \to \operatorname{SO}_{2n+1}(\mathbb{C})\} / \cong, \\ \Phi(\widetilde{\operatorname{Sp}}_n(F)) = \{\phi \colon WD_F \to \operatorname{Sp}_n(\mathbb{C})\} / \cong.$$

For $G_n = \operatorname{Sp}_n$ or $G_n = \widetilde{\operatorname{Sp}}_n$, we call an element in $\Phi(G_n(F))$ an *L*-parameter for G(F). When $G_n = \operatorname{Sp}_n$ (resp. $G_n = \widetilde{\operatorname{Sp}}_n$), any $\phi \in \Phi(G_n(F))$ can be decomposed into a direct sum

$$\phi = m_1 \phi_1 \oplus \cdots \oplus m_t \phi_t \oplus \phi' \oplus (\phi')^{\vee}$$

where ϕ_1, \ldots, ϕ_t are distinct irreducible orthogonal (resp. symplectic) representations of WD_F , m_i is the multiplicity of ϕ_i in ϕ , and ϕ' is a sum of irreducible representations of WD_F which are not orthogonal (resp. symplectic). We define the component group A_{ϕ} of ϕ by

$$A_{\phi} = \bigoplus_{i=1}^{t} (\mathbb{Z}/2\mathbb{Z})a_i.$$

Namely, A_{ϕ} is a free $\mathbb{Z}/2\mathbb{Z}$ -module of rank t, and $\{a_1, \ldots, a_t\}$ is a basis of A_{ϕ} with a_i associated to ϕ_i . For $a = a_{i_1} + \cdots + a_{i_k} \in A_{\phi}$ with $1 \leq i_1 < \cdots < i_k \leq t$, put

$$\phi^a = \phi_{i_1} \oplus \cdots \oplus \phi_{i_k}.$$

We call $z_{\phi} = \sum_{i=1}^{t} m_i a_i \in A_{\phi}$ the central element of A_{ϕ} . We denote the Pontryagin dual of A_{ϕ} by $\widehat{A_{\phi}} = \{\eta \colon A_{\phi} \to \{\pm 1\}\}.$

For $\phi \in \Phi(\mathrm{Sp}_n(F))$ (resp. $\phi \in \Phi(\widetilde{\mathrm{Sp}}_n(F))$), we say that:

- ϕ is of good parity if ϕ is a direct sum of irreducible orthogonal (resp. symplectic) representations;
- ϕ is tempered if $\phi(W_F)$ is bounded;
- ϕ is almost tempered if each irreducible constituent ϕ_i of ϕ is of the form $\phi_i = \phi'_i |\cdot|^{s_i}$ such that $\phi'_i(W_F)$ is bounded and $-1/2 < s_i < 1/2$.

The set of equivalence classes of irreducible representations of $\widetilde{\operatorname{Sp}}_n(F)$ which are genuine (resp. not genuine) is denoted by $\operatorname{Irr}(\widetilde{\operatorname{Sp}}_n(F))$ (resp. $\operatorname{Irr}(\operatorname{Sp}_n(F))$). The LLC classifies $\operatorname{Irr}(G_n(F))$ by $\Phi(G_n(F))$ for $G_n = \widetilde{\operatorname{Sp}}_n$ or $G_n = \operatorname{Sp}_n$.

Theorem B.1 ([37, 1, 3, 15]). Let $G_n = \widetilde{Sp}_n$ or $G_n = Sp_n$.

(1) There is a canonical surjective map

$$\operatorname{Irr}(G_n(F)) \twoheadrightarrow \Phi(G_n(F)).$$

For $\phi \in \Phi(G_n(F))$, we denote the inverse image of ϕ by Π_{ϕ} , and call it the L-packet of ϕ . (2) There exists an injective map

$$\Pi_{\phi} \hookrightarrow A_{\phi}.$$

This is surjective if $G_n = \widetilde{Sp}_n$. When $G_n = Sp_n$, the image of this map is equal to

$$\left\{\eta\in\widehat{A_{\phi}}\ \Big|\ \eta(z_{\phi})=1\right\}$$

When $\pi \in \Pi_{\phi}$ corresponds to $\eta \in \widehat{A_{\phi}}$, we call the pair (ϕ, η) the L-parameter for π .

- (3) When F is a non-archimedean local field of residue characteristic p > 2, an irreducible representation π is unramified if and only if its L-parameter (ϕ, η) satisfies that ϕ is unramified and $\eta = \mathbf{1}$.
- (4) $\pi \in \Pi_{\phi}$ is (almost) tempered if and only if ϕ is (almost) tempered.
- (5) If $\pi \in \Pi_{\phi}$ is discrete series, then $\phi \in \Phi(G_n(F))$ is of good parity.
- (6) If ϕ is tempered, then one can decompose

$$\phi = \phi_\tau \oplus \phi_0 \oplus \phi_\tau^\vee$$

where

- $\phi_0 \in \Phi(G_{n_0}(F))$ is of good parity;
- ϕ_{τ} is a sum of irreducible representations which are not orthogonal when $G_n = \operatorname{Sp}_n(F)$, and are not symplectic when $G_n = \widetilde{\operatorname{Sp}}_n(F)$.

Let τ be the irreducible (tempered) representation of $\operatorname{GL}_k(F)$ corresponding to ϕ_{τ} . Then for $\pi_0 \in \Pi_{\phi_0}$, the induced representation $\tau \rtimes \pi_0$ is irreducible, and the L-packet Π_{ϕ} is given by

$$\Pi_{\phi} = \{ \tau \rtimes \pi_0 \mid \pi_0 \in \Pi_{\phi_0} \}$$

If the L-parameter for π_0 is (ϕ_0, η_0) , then the one for $\tau \rtimes \pi_0$ is (ϕ, η_0) , where we regard η_0 as a character of A_{ϕ} via the canonical identification $A_{\phi} = A_{\phi_0}$.

(7) If
$$\pi = J(\tau_1|\cdot|^{s_1}, \dots, \tau_t|\cdot|^{s_t}, \pi_0)$$
, then the L-parameter (ϕ, η) for π is given by
 $\phi = \phi_1|\cdot|^{s_1} \oplus \dots \oplus \phi_t|\cdot|^{s_t} \oplus \phi_0 \oplus \phi_t^{\vee}|\cdot|^{-s_t} \oplus \dots \oplus \phi_1^{\vee}|\cdot|^{-s_1}$

and $\eta = \eta_0$, where (ϕ_0, η_0) is the L-parameter for π_0 , and ϕ_i is the (tempered) representation of WD_F corresponding to τ_i for i = 1, ..., t.

B.2. Irreducibility criterion for standard modules. In this subsection, we assume that F is non-archimedean. For each representation ϕ of WD_F , one can consider the *L*-function $L(s, \phi)$ attached to ϕ . We recall a criterion for the irreducibility of standard modules in terms of analytic properties of *L*-functions.

Proposition B.2. Let π_0 (resp. τ_i) be an irreducible tempered representation of $\operatorname{Sp}_{r_0}(F)$ (resp. $\operatorname{GL}_{k_i}(F)$) with the L-parameter (ϕ_0, η_0) (resp. with associated representation ϕ_i of WD_F), and s_1, \ldots, s_t be real numbers such that $s_1 > \cdots > s_t > 0$. Assume the following:

• If π_0 is not genuine, then

$$\left(\prod_{i=1}^{t} L(s-2s_i,\phi_i^{\vee},\wedge^2)L(s-s_i,\phi_0\otimes\phi_i^{\vee})\right)$$
$$\times \left(\prod_{1\leq i< j\leq t} L(s-s_i+s_j,\phi_i^{\vee}\otimes\phi_j)L(s-s_i-s_j,\phi_i^{\vee}\otimes\phi_j^{\vee})\right)$$

is regular at s = 1.

• If π_0 is genuine, then

$$\left(\prod_{i=1}^{t} L(s-2s_i, \phi_i^{\vee}, \operatorname{Sym}^2) L(s-s_i, \phi_0 \otimes \phi_i^{\vee})\right) \times \left(\prod_{1 \le i < j \le t} L(s-s_i+s_j, \phi_i^{\vee} \otimes \phi_j) L(s-s_i-s_j, \phi_i^{\vee} \otimes \phi_j^{\vee})\right)$$

is regular at s = 1.

Then the standard module $\tau_1 | \cdot |^{s_1} \times \cdots \times \tau_t | \cdot |^{s_t} \rtimes \pi_0$ is irreducible.

Proof. Note that our assumptions are equivalent that the adjoint *L*-function $L(s, \phi, \text{Ad})$ is regular at s = 1, where (ϕ, η) is the *L*-parameter for $J(\tau_1 | \cdot |^{s_1}, \ldots, \tau_t | \cdot |^{s_t}, \pi_0)$. The non-genuine case is a result of Heiermann [23] together with a conjecture of Gross–Prasad–Rallis (proven in [13, Appendix B]). The genuine case is [5, Theorem 3.13].

This proposition has several corollaries.

Corollary B.3. If π is irreducible and almost tempered, then π is isomorphic to an irreducible standard module.

Proof. This immediately follows from Proposition B.2.

Corollary B.4. Let μ be a unitary character of F^{\times} . Set $\delta = 0$ if $G_n = \operatorname{Sp}_n(F)$, and $\delta = 1$ if $G_n = \widetilde{\operatorname{Sp}}_n(F)$. Suppose that $\phi \in \Phi(G_n(F))$ is of good parity. Then for any $\pi \in \Pi_{\phi}$ and for any positive integer a with $a \not\equiv \delta \mod 2$, the standard module

$$\mu | \cdot |^{\frac{\alpha}{2}} \rtimes \pi$$

is irreducible.

Proof. Since ϕ is of good parity and $a \neq \delta \mod 2$, we see that $L(s-a/2, \phi \otimes \mu^{-1})$ is regular at s = 1. If $\delta = 0$, then $L(s-a, \mu^{-1}, \wedge^2) = 1$, which is entire. If $\delta = 1$, then $L(s-a, \mu^{-1}, \operatorname{Sym}^2) = L(s-a, \mu^{-2})$, which is regular at s = 1 when $a \neq 1$. By Proposition B.2, we obtain the corollary.

Corollary B.5. Let $\pi \in \Pi_{\phi}$ with ϕ of good parity, If s.s. $R_{P_1(F)}(\pi)$ contains $\chi | \cdot |^{\alpha} \boxtimes \pi_0$ for some unitary character χ and $\alpha \in \mathbb{R}$, then $2\alpha \in \mathbb{Z}$ and $2\alpha \equiv \delta \mod 2$.

Proof. First, we assume that π is discrete series. Replacing π_0 if necessary, we may assume that

$$\pi \hookrightarrow \chi | \cdot |^{\alpha} \rtimes \pi_0.$$

By Casselman's criterion (see e.g., [34, Lemma 2.4]), we see that

- $\alpha > 0;$
- π_0 is discrete series;
- the standard module $\chi | \cdot |^{\alpha} \rtimes \pi_0$ is reducible.

In particular, if $\pi_0 \in \Pi_{\phi_0}$, then $L(s - \alpha, \phi_0 \otimes \chi^{-1})$ has a pole at s = 1, or $L(s - 2\alpha, \chi^{-1}, \wedge^2)$ (resp. $L(s - 2\alpha, \chi^{-1}, \text{Sym}^2)$) has a pole at s = 1 when $\delta = 0$ (resp. $\delta = 1$). Since ϕ_0 is of good parity, this condition implies that $2\alpha \in \mathbb{Z}$ and $2\alpha \equiv \delta \mod 2$.

In general, if $\pi \in \Pi_{\phi}$ with ϕ of good parity, then

$$\pi \hookrightarrow \tau_1 \times \cdots \times \tau_r \rtimes \pi_0,$$

where

- τ_i is an irreducible discrete series representation of $\operatorname{GL}_{k_i}(F)$ to which the irreducible representation ϕ_i of WD_F corresponding is orthogonal if $\delta = 0$, and symplectic if $\delta = 1$;
- π_0 is an irreducible discrete series representation of $\text{Sp}_{n_0}(F)$.

Then the assertion follows from Tadić's formula [51] [20, Proposition 4,5].

B.3. Lowest weight modules. In this subsection, we assume that $F = \mathbb{R}$ and $\psi(x) = \exp(2\pi a \sqrt{-1}x)$ for some a > 0. Recall that we denote the irreducible lowest weight representation of $\widetilde{\mathrm{Sp}}_n(\mathbb{R})$ with lowest \widetilde{K}_{∞} -type $\det^{l/2}$ by $\mathcal{D}_{l/2}^{(n)}$. We determine the *L*-parameters for $\mathcal{D}_{l/2}^{(n)}$ when it is discrete series, which is equivalent that l > 2n. Note that the infinitesimal character of $\mathcal{D}_{l/2}^{(n)}$ is equal to

$$\left(\frac{l}{2}-1,\frac{l}{2}-2,\ldots,\frac{l}{2}-n\right).$$

Recall that the Weil group $W_{\mathbb{R}}$ of \mathbb{R} is of the form

$$W_{\mathbb{R}} = \mathbb{C}^{\times} \cup \mathbb{C}^{\times} j$$

with the group low

$$j^2 = -1, \quad jzj^{-1} = \overline{z}$$

for $z \in \mathbb{C}^{\times}$. Let sgn be the sign character of $W_{\mathbb{R}}$ defined by $\operatorname{sgn}(j) = -1$ and $\operatorname{sgn}(z) = 1$ for $z \in \mathbb{C}^{\times}$. For each positive integer k, we define a 2-dimensional irreducible representation $\rho_k \colon W_{\mathbb{R}} \to \operatorname{GL}_2(\mathbb{C})$ by

$$\rho_k(j) = \begin{pmatrix} 0 & (-1)^k \\ 1 & 0 \end{pmatrix}, \quad \rho_k(re^{\sqrt{-1}\theta}) = \begin{pmatrix} e^{k\sqrt{-1}\theta} & 0 \\ 0 & e^{-k\sqrt{-1}\theta} \end{pmatrix}$$

for r > 0 and $\theta \in \mathbb{R}/2\pi\mathbb{Z}$. Note that (a conjugate of) the image of ρ_k is contained in $O(2, \mathbb{C})$ if k is even, and in $SL_2(\mathbb{C})$ if k is odd.

For an integer l > 2n, we put

$$\phi_l^{(n)} = \begin{cases} \rho_{l-2} \oplus \rho_{l-4} \oplus \dots \oplus \rho_{l-2n} \oplus \operatorname{sgn}^n & \text{if } l \equiv 0 \mod 2, \\ \rho_{l-2} \oplus \rho_{l-4} \oplus \dots \oplus \rho_{l-2n} & \text{if } l \equiv 1 \mod 2. \end{cases}$$

Then $\phi_l^{(n)} \in \Phi(\mathrm{Sp}_n(\mathbb{R}))$ if l is even, and $\phi_l^{(n)} \in \Phi(\widetilde{\mathrm{Sp}}_n(\mathbb{R}))$ if l is odd. When $\phi = \phi_l^{(n)}$, the component group A_{ϕ} is given by

$$A_{\phi} = \left((\mathbb{Z}/2\mathbb{Z})e_{l-2} \oplus (\mathbb{Z}/2\mathbb{Z})e_{l-4} \oplus \cdots \oplus (\mathbb{Z}/2\mathbb{Z})e_{l-2n} \right) + (\mathbb{Z}/2\mathbb{Z})z_{\phi},$$

where $e_k \in A_{\phi}$ is the element associated to $\rho_k \subset \phi$.

Proposition B.6. Suppose that l > 2n. Then the L-parameter for $\mathcal{D}_{l/2}^{(n)}$ is equal to $(\phi_l^{(n)}, \eta_l^{(n)})$, where $\eta_l^{(n)}$ is determined by $\eta_l^{(n)}(e_{l-2i}) = (-1)^{i-1}$ for i = 1, ..., n.

Proof. Set $\phi = \phi_l^{(n)}$. Then the *L*-packet Π_{ϕ} is the set of discrete series representations with infinitesimal character

$$\left(\frac{l}{2}-1,\frac{l}{2}-2,\ldots,\frac{l}{2}-n\right).$$

Hence $\mathcal{D}_{l/2}^{(n)} \in \Pi_{\phi}$. The description of $\eta_l^{(n)}$ can be obtained by a similar way to [6, Appendix A] using Schmid's character identity ([46, (9.4) Theorem] and [47, Theorem (b)]). We omit the detail.

B.4. Arthur's multiplicity formula. In this subsection, we let F be a totally real number field, and ψ be a non-trivial unitary character of \mathbb{A}/F . A discrete global A-parameter for $\operatorname{Sp}_n(F)$ (resp. $\widetilde{\operatorname{Sp}}_n(F)$) is a symbol

$$\Psi = \tau_1[d_1] \boxplus \cdots \boxplus \tau_t[d_t],$$

where

- τ_i is an irreducible cuspidal unitary automorphic representation of $\operatorname{GL}_{m_i}(\mathbb{A})$;
- d_i is a positive integer such that $\sum_{i=1}^{t} m_i d_i$ is equal to 2n+1 (resp. 2n);
- if d_i is odd (resp. even), then $L(s, \tau_i, \text{Sym}^2)$ has a pole at s = 1;
- if d_i is even (resp. odd), then $L(s, \tau_i, \wedge^2)$ has a pole at s = 1;
- the central character ω_i of τ_i satisfies that $\omega_1^{d_1} \cdots \omega_l^{d_l} = 1$;
- if $i \neq j$ and $\tau_i \cong \tau_j$, then $d_i \neq d_j$.

Two A-parameters $\Psi = \bigoplus_{i=1}^{t} \tau_i[d_i]$ and $\Psi' = \bigoplus_{i=1}^{t'} \tau'_i[d'_i]$ are said to be equivalent if t = t' and there exists a permutation $\sigma \in \mathfrak{S}_t$ such that $d'_i = d_{\sigma(i)}$ and $\tau'_i \cong \tau_{\sigma(i)}$ for each i. We denote the set of equivalence classes of discrete global A-parameters for $\operatorname{Sp}_n(F)$ (resp. $\widetilde{\operatorname{Sp}}_n(F)$) by $\Psi_2(\operatorname{Sp}_n(F))$ (resp. $\Psi_2(\widetilde{\operatorname{Sp}}_n(F))$). We call an A-parameter $\Psi = \bigoplus_{i=1}^{t} \tau_i[d_i]$ tempered if $d_i = 1$ for any i. In this case, we write $\Psi = \bigoplus_{i=1}^{t} \tau_i$ for simplicity.

Let $\Psi = \bigoplus_{i=1}^{t} \tau_i$ be a tempered A-parameter for $\operatorname{Sp}_n(F)$ (resp. $\widetilde{\operatorname{Sp}}_n(F)$). For each place v of F, we denote the representation of WD_{F_v} corresponding to $\tau_{i,v}$ by $\phi_{i,v}$, and put $\Psi_v = \bigoplus_{i=1}^{t} \phi_{i,v}$. By [3, Theorem 1.4.2], we have $\Psi_v \in \Phi(\operatorname{Sp}_n(F_v))$ (resp. $\Psi_v \in \Phi(\widetilde{\operatorname{Sp}}_n(F_v))$). We define a global A-packet Π_{Ψ} by

 $\Pi_{\Psi} = \{ \pi = \bigotimes_{v}' \pi_{v} \mid \pi_{v} \in \Pi_{\Psi_{v}}, \ \pi_{v} \text{ is unramified for almost all } v \}.$

For a tempered A-parameter $\Psi = \bigoplus_{i=1}^{t} \tau_i$, the global component group A_{Ψ} of Ψ is defined by

$$A_{\Psi} = \bigoplus_{i=1}^{t} (\mathbb{Z}/2\mathbb{Z})\alpha_i.$$

Namely, A_{Ψ} is a free $\mathbb{Z}/2\mathbb{Z}$ -module of rank t, and $\{\alpha_1, \ldots, \alpha_t\}$ is a basis of A_{Ψ} with α_i associated to τ_i . There exists a localization map $A_{\Psi} \ni \alpha \mapsto \alpha_v \in A_{\Psi_v}$, so that we have a diagonal map

$$\Delta \colon A_{\Psi} \to \prod_{v} A_{\Psi_{v}}.$$

Let $\mathcal{A}_2(\widetilde{\mathrm{Sp}}_n(\mathbb{A}))$ (resp. $\mathcal{A}_2(\mathrm{Sp}_n(\mathbb{A}))$) be the set of square-integrable automorphic forms on $\widetilde{\mathrm{Sp}}_n(\mathbb{A})$ which are genuine (resp. not genuine). This is an $\widetilde{\mathrm{Sp}}_n(\mathbb{A}_{\mathrm{fin}}) \times \prod_{v \mid \infty} (\mathfrak{sp}_n(\mathbb{C}), \widetilde{K}_{\infty})$ -module. Arthur's multiplicity formula classifies $\mathcal{A}_2(\widetilde{\mathrm{Sp}}_n(\mathbb{A}))$ (resp. $\mathcal{A}_2(\mathrm{Sp}_n(\mathbb{A}))$) in terms of $\Psi_2(\widetilde{\mathrm{Sp}}_n(F))$ (resp. $\Psi_2(\mathrm{Sp}_n(F))$).

Theorem B.7 (Arthur's multiplicity formula ([3, Theorem 1.5.2], [14, Theorems 1.1, 1.3])). Let $G = \operatorname{Sp}_n$ or $G = \widetilde{\operatorname{Sp}}_n$.

(1) For each $\Psi \in \Psi_2(G(F))$, there exists an $\widetilde{\operatorname{Sp}}_n(\mathbb{A}_{\operatorname{fin}}) \times \prod_{v \mid \infty} (\mathfrak{sp}_n(\mathbb{C}), \widetilde{K}_{\infty})$ -submodule $\mathcal{A}_{2,\Psi}$ of $\mathcal{A}_2(G(\mathbb{A}))$ such that

$$\mathcal{A}_2(G(\mathbb{A})) = \bigoplus_{\Psi \in \Psi_2(G(F))} \mathcal{A}_{2,\Psi}.$$

(2) Suppose that $\pi = \bigotimes_{v}' \pi_{v}$ is an irreducible subrepresentation of $\mathcal{A}_{2,\Psi}$ with $\Psi = \boxplus_{i=1}^{t} \tau_{i}[d_{i}]$. Then for almost all $v < \infty$, the local factors π_{v} of π and $\tau_{i,v}$ of τ_{i} are unramified for any *i*. Moreover, if we denote the Satake parameter for $\tau_{i,v}$ by $\{c_{i,v,1}, \ldots, c_{i,v,m_{i}}\}$, then the Satake parameter of π_{v} is equal to

$$\bigcup_{i=1}^{t}\bigcup_{j=1}^{m_{i}}\left\{c_{i,v,j}q^{-\frac{d_{i}-1}{2}},c_{i,v,j}q^{-\frac{d_{i}-3}{2}},\ldots,c_{i,v,j}q^{\frac{d_{i}-1}{2}}\right\}$$

as multisets.

(3) When $\Psi = \bigoplus_{i=1}^{t} \tau_i$ is a tempered A-parameter, $\mathcal{A}_{2,\Psi}$ is the multiplicity-free direct sum of representations $\pi = \otimes'_v \pi_v \in \Pi_{\Psi}$ such that the character η_v of A_{Ψ_v} associated to π_v satisfies the equation

$$\left(\prod_{v} \eta_{v}\right) \circ \Delta(\alpha_{i}) = \begin{cases} \mathbf{1} & \text{if } G = \operatorname{Sp}_{n} \\ \varepsilon(\tau_{i}) & \text{if } G = \widetilde{\operatorname{Sp}}_{n} \end{cases}$$

Here, $\varepsilon(\tau_i) = \varepsilon(1/2, \tau_i, \psi) \in \{\pm 1\}$ is the global root number attached to the symplectic representation τ_i .

If a representation π of $\widetilde{\text{Sp}}_n(\mathbb{A})$ is contained in $\mathcal{A}_{2,\Psi}$, we say that π has an A-parameter Ψ , and Ψ is the A-parameter for π . We note that the following.

Lemma B.8. If Ψ is a tempered A-parameter, and $\pi = \bigotimes_{v}' \pi_{v}$ is an irreducible subrepresentation of $\mathcal{A}_{2,\Psi}$, then the local factors π_{v} are almost tempered and unitary for all v.

Proof. Since any representation π contained in $\mathcal{A}_{2,\Psi}$ is unitary, its local factors π_v are unitary for all v. By the toward Ramanujan conjecture (see [30, (2.5) Corollary] and [45, Appendix]), we see that π_v are almost tempered for all v.

Let $l = (l_v)_v \in \prod_{v \mid \infty} \mathbb{Z}$ with $l_v > 0$ and $l_v \equiv l_{v'} \mod 2$ for any $v, v' \mid \infty$. For $\Psi \in \Psi_2(\operatorname{Sp}_n(F))$ if all l_v are even, and for $\Psi \in \Psi_2(\widetilde{\operatorname{Sp}}_n(F))$ if all l_v are odd, we set

$$\mathcal{S}_{l/2,\Psi} = \mathcal{A}_{2,\Psi} \cap \mathcal{S}_{l/2} \left(\operatorname{Sp}_n(F) \setminus \widetilde{\operatorname{Sp}}_n(\mathbb{A}) \right).$$

Hence

$$\mathcal{S}_{l/2}\left(\operatorname{Sp}_n(F)\backslash \widetilde{\operatorname{Sp}}_n(\mathbb{A})\right) \cong \bigoplus_{\Psi \in \Psi_2(G(F))} \mathcal{S}_{l/2,\Psi}.$$

By Proposition B.6, if Ψ is tempered and $S_{l/2,\Psi}$ is nonzero, then $\Psi_v = \phi_{l_v}^{(n)}$ for each $v \mid \infty$. The subspace $S_{l/2,\Psi}$ is often zero.

Example B.9. Suppose that n = 1. There is a unique non-tempered A-parameter $\mathbf{1}[3]$ for $\operatorname{Sp}_1(F)$, where $\mathbf{1}$ is the trivial representation of $\operatorname{GL}_1(\mathbb{A})$. The associated subspace $\mathcal{A}_{2,\mathbf{1}[3]}$ is the space of constant functions, i.e., $\mathcal{A}_{2,\mathbf{1}[3]}$ is the trivial $\operatorname{Sp}_1(\mathbb{A}_{\operatorname{fin}}) \times \prod_{v \mid \infty} (\mathfrak{sp}_1(\mathbb{C}), K_{\infty})$ -module. On the other hand, the non-tempered A-parameters for $\widetilde{\operatorname{Sp}}_1(F)$ are of the form $\chi_{\xi}[2]$. The associated space $\mathcal{A}_{2,\chi_{\xi}[2]}$ is isomorphic to the even Weil representation $\omega_{\psi_{\xi}}^+$ as $\widetilde{\operatorname{Sp}}_1(\mathbb{A}_{\operatorname{fin}}) \times \prod_{v \mid \infty} (\mathfrak{sp}_1(\mathbb{C}), \widetilde{K}_{\infty})$ -modules. Therefore, for $\Psi = \mathbf{1}[3]$ or $\Psi = \chi_{\xi}[2]$, we have $\mathcal{S}_{l/2,\Psi} = 0$ since the trivial representation and the even Weil representations are not cuspidal. In other words, if n = 1 and $\mathcal{S}_{l/2,\Psi} \neq 0$, then Ψ is tempered.

To show Theorem 5.5, we need the following lemma.

Lemma B.10. Let $\tau = \otimes'_v \tau_v$ be an irreducible unitary cuspidal automorphic representation of $\operatorname{GL}_2(\mathbb{A})$ satisfying the conditions (A1), (A2) and (A3) in §2.5. We denote the weight of $\otimes_{v|\infty} \tau_v$ by $\pm 2k = (\pm 2k_v)_v$ with $k_v > 0$. Let $\Psi = \boxplus_{i=1}^t \tau_i$ be a tempered A-parameter for $\widetilde{\operatorname{Sp}}_r(F)$. Assume that for each infinite place $v \mid \infty$, the local factor Ψ_v is equal to $\phi_{l_v}^{(r)}$ with $l_v = 2k_v + 2r + 1$. Then $\Psi' = \Psi \boxplus \tau \chi_{-1}^r$ is a tempered A-parameter for $\widetilde{\operatorname{Sp}}_{r+1}(F)$. Moreover, for any irreducible subrepresentation $\pi = \otimes'_{v < \infty} \pi_v$ of $\mathcal{S}_{l/2,\Psi}$, there exists an irreducible subrepresentation $\pi' = \otimes'_{v < \infty} \pi'_v$ of $\mathcal{S}_{l/2,\Psi'}$ such that for each finite place $v < \infty$, the L-parameters (Ψ_v, η_v) and (Ψ'_v, η'_v) for π and π' , respectively, are related by $\eta'_v = \eta_v$ under the canonical identification $A_{\Psi'_v} = A_{\Psi_v}$.

Proof. For $v \mid \infty$, the local factor $\Psi_v = \phi_{l_v}^{(r)}$ does not contain ρ_{2k_v-1} , which is the local factor of $\tau \chi_{-1}^r$ at v. Since the central character of $\tau \chi_{-1}^r$ is trivial, the exterior square *L*-function $L(s, \tau \chi_{-1}^r, \wedge^2)$ has a pole at s = 1 by [33, Corollary 7.5]. Hence $\Psi' = \Psi \boxplus \tau \chi_{-1}^r$ is a tempered *A*-parameter for $\widetilde{\text{Sp}}_{r+1}(F)$.

Let $\pi = \otimes_{v < \infty}' \pi_v$ be an irreducible subrepresentation of $S_{l/2,\Psi}$, and define $\pi' = \otimes_{v < \infty}' \pi_v'$ so that the *L*-parameter for π_v' is equal to (Ψ_v', η_v) for each $v < \infty$, where (Ψ_v, η_v) is the *L*-parameter for π_v . To show that π' is a subrepresentation of $S_{l/2,\Psi'}$, we use Arthur's multiplicity formula (Theorem B.7 (3)). Note that the *L*-parameters $(\phi_{l_v/2}^{(r)}, \eta_{l_v})$ and $(\phi_{l_v/2}^{(r+1)}, \eta_{l_v}')$ for $\mathcal{D}_{l_v/2}^{(r)}$ and $\mathcal{D}_{l_v/2}^{(r+1)}$ are related by $\eta_{l_v}'|A_{\phi_{l_v/2}^{(r)}} = \eta_{l_v}$ and $\eta_{l_v}'(a_v) = (-1)^r$, where $a_v \in A_{\phi_{l_v/2}^{(r+1)}}$ is the element associated to ρ_{2k_v-1} . Hence for the element $\alpha_i \in A_{\Psi} \subset A_{\Psi'}$ associated to τ_i , we have

$$\left(\prod_{v<\infty}\eta'_v(\alpha_{i,v})\right)\left(\prod_{v\mid\infty}\eta'_{l_v}(\alpha_{i,v})\right) = \left(\prod_{v<\infty}\eta_v(\alpha_{i,v})\right)\left(\prod_{v\mid\infty}\eta_{l_v}(\alpha_{i,v})\right) = \varepsilon(\tau_i),$$

and for the element $\alpha \in A_{\Psi'}$ associated to $\tau \chi_{-1}^r$, we have

$$\left(\prod_{v<\infty}\eta'_v(\alpha_v)\right)\left(\prod_{v\mid\infty}\eta'_{l_v}(\alpha_v)\right) = \left(\prod_{v\mid\infty}(-1)^r\right) = \left(\prod_{v\mid\infty}\chi^r_{-1,v}(-1)\right)$$
$$= \left(\prod_{v<\infty}(\mu_v\chi^r_{-1,v})(-1)\right)\left(\prod_{v\mid\infty}(-1)^{k_v}\right) = \varepsilon(\tau\chi^r_{-1}).$$

Here, we use the condition (A3) on τ . By Arthur's multiplicity formula (Theorem B.7 (3)), we see that π' occurs in $S_{l/2,\Psi'}$.

Appendix C. Gan-Gross-Prasad conjectures

The Gan–Gross–Prasad conjecture (the GGP conjecture) predicts a relation between the nonvanishing of the Fourier–Jacobi periods with the non-vanishing of the central values of the Rankin– Selberg *L*-functions. In this appendix, we review the statements of the GGP conjecture, its refined version and its local version. As an application of local Miyawaki liftings, we prove a new case of the local GGP conjecture in $\SC.3$.

C.1. Global GGP conjectures. Let F be a totally real number field, and ψ be a non-trivial unitary character of \mathbb{A}/F . Recall that for $\xi \in F^{\times}$, the Weil representation $\omega_{\psi_{\xi}}$ of $\widetilde{J}_n(\mathbb{A}) = \widetilde{\mathrm{Sp}}_n(\mathbb{A}) \ltimes$ $V_n(\mathbb{A})$ is realized on $\mathcal{S}(X_n(\mathbb{A}))$. For $\phi \in \mathcal{S}(X_n(\mathbb{A}))$, we define the theta function by

$$\Theta_{\psi_{\xi}}^{\phi}(vg') = \sum_{x \in X_n(F)} \omega_{\psi_{\xi}}(vg')\phi(x)$$

for $v \in V_n(\mathbb{A})$ and $g' \in \widetilde{\mathrm{Sp}}_n(\mathbb{A})$. This is a genuine automorphic form on $\widetilde{J}_n(\mathbb{A})$.

Suppose that n' = n or n' = n - 1. Let φ and φ' be cusp forms on $\widetilde{\operatorname{Sp}}_n(\mathbb{A})$ and $\widetilde{\operatorname{Sp}}_{n'}(\mathbb{A})$, respectively. Assume that exactly one of φ or φ' is genuine. Then for $\phi \in \mathcal{S}(X_{n'}(\mathbb{A}))$, we define the Fourier–Jacobi period $\mathcal{P}_{n,n',\psi_{\mathcal{E}}}(\varphi,\varphi',\phi)$ by

$$\mathcal{P}_{n,n,\psi_{\xi}}(\varphi,\varphi',\phi) = \int_{\operatorname{Sp}_{n}(F)\backslash\operatorname{Sp}_{n}(\mathbb{A})} \varphi(g,\zeta)\varphi'(g,\zeta)\overline{\Theta_{\psi_{\xi}}^{\phi}(g,\zeta)}dg,$$
$$\mathcal{P}_{n,n-1,\psi_{\xi}}(\varphi,\varphi',\phi) = \int_{V_{n-1}(F)\backslash V_{n-1}(\mathbb{A})} \int_{\operatorname{Sp}_{n-1}(F)\backslash\operatorname{Sp}_{n-1}(\mathbb{A})} \varphi(v(g,\zeta))\varphi'(g,\zeta)\overline{\Theta_{\psi_{\xi}}^{\phi}(v(g,\zeta))}dgdv$$

The GGP conjecture is stated as follows:

Conjecture C.1 (Gan–Gross–Prasad conjecture [11, Conjecture 24.1]). Suppose that n' = n or n' = n - 1. Let $\pi = \otimes'_v \pi_v$ and $\pi' = \otimes'_v \pi'_v$ be irreducible cuspidal automorphic representations of $\widetilde{\operatorname{Sp}}_n(\mathbb{A})$ and $\widetilde{\operatorname{Sp}}_{n'}(\mathbb{A})$, respectively. Assume that exactly one of π or π' is genuine, and that the A-parameters for π and π' are tempered. Then the following are equivalent:

- (1) The Fourier-Jacobi period $\mathcal{P}_{n,n',\psi_{\xi}}$ is not identically zero on $\pi \times \pi' \times \mathcal{S}(X_{n'}(\mathbb{A}))$.
- (2) The central value $L(1/2, \pi \times \pi' \times \chi_{\xi})$ is nonzero and the local Hom-space

$$\operatorname{Hom}_{\operatorname{Sp}_{n'}(F_v)}(\pi_v \otimes \pi'_v \otimes \overline{\omega_{\psi_{\xi,v}}}, \mathbb{C})$$

is nonzero for any place v of F.

For the case where n' = n, Yamana [58, Theorem 1.1] proved that (1) implies (2). See also [58, Remark 6.6]. Conjecture C.1 implies Conjecture 5.2 (1) since the A-parameter for $\overline{\pi'}$ is given by

$$\begin{cases} \Psi' = \boxplus_{j=1}^{t'} \tau'_j & \text{if } \pi' \text{ is not genuine,} \\ \Psi' \otimes \chi_{-1} = \boxplus_{j=1}^{t'} (\tau'_j \otimes \chi_{-1}) & \text{if } \pi' \text{ is genuine,} \end{cases}$$

where $\Psi' = \bigoplus_{j=1}^{t'} \tau'_j$ is the A-parameter for π' .

To obtain Conjecture 5.2 (2), we need a refined version of Conjecture C.1. To state this, we define a local period integral. For each place v, we fix a Haar measure on $\operatorname{Sp}_{n'}(F_v)$ such that $\operatorname{vol}(\operatorname{Sp}_{n'}(\mathfrak{o}_v); dg'_v) = 1$ for almost all v, and that $dg' = \prod_v dg'_v$ is the Tamagawa measure on $\operatorname{Sp}_{n'}(\mathbb{A})$. Fix an $\operatorname{Sp}_n(F_v)$ -invariant inner product $\langle , \rangle_{\pi_v}$ of $\pi_v \times \pi_v$ such that

$$\int_{\operatorname{Sp}_n(F)\backslash \operatorname{Sp}_n(\mathbb{A})} \varphi_1(g) \overline{\varphi_2(g)} dg = \prod_v \langle \varphi_{1,v}, \varphi_{2,v} \rangle_{\pi_v}$$

for $\varphi_1 = \otimes_v \varphi_{1,v}, \varphi_2 = \otimes_v \varphi_{2,v} \in \pi$. Similarly, fix an $\operatorname{Sp}_{n'}(F_v)$ -invariant inner product $\langle, \rangle_{\pi'_v}$ of $\pi'_v \times \pi'_v$. Recall that $\omega_{\psi_{\xi,v}}$ has a $\widetilde{J}_{n'}(F_v)$ -invariant pairing

$$(\phi_{1,v},\phi_{2,v})_v = \int_{X_{n'}(F_v)} \phi_{1,v}(x_v) \overline{\phi_{2,v}(x_v)} dx_v,$$

which satisfies that $(\phi_1, \phi_2) = \prod_v (\phi_{1,v}, \phi_{2,v})_v$ for $\phi_1 = \bigotimes_v \phi_{1,v}, \phi_2 = \bigotimes_v \phi_{2,v} \in \mathcal{S}(X_{n'}(\mathbb{A}))$. For $\varphi_v \in \pi_v, \varphi'_v \in \pi'_v$ and $\phi_v \in \mathcal{S}(X_{n'}(F_v))$, we define a local period integral by $\alpha_v(\varphi_v, \varphi'_v, \phi_v)$

$$= \begin{cases} \int_{\operatorname{Sp}_{n}(F_{v})} \langle \pi_{v}(g)\varphi_{v},\varphi_{v}\rangle_{\pi_{v}} \langle \pi'_{v}(g)\varphi'_{v},\varphi'_{v}\rangle_{\pi'_{v}} \overline{\langle \omega_{\psi_{\xi,v}}(g)\phi_{v},\phi_{v}\rangle_{v}} dg & \text{if } n' = n, \\ \int_{V_{n-1}(F_{v})} \int_{\operatorname{Sp}_{n-1}(F_{v})} \langle \pi_{v}(ug)\varphi_{v},\varphi_{v}\rangle_{\pi_{v}} \langle \pi'_{v}(g)\varphi'_{v},\varphi'_{v}\rangle_{\pi'_{v}} \overline{\langle \omega_{\psi_{\xi,v}}(ug)\phi_{v},\phi_{v}\rangle_{v}} dg du & \text{if } n' = n-1 \end{cases}$$

This integral is absolutely convergent by [56, Proposition 2.2.1].

The refined version of the GGP conjecture is stated as follows:

Conjecture C.2 (Xue [56, Conjecture 2.3.1]). Let $\pi = \bigotimes'_v \pi_v$ and $\pi' = \bigotimes'_v \pi'_v$ be irreducible cuspidal automorphic representations of $\widetilde{\operatorname{Sp}}_n(\mathbb{A})$ and $\widetilde{\operatorname{Sp}}_{n'}(\mathbb{A})$ with tempered A-parameters Ψ and Ψ' , respectively. Assume that exactly one of π or π' is genuine. Fix a sufficiently large finite set S of places of F containing all "bad" places.

- (1) For each place v, the local Hom-space Hom_{Sp_n'(F_v)} $(\pi_v \otimes \pi'_v \otimes \overline{\omega}_{\psi_{\xi,v}}, \mathbb{C})$ is nonzero if and only if $\alpha_v(\varphi_v, \varphi'_v, \phi_v) \neq 0$ for some $\varphi_v \in \pi_v$, $\varphi'_v \in \pi'_v$ and $\phi_v \in \omega_{\psi_{\xi,v}}$.
- (2) Let $\varphi = \otimes_v \varphi_v \in \pi$, $\varphi' = \otimes_v \varphi'_v \in \pi'$, and $\phi = \otimes_v \phi_v \in \mathcal{S}(X_{n'}(\mathbb{A}))$ be factorizable elements. Then there exists a constant Δ^S , which is an explicit product of partial zeta values, such that

$$\begin{aligned} |\mathcal{P}_{n,n',\psi_{\xi}}(\varphi,\varphi',\phi)|^{2} \\ &= \frac{2\Delta^{S}}{|A_{\Psi}||A_{\Psi'}|} \left. \frac{L^{S}(s,\pi\times\pi'\times\chi_{\xi})}{L^{S}(s+\frac{1}{2},\pi,\operatorname{Ad})L^{S}(s+\frac{1}{2},\pi,\operatorname{Ad})} \right|_{s=\frac{1}{2}} \times \prod_{v\in S} \alpha_{v}(\varphi_{v},\varphi'_{v},\phi_{v}) \end{aligned}$$

Here, $L^{S}(s, \pi \times \pi' \times \chi_{\xi})$ is the partial Rankin–Selberg L-function, and $L^{S}(s, \pi, \text{Ad})$ and $L^{S}(s, \pi', \text{Ad})$ are the partial adjoint L-functions of π and π' , respectively.

Conjecture C.2 together with the following lemma implies Conjecture 5.2 (2).

Lemma C.3. Assume that n' = n. Let v be a real place of F, and ξ be a positive real number (in F_v). Then for nonzero lowest weight vectors $\varphi_{l/2} \in \mathcal{D}_{l/2}^{(n)}$, $\varphi_{(l-1)/2} \in \mathcal{D}_{(l-1)/2}^{(n)}$, and $\phi_{\xi}^0 \in \omega_{\psi_{\xi,v}}$, we have $\alpha_v(\varphi_{l/2}, \overline{\varphi_{(l-1)/2}}, \phi_{\xi}^0) \neq 0$.

Proof. Set $\pi_v = \mathcal{D}_{l/2}^{(n)}$ and $\pi'_v = \mathcal{D}_{(l-1)/2}^{(n)}$. Note that the subspace of $\mathcal{D}_{(l-1)/2}^{(n)} \otimes \omega_{\psi_{\xi,v}}$ on which \widetilde{K}_{∞} acts by det^{l/2} is one dimension, and is spanned by $\varphi_{(l-1)/2} \otimes \phi_{\xi}^0$. Since $\operatorname{Hom}_{\widetilde{\operatorname{Sp}}_n(\mathbb{R})}(\mathcal{D}_{l/2}^{(n)}, \mathcal{D}_{(l-1)/2}^{(n)} \otimes \omega_{\psi_{\xi,v}}) \neq$ 0, we may regard $\varphi_{(l-1)/2} \otimes \phi_{\xi}^0$ as a lowest weight vector of $\mathcal{D}_{l/2}^{(n)} \subset \mathcal{D}_{(l-1)/2}^{(n)} \otimes \omega_{\psi_{\xi,v}}$. Hence we can regard

$$\langle \pi'_{v}(g)\varphi_{(l-1)/2},\varphi_{(l-1)/2}\rangle_{\pi'_{v}}(\omega_{\psi_{\xi,v}}(g)\phi^{0}_{\xi},\phi^{0}_{\xi})_{v}$$

as a (nonzero) matrix coefficient of $\mathcal{D}_{l/2}^{(n)}$ on which the left-right translation of $\widetilde{K}_{\infty} \times \widetilde{K}_{\infty}$ is equal to $\det^{l/2} \boxtimes \det^{l/2}$. Since such a matrix coefficient is a scalar multiple of $\langle \pi_v(g)\varphi_{l/2},\varphi_{l/2}\rangle_{\pi_v}$, we have $\alpha_v(\varphi_{l/2},\overline{\varphi_{(l-1)/2}},\phi_{\xi}^0) \neq 0$.

C.2. Local GGP conjecture. In this subsection, we let F be a non-archimedean local field of characteristic zero. Fix a non-trivial additive character ψ of F and an element $\xi \in F^{\times}$. Let π_1 and π_2 be irreducible representations of $\widetilde{\operatorname{Sp}}_r(F)$ and $\widetilde{\operatorname{Sp}}_{r-1}(F)$, respectively. Assume that exactly one of them is genuine. We set

$$d_{r,r-1,\xi}(\pi_1,\pi_2) = \dim \operatorname{Hom}_{J_{r-1}(F)}(\pi_1 \otimes \pi_2 \otimes \overline{\omega_{\psi_{\xi}}^{(r-1)}}, \mathbb{C}).$$

Similarly, let π'_1 and π'_2 be irreducible representations of $\widetilde{\operatorname{Sp}}_n(F)$. Assume that exactly one of them is genuine. We set

$$d_{n,n,\xi}(\pi'_1,\pi'_2) = \dim \operatorname{Hom}_{\operatorname{Sp}_n(F)}(\pi'_1 \otimes \pi'_2 \otimes \omega_{\psi_{\xi}}^{(n)}, \mathbb{C}).$$

The multiplicity one theorem proven by Sun [48] and Gan–Gross–Prasad [11, Corollary 16.2] asserts that

$$d_{r,r-1,\xi}(\pi_1,\pi_2) \le 1, \quad d_{n,n,\xi}(\pi'_1,\pi'_2) \le 1$$

for any $\pi_1, \pi_2, \pi'_1, \pi'_2$.

For a symplectic representation ϕ of WD_F , we denote the root number attached to ϕ by $\varepsilon(\phi) = \varepsilon(1/2, \phi, \psi)$. This value does not depend on ψ , and is in $\{\pm 1\}$. The local GGP conjecture for the symplectic-metaplectic case (proven by the author [5]) gives complete descriptions of $d_{r,r-1,\xi}(\pi_1, \pi_2)$ and $d_{n,n,\xi}(\pi'_1, \pi'_2)$ in terms of internal structures of *L*-packets when all $\pi_1, \pi_2, \pi'_1, \pi'_2$ are almost tempered.

Theorem C.4 ([5, Theorem 1.3, Corollary 1.4], [11, Proposition 18.1]). Assume that all $\pi_1, \pi_2, \pi'_1, \pi'_2$ are almost tempered. Let (ϕ_1, η_1) , (ϕ_2, η_2) , (ϕ'_1, η'_1) , (ϕ'_2, η'_2) be the L-parameters for $\pi_1, \pi_2, \pi'_1, \pi'_2$, respectively. Then $d_{r,r-1,\xi}(\pi_1, \pi_2) \neq 0$ if and only if

$$\begin{cases} \eta_1(a) = \varepsilon(\phi_1^a \otimes \phi_2 \otimes \chi_{\xi})\varepsilon(\phi_1 \otimes \phi_2 \otimes \chi_{\xi})^{\det(a)} \det(\phi_1^a)((-1)^{\frac{1}{2}\dim(\phi_2)}\xi) \\ \eta_2(b) = \varepsilon(\phi_1 \otimes \phi_2^b \otimes \chi_{\xi})\varepsilon(\phi_2^b)\chi_{\xi}(-1)^{\frac{1}{2}\dim(\phi_2^b)} \end{cases}$$

for $a \in A_{\phi_1}$ and $b \in A_{\phi_2}$, and $d_{n,n,\xi}(\pi'_1,\pi'_2) \neq 0$ if and only if

$$\begin{cases} \eta_1'(a) = \varepsilon(\phi_1'^a \otimes \phi_2' \otimes \chi_{\xi})\varepsilon(\phi_1' \otimes \phi_2' \otimes \chi_{\xi})^{\det(a)} \det(\phi_1'^a)((-1)^{\frac{1}{2}\dim(\phi_2')}\xi), \\ \eta_2'(b) = \varepsilon(\phi_1' \otimes \phi_2'^b \otimes \chi_{\xi})\varepsilon(\phi_2'^b)\chi_{\xi}(-1)^{\frac{1}{2}\dim(\phi_2'^b)} \end{cases}$$

for $a \in A_{\phi'_1}$ and $b \in A_{\phi'_2}$.

C.3. Application of local Miyawaki liftings. As in Proposition 3.10, local Miyawaki liftings satisfy a seesaw identity. Using this, one can describe $d_{n,n,\xi}(\pi'_1,\pi'_2)$ for a new case. Recall that the unique irreducible algebraic representation of $SL_2(\mathbb{C})$ of dimension d is denoted by S_d .

Theorem C.5. Let π_1 and π_2 be irreducible almost tempered unitary representations of $\widetilde{\operatorname{Sp}}_r(F)$ and $\widetilde{\operatorname{Sp}}_{r-1}(F)$, on which $\{\pm 1\}$ acts by $(\pm 1)^{n+r}$ and $(\pm 1)^{n+r-1}$, respectively. Fix a unitary character μ of F^{\times} .

- (1) Assume one of the following conditions:
 - The L-parameter ϕ_1 for π_1 does not contain $\mu^{\pm 1}S_d$ for any $d \ge n r$ with $d \equiv n r \mod 2$;
 - n = r or n = r + 1.

Then we have

$$d_{n,n,\xi}((\mu\chi_{-1}^{n+r-1} \circ \det_{n-r}) \rtimes \pi_1, (\mu\chi_{\xi} \circ \det_{n-r+1}) \rtimes \pi_2) = d_{r,r-1,\xi}(\pi_1, \pi_2).$$

In particular, $d_{n,n,\xi}((\mu\chi_{-1}^{n+r-1}\circ\det_{n-r})\rtimes\pi_1,(\mu\chi_{\xi}\circ\det_{n-r+1})\rtimes\pi_2)$ can be described in terms of internal structures of the L-packets for π_1 and π_2 .

(2) Set $\mu' = \mu \chi_{-1}^{[(n+r-1)/2]}$ and $\tau' = \mu' \times \mu'^{-1}$. If an irreducible representation π'_1 of $\widetilde{\operatorname{Sp}}_n(F)$ satisfies that $d_{n,n,\xi}(\pi'_1, (\mu\chi_{\xi} \circ \det_{n-r+1}) \rtimes \pi_2) \neq 0$, then $\mathcal{M}_{\psi,\tau'}^{(r)}(\pi'_1) \neq 0$.

Proof. Note that $\mu \chi_{-1}^{n+r-1} = \mu' \chi_{-1}^{[(n+r)/2]}$. By Theorem 3.1 (2), (4) and the seesaw identity (Proposition 3.10), we have

$$d_{n,n,\xi}((\mu\chi_{-1}^{n+r-1} \circ \det_{n-r}) \rtimes \pi_{1}, (\mu\chi_{\xi} \circ \det_{n-r+1}) \rtimes \pi_{2}) \neq 0$$

$$\iff d_{n,n,\xi}\left(\mathcal{M}_{\psi,\tau'}^{(n)}(\pi_{1}), \mathcal{M}_{\psi,\tau'\chi_{\xi}}^{(n)}(\pi_{2})\right) \neq 0$$

$$\iff \operatorname{Hom}_{\widetilde{\operatorname{Sp}}_{n}(F)}\left(\mathcal{M}_{\psi,\tau'\chi_{\xi}}^{(n)}(\pi_{2}^{\vee}) \otimes \omega_{\psi_{\xi}}^{(n)}, \mathcal{M}_{\psi,\tau'}^{(n)}(\pi_{1})\right) \neq 0$$

$$\iff \operatorname{Hom}_{\widetilde{J}_{r-1}}\left(\mathcal{M}_{\psi,\tau'}^{(r)}\left(\mathcal{M}_{\psi,\tau'}^{(n)}(\pi_{1})\right), \pi_{2}^{\vee} \otimes \omega_{\psi_{\xi}}^{(r-1)}\right) \neq 0$$

$$\iff \operatorname{Hom}_{\widetilde{J}_{r-1}}\left(\pi_{1}, \pi_{2}^{\vee} \otimes \omega_{\psi_{\xi}}^{(r-1)}\right) \neq 0$$

$$\iff d_{r,r-1,\xi}(\pi_{1}, \pi_{2}) \neq 0.$$

Hence we obtain (1).

When $d_{n,n,\xi}(\pi'_1, (\mu\chi_{\xi} \circ \det_{n-r+1}) \rtimes \pi_2) \neq 0$, the seesaw identity implies that

$$\operatorname{Hom}_{\widetilde{J}_{r-1}}\left(\mathcal{M}_{\psi,\tau'}^{(r)}(\pi_1'),\pi_2^{\vee}\otimes\omega_{\psi}^{(r-1)}\right)\neq 0.$$

In particular, $\mathcal{M}_{\psi,\tau'}^{(r)}(\pi_1') \neq 0$. Hence we have (2).

References

- J. Adams and D. Barbasch, Genuine representations of the metaplectic group, Compositio Math. 113 (1998), no. 1, 23–66.
- H. Aoki, Estimating Siegel modular forms of genus 2 using Jacobi forms, J. Math. Kyoto Univ. 40 (2000), no. 3, 581–588.
- [3] J. Arthur, The endoscopic classification of representations. Orthogonal and symplectic groups, American Mathematical Society Colloquium Publications, 61. American Mathematical Society, Providence, RI, 2013. xviii+590 pp.
- M. Asgari and R. Schmidt, Siegel modular forms and representations, Manuscripta Math. 104 (2001), no. 2, 173–200.
- [5] H. Atobe, The local theta correspondence and the local Gan-Gross-Prasad conjecture for the symplecticmetaplectic case, Math. Ann. 371 (2018), no. 1-2, 225–295.
- [6] H. Atobe, On the non-vanishing of theta liftings of tempered representations of U(p,q), arXiv:1610.07794v1.
- [7] H. Atobe and W. T. Gan, Local theta correspondence of tempered representations and Langlands parameters, Invent. Math. 210 (2017), no. 2, 341–415.
- [8] D. Blasius, Hilbert modular forms and the Ramanujan conjecture, Noncommutative geometry and number theory, 35–56, Aspects Math., E37, Friedr. Vieweg, Wiesbaden, 2006.
- C. Bushnell, Representations of reductive p-adic groups: localization of Hecke algebras and applications, J. London Math. Soc. (2) 63 (2001), no. 2, 364–386.
- [10] A. Caraiani, Local-global compatibility and the action of monodromy on nearby cycles, Duke Math. J. 161 (2012), no. 12, 2311–2413.
- [11] W. T. Gan, B. H. Gross and D. Prasad, Symplectic local root numbers, central critical L-values, and restriction problems in the representation theory of classical groups, Astérisque. No. 346 (2012), 1–109.
- [12] W. T. Gan and N. Gurevich, Restrictions of Saito-Kurokawa representations. With an appendix by Gordan Savin, Contemp. Math., 488, Israel Math. Conf. Proc., Automorphic forms and L-functions I. Global aspects, 95–124, Amer. Math. Soc., Providence, RI, 2009.
- [13] W. T. Gan and A. Ichino, The Gross-Prasad conjecture and local theta correspondence, Invent. Math. 206 (2016), no. 3, 705–799.

52

- [14] W. T. Gan and A. Ichino, The Shimura–Waldspurger correspondence for Mp_{2n} , Ann. of Math. (2) 188 (2018), no. 3, 965–1016.
- [15] W. T. Gan and G. Savin, Representations of metaplectic groups I: epsilon dichotomy and local Langlands correspondence, Compos. Math. 148 (2012), no. 6, 1655–1694.
- [16] W. T. Gan and S. Takeda, On the Howe duality conjecture in classical theta correspondence, Advances in the theory of automorphic forms and their L-functions, 105–117, Contemp. Math., 664, Amer. Math. Soc., Providence, RI, 2016.
- [17] P. Garrett and B. Heim, Hecke duality of Ikeda lifts, J. Number Theory 146 (2015), 171–186.
- [18] D. Ginzburg, S. Rallis and D. Soudry, On explicit lifts of cusp forms from GL_m to classical groups, Ann. of Math. (2) **150** (1999), no. 3, 807–866.
- [19] D. Ginzburg, S. Rallis and D. Soudry, On Fourier coefficients of automorphic forms of symplectic groups, Manuscripta Math. 111 (2003), no. 1, 1–16.
- [20] M. Hanzer and G. Muić, Parabolic induction and Jacquet functors for metaplectic groups, J. Algebra 323 (2010), no. 1, 241–260.
- M. Hanzer and G. Savin, Eisenstein series arising from Jordan algebras, Canadian Journal of Mathematics, http://dx.doi.org/10.4153/CJM-2018-033-2.
- [22] S. Hayashida, Lifting from two elliptic modular forms to Siegel modular forms of half-integral weight of even degree, Doc. Math. 21 (2016), 125–196.
- [23] V. Heiermann, A note on standard modules and Vogan L-packets, Manuscripta Math. 150 (2016), no. 3-4, 571–583.
- [24] T. Ibukiyama, C. Poor, and D. S. Yuen, Jacobi forms that characterize paramodular forms, Abh. Math. Semin. Univ. Hambg. 83 (2013), no. 1, 111–128.
- [25] A. Ichino, Pullbacks of Saito-Kurokawa lifts, Invent. Math. 162 (2005), no. 3, 551–647.
- [26] T. Ikeda, On the theory of Jacobi forms and Fourier-Jacobi coefficients of Eisenstein series, J. Math. Kyoto Univ. 34 (1994), no. 3, 615–636.
- [27] T. Ikeda, On the lifting of elliptic cusp forms to Siegel cusp forms of degree 2n, Ann. of Math. (2) 154 (2001), no. 3, 641–681.
- [28] T. Ikeda, Pullback of the lifting of elliptic cusp forms and Miyawaki's conjecture, Duke Math. J. 131 (2006), no. 3, 469–497.
- [29] T. Ikeda and S. Yamana, On the lifting of Hilbert cusp forms to Hilbert-Siegel cusp forms, Ann. Sci. Ec. Norm. Sup. (4), to appear.
- [30] H. Jacquet and J. A. Shalika, On Euler products and the classification of automorphic representations. I, Amer. J. Math. 103 (1981), no. 3, 499–558.
- [31] C. Jantzen, Degenerate principal series for symplectic and odd-orthogonal groups, Mem. Amer. Math. Soc. 124 (1996), no. 590, viii+100 pp.
- [32] D. Jiang and L. Zhang, Arthur parameters and cuspidal automorphic modules of classical groups, arXiv:1508.03205v3.
- [33] P. K. Kewat and R. Raghunathan, On the local and global exterior square L-functions of GL_n, Math. Res. Lett. 19 (2012), no. 4, 785–804.
- [34] T. Konno, A note on the Langlands classification and irreducibility of induced representations of p-adic groups, Kyushu J. Math. 57 (2003), no. 2, 383–409.
- [35] S. S. Kudla and S. Rallis, Ramified degenerate principal series representations for Sp(n), Israel J. Math. **78** (1992), no. 2-3, 209–256.
- [36] S. S. Kudla and S. Rallis, On first occurrence in the local theta correspondence, Automorphic representations, L-functions and applications: progress and prospects, 273–308, Ohio State Univ. Math. Res. Inst. Publ., 11, de Gruyter, Berlin, 2005.
- [37] R. P. Langlands, On the classification of irreducible representations of real algebraic groups, Representation theory and harmonic analysis on semisimple Lie groups, 101–170, Math. Surveys Monogr., 31, Amer. Math. Soc., Providence, RI, 1989.
- [38] E. Lapid and S. Rallis, On the local factors of representations of classical groups, Automorphic representations, L-functions and applications: progress and prospects, 309–359, Ohio State Univ. Math. Res. Inst. Publ., 11, de Gruyter, Berlin, 2005.
- [39] I. Miyawaki, Numerical examples of Siegel cusp forms of degree 3 and their zeta-functions, Mem. Fac. Sci. Kyushu Univ. Ser. A 46 (1992), no. 2, 307–339.
- [40] C. Mœglin, M.-F. Vignéras and J.-L. Waldspurger, Correspondances de Howe sur un corps p-adique, Lecture Notes in Mathematics, 1291. Springer-Verlag, Berlin, 1987. viii+163 pp.

- [41] C. Mœglin and J.-L. Waldspurger, Le spectre résiduel de GL(n), Ann. Sci. École Norm. Sup. (4) 22 (1989), no. 4, 605–674.
- [42] P. D. Nelson and A. Venkatesh, The orbit method and analysis of automorphic forms, arXiv:1805.07750v2.
- [43] I. I. Piatetski-Shapiro and S. Rallis, L-functions for the classical groups, Explicit constructions of automorphic L-functions, Lecture Notes in Mathematics 1254, Springer-Verlag, Berlin, 1987, pp. 1–52.
- [44] Y. Qiu, Periods of Saito-Kurokawa representations, Int. Math. Res. Not. IMRN 2014 (2014), no. 24, 6698–6755.
- [45] Z. Rudnick and P. Sarnak, Zeros of principal L-functions and random matrix theory, Duke Math. J. 81 (1996), no. 2, 269–322.
- [46] W. Schmid, On the characters of the discrete series. The Hermitian symmetric case, Invent. Math. 30 (1975), no. 1, 47–144.
- [47] W. Schmid, Two character identities for semisimple Lie groups, Non-commutative harmonic analysis (Actes Colloq., Marseille-Luminy, 1976), pp. 196–225. Lecture Notes in Math., Vol. 587, Springer, Berlin, 1977.
- [48] B. Sun, Multiplicity one theorems for Fourier-Jacobi models, Amer. J. Math. 134 (2012), no. 6, 1655– 1678.
- [49] B. Sun, On representations of real Jacobi groups, Sci. China Math. 55 (2012), no. 3, 541–555.
- [50] W. J. Sweet, Jr., Functional equations of p-adic zeta integrals and representations of the metaplectic group, preprint (1995).
- [51] M. Tadić, Structure arising from induction and Jacquet modules of representations of classical p-adic groups, J. Algebra 177 (1995), no. 1, 1–33.
- [52] M. Tadić, On reducibility of parabolic induction, Israel J. Math. 107 (1998), 29–91.
- [53] J.-L. Waldspurger, Correspondances de Shimura et quaternions, Forum Math. 3 (1991), no. 3, 219–307.
- [54] N. R. Wallach, On the constant term of a square integrable automorphic form, Operator algebras and group representations, Vol. II (Neptun, 1980), 227–237, Monogr. Stud. Math., 18, Pitman, Boston, MA, 1984.
- [55] M. H. Weissman, The Fourier-Jacobi map and small representations, Represent. Theory 7 (2003), 275– 299.
- [56] H. Xue, Refined global Gan-Gross-Prasad conjecture for Fourier-Jacobi periods on symplectic groups, Compos. Math. 153 (2017), no. 1, 68–131.
- [57] H. Xue, Central values of degree six L-functions, J. Number Theory 203 (2019), 350–359.
- [58] S. Yamana, Periods of automorphic forms: the trilinear case, J. Inst. Math. Jussieu 17 (2018), no. 1, 59–74.
- [59] A. V. Zelevinsky, Induced representations of reductive p-adic groups. II. On irreducible representations of GL(n), Ann. Sci. École Norm. Sup. (4) 13 (1980), no. 2, 165–210.

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, KITA 10, NISHI 8, KITA-KU, SAPPORO, HOKKAIDO, 060-0810, JAPAN

Email address: atobe@math.sci.hokudai.ac.jp