Title	Melatonin suppression during a simulated night shift in medium intensity light is increased by 10-minute breaks in dim light and decreased by 10-minute breaks in bright light
Author(s)	Lee, Sang-II; Kinoshita, Saki; Noguchi, Anna; Eto, Taisuke; Ohashi, Michihiro; Nishimura, Yuki; Maeda, Kaho; Motomura, Yuki; Awata, Yasuhiro; Higuchi, Shigekazu
Citation	Chronobiology International, 37(6), 897-909 https://doi.org/10.1080/07420528.2020.1752704
Issue Date	2020-06
Doc URL	http://hdl.handle.net/2115/81619
Rights	This is an Accepted Manuscript of an article published by Taylor & Francis in Chronobiology International on June 2020, available online: http://www.tandfonline.com/10.1080/07420528.2020.1752704 .
Type	article (author version)
File Information	LCBI-2019-0234.R2_Proof_hi2.pdf

[Instructions for use](#)

Melatonin suppression during a simulated night shift in medium intensity light is increased by 10-minute breaks in dim light and decreased by 10-minute breaks in bright light

Journal:	<i>Chronobiology International</i>
Manuscript ID	LCBI-2019-0234.R2
Manuscript Type:	Original Reports
Date Submitted by the Author:	n/a
Complete List of Authors:	LEE, SANG-IL; Kyushu University Graduate School of Design Faculty of Design, Department of Human Science; Hokkaido University Faculty of Engineering, Division on Human Environmental Systems Kinoshita, Saki; Kyushu University, Graduate School of Integrated Frontier Science, Department of Kansei Science Noguchi, Anna; Kyushu University, Graduate School of Integrated Frontier Science, Department of Kansei Science Eto, Taisuke; Kyushu University, Graduate School of Integrated Frontier Science, Department of Kansei Science Ohashi, Michihiro; Kyushu University, Graduate School of Integrated Frontier Science, Department of Kansei Science Email address Nishimura, Yuki; Kyushu University, Graduate School of Integrated Frontier Science, Department of Kansei Science Email address; National Institute of Occupational Safety and Health Japan Noborito District Maeda, Kaho; Japan Aerospace Exploration Agency Institute of Aerospace Technology Tsukuba Space Center, Ground Facilities Department Motomura, Yuki; Kyushu University Graduate School of Design Faculty of Design, Department of Human Science Awata, Yasuhiro; Japan Aerospace Exploration Agency Institute of Aerospace Technology Tsukuba Space Center, Ground Facilities Department Higuchi, Shigekazu; Kyushu University Graduate School of Design Faculty of Design, Department of Human Science
Keywords:	Humans, Night shift work, Intermittent light at night exposure, Light adaptation, Short duration, Melatonin suppression, Subjective sleepiness, Performance

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

SCHOLARONE™
Manuscripts

**Melatonin suppression during a simulated night shift in medium
intensity light is increased by 10-minute breaks in dim light and
decreased by 10-minute breaks in bright light**

Sang-il Lee^{a, b}, Saki Kinoshita^c, Anna Noguchi^c, Taisuke Eto^c, Michihiro
Ohashi^c, Yuki Nishimura^{c, e}, Kaho Maeda^d, Yuki Motomura^a, Yasuhiro
Awata^d, and Shigekazu Higuchi^{a*}

*^aDepartment of Human Science, Faculty of Design, Kyushu University, Fukuoka,
Japan; ^bDivision of Human Environmental Systems, Faculty of Engineering, Hokkaido
University, Sapporo, Japan; ^cDepartment of Kansei Science, Graduate School of
Integrated Frontier Science, Kyushu University, Fukuoka, Japan; ^dGround Facilities
Department, Japan Aerospace Exploration Agency, Tsukuba, Japan; ^eNational Institute
of Occupational Safety and Health, Kawasaki, Japan*

*corresponding author:

Shigekazu Higuchi, Ph.D
Department of Human Science
Faculty of design, Kyushu University
4-9-1 Shiobaru, Minami-ku, Fukuoka, 815-8540, Japan

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Exposure to light at night results in disruption of endogenous circadian rhythmicity and/or suppression of pineal melatonin, which can consequently lead to acute or chronic adverse health problems. In the present study, we investigated whether exposure to very dim light or very bright light for a short duration influences melatonin suppression, subjective sleepiness, and performance during exposure to constant moderately bright light. Twenty-four healthy male university students were divided into two experimental groups: Half of them (mean age: 20.0 ± 0.9 years) participated in an experiment for short-duration (10 min) light conditions of medium intensity light (430 lx, medium breaks) vs. very dim light (< 1 lx, dim breaks) and the other half (mean age: 21.3 ± 2.5 years) participated in an experiment for short-duration light conditions of medium intensity light (430 lx, medium breaks) vs. very bright light (4700 lx, bright breaks). Each simulated night shift consisting of 5 sets (each including 50-minute night work and 10-minute break) was performed from 01:00 to 06:00h. The subjects were exposed to medium intensity light (550 lx) during the night work. Each 10-minute break was conducted every hour from 02:00 to 06:00h. Salivary melatonin concentrations were measured, subjective sleepiness was assessed, the psychomotor vigilance task was performed at hourly intervals from 21:00h until the end of the experiment. Compared to melatonin suppression between 04:00 and 06:00h in the condition of medium breaks, the condition of dim breaks significantly promoted melatonin suppression and the condition of bright breaks significantly diminished melatonin suppression. However, there was no remarkable effect of either dim breaks or bright breaks on subjective sleepiness and performance of the psychomotor vigilance task. Our findings suggest that periodic exposure to light for short durations during exposure to a constant light environment affects the sensitivity of pineal melatonin to constant light depending on the difference between light intensities in the two light conditions (i.e., short light exposure vs. constant light exposure). Also, our findings indicate that exposure to light of various intensities at night could be a factor influencing the light-induced melatonin suppression in real night work settings.

Keywords: Humans; Night shift work; Intermittent light at night exposure; Light adaptation; Short duration; Melatonin suppression; Subjective sleepiness; Performance

35 Introduction

36 The 24-h light-dark cycle in nature is known as the strongest zeitgeber (i.e., time giver)
37 for almost all mammals. In humans, the suprachiasmatic nuclei (SCN), i.e., the
38 circadian pacemaker, regulates circadian rhythmicity throughout the body via
39 phototransduction input from ganglion cells in the retina (Weaver 1998). For instance,
40 the SCN restrains pineal melatonin production during the daytime and allows melatonin
41 secretion during the night. It has been believed that the release timing and the amount of
42 melatonin secretion are related to the regulation of physiological and behavioral
43 circadian rhythm (Macchi & Bruce 2004). However, the extension of daylight duration
44 in modern life by using artificial lighting at night is likely to lead to acute melatonin
45 suppression and circadian disturbances, which are partially responsible for some health
46 problems (Smolensky et al. 2015; Smolensky et al. 2016; Lunn et al. 2017; Touitou et
47 al. 2017).

48 The magnitude of melatonin suppression varies depending on light intensity
49 (Zeitzer et al. 2000), exposure duration (Aoki et al. 1998), and wavelength composition
50 (Brainard et al. 2001). Recently, however, there is growing evidence that prior light
51 history has an impact on the magnitude of melatonin suppression in response to a
52 subsequent light stimulus at night. Several studies have suggested that less daylight can
53 increase melatonin sensitivity to light at night (Hebert et al. 2002; Smith et al. 2004). A
54 field study with human subjects showed that melatonin suppression in response to 500
55 lx light at night was greater following exposure to dim light (wearing dark goggles with
56 2% transmission lenses) for one week than following exposure to bright light for one
57 week (Hebert et al. 2002). Similar results were obtained in laboratory studies on
58 melatonin suppression (Smith et al. 2004) and circadian phase shift (Chang et al. 2011).
59 The results of those studies, however, were likely to have been affected by great
60 differences in the intensities of prior light conditions. One study, however, showed
61 significantly dampened melatonin suppression in response to blue light (460 nm
62 monochromatic light) following 2-h exposure to dim white light (18 lx) compared to
63 that following 2-h dark adaptation (Jasser et al. 2006). Taken together, the results
64 suggest that melatonin sensitivity to a light stimulus can be increased or decreased
65 depending on the relative intensity of a prior light stimulus to the target light stimulus.

66 Although the mechanism involved in the effect of prior light history is not
67 clear, it seems that a long period of photic adaptation alters the absolute response
68 threshold of photoreceptors and/or photosensory inputs from the photoreceptors to the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

SCN. In humans, the visual photoreceptors (i.e., rods and cones) and especially a small subset of retinal ganglion cells expressing melanopsin (mRGC) contribute to non-visual effects on the circadian system (Gooley et al. 2012); those are designed to adapt to various changes in ambient light in a short time. For instance, exposure of the eyes to bright light desensitizes the visual photoreceptors to facilitate response to intensity increment (i.e., light adaptation) (Fain et al. 2001). Conversely, darkness fully recovers the photoreceptors from the desensitized state (i.e., dark adaptation) (Lamb & Pugh 2004). It should be noted, however, that the time required for photo-regeneration becomes longer as the photopigment bleaches more. Recently, evidence of the adaptation capacity of mRGC to light has been provided (Wong et al. 2005). One study with young human subjects, for example, showed that blocking short-wavelength light, which dramatically activates the mRGC, by using orange-colored contact lenses (i.e., blue light-filtering lenses) immediately reduced melatonin suppression but that the reduction in melatonin suppression disappeared 16 days after wearing the blue light-filtering lenses (Gimenez et al. 2014).

          If two lights of different intensities are emitted alternately for long or short durations, can the effects of the short duration exposure be ignored? Some researchers have demonstrated that using intermittent light (e.g., alternate exposure to bright light and darkness) during the night for multiple days is effective for shifting the phase of human circadian rhythm (Baehr et al. 1999; Rimmer et al. 2000; Crowley et al. 2003; Gronfier et al. 2004; Smith et al. 2009). A mathematical model of the effects of brief light on the human circadian pacemaker has been proposed for explaining the results of previous investigations (Kronauer et al. 1999). Recent studies, however, showed that even brief (12 minutes or shorter) exposure to bright light was able to elicit circadian phase delay and melatonin suppression after previous adaptation to dim light during a constant routine (Chang et al. 2012; Rahman et al. 2017).

          Although there is a possibility that exposure to darkness or dimmer light or brighter light can alter (at least temporarily) the sensitivity of non-visual responses to a subsequent light stimulus, the intermittent light conditions used in most previous studies are far from real working conditions. Furthermore, the effects of short exposure to very dim light during night work under light of constant medium intensity are unknown. In the present study, we therefore investigated whether short-duration exposure to very dim light or very bright light influences melatonin suppression, subjective sleepiness, and performance during exposure to constant light of medium intensity.

103

104 **Materials and Methods**

105 *Subjects*

106 Twenty-four healthy male university students participated in this study. Twelve subjects
107 (mean $\pm$ SD age: 20.0 $\pm$ 0.9 years) participated in Experiment 1 and the remaining 12
108 subjects (mean $\pm$ SD age: 21.3 $\pm$ 2.5 years) participated in Experiment 2. None of the
109 participants showed extreme morningness or extreme eveningness as assessed by a
110 Japanese version of the Morningness-Eveningness Questionnaire (Ishihara et al. 1984).
111 Subjects who had engaged in night shift work or who had experienced time zone travel
112 (i.e., at least > 1 time zone) in the previous three months were excluded from the study.
113 Signed written informed consent to take part in the research study, which was approved
114 by the Ethical Committee of Kyushu University, was obtained from all participants. The
115 experiments were conducted in accordance with the Declaration of Helsinki.

117 *Experimental light conditions*

118 The vertical illuminance and irradiance of each light condition were measured at eye
119 level in a sitting position using an illuminance spectroradiometer (CL-500A, KONICA
120 MINOLTA INC., Japan). Two experimental chambers were used for the experiment
121 (chamber 1) and for 10 min breaks (chamber 2). In chamber 1, light-emitting diode
122 (LED) ceiling lights (HH-LC569A, Panasonic Inc., Japan) were set up for exposure to
123 constant medium intensity light (~550 lx). Fluorescent ceiling lights (FPL36CW, Panasonic
124 Inc., Japan) were used for medium breaks (~430 lx) or bright breaks (~4700 lx). For dim
125 breaks (<1 lx), incandescent bulbs were installed on the floor as indirect lighting.
126 Detailed information on each light condition is given in Table 1. Melanopic lux was
127 calculated using an excel-based toolbox provided by the Lucas Group at the University
128 of Manchester (Enezi et al. 2011).
129 (<http://lucasgroup.lab.manchester.ac.uk/measuringmelanopicilluminance/>).

131 *Procedure*

132 Two experiments with a crossover design, including dim breaks vs. medium breaks
133 (Experiment 1) and bright breaks vs. medium breaks (Experiment 2), were conducted to
134 investigate the effects of each short-duration (10 min) light exposure condition on
135 melatonin suppression, subjective sleepiness, and performance during exposure to
136 constant bright light. Each participant was therefore required to visit our laboratory

twice with an interval of 2 weeks. Six participants simultaneously participated in each experiment. Prior to the experiment, participants were instructed to sleep for more than 7 hours between 00:00 and 08:00h for one week. An accelerometry-based activity monitor (Lifecorder plus, Suzuken Co Ltd, Japan) and daily sleep diary were used to confirm the implementation of sleep intervention during the control period.

Furthermore, each participant sent us a message via a mobile phone shortly before bedtime and shortly after waking up. If there was no message from a participant, we called the participant to confirm his situation. The participants were instructed not to drink alcohol from three days before the experiment. Also, excessive exercise, napping, and caffeine consumption were not allowed commencing the day before the experiment.

The procedures used for Experiment 1 and Experiment 2 were same (Figure 1). The participants arrived at the experimental facility at about 12:00h and dressed into the experimental clothes (short-sleeved T-shirt, short pants, and no socks) after receiving brief instructions for the experiment. The participants stayed in chamber 1 in a sitting position from 13:00h until the end of the experiment. The room illuminance was 275 lx (vertical illuminance at eye level) from 13:00 to 19:00h. The light illuminance was then changed into a dim light (< 10 lx) from 19:00 to 01:00h. The participants were allowed to use portable devices (e.g., smartphone, tablet pc, laptop, set to minimum brightness) and read books between 19:00 and 00:00h. For reference, the illuminances of the self-illuminating portable devices were measured under 2 lx at a distance of 20 cm from the center of each screen. The participants started simulated night work in a sitting position from 00:00 to 06:00h (practice session between 00:00 and 01:00h) in chamber 1. The light illuminance was changed to medium intensity light (550 lx) from 01:00 until 06:00h. Each break condition was conducted every hour from 02:00 to 06:00h in chamber 2 and lasted 10 min. During each break, all of the participants performed light stretching (~2 min in a standing position) and a word chain game (Shiritori, ~5 min in a sitting position) and answered questionnaires (~3 min in a sitting position) that addressed subjective sleepiness (Karolinska sleepiness scale, KSS) and 'dummy variables' (e.g., physical fatigue and mood state). Each break was followed by 50-min of simulated night work in which the participants conducted PVT and answered the KSS and dummy questionnaires for 10 min followed by card games for 20 min. After a 5-min rest period, the participants conducted PVT, collected saliva, and answered the KSS for 15 min (Figure 1b). The order of the break conditions in each experiment (i.e., Experiment 1 or Experiment 2) was random for each participant. For example, some participants

performed the medium breaks on the first visit and the bright breaks (or dim breaks) on the next visit, and vice versa for other participants. Participants had dinner at 19:00h (typical Japanese food, the same dishes for all participants in every experiment) and a late-night snack at 23:30h (rice, miso soup). No drinks except for water were provided throughout each experiment.

Salivary samples were collected hourly using a plain cotton plug (Salivette Sarstedt, Germany) from 21:00h until the end of the experiment. Participants did not drink any water for 15 min prior to each salivary sample collection. Subjective sleepiness was evaluated using the KSS at 1-hour intervals from 21:00 to 06:00h, 5 min before collecting the salivary sample. Subjective sleepiness during each break was also evaluated to confirm the acute effects of the break on sleepiness. Each participant performed the Psychomotor Vigilance Task (PVT) twice with a 1-hour interval, soon after the break and shortly before the next break. The PVT was performed for 5 min using Presentation (Neurobehavioral Systems Inc., Albany, CA). A visual stimulus was displayed randomly on the PC display at intervals of 2 to 10 sec. Participants were instructed to press the space key on the keyboard as soon as possible after the appearance of the visual stimulus. A beep sound was emitted from each earphone if the participants did not react within 3000 msec.

Sample analysis

Salivary melatonin concentrations were measured by radioimmunoassay kit (RK-DSM; Buhlmann Laboratories AG, Allschwil, Switzerland). Melatonin area under the curve (AUC; trapezoidal approximation) between 21:00 and 06:00h was calculated to evaluate the overall effect of each break condition on melatonin suppression. Data of three participants were excluded from analysis of the results for bright breaks, since each participant showed a gap longer than 1 hour in the time of dim light melatonin onset (DLMO) between the conditions (medium breaks vs. bright breaks). DLMO was determined by linear interpolation between two time points at which melatonin concentration crossed the 3.0 pg/ml threshold (Benloucif et al. 2008).

Statistical analysis

In statistical comparisons between the conditions (medium break vs. bright break or medium break vs. dim break) for the melatonin profile, subjective sleepiness (KSS), performance (PVT), repeated-measures two-way ANOVA (SPSS 23.0, IBM® SPSS®

Statistics) with light conditions and time (during the simulated night work) as independent factors was conducted. Greenhouse-Geisser correction was performed when Mauchly's sphericity assumption was largely violated. A two-sided, paired Student's t-test was used for planned comparisons between the light conditions during the night work when a significant interaction between the independent factors was found. For the comparison of numbers of PVT lapses between the conditions, the Wilcoxon signed-rank test was conducted. A P-value of less than 0.05 was considered statistically significant.

Results

Melatonin suppression

Figure 2 shows the melatonin profiles obtained from each experiment. In both experiments, melatonin gradually increased under the dim light condition (21:00-01:00h) but was immediately attenuated by light exposure (550 lx) from 01:00h in both experiments. However, the aspects of melatonin suppression were different with the light conditions during breaks.

In Experiment 1 (Figure 2a), repeated-measures two-way ANOVA with light condition (medium break vs. dim break) and time (01:00~06:00h) showed a main effect in light condition ($F_{1, 11} = 6.966, P = 0.027$) but not in time ($F_{1.951, 21.463} = 0.511, P = 0.603, ns$). There was a significant interaction between condition and time ($F_{2.827, 31.095} = 3.826, P = 0.021$). A paired t-test for melatonin concentrations at each time point showed that the dim break resulted in greater melatonin suppression than did the medium break at 04:00, 05:00, and 06:00h ($P = 0.011, P = 0.007$, and $P = 0.001$, respectively). A comparison of the melatonin AUCs between the conditions showed that there was a significant tendency for lower melatonin concentration in the dim break condition compared to that in the medium break condition.

Similarly, in Experiment 2 (Figure 2b), repeated-measures two-way ANOVA with light condition (medium break vs. bright break) and time (01:00~06:00h) showed a main effect in light condition ($F_{1, 8} = 9.837, P = 0.014$) but not in time ($F_{5, 40} = 0.981, P = 0.441, ns$). A significant interaction between condition and time was found ($F_{5, 40} = 4.484, P = 0.002$). A paired t-test for melatonin concentrations at each time point showed that the bright break resulted in lower melatonin suppression than did the medium break at 04:00, 05:00, and 06:00h ($P = 0.009, P = 0.038$, and $P = 0.001$, respectively). There was also a significant tendency for higher melatonin concentration

in the bright breaks than in the medium breaks at 05:00h (paired t-test, $P = 0.050$). The melatonin AUC in the bright breaks was significantly greater than that in the medium breaks.

Subjective sleepiness (KSS)

The results for subjective sleepiness are shown in Figure 3. In Experiment 1 (dim breaks), subjective sleepiness gradually increased over time during the simulated night work span ($F_{8, 34.031} = 9.066$; $p < 0.001$), but it showed almost the same pattern in the conditions (medium vs. dim) ($F_{1, 11} = 0.723$; $p = 0.413$). In addition, no significant interaction was found between conditions and time ($F_{8, 42.824} = 1.709$; $p = 0.167$). Similarly, in Experiment 2 (bright breaks), there was a significant main effect of time ($F_{8, 64} = 14.030$; $p < 0.001$), but no main effect of condition ($F_{1, 8} = 0.018$; $p = 0.897$), and no interaction between condition and time ($F_{8, 64} = 1.195$; $p = 0.316$) were found.

PVT

Figure 4 shows the results for reaction speed (mean reciprocal reaction time: mean 1/RT) (Basner & Dinges 2011) in the two experiments. In ANOVA analysis for mean 1/RT, there were significant main effects of time in Experiment 1 (dim breaks) ($F_{8, 27.005} = 8.729$; $p = 0.001$) and in Experiment 2 (bright breaks) ($F_{8, 16.421} = 11.593$; $p = 0.001$). However, mean 1/RT was not significantly different between conditions in both experiments ($F_{1, 11} = 0.0004$; $p = 0.984$ in Experiment 1, $F_{1, 8} = 0.018$; $p = 0.897$ in Experiment 2). Also, no interactions between conditions and time were found ($F_{8, 88} = 0.927$; $p = 0.499$ in Experiment 1, $F_{8, 22.129} = 1.073$; $p = 0.377$ in Experiment 2).

A comparison of the numbers of lapses at each time point (Wilcoxon signed-rank test) showed that there was no significant difference at any time points between medium breaks and dim breaks and between medium breaks and bright breaks (Figure 4).

Discussion

In the present study, we investigated whether periodic short-duration exposures (for 10 min at hourly intervals) to very bright (bright breaks) or very dim light (dim breaks) affect physiological responses including melatonin suppression, subjective sleepiness, and performance during exposure to constant medium intensity light. We found that both the dim breaks and bright breaks indirectly, rather than directly, affected melatonin suppression

273 during the experiments: the dim breaks promoted melatonin suppression, whereas the
274 bright breaks diminished melatonin suppression. A possible reason for these results is
275 that the dim breaks or the bright breaks sensitized or desensitized pineal melatonin to
276 the subsequent constant light exposure during the experiment.

277 The effect of brief light exposure during each break on melatonin synthesis was
278 first observed at the 04:00h time point, just before the 3rd break in both experiments,
279 and the sensitized or desensitized states lasted until the end of the experiment (i.e.,
280 06:00h). However, this does not mean that the first break had no effect at all, but rather
281 it seems that adaptation to dim light for 4 hours before the start of the experiment
282 caused strong melatonin suppression by sensitizing melatonin responsiveness to light
283 (see melatonin suppressions at 02:00h in Figure 2). This may indicate that melatonin
284 sensitivity can be changed shortly after short-duration exposure to light. Our results
285 might suggest that using brief photo-adaptation probably enables real-time adjustment
286 of melatonin sensitivity against a current light stimulus depending on relative photic
287 intensity.

288 Although the results of the present study are similar to previous findings of prior
289 light history having an effect on melatonin (see Introduction section), the underlying
290 mechanisms responsible for the results appear to be different. In previous studies, it is
291 more likely to be a compensatory adaptation by long-term adaptation of photoreceptors
292 to a given photo-environment. On the other hand, although the mechanisms by which
293 short adaptations during breaks contribute to melatonin sensitivity are not known, they
294 seem to be associated with bleaching and recovery mechanism of photoreceptors in the
295 retina. For instance, the bleached photopigments might be partially recovered by the
296 dim breaks, leading to an increase in gain of the phototransduction cascade. Likewise,
297 profound bleaching of a substantial fraction of the photopigments due to exposure to
298 very bright light during the bright break might lead to attenuation of melatonin
299 sensitivity to the subsequent light (Fain et al. 2001). On the other hand, visual
300 photoreceptors, including rods and cones, saturate at a relatively low-intensity level of
301 light (Lucas et al. 2003). Given that mRGC compensate the functional limitations of the
302 visual photoreceptors for higher light intensities (Gooley et al. 2012), it might be more
303 important to understand whether the mRGC has such capacity of light adaptation. The
304 photopigment melanopsin has been shown to be homologous to invertebrate opsin
305 (rhododermic opsin) (Shichida & Matsuyama 2009), and it has therefore been
306 hypothesized that melanopsin uses the rhododermic phototransduction cascade (Hillman

et al. 1983). Although controversial, several previous studies have provided evidence of a bi- or tri-stable signaling state in mammalian melanopsin including, for example, red-light enhancement for pupil response to blue light (Graham et al. 2008; Mure et al. 2009; Emanuel & Do 2015).

Another in vitro study demonstrated that prior light stimulus alters the sensitivity of rat mRGC to subsequent light exposure in a way similar to that of photoreceptor adaptation, rather than neural network adaptation: a brief flash desensitized the cells whereas darkness re-sensitized the cells without synaptic inputs from rods and cones (Wong et al. 2005). According to the study, mRGC completed light adaptation (i.e., desensitization) within 5 min. On the other hand, the kinetics of dark adaptation appeared to be even slower for mRGC than for rods, as the cell showed a striking increase in sensitivity after 30–40 min of dark adaptation and kept increasing for at least 2 h and 40 min. More recent studies, however, have shown that synaptic inputs from the classical photoreceptors (i.e., rods and cones) via the inner plexiform layer to mRGC increase the sensitivity of mRGC to light (Wong et al. 2007). Based on these results, exposure to very bright light for 10 min (i.e., bright breaks) might be sufficient to cause a massive decrease in the photosensitivity of mRGC, and this phenomenon likely lead to the attenuation of melatonin suppression in Experiment 1. However, recovery in near darkness for 10 min (i.e., dim breaks) was probably not sufficient to elicit a significant increase in the photosensitivity of mRGC; rather, synaptically mediated signals from partially dark-adapted classical photoreceptors might be more responsible for the promotion of melatonin suppression in Experiment 2.

There were no remarkable effects of dim breaks or bright breaks on alertness, i.e., subjective sleepiness and performance of the psychomotor vigilance task. As an indirect alerting effect of light via retinal projection to the SCN, the magnitude of melatonin suppression was thought to be involved in subjective sleepiness (Cajochen 2007). Nonetheless, subjective sleepiness or reaction speed (i.e., mean 1/RT) was consistently increased or decreased over time in a similar pattern regardless of the break conditions. Similarly, unlike medium breaks, bright breaks and dim breaks did not have an additional effect on the number of lapses. On the other hand, as a direct alerting effect of light (Souman et al. 2018), the bright breaks were expected to be able to delay the decrease in alertness. However, we could not find such a beneficial effect even when we compared the reaction velocities or numbers of lapses before and after the bright breaks (Figure 4b and Figure 5b). Moreover, although acute reduction of

1
2
3 341 subjective sleepiness tended to emerge during each break session, it seems to be a
4 342 temporal effect associated with moving to a break room or a light stretch during each
5 343 break session, rather than than a direct alerting effect of light during each break.
6
7 344 Nevertheless, it should be noted that the participants were continuously exposed to
8 345 medium intensity light during the simulated night work, and this might have diluted the
9 346 additional effects of bright breaks or dim breaks on alertness considering the dose-response
10 347 relationship between light intensity and alertness (Cajochen et al. 2000). Also, high
11 348 sleep pressure due to prior wakefulness might be partially responsible for the results,
12 349 since the participants did not take a nap before the start of the experiments.
13
14 350 Epidemiological studies conducted over the past few decades have suggested
15 351 adverse relationships of night shift work with acute and chronic adverse health
16 352 problems (Kantermann et al. 2010; Parent et al. 2012; Evans & Davidson 2013; Kamdar
17 353 et al. 2013). Direct effects of exposure to bright white light, especially blue-enriched
18 354 light, on the circadian system, such as melatonin suppression and circadian
19 355 misalignment, for example, between the biological clock and the social-behavioral cycle
20 356 have been suspected as factors involved in the risks (Wittmann et al. 2006; Touitou et
21 357 al. 2017). Nonetheless, in some ways, exposure to bright white light (i.e., blue-enriched
22 358 light) is also helpful for keeping night workers awake and providing better visibility,
23 359 leading to better performance and fewer accidents due to human errors (Cajochen 2007;
24 360 Chellappa et al. 2011; Kraneburg et al. 2017). For attenuation of melatonin suppression
25 361 without a negative effect on performance, the use of lighting with less short-wavelength
26 362 components (Kozaki et al. 2008) and wearing blue light-filtering goggles (Kayumov et
27 363 al. 2005), or a red-visor cap (Higuchi et al. 2011), have been proposed. In addition to
28 364 these proposals, a countermeasure for night shift workers is also suggested by our
29 365 findings that bright breaks can reduce melatonin suppression by light without having
30 366 adverse effects on sleepiness or performance.
31
32 367 However, a field study in which the effect of bright light exposure during a short
33 368 break (~20 min) in night work on melatonin was investigated showed no such
34 369 desensitization in melatonin suppression. In that field study, the subjects showed greater
35 370 melatonin suppression in night work when they took a break with exposure to bright
36 371 light (2500 lx) than when they took a break with exposure to normal light (300 lx)
37 372 (Lowden et al. 2004). However, a limitation of that field study is that the timing and
38 373 duration of the breaks were not strictly controlled. Another limitation is that the subjects
39 374 were allowed to leave the workplace for a short period. These limitations, however,

rather remind us about a question if similar results could be obtained by conducting bright breaks in a real night workplace. Additionally, it is uncertain in the present study whether the hourly repetitive execution of breaks was essential to achieve persistent effects on melatonin suppression. Although taking rest breaks is known to be effective for decreasing accident risks, recovering from physical fatigue and maintaining arousal level, taking breaks more than once per hour tends to disturb work (Tucker 2003). In this regard, it is essential to clarify the minimum number of breaks that is necessary to obtain the same results as those in the present study.

In the present study, although the pace of melatonin synthesis was remarkably diminished after light exposure (~550 lx), we did not observe dramatic melatonin suppression as found in some previous studies using a protocol and illuminance level similar to the present study. For example, McIntyre et al. (1989) reported that 1-h light exposure (500 lx) from midnight caused about 40% suppression of melatonin compared to the melatonin concentration just before light exposure (Mcintyre et al. 1989). Laakso et al. (1993) reported that melatonin suppression following 1-h light exposure from 23:00h amounted to as much as 53% (Laakso et al. 1993). Ethnicity might be partially responsible for the inconsistency in melatonin suppression induced by nocturnal light exposure. Higuchi et al. (2007) reported that melatonin suppression following 2-h light exposure (1000 lx) was greater in Caucasian than Asian subjects (Higuchi et al. 2007a). Although Aoki et al. (1998) also found that melatonin suppression amounted to as much as 40.1% following 2-h light exposure (500 lx) in Asian subjects (Aoki et al. 1998), dark adaptation by 5-h sleep before the light exposure possibly influenced the result (Jasser et al. 2006). Subjects in the previous study were directly exposed to a specially designed light source in a fixed position. However, in the present study, we used ceiling light, and the gaze of each participant was not strictly fixed; hence, the light intensity reaching the retina might have been less than 550 lx in the present study.

There is a question that remains unanswered: Can bright breaks or dim breaks modify the circadian phase shift caused by light exposure? There have been practical interventions using intermittent light to entrain the circadian clock of shift workers to long-term night shift duty (Baehr et al. 1999; Crowley et al. 2003; Smith et al. 2009; Smith & Eastman 2012). Intermittent light was used for multiple days in those previous studies, mainly to delay the phase of the circadian pacemaker. However, the results for melatonin in the present study indicate the possibility that circadian phase delay during night work can not only be promoted by conducting dim breaks, but it can also be

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

attenuated by conducting bright breaks. Given the greater melatonin suppression in the condition of dim breaks than in the condition of medium breaks, conducting dim breaks during night work can probably cause a larger phase delay than can continuous exposure to medium intensity light. Modulation of circadian phase to both advance and delay might be easier by conducting bright or dim breaks based on the human phase response curve (St Hilaire et al. 2012).

This study has several limitations. We measured salivary melatonin levels at hourly intervals; hence, acute effects of breaks on melatonin could not be determined in the present study. All of the participants in this study were healthy young male adults. However, inter-individual differences in the sensitivity of pineal melatonin have been shown in previous studies. Although there is still lack of agreement, there has been an accumulation of evidence indicating an age-dependent difference in pineal melatonin sensitivity (Charman 2003; Higuchi et al. 2014; Lee et al. 2018). Also, one study has suggested greater sensitivity in females than in males (Monteleone et al. 1995). The experiments in this study were conducted in different seasons: The experiment for dim breaks was conducted in summer (July), while the experiment for bright breaks was conducted in winter (from January to February). It has been reported that melatonin suppression by light at night is greater in winter than in summer (Higuchi et al. 2007b). Therefore, it is necessary to verify the reproducibility of our findings for different seasons.

In addition, inter-individual differences in the photo-sensitivity of pineal melatonin have been shown in previous studies (Higuchi et al. 2008; Santhi et al. 2012; Phillips et al. 2019). Indeed, in the present study, some participants showed strong melatonin suppression during light exposure, while others, especially participants who had a relatively low melatonin level at 01:00h (e.g., below 10 pg/ml), showed weak melatonin suppression. Furthermore, some participants showed quick recovery from the melatonin suppression and an increase in melatonin concentration over time. It remains unclear what causes the individual differences, but several recent studies have suggested that genetic variations in the clock genes are associated with inter-individual differences in melatonin suppression (Chellappa et al. 2012; Akiyama et al. 2017). Further investigation should be carried out to identify the individual differences in non-visual photo-sensitivity.

Our findings suggest that periodic exposure to light for a short duration during exposure to constant light affects melatonin sensitivity to the constant light depending

on the difference between light intensities in the light conditions (i.e., exposure to short light vs. exposure to constant light). In most previous studies, the effects of light with fixed intensity and/or spectral composition on the circadian system were investigated. However, humans generally do not stay at the same place for long duration; the light environment surrounding us frequently changes in real life. In this regard, the findings in the present study suggest that exposure to light of various intensities at night could be a factor influencing the light-induced melatonin suppression in real life.

Acknowledgments

We thank Takuya Tsuyama at Kyushu University for his help in the preparation of this study. This work was supported by the Japan Society for the Promotion of Science under Grant-in-Aid for Scientific Research (KAKENHI) number 19H03316.

Declaration of interest statement

The authors report no conflict of interest.

References

- Akiyama T, Katsumura T, Nakagome S, Lee SI, Joh K, Soejima H, Fujimoto K, Kimura R, Ishida H, Hanihara T, Yasukouchi A, Satta Y, Higuchi S, Oota H. 2017. An ancestral haplotype of the human PERIOD2 gene associates with reduced sensitivity to light-induced melatonin suppression. *Plos One*. 12.
- Aoki H, Yamada N, Ozeki Y, Yamane H, Kato N. 1998. Minimum light intensity required to suppress nocturnal melatonin concentration in human saliva. *Neurosci Lett*. 252:91-94.
- Baehr EK, Fogg LF, Eastman CI. 1999. Intermittent bright light and exercise to entrain human circadian rhythms to night work. *Am J Physiol-Reg I*. 277:R1598-R1604.
- Basner M, Dinges DF. 2011. Maximizing Sensitivity of the Psychomotor Vigilance Test (PVT) to Sleep Loss. *Sleep*. 34:581-591.
- Benloucif S, Burgess HJ, Klerman EB, Lewy AJ, Middleton B, Murphy PJ, Parry BL, Revell VL. 2008. Measuring Melatonin in Humans. *J Clin Sleep Med*. 4:66-69.
- Brainard GC, Hanifin JP, Greeson JM, Byrne B, Glickman G, Gerner E, Rollag MD. 2001. Action spectrum for melatonin regulation in humans: evidence for a novel circadian photoreceptor. *J Neurosci*. 21:6405-6412.
- Cajochen C. 2007. Alerting effects of light. *Sleep Med Rev*. 11:453-464.
- Cajochen C, Zeitzer JM, Czeisler CA, Dijk DJ. 2000. Dose-response relationship for light intensity and ocular and electroencephalographic correlates of human alertness. *Behav Brain Res*. 115:75-83.
- Chang AM, Santhi N, St Hilaire M, Gronfier C, Bradstreet DS, Duffy JF, Lockley SW, Kronauer RE, Czeisler CA. 2012. Human responses to bright light of different durations. *J Physiol-London*. 590:3103-3112.
- Chang AM, Scheer FAJL, Czeisler CA. 2011. The human circadian system adapts to prior photic history. *J Physiol-London*. 589:1095-1102.
- Charman WN. 2003. Age, lens transmittance, and the possible effects of light on melatonin suppression. *Ophthalmic Physiol Opt*. 23:181-187.
- Chellappa SL, Steiner R, Blattner P, Oelhafen P, Gotz T, Cajochen C. 2011. Non-Visual Effects of Light on Melatonin, Alertness and Cognitive Performance: Can Blue-Enriched Light Keep Us Alert? *PLoS One*. 6.
- Chellappa SL, Viola AU, Schmidt C, Bachmann V, Gabel V, Maire M, Reichert CF, Valomon A, Gotz T, Landolt HP, Cajochen C. 2012. Human Melatonin and Alerting Response to Blue-Enriched Light Depend on a Polymorphism in the Clock Gene PER3. *J Clin Endocrinol Metab*. 97:E433-E437.
- Crowley SJ, Lee C, Tseng CY, Fogg LF, Eastman CI. 2003. Combinations of bright light, scheduled dark, sunglasses, and melatonin to facilitate circadian entrainment to night shift work. *J Biol Rhythms*. 18:513-523.
- Emanuel AJ, Do MTH. 2015. Melanopsin Tristability for Sustained and Broadband Phototransduction. *Neuron*. 85:1043-1055.
- Enezi J, Revell V, Brown T, Wynne J, Schlangen L, Lucas R. 2011. A "melanopic" spectral efficiency function predicts the sensitivity of melanopsin photoreceptors to polychromatic lights. *J Biol Rhythms*. 26:314-323.
- Evans JA, Davidson AJ. 2013. Health Consequences of Circadian Disruption in Humans and Animal Models. *Prog Mol Biol Transl*. 119:283-323.
- Fain GL, Matthews HR, Cornwall MC, Koutalos Y. 2001. Adaptation in vertebrate photoreceptors. *Physiol Rev*. 81:117-151.

- 507 Gimenez MC, Beersma DG, Bollen P, van der Linden ML, Gordijn MC. 2014. Effects
508 of a chronic reduction of short-wavelength light input on melatonin and sleep
509 patterns in humans: evidence for adaptation. *Chronobiol Int.* 31:690-697.
- 510 Gooley JJ, Mien IH, St Hilaire MA, Yeo SC, Chua ECP, van Reen E, Hanley CJ, Hull
511 JT, Czeisler CA, Lockley SW. 2012. Melanopsin and Rod-Cone Photoreceptors
512 Play Different Roles in Mediating Pupillary Light Responses during Exposure to
513 Continuous Light in Humans. *J Neurosci.* 32:14242-14253.
- 514 Graham DM, Wong KY, Shapiro P, Frederick C, Pattabiraman K, Berson DM. 2008.
515 Melanopsin ganglion cells use a membrane-associated rhabdomic
516 phototransduction cascade. *J Neurophysiol.* 99:2522-2532.
- 517 Gronfier C, Wright KP, Kronauer RE, Jewett ME, Czeisler CA. 2004. Efficacy of a
518 single sequence of intermittent bright light pulses for delaying circadian phase in
519 humans. *Am J Physiol-Endoc M.* 287:E174-E181.
- 520 Hebert M, Martin SK, Lee C, Eastman CI. 2002. The effects of prior light history on the
521 suppression of melatonin by light in humans. *J Pineal Res.* 33:198-203.
- 522 Higuchi S, Fukuda T, Kozaki T, Takahashi M, Miura N. 2011. Effectiveness of a Red-
523 visor Cap for Preventing Light-induced Melatonin Suppression during
524 Simulated Night Work. *J Physiol Anthropol.* 30:251-258.
- 525 Higuchi S, Ishibashi K, Aritake S, Enomoto M, Hida A, Tamura M, Kozaki T,
526 Motohashi Y, Mishima K. 2008. Inter-individual difference in pupil size
527 correlates to suppression of melatonin by exposure to light. *Neurosci Lett.*
528 440:23-26.
- 529 Higuchi S, Motohashi Y, Ishibashi K, Maeda T. 2007a. Influence of eye colors of
530 Caucasians and Asians on suppression of melatonin secretion by light. *Am J*
531 *Physiol-Reg I.* 292:R2352-R2356.
- 532 Higuchi S, Motohashi Y, Ishibashi K, Maeda T. 2007b. Less exposure to daily ambient
533 light in winter increases sensitivity of melatonin to light suppression.
534 *Chronobiol Int.* 24:31-43.
- 535 Higuchi S, Nagafuchi Y, Lee S, Harada T. 2014. Influence of Light at Night on
536 Melatonin Suppression in Children. *J Clin Endocrinol Metab.* 99:3298-3303.
- 537 Hillman P, Hochstein S, Minke B. 1983. Transduction in Invertebrate Photoreceptors -
538 Role of Pigment Bistability. *Physiol Rev.* 63:668-772.
- 539 Ishihara K, Saitoh T, Inoue Y, Miyata Y. 1984. Validity of the Japanese Version of the
540 Morningness-Eveningness Questionnaire. *Percept Mot Skills.* 59:863-866.
- 541 Jasser SA, Hanifin JP, Rollag MD, Brainard GC. 2006. Dim light adaptation attenuates
542 acute melatonin suppression in humans. *J Biol Rhythms.* 21:394-404.
- 543 Kamdar BB, Tergas AI, Mateen FJ, Bhayani NH, Oh J. 2013. Night-shift work and risk
544 of breast cancer: a systematic review and meta-analysis. *Breast Cancer Res*
545 *Treat.* 138:291-301.
- 546 Kantermann T, Juda M, Vetter C, Roenneberg T. 2010. Shift-work research: Where do
547 we stand, where should we go? *Sleep and Biological Rhythms.* 8:95-105.
- 548 Kayumov L, Casper RF, Hawa RJ, Perelman B, Chung SA, Sokalsky S, Shapiro CM.
549 2005. Blocking low-wavelength light prevents nocturnal melatonin suppression
550 with no adverse effect on performance during simulated shift work. *J Clin*
551 *Endocrinol Metab.* 90:2755-2761.
- 552 Kozaki T, Koga S, Toda N, Noguchi H, Yasukouchi A. 2008. Effects of short
553 wavelength control in polychromatic light sources on nocturnal melatonin
554 secretion. *Neurosci Lett.* 439:256-259.

- Kraneburg A, Franke S, Methling R, Griefahn B. 2017. Effect of color temperature on melatonin production for illumination of working environments. *Appl Ergon.* 58:446-453.
- Kronauer RE, Forger DB, Jewett ME. 1999. Quantifying human circadian pacemaker response to brief, extended, and repeated light stimuli over the photopic range. *J Biol Rhythms.* 14:500-515.
- Laakso ML, Hatonen T, Stenberg D, Alila A, Smith S. 1993. One-Hour Exposure to Moderate Illuminance (500-Lux) Shifts the Human Melatonin Rhythm. *J Pineal Res.* 15:21-26.
- Lamb TD, Pugh EN. 2004. Dark adaptation and the retinoid cycle of vision. *Prog Retin Eye Res.* 23:307-380.
- Lee SI, Matsumori K, Nishimura K, Nishimura Y, Ikeda Y, Eto T, Higuchi S. 2018. Melatonin suppression and sleepiness in children exposed to blue-enriched white LED lighting at night. *Physiol Rep.* 6.
- Lowden A, Akerstedt T, Wibom R. 2004. Suppression of sleepiness and melatonin by bright light exposure during breaks in night work. *J Sleep Res.* 13:37-43.
- Lucas RJ, Hattar S, Takao M, Berson DM, Foster RG, Yau KW. 2003. Diminished pupillary light reflex at high irradiances in melanopsin-knockout mice. *Science.* 299:245-247.
- Lunn RM, Blask DE, Coogan AN, Figueiro M, Gorman MR, Hall JE, Hansen J, Nelson RJ, Panda S, Smolensky MH, Stevens RG, Turek FW, Vermeulen R, Carreon T, Caruso CC, Lawson CC, Thayer KA, Twery MJ, Ewens AD, Garner SC, Schwingl PJ, Boyd WA. 2017. Health consequences of electric lighting practices in the modern world: A report on the National Toxicology Program's workshop on shift work at night, artificial light at night, and circadian disruption. *Sci Total Environ.* 607:1073-1084.
- Macchi MM, Bruce JN. 2004. Human pineal physiology and functional significance of melatonin. *Front Neuroendocrinol.* 25:177-195.
- Mcintyre IM, Norman TR, Burrows GD, Armstrong SM. 1989. Human Melatonin Suppression by Light Is Intensity Dependent. *J Pineal Res.* 6:149-156.
- Monteleone P, Esposito G, Larocca A, Maj M. 1995. Does Bright Light Suppress Nocturnal Melatonin Secretion More in Women Than Men. *J Neural Transm-Gen.* 102:75-80.
- Mure LS, Cornut PL, Rieux C, Drouyer E, Denis P, Gronfier C, Cooper HM. 2009. Melanopsin Bistability: A Fly's Eye Technology in the Human Retina. *PLoS One.* 4.
- Parent ME, El-Zein M, Rousseau MC, Pintos J, Siemiatycki J. 2012. Night Work and the Risk of Cancer Among Men. *Am J Epidemiol.* 176:751-759.
- Phillips AJK, Vidadar P, Burns AC, McGlashan EM, Anderson C, Rajaratnam SMW, Lockley SW, Cain SW. 2019. High sensitivity and interindividual variability in the response of the human circadian system to evening light. *Proc Natl Acad Sci U S A.* 116:12019-12024.
- Rahman SA, Hilaire MAS, Chang AM, Santhi N, Duffy JF, Kronauer RE, Czeisler CA, Lockley SW, Klerman EB. 2017. Circadian phase resetting by a single short-duration light exposure. *Jci Insight.* 2.
- Rimmer DW, Boivin DB, Shanahan TL, Kronauer RE, Duffy JF, Czeisler CA. 2000. Dynamic resetting of the human circadian pacemaker by intermittent bright light. *Am J Physiol-Reg I.* 279:R1574-R1579.
- Santhi N, Thorne HC, van der Veen DR, Johnsen S, Mills SL, Hommes V, Schlangen LJM, Archer SN, Dijk DJ. 2012. The spectral composition of evening light and

- individual differences in the suppression of melatonin and delay of sleep in humans. *J Pineal Res.* 53:47-59.
- Shichida Y, Matsuyama T. 2009. Evolution of opsins and phototransduction. *Philos T R Soc B.* 364:2881-2895.
- Smith KA, Schoen MW, Czeisler CA. 2004. Adaptation of human pineal melatonin suppression by recent photic history. *J Clin Endocrinol Metab.* 89:3610-3614.
- Smith MR, Eastman CI. 2012. Shift work: health, performance and safety problems, traditional countermeasures, and innovative management strategies to reduce circadian misalignment. *Nat Sci Sleep.* 4:111-132.
- Smith MR, Fogg LF, Eastman CI. 2009. Practical Interventions to Promote Circadian Adaptation to Permanent Night Shift Work: Study 4. *J Biol Rhythms.* 24:161-172.
- Smolensky MH, Hermida RC, Reinberg A, Sackett-Lundeen L, Portaluppi F. 2016. Circadian disruption: New clinical perspective of disease pathology and basis for chronotherapeutic intervention. *Chronobiol Int.* 33:1101-1119.
- Smolensky MH, Sackett-Lundeen LL, Portaluppi F. 2015. Nocturnal light pollution and underexposure to daytime sunlight: Complementary mechanisms of circadian disruption and related diseases. *Chronobiol Int.* 32:1029-1048.
- Souman JL, Tinga AM, te Pas SF, van Ee R, Vlaskamp BNS. 2018. Acute alerting effects of light: A systematic literature review. *Behav Brain Res.* 337:228-239.
- St Hilaire MA, Gooley JJ, Khalsa SBS, Kronauer RE, Czeisler CA, Lockley SW. 2012. Human phase response curve to a 1 h pulse of bright white light. *J Physiol-London.* 590:3035-3045.
- Touitou Y, Reinberg A, Touitou D. 2017. Association between light at night, melatonin secretion, sleep deprivation, and the internal clock: Health impacts and mechanisms of circadian disruption. *Life Sci.* 173:94-106.
- Tucker P. 2003. The impact of rest breaks upon accident risk, fatigue and performance: a review. *Work Stress.* 17:123-137.
- Weaver DR. 1998. The suprachiasmatic nucleus: A 25-year retrospective. *J Biol Rhythms.* 13:100-112.
- Wittmann M, Dinich J, Meroow M, Roenneberg T. 2006. Social jetlag: Misalignment of biological and social time. *Chronobiol Int.* 23:497-509.
- Wong KY, Dunn FA, Berson DM. 2005. Photoreceptor adaptation in intrinsically photosensitive retinal ganglion cells. *Neuron.* 48:1001-1010.
- Wong KY, Dunn FA, Graham DM, Berson DM. 2007. Synaptic influences on rat ganglion-cell photoreceptors. *J Physiol-London.* 582:279-296.
- Zeitzer JM, Dijk DJ, Kronauer RE, Brown EN, Czeisler CA. 2000. Sensitivity of the human circadian pacemaker to nocturnal light: melatonin phase resetting and suppression. *J Physiol-London.* 526:695-702.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

646 Table 1. Light conditions used in this study

	Night Shift Light (01:00-06:00h)		10 Min Breaks (4 per night, 1/h)	
	Medium Intensity Light	Medium Intensity Light	Bright Light	Dim Light
Illuminance (lx)	550	430	4700	1
Color temperature (K)	4500	3850	5000	-
Photon flux (log ₁₀ 1/cm ² /sec)	14.70	14.54	15.62	12.63
Melanopic lux	83.37	52.86	787.84	0.09

647

Figure 1. Experimental protocol (a) and details of the experimental tasks during the break and during the simulated night work (b). Participants always stayed in chamber 1 from 13:00h to the end of the experiment (i.e., 06:00h the next morning) except for when they took breaks in chamber 2.

Figure 2. Melatonin profiles (means $\pm$ standard error) and AUCs (means + standard error) in Experiment 1 (a: medium breaks vs. dim breaks) and Experiment 2 (b: medium breaks vs. bright breaks). The black arrows indicate the times when breaks were conducted. **: $p < 0.01$, *: $p < 0.05$

Figure 3. Subjective sleepiness (means $\pm$ standard error) in Experiment 1 (a: medium breaks vs. dim breaks) and Experiment 2 (b: medium breaks vs. bright breaks). The black arrows indicate the times when breaks were conducted.

Figure 4. Cognitive performance (i.e., reaction speed [mean 1/RT]; a, b) and number of lapses (c, d) in Experiment 1 (medium breaks vs. dim breaks; left columns) and Experiment 2 (medium breaks vs. bright breaks; right columns). The black arrows indicate the times when breaks were conducted.

Figure 1

Figure 1. Experimental protocol (a) and details of the experimental tasks during the break and during the simulated night work (b). Participants always stayed in chamber 1 from 13:00h to the end of the experiment (i.e., 06:00h the next morning) except for when they took breaks in chamber 2.

338x190mm (600 x 600 DPI)

Figure 2

Figure 2. Melatonin profiles (means $\pm$ standard error) and AUCs (means $\pm$ standard error) in Experiment 1 (a: medium breaks vs. dim breaks) and Experiment 2 (b: medium breaks vs. bright breaks). The black arrows indicate the times when breaks were conducted. **: $p < 0.01$, *: $p < 0.05$

190x275mm (300 x 300 DPI)

Figure 3

Figure 3. Subjective sleepiness (means $\pm$ standard error) in Experiment 1 (a: medium breaks vs. dim breaks) and Experiment 2 (b: medium breaks vs. bright breaks). The black arrows indicate the times when breaks were conducted.

190x275mm (300 x 300 DPI)

Figure 4

Figure 4. Cognitive performance (i.e., reaction speed [mean 1/RT]; a, b) and number of lapses (c, d) in Experiment 1 (medium breaks vs. dim breaks; left columns) and Experiment 2 (medium breaks vs. bright breaks; right columns). The black arrows indicate the times when breaks were conducted.

190x275mm (300 x 300 DPI)