



Title	Coupled nitrogen and oxygen isotope effects of anaerobic ammonium oxidation (anammox) [an abstract of dissertation and a summary of dissertation review]
Author(s)	小林, 香苗
Citation	北海道大学. 博士(工学) 甲第14449号
Issue Date	2021-03-25
Doc URL	<a href="http://hdl.handle.net/2115/81648">http://hdl.handle.net/2115/81648</a>
Rights(URL)	<a href="https://creativecommons.org/licenses/by/4.0/">https://creativecommons.org/licenses/by/4.0/</a>
Type	theses (doctoral - abstract and summary of review)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Kanae_Kobayashi_abstract.pdf (論文内容の要旨)



[Instructions for use](#)

## 学位論文内容の要旨

博士の専攻分野の名称 博士（工学） 氏名 小林 香苗

### 学位論文題名

Coupled nitrogen and oxygen isotope effects of anaerobic ammonium oxidation (anammox)

(アナモックス細菌の窒素および酸素同位体分別の解析)

Natural abundance of stable nitrogen (N) and oxygen (O) isotopes ( $\delta^{15}\text{N}$  and  $\delta^{18}\text{O}$ ) are invaluable biogeochemical tracers for assessing the N transformations in the environment. To fully exploit these tracers, the N and O isotope effects ( $^{15}\epsilon$  and  $^{18}\epsilon$ ) associated with the respective N transformation processes must be known. Anaerobic ammonium oxidation (anammox) and denitrification are the two major sinks of fixed N. In addition, anammox bacteria contribute to re-oxidation of nitrite to nitrate, because they fix  $\text{CO}_2$  into biomass with reducing equivalents generated from oxidation of nitrite to nitrate. Nitrate production by anammox bacteria influences the nitrite and nitrate N and O isotope effects in freshwater and marine systems. Despite the significant importance of anammox bacteria in the global N cycle,  $^{15}\epsilon$  and  $^{18}\epsilon$  of anammox are not well known. Therefore, the never yet determined  $^{15}\epsilon$  and  $^{18}\epsilon$  associated with anammox were investigated in this study.

Firstly, the  $^{15}\epsilon$  were determined for ‘*Ca. Scalindua sp.*’, ‘*Ca. Jettenia caeni*’, and ‘*Ca. Brocadia sinica*’ growing in continuous enrichment cultures. All three anammox species yielded similar  $^{15}\epsilon$  values of  $\text{NH}_4^+$  oxidation to  $\text{N}_2$  ( $^{15}\epsilon_{\text{NH}_4^+ \rightarrow \text{N}_2} = 30.9 \sim 32.7 \text{ ‰}$ ) and inverse kinetic isotope effects of  $\text{NO}_2^-$  oxidation to  $\text{NO}_3^-$  ( $^{15}\epsilon_{\text{NO}_2^- \rightarrow \text{NO}_3^-} = -45.3 \text{ ‰} \sim -30.1 \text{ ‰}$ ). In contrast, the values of  $\text{NO}_2^-$  reduction to  $\text{N}_2$  was significantly different among three species ( $^{15}\epsilon_{\text{NO}_2^- \rightarrow \text{N}_2} = 5.9 \sim 29.5 \text{ ‰}$ ), which is probably because individual anammox bacteria species might possess different types of nitrite reductase.

Secondly, the  $^{18}\epsilon$  were determined for ‘*Ca. Scalindua sp.*’, which is a putative marine species. Determination of  $^{18}\epsilon$  of anammox is more challenging because the  $\delta^{18}\text{O}_{\text{NO}_2^-}$  value is affected by abiotic O isotope exchange between  $\text{NO}_2^-$  and  $\text{H}_2\text{O}$  ( $k_{eq}$ ,  $^{18}\epsilon_{eq}$ ) and incorporation of a water-derived O atom into  $\text{NO}_3^-$  during  $\text{NO}_2^-$  oxidation to  $\text{NO}_3^-$  ( $^{18}\epsilon_{\text{H}_2\text{O}}$ ). In order to determine abiotic  $k_{eq}$ ,  $^{18}\epsilon_{eq}$ , and  $^{18}\epsilon_{\text{H}_2\text{O}}$ , batch experiments with different  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  values of medium were conducted. Oxygen isotope ratio measurements of  $\text{NO}_2^-$  and  $\text{NO}_3^-$  by the azide method and denitrifier method are sensitive to the  $\delta^{18}\text{O}$  of sample water. However, the influence of  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  on those measurements has not been quantitatively evaluated and documented so far. Therefore, the influence of  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  of sample on  $\delta^{18}\text{O}$  analysis of  $\text{NO}_2^-$  and  $\text{NO}_3^-$  were quantitatively evaluated. Then, the rate of abiotic O isotope exchange between  $\text{NO}_2^-$  and  $\text{H}_2\text{O}$ :  $k_{eq} = 1.13 \times 10^{-2} \text{ (h}^{-1}\text{)}$ , as well as equilibrium isotope effects:  $^{18}\epsilon_{eq} = 11.9 \text{ ‰}$  were experimentally determined. To determine  $^{18}\epsilon$  of each reaction, batch culture experiments with different  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  values of medium ( $\delta^{18}\text{O}_{\text{H}_2\text{O}} = -12.6 \sim 110.1 \text{ ‰}$ ) were conducted for ‘*Ca. Scalindua sp.*’. A numerical model was developed for estimation of respective  $^{18}\epsilon$  of anammox reaction, resulting in  $^{18}\epsilon_{\text{NO}_2^- \rightarrow \text{N}_2} = 8.4 \sim 10.0 \text{ ‰}$ ,  $^{18}\epsilon_{\text{NO}_2^- \rightarrow \text{NO}_3^-} = -3.0 \sim -1.2 \text{ ‰}$ , and  $^{18}\epsilon_{\text{H}_2\text{O}} = 25.8 \sim 27.8 \text{ ‰}$ , respectively. These  $^{18}\epsilon$  values were determined for the first time in the world.

These obtained dual N and O isotopic effects could provide significant insights into the contribution

of anammox bacteria to the fixed N loss and nitrite reoxidation in (recycling N) in various natural environments.