Title	Disinfectant potential in inactivation of epidemic keratoconjunctivitis-related adenoviruses by potassium peroxymonosulfate
Author(s)	Hashizume, Mei; Aoki, Koki; Ohno, Shigeaki; Kitaichi, Nobuyoshi; Yawata, Nobuyo; Gonzalez, Gabriel; Nonaka, Hirotaka; Sato, Seiichi; Takaoka, Akinori
Citation	European journal of ophthalmology, 31(2), 379-384 https://doi.org/10.1177/1120672119891408
Issue Date	2021-03
Doc URL	http://hdl.handle.net/2115/81697
Rights	Mei Hashizume, Koki Aoki, Shigeaki Ohno, Nobuyoshi Kitaichi, Nobuyo Yawata, Gabriel Gonzalez, Hirotaka Nonaka, Seiichi Sato, Akinori Takaoka, Disinfectant potential in inactivation of epidemic keratoconjunctivitis-related adenoviruses by potassium peroxymonosulfate Show less, European Journal of Ophthalmology Vol. 31(2) pp. 379-384. Copyright ©2021 (Copyright Holder). DOI:10.1177/1120672119891408
Type	article (author version)
File Information	TEXT_MH_EJO_final.pdf

[Instructions for use](#)

Disinfectant potential in inactivation of epidemic keratoconjunctivitis (EKC)-related adenoviruses by potassium peroxymonosulfate

Mei Hashizume^{1,2}, Koki Aoki^{1,3,4}, Shigeaki Ohno^{3,4}, Nobuyoshi Kitaichi^{3,4}, Nobuyo Yawata⁵, Gabriel Gonzalez⁶, Hirotaka Nonaka^{1,2}, Seiichi Sato^{1,2} and Akinori Takaoka^{1,2}

¹ Division of Signalling in Cancer and Immunology, Institute for Genetic Medicine, Hokkaido University, Sapporo, Hokkaido, Japan

² Molecular Medical Biochemistry Unit, Biological Chemistry and Engineering Course, Graduate School of Chemical Sciences and Engineering, Hokkaido University, Sapporo, Hokkaido, Japan

³ Department of Ophthalmology, Health Sciences University of Hokkaido, Sapporo, Hokkaido, Japan

⁴ Department of Ophthalmology, Faculty of Medicine and Graduate School of Medicine, Hokkaido University, Sapporo, Hokkaido, Japan

⁵ Department of Ocular Pathology and Imaging Science, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan

⁶ Division of Bioinformatics, Research Center for Zoonosis Control, Hokkaido University, Sapporo, Hokkaido, Japan

Corresponding author:

Akinori Takaoka

takaoka@igm.hokudai.ac.jp

Tel; +81 11-706-5020

Fax; +81 11-706-7541

Short title: Inactivation of EKC-related HAdV

ABSTRACT

Purpose: To be tested the antiviral effectivity of potassium peroxymonosulfate (RUBYSTA[®], KYORIN) against five EKC-related types of *Human adenovirus D* (HAdV-D) *in vitro*.

Methods: Five types of HAdV-D (8, 37, 53, 54 and 56) were incubated with 1% potassium peroxymonosulfate, 0.1% sodium hypochlorite (NaClO) or alcohol-based disinfectant for 30 sec or 1 min. These solutions were subjected to measurements of viral titers by infection assays in A549 cells. At day 6 post-infection, both, supernatants and cells, were collected and the viral genome was assessed by real-time PCR analysis.

Results: Treatments with 1% potassium peroxymonosulfate led to significant reduction of all tested HAdV-D types comparable to disinfecting effects by 0.1% NaClO. Overall, potassium peroxymonosulfate demonstrated sufficient inactivation of the major EKC-causing HAdV-D to be considered for disinfection and prevention purposes in ophthalmological clinics and hospitals.

Conclusions: This study demonstrated that potassium peroxymonosulfate is a promising disinfectant for the prevention of EKC nosocomial infections in ophthalmological clinics.

Keywords: HAdV, EKC, disinfectants, prevention, RUBYSTA, Virkon

Introduction

Human adenoviruses (HAdVs) are currently classified into over 85 accepted human adenovirus types which can be divided into seven species (A to G). These types are classified by phylogenetic analysis of three genes encoding capsid proteins, which have large difference among species or types. HAdV infections are associated with respiratory diseases, conjunctivitis, obesity, gastroenteritis and even cancer.¹ HAdVs are highly contagious and resistant to multiple disinfectants due to physical properties of their capsid proteins.^{1,2} Since HAdVs lack envelopes like other viruses such as norovirus, they are highly resistant to lipid disinfectants including alcohol and soap.^{3,4} The resistance of adenoviral capsid proteins varies according to environmental conditions such as temperature and relative humidity; furthermore, virions have been shown to survive for many days on plastic or metal surfaces and eyedrop bottles.⁵⁻⁷ These HAdV features are closely related to the spread of infections such as nosocomial outbreaks and outbreaks in schools, military bases, and swimming pools.^{1,8,9}

HAdVs cause multiple clinical diseases, including pneumonia, conjunctivitis, gastroenteritis, hepatitis, and myocarditis. Among such HAdV-associated diseases, epidemic keratoconjunctivitis (EKC) is an ocular infection, which is caused mainly by HAdV-D types and commonly transmitted from hands to eyes. In particular, infections by HAdV-D8, -D19a, -D37, -D53, -D54 and -D56 are more frequently reported with the etiology of EKC, which can develop into severe cases with impaired vision.^{2,10} A number of adenoviral EKC outbreaks are seen in ophthalmological clinics and hospitals worldwide.¹¹⁻¹³ In this regard, disinfection against viable HAdVs is critical for the prevention of such outbreaks. The most frequently used disinfectant solutions against HAdVs in clinic are ethanol (80%), popidone iodine (0.2%), and sodium hypochlorite (NaClO; 0.1%).¹⁴ Although NaClO has been reported as the most effective disinfectant against HAdV-D,¹⁴ it also presents problems such as metal corrosion and irritating smell, which hinder its usage in healthcare facilities.¹⁵ Recently a disinfectant including potassium peroxymonosulfate “Multi-purpose disinfectant cleaner RUBYSTA®” (KYORIN Medical Supply, Tokyo, Japan) , which is also known as Virkon® in other countries, has been officially approved for various pathogenic agents including non-enveloped virus such as norovirus:¹⁶ it is odourless and less metal corrosive

than NaClO because of the potassium peroxymonosulfate monohydrate as oxidizing agent.¹⁷ In the United States, it is registered as antimicrobial products for cleaning and disinfecting hard surfaces in hospital by the US Environmental Protection Agency (EPA). Moreover, a report showed that potassium peroxymonosulfate can inactivate HAdV-C5 and -C6,¹⁷ however, it remains unclear whether potassium peroxymonosulfate also shows effectiveness against EKC-related HAdVs. In this study, to assess the potential of potassium peroxymonosulfate as a disinfectant against EKC-related HAdV-D strains, we compared the effects of potassium peroxymonosulfate and NaClO on the viability of these HAdVs, by measuring the viral titers after treatments with them.

Methods

Reagents

Potassium peroxymonosulfate solution (RUBYSTA[®]) was obtained from KYORIN Medical Supply (Tokyo, Japan). Sodium hypochlorite (NaClO) and sodium thiosulfate were obtained from Wako (Osaka, Japan). Antiseptic Solution WELPAS[®] for Hand 0.2% (alcoholic chlorbenzarconium) was purchased from Maruishi pharmaceutical (Osaka, Japan).

Cell culture

A549 cells were purchased from ATCC and cultured in DMEM medium supplemented with 4 mM L-glutamine and 10% heat-inactivated fetal bovine serum (FBS) (Gibco). Also, DMEM medium containing 5% FBS was used for culture of infected A549 cells.

Virus

HAdV-B3 and HAdV-D8 were provided from the Department of Medicine, Ophthalmology, Fukuoka Dental College. HAdV-D37, -D53 and -D56 were obtained from Kumamoto Institute of Public-Health and Environmental. HAdV-D54 was provided from the Department of Ophthalmology, Hokkaido University. For virus propagation, A549 cells were infected with each virus solution and incubated. After observation of cytopathic effects (CPE) in the infected culture, cells and culture media were collected followed by 3 times of freeze and thaw cycles. The supernatant including virus was collected by centrifuge at 3,000 rpm for 10 min. All viral titers were determined by TCID₅₀.

Examination of HAdV replication

A549 cells cultured in 24 well-plate were infected with HAdV at different concentrations (multiplicity of infection, MOI = 0.1, 1, 10). Then, infected cells were collected at 1, 3, 6 and 9 days after infection and the copy number was determined using real-time PCR.

Effects of disinfectants against HAdV were assessed using real-time polymerase chain reaction

A 0.02%, 0.2%, and 2% (w/v) potassium peroxymonosulfate solution and 0.002%, 0.02%, and 0.2% (w/v) NaClO solution were prepared. HAdV solutions were incubated with 0.01%, 0.1%, and 1% (w/v) potassium peroxymonosulfate solution and 0.001%, 0.01%, and 0.1% (w/v) NaClO solution at final concentration (the effective chlorine concentration is 10 ppm, 100 ppm and 1,000 ppm, respectively) for 30 sec or 1 min. After incubation, 0.1% (w/v) sodium thiosulfate at final concentration was added. The alcoholic chlorbenzarconium also reacted with HAdV for 1 min, then diluted 260 times with DMEM. Next, the HAdV solutions after treatment by three different disinfectants were used to infect A549 cells. After 6 days, viral DNAs were isolated from culture cells and their medium. Real-time PCR was performed using SYBR Premix Ex Taq (TOYOBO) and analysed on a StepOnePlus real-time PCR system (Applied Biosystems). Primers for amplification of HAdV DNA by real-time PCR were as follows. Forward primer: 5'-TTCCCCATGGC(A/T/C/G)CACAA(C/T)AC-3', reverse primer: 5'-TGCC(T/G)(A/G)CTCAT(A/G)GGCTG(A/G)AAGTT-3' (product length: 554 bp, product Tm: 88.0 °C). The real-time PCR condition included a first step at 98 °C, followed by 45 cycles of 10 sec at 98 °C, 10 sec at 50 °C and 45 sec at 68 °C. Copy numbers were calculated using a standard curve method.

Cell growth assay

A549 cells seeded in 96 well-plate were treated with potassium peroxymonosulfate at the concentration with 1%, 0.1% and 0.01%. After 24 h we used Cell Counting Kit-8 (DOJINDO, Japan) and measured absorbance following the manufacturer's protocol.

Statistical analysis

Values are shown as mean $\pm$ standard deviation. Statistical significance between two samples was determined with Student's *t*-test.

Results

To determine an appropriate timepoint and MOI of HAdVs for measurement of viral titers, we first conducted time-course analyses of viral propagation in A549 cells. This cell line is known to provide an advantage of viral growth since it lacks expression of cGAS and STING, which are reported to be innate sensors for HAdVs.¹⁸ As shown in Figure 1, HAdV-B3, HAdV-D8, HAdV-D37, HAdV-D53, HAdV-D54, and HAdV-D56, which are known to cause EKC, were successfully replicated in A549 cells in a dose-dependent manner. The levels of DNA copy number became saturated at day 6 after infection with any of the tested HAdVs. Therefore, we decided to assess viral titers at day 6 after infection with each type of HAdV-D.

To evaluate the effects of potassium peroxymonosulfate on the viability of EKC-related HAdV-D types, we tested five HAdV-D types: 8, 37, 53, 54, and 56, together with HAdV-B3 as a control, which does not directly cause EKC. We also analysed the effects of NaClO and alcoholic chlorbenzarconium for comparison. First we checked cytotoxic effect of potassium peroxymonosulfate, and confirmed that it has no effect on A549 cell growth in our experimental condition (data not shown). Thus we used 1% potassium peroxymonosulfate which is recommended concentration for using for clinical setting and compare with NaClO. As shown in Figure 2, the viral replication of all the types of HAdV-D tested were significantly suppressed by more than 4, which was comparable or more effective than NaClO in HAdV-D8, -D37 and -D54. Next, we examined the antiviral effect of several concentration disinfectants. Surprisingly, the 100 ppm (0.1%w/v) potassium peroxymonosulfate still had sufficient antiviral effect (Figure 3). In contrast low concentration of NaClO couldn't reduce infectivity of HAdV-D as previously reported.¹⁴ These data indicated that potassium peroxymonosulfate sufficiently reduced the infectivity of EKC-related HAdV types.

Additionally, we also checked whether alcoholic chlorbenzarconium, which is commonly used for hand wash in hospital, could inactivate EKC-caused HAdV-D. For 1 min reaction, treatment with alcoholic chlorbenzarconium, did not affect the viabilities of all tested types of HAdV (Figure 2b).

Discussion

EKC is typically caused by HAdV-D8, -D19a, -D37, and in recent years new types of EKC-related HAdV, HAdV-D53, -D54 and -D56, have emerged widely in Japan.^{2,19} Since these types are highly infectious and often spread in hospitals, control of such nosocomial infections is recognized as an important issue. This study demonstrated that potassium peroxymonosulfate is a promising disinfectant for the prevention of EKC nosocomial infections in ophthalmological clinics. We also showed that comparable to 0.1% NaClO, 1% potassium peroxymonosulfate sufficiently reduced the yield of infectious virions of all tested HAdV types, supporting the usefulness of potassium peroxymonosulfate against EKC-related HAdV-D types.

0.1% Hypochlorous acid NaClO is widely known as the most effective in mid-level disinfectants¹⁴ despite drawbacks such as corrosive properties that limit its use with medical devices and the irritating smell that is inappropriate to use in clinical settings.¹⁵ In contrast, potassium peroxymonosulfate produce few effects on metals, especially non-reactive to stainless steel.²⁰ Furthermore, potassium peroxymonosulfate is odourless, easy to prepare and store. Our current study indicated that the use of potassium peroxymonosulfate is as effective as NaClO against HAdV-D types, without the main drawbacks of the former. These results support the use of potassium peroxymonosulfate for disinfection of medical devices such as tonometer, slit lamp microscope and eyedrop bottle in healthcare facilities, in particular, at eye clinics.

It is noteworthy that potassium peroxymonosulfate has been shown to have antimicrobial activity against different types of pathogenic agents other than adenovirus, including hepatitis B and C viruses, poliovirus, enterovirus, feline calicivirus, several strains of different bacteria, spores, and fungi.^{16,21-25} In the case of adenovirus, it was reported that potassium peroxymonosulfate inactivated HAdV-C5 and -C6,¹⁷ which are known as respiratory pathogens. The chlorine-based disinfectants seem to disrupt capsid proteins of HAdV. Among various types of HAdV, there is a large difference in their amino acid composition especially in capsid proteins. Recent long term surveillance suggested that not only different type but also a genetic variation is associated with the epidemics of adenoviral keratoconjunctivitis.²⁶ The information was limited regarding the availability of potassium

200 peroxymonosulfate as a disinfectant against EKD-related types of HAdV. Our results
201 extended such results to effective inactivation of HAdV-B3, -D8, -D37, -D53, -D54 and -
202 D56.

203 We demonstrated that 1% potassium peroxymonosulfate was more effective than
204 0.1% NaClO against HAdV-D54. HAdV-D54 was firstly reported and isolated from a
205 nosocomial outbreak in a Japanese hospital in 2000.²⁷ HAdV-D54 has been often isolated
206 since 2011, according to surveillance of EKC causative agents in Japan, and has unique
207 genome sequences encoded capsid protein,² however, the detailed virological features of
208 HAdV-D54 remain poorly understood. *Kaneko et al.* reported that HAdV-D54 is
209 characterized by cytopathic effects (CPE) harder to observe than other EKC-related types.²⁸
210 In relation to this report, our data showed that the replication rate of HAdV-D54 in A549
211 cells is the lowest among all HAdVs we tested (Figure 1e). In this regard, HAdV-D54 or its
212 interaction with host cells may have some factors behind the slow replication rate and CPE
213 due to its gene polymorphism or interaction with host factors. Despite these conditions, this
214 type is an important infectious EKC agent in Japan with severe cases leading to corneal
215 complications.^{2,28}

216 In conclusion, we showed that 1% potassium peroxymonosulfate has a strong
217 disinfectant activity against EKC-related types of HAdV. Considering that negligible odor
218 and corrosive effect, this study suggests that potassium peroxymonosulfate is a superior
219 disinfectant in as a prevention measure against nosocomial infections and accidental
220 spreading of infectious agents.

221 **Acknowledgements**

222 We thank R. Hinokuma and S. Hashimoto for HAdV, and H. Watanabe for helpful advice.

223 This work was supported by financial support from KYORIN Medical Supply Co., Ltd
224 (Tokyo, Japan).

225

226 **Declaration of conflicting interests**

227 None.

References

1. Hall K, Blair Zajdel ME, Blair GE. Unity and diversity in the human adenoviruses: exploiting alternative entry pathways for gene therapy. *Biochem J* 2010; 431: 321–336.
2. Gonzalez G, Yawata N, Aoki K, et al. Challenges in management of epidemic keratoconjunctivitis with emerging recombinant human adenoviruses. *J Clin Virol* 2019; 112: 1–9.
3. Wutzler P, Sauerbrei A. Virucidal efficacy of a combination of 0.2% peracetic acid and 80% (v/v) ethanol (PAA-ethanol) as a potential hand disinfectant. *J Hosp Infect* 2000; 46: 304–308.
4. Uzuner H, Karadenizli A, Er DK, et al. Investigation of the efficacy of alcohol-based solutions on adenovirus serotypes 8, 19 and 37, common causes of epidemic keratoconjunctivitis, after an adenovirus outbreak in hospital. *J Hosp Infect* 2018; 100: e30–e36.
5. Nauheim RC, Romanowski EG, Araullo-Cruz T, et al. Prolonged recoverability of desiccated adenovirus type 19 from various surfaces. *Ophthalmology* 1990; 97: 1450–3.
6. Gordon YJ, Gordon RY, Romanowski E, et al. Prolonged recovery of desiccated adenoviral serotypes 5, 8, and 19 from plastic and metal surfaces in vitro. *Ophthalmology* 1993; 100: 1835–9; discussion 1839–40.
7. Uchio E, Ishiko H, Aoki K, et al. Adenovirus detected by polymerase chain reaction in multidose eyedrop bottles used by patients with adenoviral keratoconjunctivitis. *Am J Ophthalmol* 2002; 134: 618–9.
8. Ghebremedhin B. Human adenovirus: Viral pathogen with increasing importance. *Eur J Microbiol Immunol* 2014; 4: 26–33.
9. Hamada N, Gotoh K, Hara K, et al. Nosocomial outbreak of epidemic keratoconjunctivitis accompanying environmental contamination with adenoviruses. *J Hosp Infect* 2008; 68: 262–268.
10. Tabbara KF, Omar N, Hammouda E, et al. Molecular epidemiology of adenoviral keratoconjunctivitis in Saudi Arabia. *Mol Vis* 2010; 16: 2132–6.

- 258 11. Warren D, Nelson KE, Farrar JA, et al. A large outbreak of epidemic
259 keratoconjunctivitis: problems in controlling nosocomial spread. *J Infect Dis* 1989;
260 160: 938–43.
- 261 12. Takeuchi R, Nomura Y, Kojima M, et al. A nosocomial outbreak of epidemic
262 keratoconjunctivitis due to adenovirus type 37. *Microbiol Immunol* 1990; 34: 749–
263 54.
- 264 13. Alai N. Enhancing best practices in ophthalmology for prevention of nosocomial
265 epidemic keratoconjunctivitis infections. *Curr Med Res Opin* 2016; 32: 1757–1758.
- 266 14. Akanuma M. Evaluation of Disinfectant against Adenovirus by Real-time
267 Polymerase Chain Reaction. *J japanese ophthalmological Soc* 2007; 111: 384–390.
- 268 15. Fukuzaki S. Mechanisms of Actions of Sodium Hypochlorite in Cleaning and
269 Disinfection Processes. *Biocontrol Sci* 2006; 11: 147–157.
- 270 16. Eleraky NZ, Potgieter LND, Kennedy MA. Virucidal Efficacy of Four New
271 Disinfectants. *J Am Anim Hosp Assoc* 2002; 38: 231–234.
- 272 17. McCormick L, Maheshwari G. Inactivation of adenovirus types 5 and 6 by Virkon®
273 S. *Antiviral Res* 2004; 64: 27–33.
- 274 18. Lam E, Falck-Pedersen E. Unabated Adenovirus Replication following Activation of
275 the cGAS/STING-Dependent Antiviral Response in Human Cells. *J Virol* 2014; 88:
276 14426–14439.
- 277 19. Ishiko H, Aoki K. Spread of Epidemic Keratoconjunctivitis Due to a Novel Serotype
278 of Human Adenovirus in Japan. *J Clin Microbiol* 2009; 47: 2678–2679.
- 279 20. Okaue A, Ozawa T, Ogura Y, et al. Evaluation of Corrosiveness to Various Surface
280 Materials of Complex Type Chlorine-Based Disinfectant Cleaner. *Japanese J Infect*
281 *Prev Control* 2015; 30: 325–330.
- 282 21. Gasparini R, Pozzi T, Magnelli R, et al. Evaluation of in vitro efficacy of the
283 disinfectant Virkon. *Eur J Epidemiol* 1995; 11: 193–7.
- 284 22. Hernandez A, Martró E, Matas L, et al. Assessment of in-vitro efficacy of 1%
285 Virkon® against bacteria, fungi, viruses and spores by means of AFNOR guidelines.
286 *J Hosp Infect* 2000; 46: 203–209.

- 287 23. Chan YF, Abu Bakar S. Virucidal activity of Virkon S on human enterovirus. *Med J*
288 *Malaysia* 2005; 60: 246–8.
- 289 24. Roongruangchai K, Kummalue T, Sookkua T, et al. Comparison of Pouzolzia indica
290 methanolic extract and Virkon against cysts of Acanthamoeba spp. *Southeast Asian J*
291 *Trop Med Public Health* 2010; 41: 776–84.
- 292 25. Ogura Y, Ozawa T, Nojima Y, et al. Antimicrobial Efficacy of Complex-Type
293 Chlorine-Based Disinfectant Cleaner Against Several Pathogenic Microorganisms.
294 *Japanese J Infect Prev Control* 2015; 30: 391–398.
- 295 26. Ariga T, Shimada Y, Shiratori K, et al. Five New Genome Types of Adenovirus
296 Type 37 Caused Epidemic Keratoconjunctivitis in Sapporo, Japan, for More Than 10
297 Years. *J Clin Microbiol* 2005; 43: 726–732.
- 298 27. Ishiko H, Shimada Y, Konno T, et al. Novel Human Adenovirus Causing
299 Nosocomial Epidemic Keratoconjunctivitis. *J Clin Microbiol* 2008; 46: 2002–2008.
- 300 28. Kaneko H, Suzutani T, Aoki K, et al. Epidemiological and virological features of
301 epidemic keratoconjunctivitis due to new human adenovirus type 54 in Japan. *Br J*
302 *Ophthalmol* 2011; 95: 32–36.

Figure 1. Time course analyses of HAdV DNA copy number at the indicated times by real-time PCR (1, 3, 6 and 9 days) after HAdV-B3 (a), HAdV-D8 (b), HAdV-D37 (c), HAdV-D53 (d), HAdV-D54 (e) and HAdV-D56 (f) infection (multiplicity of infection = 0.1; ●, 1; ▲, 10; ■) in A549 cells.

308

309 **Figure 2.** Effect of 1% potassium peroxymonosulfate (blank bar), 0.1% sodium hypochlorite
 310 (black bar) and alcoholic chlorbenzarconium (grey bar) on the inactivation of HAdV-D for
 311 30 sec (a) or 1min (b). ** $P < 0.01$ and * $P < 0.05$ vs potassium peroxymonosulfate. NS, not
 312 significant. Data are presented as mean and standard deviation (n=3).

Figure 3. Effect of various concentration potassium peroxymonosulfate (circle) and sodium hypochlorite (square) on the inactivation of HAdV-B3 (a), HAdV-D8 (b), HAdV-D37 (c), HAdV-D53 (d), HAdV-D54 (e) and HAdV-D56 (f) for 1min. ** $P < 0.01$ vs potassium peroxymonosulfate. Data are presented as mean and standard deviation (n=3).