Title	Review of the Taxonomy, Diversity, Ecology, and Other Biological Aspects of Order Tanaidacea from Japan and Surrounding Waters
Author(s)	Kakui, Keiichi
Citation	Species Diversity of Animals in Japan, 603-627 https://doi.org/10.1007/978-4-431-56432-4_23
Issue Date	2017
Doc URL	http://hdl.handle.net/2115/81780
Type	bookchapter (author version)
File Information	Kakui_Chapter_603-627.pdf

[Instructions for use](#)

This is a post-peer-review, pre-copyedit version of a chapter published in “Species Diversity of Animals in Japan” (Motokawa M, Kajihara H, editors). The final authenticated version is available online at: https://doi.org/10.1007/978-4-431-56432-4_23.

Chapter 23

Review of the Taxonomy, Diversity, Ecology, and Other Biological Aspects of Order Tanaidacea from Japan and Surrounding Waters

Keiichi Kakui

Faculty of Science, Hokkaido University, Sapporo, Hokkaido 060-0810, Japan

e-mail: kakui@mail.sci.hokudai.ac.jp

telephone: +81-11-706-2750

Abstract The order Tanaidacea is a group of benthic crustaceans, most of which are small, up to a few millimeters long. Tanaidaceans are distributed worldwide, with more than 1,200 described species. Following the first taxonomic paper on a Japanese tanaidacean in 1936, many researchers have studied their taxonomy, morphology, reproductive biology, or ecology in the waters around Japan. This chapter presents a brief introduction to tanaidaceans and then reviews what is known of their systematics (taxonomy and phylogeny), biology (including feeding habits, phenology, morphology, reproductive modes, parasites, predators), and ecology in Japan. The chapter ends with a summary and prospects for future research. The general conclusion is that tanaidaceans have been under-studied, both globally and within Japan; the 104 nominal species reported from around Japan and the 1,200 species reported globally likely represent a fraction of the actual diversity. The phylogeny of tanaidaceans is largely unresolved at all taxonomic levels. Recent, significant new discoveries dealing with herbivory, selfing, skin-digging activity in holothuroid hosts, possible sound production, and tube building suggest that much remains to be learned about their general biology.

Keywords Crustacea • Peracarida • Tanaidacea • Taxonomy • Biology • Ecology • Japan

23.1 Introduction

The order Tanaidacea is a group of aquatic crustaceans in the superorder Peracarida, which includes Isopoda, Amphipoda, Cumacea, and some other groups. More than 1,200 living tanaidacean species have been described worldwide, comprising 33 families, with several dozens of species regarded as family *incertae sedis* (Anderson 2013); this is likely only a small fraction of the actual species diversity (cf. Błażewicz-Paszkowycz 2014). Tanaidacea also includes a few extinct species; 16 fossil taxa have been recorded from the Lower Carboniferous to the Middle Cretaceous (see

Sánchez-García et al. 2015). Tanaidaceans differ from other crustaceans in the following combination of characters: 1) a cephalon and the two most anterior thoracic somites form a cephalothorax, resulting in a pereon consisting of six pereonites; 2) the cephalothorax is covered with a carapace; 3) the cephalothorax has a branchial chamber posterolaterally on each side, each containing an epignath (an organ related to respiration); 4) the appendage on the second thoracic somite (seventh segment in the cephalothorax) is a cheliped (it is subchelate in a few groups; cf. Kakui and Angsupanich 2012); and 5) there is one pair of narrow, biramous or uniramous uropods.

Tanaidaceans are typically a few millimeters long, though the largest reported is 75 mm long (Gamo 1984). While several groups inhabit non-marine environments such as brackish lakes, freshwater systems, and anchialine caves (Kakui et al. 2010; Chapman et al. 2011; Guţu and Iliffe 2011; Błazewicz-Paszkowycz et al. 2012), most species are marine. Marine species have been reported worldwide, ranging in depth from the intertidal zone to bathyal depths at around 9,000 m. Aside from a few parasitic or symbiotic species such as *Exspina typica* Lang, 1968 and *Terebellatanais floridanus* Suárez-Morales et al., 2011 (Alvaro et al. 2011; Suárez-Morales et al. 2011), tanaidaceans are free living (cf. Błazewicz-Paszkowycz 2014). They show diverse modes of life (Fig. 23.1); for example, most apseudomorph tanaidaceans are burrowers; some species in the family Pagurapseudidae utilize empty gastropod shells as a mobile shelter, like hermit crabs; and many species use threads to construct tube-like nests in the sea bottom, or on the surfaces of seaweeds, seagrasses, other animals, or abiotic substrata (Hassack and Holdich 1987; Kakui and Hiruta 2014; hereafter, ‘tube-dwellers’). Tanaidaceans are typical peracaridans, in that they lack a planktonic larval stage and brood their young, and release benthic offspring directly (so-called ‘direct development’), a feature that might be responsible for the occasionally extremely high local densities reported, e.g., 146,000 individuals/m² in *Allotanaïs hirsutus* (Beddard, 1886) and 140,000 individuals/m² in *Mesokalliapseudes crassus* (Menzies, 1953) (Barnard 1970; Delille et al. 1985).

The earliest report of tanaidaceans from the waters surrounding Japan was Stephensen (1936), dealing with a tanaidid species collected in the Kurile Islands. There have been many subsequent studies, continuing to the present, that deal with taxonomy, phylogeny, biology, and ecology. In this chapter, I review previous studies on the systematics, biology, and ecology of tanaidaceans around Japan.

23.2 Systematics

Tanaidacean systematics (including taxonomy) remains active, with recent discoveries

of taxa even at the family level (e.g., Larsen 2012). In this section, I review tanaidacean systematics around Japan.

23.2.1 *Taxonomic and Phylogenetic Studies of Tanaidaceans Around Japan*

Knud Hensch Stephensen, a Danish zoologist, published the first report specifically dealing with a tanaidacean from Japanese waters. This study (Stephensen 1936), based on specimens collected by Denzaburo Miyadi from a lake on Kunashiri Island, Kurile Islands, reported and redescribed *Sinelobus stanfordi* (Richardson, 1901).

The first Japanese specialist on tanaidacean taxonomy was Sueo Shiino, who also actively studied other crustaceans, including isopods, copepods, and rhizocephalans. His publications include five papers on the Japanese tanaidacean fauna (Shiino 1937, 1951a, b, 1952, 1966), with descriptions of seven new species, as well as five papers on tanaidaceans outside Japan (e.g., Shiino 1963).

Another relevant specialist was the Soviet researcher Rosalia Konstantinovna Kudinova-Pasternak, who studied deep-sea tanaidaceans collected from the Sea of Japan, the Kurile–Kamchatka Trench, the Japan Trench, the Izu–Ogasawara Trench, and the Pacific abyssal plane, and described more than 25 new species from these areas (Kudinova-Pasternak 1966a, b, 1970, 1976, 1978, 1984). Other researchers were active around the same time. Kussakin and Tzareva (1972) described several tanaidomorphs from the Kurile Islands. Sieg (1980) described two new tanaidids, *Zeuxo coralensis* Sieg, 1980 and *Z. maledivensis* Sieg, 1980; although the type locality of these species is the Maldives Islands, Sieg included Japanese specimens in the material examined for the original descriptions. Ishimaru (1985) described two leptocheliids, one of which was new to science.

After a hiatus of nearly two decades in Japanese tanaidacean studies, Nunomura (2005) described a new brackish apseudomorph from Okinawa. A further boost to tanaidacean taxonomy came in 2007, when the Danish researcher Kim Richard Larsen came to Japan for a 2-year stay. He and his host, Michitaka Shimomura, studied both shallow-water and deep-sea tanaidaceans, described 12 new species in four papers, and edited a monograph (with the participation of four other researchers) that included descriptions of 15 new species (Larsen and Shimomura 2006, 2007a, b, 2008, 2009; Bamber 2007; Bird 2007a, b; Błażewicz-Paszkowycz 2007; Larsen 2007; McLelland 2007). Around the same time, colleagues and I began studies that resulted in eight new species described from Japan, with several redescrptions (Kakui et al. 2007, 2010, 2012; Kakui and Kajihara 2011; Kakui and Yamasaki 2013; Kakui and Kohtsuka 2015; Kakui and Naruse 2015). Another currently active specialist is the Polish researcher

Magdalena Błażewicz-Paszkowycz, who has published descriptions of nine new species collected by two joint Russian–German expeditions, SoJaBio (Sea of Japan) and KuramBio (Kurile–Kamchatka Trench) (Błażewicz-Paszkowycz et al. 2013, 2015a, b).

The known tanaidacean fauna around Japan currently stands at 104 species (Table 23.1; Fig. 23.2), although some of these records may be in error due to misidentification (cf. Błażewicz-Paszkowycz et al. 2013). Some papers have only presented new distributional records or partial descriptions (e.g., Nunomura 1985; Ariyama and Ohtani 1990; Saito 1996; Kakui et al. 2014).

Since the pioneering work by Drumm (2010), three molecular phylogenetic studies on tanaidaceans have been published, all of which have included representative species collected around Japan. Kakui et al. (2011) focused on the relationships among families or higher taxa and obtained the first sequence data from neotanaids; based on the results, these authors proposed a two-suborder, four-superfamily classification for Recent tanaidaceans. Kakui et al. (2012) and Błażewicz-Paszkowycz et al. (2015b) studied the phylogenetic position of *Arctotanais* and *Protanais*, respectively, within Tanaidomorpha.

23.2.2 *Some Remarks on the Tanaidacean Fauna Around Japan*

Japan extends across several climatic zones and is affected by two warm and two cold currents, and thus the tanaidacean fauna in coastal area shows some regionality (Kakui unpubl. data from several years of sampling, including by snorkeling and SCUBA). At the family level, for example, Kalliapseudidae, Parapseudidae, Apseudidae, and Pagurapseudidae have been detected only from Honshu to the Nansei Islands, but not around Hokkaido. Metapseudidae from coastal regions includes species in genera previously reported (*Apseudomorpha* and *Synapseudes*; Shiino 1951b) as well as representatives of other genera such as *Curtipleon* (Kakui unpubl. data), with differences in distributional ranges among genera. To date, *Apseudomorpha* has been collected from Hokkaido to the Ryukyu Islands; *Synapseudes* from central Honshu to the Ryukyu Islands; and *Curtipleon* only in the vicinity of the Ogasawara and Ryukyu Islands (Kakui unpubl. data). Nototanaididae is presently known from only around the Ryukyu Islands (Kakui and Yamasaki 2013). These distributional differences may result from differences in habitat preference (e.g., brown alga bed, mudflat, coral reef, etc.) as well as differences in tolerance across a wide range of water temperature related to latitude and ocean currents. In addition, mangrove forests extend farther north in Japan than anywhere else in the world except Bermuda (Giri et al. 2011), contributing to tanaidacean diversity—as evidenced, for example, by the nototanaid *Nesotanais*

ryukyuensis Kakui et al., 2010, so far known only from brackish waters around islands with mangroves (Kakui and Yamasaki 2013). Likewise, the parapseudid *Longiflagrum nasutus* (Nunomura, 2005) is known only from the Manko mangrove wetland on Okinawa Island. It is presently unknown whether these species simply inhabit tropical/subtropical brackish waters, or are specifically associated with mangrove environments. Interestingly, both species appear to be distributional outliers, with the nearest neighboring congener thousands of kilometers away: for *Nesotanaïs*, *N. macLaughlinae* Guțu and Iliffe, 1989 from Palau; for *Longiflagrum*, *L. koyonense* Angsupanich, 2004 from Thailand (Guțu and Iliffe 1989; Angsupanich 2004).

Tanaididae is one of two families commonly occurring in nearshore and brackish environments in Japan. *Zeuxo*, the most species-rich genus in this family, is distributed from Hokkaido to the Ryukyu Islands, with high population abundances from Hokkaido to Kyushu, and can be collected from many microhabitats such as laminarian holdfasts, as epibionts on seagrasses or algae (Fig. 23.1c), or sandy bottoms. Pairs of sympatric species are known, from several areas (e.g., around the Misaki Marine Biological Station), that show different pigmentation patterns (Fig. 23.3) and different frequencies of occurrence among various microhabitats, which implies habitat segregation. Many different pigmentation patterns are evident in *Zeuxo* specimens collected in Japan, each likely corresponding to an independent species (Kakui unpubl. data). To date, three species have been reported in Japan (Table 23.1), though their type localities are far away: Addu Atoll (Maldives) for *Z. coralensis*; Fadiffolu Atoll (Maldives) for *Z. maledivensis*; Monterey Bay (USA) for *Z. normani* (Richardson, 1905) (Sieg 1980). The validity of the previous identifications of *Zeuxo* species in Japan are thus suspect and need to be checked.

The confamilial genus *Tanaïs* is also distributed throughout Japan, with high population abundances in southern Japan (in contrast to *Zeuxo*). Species in *Tanaïs* are positively phototactic and can be collected at night with a light trap (Naoto Jimi pers. comm.). *Sinelobus* is distributed in brackish lower river regions and estuaries from northern to southern Japan. Japanese material currently treated as *Sinelobus stanfordi* actually contains at least two species, one in northern and the other in southern Japan, but they are sympatric at least in Kochi Bay, Shikoku (Kakui unpubl. data).

The other common family in coastal regions is Leptocheliidae, species of which occur abundantly intertidally in coral-rubble beds, and throughout Japan on sandy or muddy bottoms, and are also common subtidally in the shelf zone. Despite their abundance, leptocheliids have been poorly studied in Japan (Kussakin and Tzareva 1972; Ishimaru 1985; Larsen and Shimomura 2007a), with only one subtidal record

(Larsen and Shimomura 2007a), and many species likely remain to be discovered.

Tanaidacean samples from the deep sea around Japan contain numerous paratanaoid species; among 15 paratanaoid families with deep-sea records (cf. Suárez-Morales et al. 2011; Bird 2012; Blazewicz-Paszkowycz et al. 2012), only Mirandotanaidae has not been reported. One neotanaoid and three apseudoid families (Neotanaidae; Apseudidae, Parapseudidae, and Sphyrapodidae) are also represented. In addition, the tanaidoid *Protanais birsteini* (Kudinova-Pasternak, 1970) has been reported from hadal depths along the Kurile–Kamchatka Trench (Kudinova-Pasternak 1970; Blazewicz-Paszkowycz et al. 2015b). Sphyrapodidae is especially well represented; to date, two genera and four species have been reported from the deep sea around Japan [excluding doubtful records of *Pseudosphyrapus serratus* (G. O. Sars, 1882)] (Table 23.1). Sphyrapodids have been detected all around Japan, except off the Pacific coast from eastern Hokkaido to central Honshu (Fig. 23.4) (Kakui unpubl. data). However, the distributions of the four species are more limited, as indicated in Fig. 23.4. Unidentified specimens in *Pseudosphyrapus* (*P. sp.* in Fig. 23.4) that are closely related to or possibly conspecific with *P. quintolongus* occur off the coasts of Shikoku and Honshu. The congener *P. anomalus* (G. O. Sars, 1969), with a circumpolar Arctic distribution, occurs to the north of Japan, but no other congeners have been reported from Pacific except for those around Japan. A phylogeographic study would shed light on this interesting distributional pattern.

23.3 Biology and Ecology

The biology of tanaidaceans has long interested biologists; Darwin (1872), for example, referred to their male dimorphism in “The Origin of Species”. In addition, their high abundance in bottom habitats suggests tanaidaceans might play an important role in ecosystems (Larsen et al. 2015). However, there have been relatively few studies in tanaidacean biology and ecology, perhaps due to the small size of individuals and the difficulty of identification, which relies on microscopic characters. In this section, I review knowledge on several aspects of tanaidacean biology and ecology derived from studies in Japan.

23.3.1 Morphology

Tanaidacean morphology has been studied for well over a century, as the following examples illustrate. Blanc (1884) made detailed observations on the thoracic glands associated with thread production in a tube dweller, *Heterotanaïs oerstedii* (Krøyer, 1842). Dennell (1937) illustrated feeding and respiratory structures in *Apseudes talpa*

(Montagu, 1808). Johnson and Attramadal (1982a) studied morphological structures and behavior related to tube dwelling in *Tanais dulongii* (Audouin, 1826). Drumm (2005) examined feeding, respiratory, and cleaning behaviors and their related morphology in two kalliapseudid species, and Wirkner and Richter (2008) studied the morphology of the haemolymph vascular system among three superfamilies [*T. dulongii*, *Paradoxapseudes bermudeus* (Băcescu, 1980), and *Neotanais* sp.].

Several studies in Japan have also dealt with morphology. In a study of the ultrastructure of the branchial chamber and epignath in *Sinelobus stanfordi*, Kikuchi and Matsumasa (1993) showed that the epithelia of both the epignath and the inner wall of the carapace (branchiostegite) are the ion-transporting type but differ in structure; the authors suggested that the latter epithelium plays a major role in osmoregulation and respiratory gas exchange. Kakui et al. (2010) observed opposing serial ridges on the inner surface of the left and right chelipeds in two *Nesotanais* species. Through behavioral observations and a comparison with similar structures in sesamid crabs used to produce sound, that study suggested that the paired ridges in *Nesotanais* might likewise function to produce sound. In the tanaidid *Arctotanais alascensis* (Richardson, 1899), Kakui et al. (2012) discovered a unique semicircular ischium on all pereopods and speculated that this character might represent a transitional stage between the circular-shaped ischium present in most tanaidaceans and complete loss of that article in tanaidids other than *A. alascensis*. Finally, in a study of the thread-producing system in the kalliapseudid apseudomorph *Phoxokalliapseudes tomiokaensis* (Shiino, 1966), Kakui and Hiruta (2014) reported four distinct types of secretory system in the pereopods, with the type in pereopod 1 possibly involving secretory setae.

23.3.2 Reproductive Modes

Tanaidaceans are highly diverse in reproductive mode. Most species are gonochoristic, like humans, with males and females engaging in sexual reproduction. However, simultaneous hermaphroditism also occurs, first reported in a species in *Apseudes* (Lang 1953). Only recently, Kakui and Hiruta (2013) showed that the simultaneously hermaphroditic *Apseudes* sp., collected from a public aquarium in Japan, can self-fertilize, the first example of self-fertilization in the crustacean class Malacostraca. The same showed that *Apseudes* sp. becomes simultaneously hermaphroditic via a male-like phase in which only the testes are mature, which means that this species is a protandric simultaneous hermaphrodite. Specimens bearing both male and female external reproductive features have been reported in several species in genera other than *Apseudes* (cf. Rumbold et al. 2015), and some of these may likewise be functional

simultaneous hermaphrodites.

Some tanaidaceans are sequential hermaphrodites. Bückle Ramírez (1965) demonstrated that *Heterotanaïs oerstedii* is diandrous, producing both primary males (developing directly from juveniles) and secondary males (developing by sex change from females). *Nototanaïs* and subsequently several other tanaidomorphs have proven similarly to be protogynous sequential hermaphrodites, changing in sex from female to male during the life cycle (cf. Rumbold et al. 2015). Although protandric sequential hermaphrodites (changing sex from male to female) have also been suspected among tanaidaceans (Wolff 1956), no case has been confirmed. Błażewicz-Paszkowycz et al. (2014) demonstrated that a single species (*Agathotanaïs ingolfi* Hansen, 1913) can produce two different male morphs (a sedentary form and a swimming form) (see also Bird and Holdich 1988), indicating there is much left to learn about tanaidacean life cycles; future studies of their reproductive systems will help elucidate the evolution of reproduction in Tanaidacea, and even in Malacostraca.

23.3.3 Phenology

Tanaidacean phenology has not been well studied, though there are a few reports dealing with several families (e.g., Aboul-Dahab et al. 2001; Fonseca and D’Incao 2003; Pennafirme and Soares-Gomes 2009). Several species appear to have an annual breeding period (Johnson and Attramadal 1982b; Fonseca and D’Incao 2003), whereas others reproduce year-around (Aboul-Dahab et al. 2001). Pennafirme and Soares-Gomes (2009) showed that the life cycle can vary (e.g., presence or absence of a distinct breeding period) between tropical and subtropical populations of a single species.

Several papers have described seasonal changes in tanaidacean density in Japan. Mukai (1971) and Kito (1975) documented seasonal fluctuations associated with brown macroalgae in the genus *Sargassum*. While the former study just identified tanaidaceans to the ordinal level (Tanaidacea), the population involved probably belonged to Tanaididae; the latter study identified the populations studied as “*Anatanaïs normani* (Richardson).” Both studies detected an acute annual density peak of tanaidaceans during the withering season for *Sargassum* seaweeds, i.e., around 2–4 months after the peak of the mean wet weight of the seaweeds. These results suggest that the tanaidaceans studied have a breeding period, and that this period partly overlaps the season of maximum growth and biomass accumulation of the seaweed. In a study on the distribution patterns of five peracaridans in relation to environmental factors in a shallow, brackish lagoon, Matsumasa and Kurihara (1988) included phenological data

from April 1984 to May 1985 for the tanaidid *Sinelobus stanfordi*, which inhabited the concrete wall of the lagoon embankment. The tanaidid density was quite low from April to June, increased in July, remained relatively constant until January, and then gradually decreased from February to May. As above, these data may indicate a breeding period for the *Sinelobus* population in the study area, possibly from June to November or December. Nakaoka et al. (2001) studied seasonal and between-substrate variation in the mobile epibiont community (including *Zeuxo* sp., discussed under 'Feeding habits' below) in a multi-species seagrass bed. Although the density of *Zeuxo* sp. was close to zero most months, an acute peak occurred in July (on *Zostera marina* L.) or August (on *Zostera caulescens* Miki), presumably associated with spathe use by the tanaidid. That study also found that *Zeuxo* sp. utilizes the spathes of both *Zostera* seagrass species from June to August, but detected juveniles only in August. These data indicate seasonal breeding, but the duration remains unknown.

23.3.4 Feeding Habits

Tanaidaceans show diverse feeding modes and include detritivores, scavengers, browsers, raptorial carnivores, opportunists, and filter-feeders (see Larsen et al. 2015). Nakaoka (2002) discovered an interesting mode in a *Zeuxo* species in Japan. Nakaoka (2002) first found *Zeuxo* sp. inside spathes of the seagrasses *Zostera marina* and *Z. caulescens*, and observed bored seeds. Through laboratory experiments and a monthly field survey, he showed that *Zeuxo* sp. consumes seagrass seeds by boring, constructs dwelling tubes in the spathes, and utilizes the spathes as the site of reproduction and as a nursery. However, *Zeuxo* sp. only seasonally utilizes the spathes, because flowering and fruiting by the seagrasses are restricted to a few summer months; during the remainder of the year, this tanaidid is nearly absent from the plants (Nakaoka et al. 2001).

Species in *Protanais* are one of the main indicators of successional stages in deep-sea wood-fall communities (McClain and Barry 2014), and are suspected of preying on *Xylophaga* wood-boring bivalves (another of the main indicators), their fecal pellets, or associated fecal bacteria (Larsen 2006; McClain and Barry 2014). Błażewicz-Paszkowycz et al. (2015b), however, found wood tissue and bacterial cells in the tanaidacean digestive tract, and nematodes and protists on the mouthparts, but no residues from bivalves. This suggests that, as with the hadal amphipod *Hirondellea gigas* (Birstein and Vinogradov, 1955) (Kobayashi et al. 2012), *Protanais birsteini* may be xylophagous and contain wood-digesting enzymes.

23.3.5 Parasites

Despite high local abundances of tanaidaceans, there are only a few reports of associated parasites or commensal animals. Peritrich ciliates, foraminiferans, lasaeid bivalves, tantulocarideans, and nicothoid copepods have been reported as ecto- or mesoparasites on tanaidaceans (Warén and Carrozza 1994; Fernandez-Leborans and Sorbe 2003; Larsen 2005; Oliver et al. 2010; Błażewicz-Paszkowycz 2014), and mites, harpacticoid copepods, and ostracods have been reported as commensals in their burrows or tubes (Johnson and Attramadal 1982b; Drumm 2005). A few ecto- and mesoparasites are known from Japanese waters. Kakui and Kohtsuka (2015: fig. 8) presented an image of unidentified peritrich ciliates parasitic on *Agathotanaïs misakiensis* Kakui and Kohtsuka, 2015 and Błażewicz-Paszkowycz et al. (2015b) reported fibrous filamentous bacteria attached to appendages of *Protanaïs birsteini*. In addition, Kakui (2015b) reported copepods parasitic on the tanaidaceans *Fageapseudes* sp. and *Leptochelia* sp. collected from the East China Sea and Pacific Ocean. The copepods proved to represent two undescribed species in the nicothoid genus *Rhizorhina*—the first report of copepods parasitic on tanaidaceans from the North Pacific, and only the third worldwide.

Until recently, only nematodes and acanthocephalan larvae had been reported as endoparasites of tanaidaceans (cf. Larsen 2005). Kakui (2014) added trematodes to this list, after discovering the encysted metacercariae of a trematode inside the body cavity of the parapseudid *Longiflagrum nasutus*. Morphology and DNA sequence data placed the trematode in the family Microphallidae, species of which chiefly utilize birds as the definitive host. The true diversity of animals parasitic on tanaidaceans may be greatly underestimated.

23.3.6 Predators

Tanaidaceans have been reported in stomach contents from diverse animal groups, including crustaceans, polychaetes, sea anemones, wading birds, and fishes (Shaffer 1979; Oliver et al. 1982; Larsen 2005). In Japan, Sano et al. (1984) found that 14 species of coral-reef fishes in 10 families consume tanaidaceans. Other studies (Nakagami et al. 2000; Nakamura et al. 2003; Inoue et al. 2005; Abe 2007) have found tanaidacean prey in shallow-water fishes. Tsubaki et al. (2012) reported tanaidacean as a rare prey component in stomach contents of the deep-sea fish *Caelorinchus anatirostris* Jordan and Gilbert, 1904 (Macrouridae). Kakui (2015a) reported *Zeuxo* sp. from the stomach of an intertidal goby (*Chaenogobius annularis* Gill, 1859) and compared the sex ratio and size-distribution of this tanaidacean in the stomach with those in an environmental substratum sample. The sex ratio in the tanaidacean prey was

significantly male biased, and males in the stomach were significantly larger than those in the environmental sample. Kakui (2015a) concluded that the observed sex and size bias of the prey was likely related to a prey-size preference by the fish, and perhaps to behavioral differences between male and female *Zeuxo* sp., as Johnson and Attramadal (1982b) had previously suggested for a confamilial species.

23.3.7 Ecological Surveys

Tanaidaceans are sometimes reported in ecological surveys documenting faunal assemblages. In Japan, a few such surveys have included this crustacean, including the four studies (Mukai 1971; Kito 1975; Matsumasa and Kurihara 1988; Nakaoka et al. 2001) mentioned above under ‘Phenology’; studies on temporal and spatial variation in community structure and the effects of environmental factors in a seagrass ecosystem (Yamada et al. 2007, 2014); studies of differences in the faunal assemblage among habitats (seagrass bed–coral surface–sandy bottom) or along a depth gradient (Nakamura and Sano 2005; Nakamura et al. 2012; Takada et al. 2012); and a study comparing tanaidacean diversity and abundance among deep-sea sampling sites (Błażewicz-Paszkowycz et al. 2015a). By using experimental substrata in a brackish lagoon, Matsumasa (1994) examined the effects of secondary substratum type on small crustacean assemblages, focusing on three amphipods and a tanaidid, *Sinelobus stanfordi*, and found that the type of secondary substratum directly determined the dominant species. Larsen and Shimomura (2008) investigated the recolonization ability of tanaidaceans on a shallow sandy bottom, and found that a burrower (*Parapseudes arenamans* Larsen and Shimomura, 2008) had greater recolonizing ability than a tube dweller (*Paranesotanais longicephala* Larsen and Shimomura, 2008).

23.4 Summary and Future Perspectives

As mentioned, the numbers of tanaidacean species hitherto reported both globally and from around Japan are likely a small fraction of the actual diversities. This is true even among putatively common shallow-water species; for example, even well-surveyed faunas in the vicinity of marine biological stations harbor many species new to Japan or previously undetected (Fig. 23.5) (Kakui and Kohtsuka 2015; Kakui and Naruse 2015; Nakano et al. 2015). Deep-sea tanaidaceans are even more poorly known.

Błażewicz-Paszkowycz et al. (2015a) detected at least 48 species in the KuramBio collection, none of which has yet been identified to the species level, and which likely include many undescribed species. Likewise, collections made during recent cruises by research/training vessels (e.g., *Nagasaki-maru*, *Seisui-maru*, *Soyo-maru*, *Tansei-maru*,

Toyoshio-maru) around Japan include many deep-sea and some shallow-water tanaidaceans (Fig. 23.5), with a high proportion of species new to Japan or undescribed (Kakui unpubl. data). Tanaidacean systematics is also in its infancy. Further molecular studies are badly needed at all taxonomic levels; even the monophyly of and relationships among the four “superfamilies” sensu Kakui et al. (2011) remain open to question.

Observations of living animals have recently elucidated various new aspects of tanaidacean biology, including a novel feeding mode (seed consumption; Nakaoka 2002), selfing (Kakui and Hiruta 2013), “skin-digging” activity in the holothuroid host by *Exspina typica* (Alvaro et al. 2011), the possibility of producing sound (Kakui et al. 2010), and thread production in apseudomorphs (Drumm 2005; Kakui and Hiruta 2014). This high rate of recent discoveries suggests that many other important discoveries remain to be made, especially with the application of approaches (e.g., breeding experiments) or techniques (e.g., bioimaging, microCT, volume electron microscopy, next-generation sequencing, micromanipulation, genome editing) not previously applied to tanaidaceans. New discoveries will not only broaden our knowledge of tanaidaceans, but will also better place this minor but attractive group in the broader contexts of crustacean biology and general biology.

Acknowledgments

I thank Ryutaro Goto, Masatoshi Matsumasa, and Daisuke Shimada for help in literature surveys; Graham J. Bird, Magdalena Błażewicz-Paszkowycz, and Richard W. Heard for critical comments on the manuscript; and Matthew H. Dick for reviewing and editing the manuscript.

References

- Abe F (2007) Studies on the use of eelgrass-shaped substrate reef by the juvenile red sea bream, *Pagrus major*, and other organisms. Bull Mar Sci Fish Kochi Univ 25:5–57 [in Japanese with English abstract]
- Aboul-Dahab HM, Hussein MA, Soliman FE, Ismail TG (2001) Reproductive aspects on the peracarid crustaceans *Leptochelia erythraea* and *Cyproidea ornata*, from the Red Sea, Egypt. Bull Fac Sci Assiut Univ 30(1-E):23–36
- Alvaro MC, Błażewicz-Paszkowycz M, Davey M, Schiaparelli S (2011) Skin-digging tanaids: the unusual parasitic behaviour of *Exspina typica* in Antarctic waters and worldwide deep basins. Antarct Sci 23:343–348
- Amante C, Eakins BW (2009) ETOPO1 1 Arc-Minute Global Relief Model: Procedures,

- Data Sources and Analysis. NOAA Technical Memorandum NESDIS NGDC-24. National Geophysical Data Center, NOAA. doi:10.7289/V5C8276M. Accessed 15 July 2015
- Anderson G (2013) Tanaidacea—thirty years of scholarship (vers. 1.1, May 2013). <http://peracarida.usm.edu/TanaidaceaText.pdf>. Accessed 1 Dec 2014
- Angsupanich S (2004) A new species of *Longiflagrum* (Tanaidacea, Parapseudidae), from Songkhla Lagoon, Thailand. *Crustaceana* 77:849–860
- Ariyama H, Ohtani M (1990) Record of tanaidacean Crustacea *Sinelobus stanfordi* (Richardson) from the Yodo River in Osaka. *Nanki Seibutsu* 32:23–26 [in Japanese with English abstract]
- Bamber RN (2007) Suborder Apseudomorpha Sieg, 1980 and Neotanaidomorpha Sieg, 1980. In: Larsen K, Shimomura M (eds) Tanaidacea (Crustacea: Peracarida) from Japan III. The deep trenches; the Kurile–Kamchatka Trench and Japan Trench. *Zootaxa* 1599. Magnolia Press, Auckland, pp 13–40
- Barnard JL (1970) Benthic ecology of Bahia de San Quintin Baja California. *Smithson Contrib Zool* 44:1–60
- Bird GJ (2007a) Families Anarthruridae Lang, 1971, Colletteidae Larsen & Wilson, 2002, and Leptognathiidae Sieg, 1976. In: Larsen K, Shimomura M (eds) Tanaidacea (Crustacea: Peracarida) from Japan III. The deep trenches; the Kurile–Kamchatka Trench and Japan Trench. *Zootaxa* 1599. Magnolia Press, Auckland, pp 61–85
- Bird GJ (2007b) Family *incertae sedis*. In: Larsen K, Shimomura M (eds) Tanaidacea (Crustacea: Peracarida) from Japan III. The deep trenches; the Kurile–Kamchatka Trench and Japan Trench. *Zootaxa* 1599. Magnolia Press, Auckland, pp 121–149
- Bird GJ (2012) A new leptocheiloid family, Heterotanoididae (Crustacea: Peracarida: Tanaidacea), and a new species of *Heterotanoides* from New Zealand. *Zootaxa* 3481:1–26
- Bird GJ, Holdich DM (1988) Deep-sea Tanaidacea (Crustacea) of the North-east Atlantic: the tribe Agathotanaini. *J Nat Hist* 22:1591–1621
- Blanc H (1884) Contribution a l’histoire naturelle des asellotes hétéropodes observations faites sur la *Tanais ærstedii*, Krøyer. *Recl Zool Suisse Sér I* 1:189–258 + pls X–XII
- Błażewicz-Paszkowycz M (2007) Family Nototanaidae Sieg, 1976 and Typhlotanaidae Sieg, 1984. In: Larsen K, Shimomura M (eds) Tanaidacea (Crustacea: Peracarida) from Japan III. The deep trenches; the Kurile–Kamchatka Trench and Japan Trench. *Zootaxa* 1599. Magnolia Press, Auckland, pp 101–120

- Błażewicz-Paszkowycz M (2014) 5.19. Tanaidacea. In: De Broyer C, Koubbi P, Griffiths HJ, Raymond B, Udekem d'Acoz C d', Van de Putte AP, Danis B, David B, Grant S, Gutt J, Held C, Hosie G, Huettmann F, Post A, Ropert-Coudert Y (eds) Biogeographic atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, pp 173–180
- Błażewicz-Paszkowycz M, Bamber R, Anderson G (2012) Diversity of Tanaidacea (Crustacea: Peracarida) in the world's oceans—how far have we come? PLoS ONE 7. doi:10.1371/journal.pone.0033068
- Błażewicz-Paszkowycz M, Bamber RN, Jóźwiak P (2013) Tanaidaceans (Crustacea: Peracarida) from the SoJaBio joint expedition in slope and deeper waters in the Sea of Japan. Deep-Sea Res II 86–87:181–213
- Błażewicz-Paszkowycz M, Jennings RM, Jeskulke K, Brix S (2014) Discovery of swimming males of Paratanaoidea (Tanaidacea). Pol Polar Res 35:415–453
- Błażewicz-Paszkowycz M, Pabis K, Jóźwiak P (2015a) Tanaidacean fauna of the Kuril–Kamchatka Trench and adjacent abyssal plain—abundance, diversity and rare species. Deep-Sea Res II 111:325–332
- Błażewicz-Paszkowycz M, Kobylecka E, Jennings RN (2015b) Redescription of wood-associated tanaidacean *Protanais birsteini* (Kudinova-Pasternak, 1970) and its relationship within the Tanaididae. Deep-Sea Res II 111:333–342
- Bückle Ramírez LF (1965) Untersuchungen über die Biologie von *Heterotanaïs oerstedii* Krøyer (Crustacea, Tanaidacea). Z Morph Ökol Tiere 55:714–782
- Chapman MA, Lewis MH, Winterbourn MJ (2011) Guide to the freshwater Crustacea of New Zealand. New Zealand Freshwater Sciences Society, Christchurch
- Darwin CR (1872) The origin of species by means of natural selection, or the preservation of favoured races in the struggle for life. 6th edn. John Murray, London
- Delille D, Guidi LD, Soyer J (1985) Nutrition of *Allotanaïs hirsutus* (Crustacea: Tanaidacea) at Kerguelen Island. In: Siegfried WR, Condy PR, Laws RM (eds) Antarctic nutrient cycles and food webs. Springer, Berlin, pp 378–380
- Dennell R (1937) On the feeding mechanism of *Apseudes talpa*, and evolution of the peracaridan feeding mechanisms. T Roy Soc Edinb 59:57–78
- Drumm DT (2005) Comparison of feeding mechanisms, respiration, and cleaning behavior in two kalliapseudids, *Kalliapseudes macsweenyi* and *Psammokalliapseudes granulatus* (Peracarida: Tanaidacea). J Crustacean Biol 25:203–211
- Drumm DT (2010) Phylogenetic relationships of Tanaidacea (Eumalacostraca:

- Peracarida) inferred from three molecular loci. *J Crustacean Biol* 30:692–698
- Fernandez-Leborans G, Sorbe J-C (2003) A new protozoan of the genus *Cothurnia* (Ciliophora, Peritrichia), epibiont on the marine tanaid *Halmyrapseudes spaansi* (Crustacea Peracarida) from the French Guiana (SE America). *Zool Anz* 242:267–275
- Fonseca DB, D’Incao F (2003) Growth and reproductive parameters of *Kalliapseudes schubartii* in the estuarine region of the Lagoa dos Patos (southern Brazil). *J Mar Biol Assoc UK* 83:931–935
- Gamo S (1984) A new remarkably giant tanaid, *Gigantapseudes maximus* sp. nov. (Crustacea) from the abyssal depths far off southeast of Mindanao, the Philippines. *Sci Rep Yokohama Natl Univ Sect II* 31:1–12
- Giri C, Ochieng E, Tieszen LL, Zhu Z, Singh A, Loveland T, Masek J, Duke N (2011) Status and distribution of mangrove forests of the world using earth observation satellite data. *Global Ecol Biogeogr* 20:154–159
- Guṭu M, Iliffe TM (1989) Description of two new species of Tanaidacea (Crustacea) from the marine water caves of the Palau Islands (Pacific Ocean). *Trav Mus Hist Nat Grigore Antipa* 30:169–180
- Guṭu M, Iliffe TM (2011) *Leptochelia vatulelensis* (Crustacea: Tanaidacea), a new species from anchialine caves of the south-western Pacific. *Trav Mus Natl Hist Nat Grigore Antipa* 54:351–363
- Hassack E, Holdich DM (1987) The tubicolous habit amongst the Tanaidacea (Crustacea, Peracarida) with particular reference to deep-sea species. *Zool Scr* 16:223–233
- Inoue T, Suda Y, Sano M (2005) Food habits of fishes in the surf zone of a sandy beach at Sanrimatsubara, Fukuoka Prefecture, Japan. *Ichthyol Res* 52:9–14
- Ishimaru S (1985) A new species of *Leptochelia* (Crustacea, Tanaidacea) from Japan, with a redescription of *L. savignyi* (Krøyer, 1842). *Publ Seto Mar Biol Lab* 30:241–267
- Johnson SB, Attramadal YG (1982a) A functional-morphological model of *Tanais cavolinii* Milne-Edwards (Crustacea, Tanaidacea) adapted to a tubicolous life-strategy. *Sarsia* 67:29–42
- Johnson SB, Attramadal YG (1982b) Reproductive behaviour and larval development of *Tanais cavolinii* (Crustacea: Tanaidacea). *Mar Biol* 71:11–16
- Kakui K (2014) A novel transmission pathway: first report of a larval trematode in a tanaidacean crustacean. *Fauna Ryukyuana* 17:13–22
- Kakui K (2015a) First report of *Zeuxo* sp. (Crustacea: Tanaidacea) as prey for the

- forktongue goby, *Chaenogobius annularis* Gill, 1859. Rishiri Stud 34:1–6 [in Japanese with English abstract]
- Kakui K (2015b) Descriptions of two new species of *Rhizorhina* Hansen, 1892 (Copepoda: Siphonostomatoida: Nicothoidae) parasitic on tanaidacean crustaceans, with a note on their phylogenetic position. Syst Parasitol (in press)
- Kakui K, Angsupanich S (2012) *Birdotanais songkhlaensis*, a new genus and species of Nototanaidae (Crustacea: Tanaidacea) from Thailand. Raffles B Zool 60:421–432
- Kakui K, Hiruta C (2013) Selfing in a malacostracan crustacean: why a tanaidacean but not decapods. Naturwissenschaften 100:891–894
- Kakui K, Hiruta C (2014) Diverse pereopodal secretory systems implicated in thread production in an apseudomorph tanaidacean crustacean. J Morphol 275:1041–1052
- Kakui K, Kajihara H (2011) *Pseudosphyrapus cuspidiger* sp. nov. from the Nansei Islands, Japan, with a supplementary description of *Kudinopasternakia balanorostrata* Kakui, Kajihara and Mawatari, 2007 (Crustacea: Tanaidacea: Apseudomorpha: Sphyrapodidae). Bull Natl Mus Nat Sci Ser A Suppl 5:53–70
- Kakui K, Kohtsuka H (2015) Two new shallow-water species of *Agathotanaeis* (Crustacea: Tanaidacea) from Japan. Spec Divers 20:45–58
- Kakui K, Naruse T (2015) *Indoapseudes bamberi* sp. nov. (Crustacea: Tanaidacea: Pagurapseudidae: Hodometricinae) from Iriomote Island, Ryukyu Islands, southwestern Japan. Zootaxa 3995:138–148
- Kakui K, Yamasaki H (2013) Nototanaids (Crustacea: Tanaidacea) from Japan, with the description of a new species of *Nototanoides*. Spec Divers 18:245–254
- Kakui K, Kajihara H, Mawatari SF (2007) Two new sphyrapodid species (Crustacea: Tanaidacea: Apseudomorpha) from southwestern Japan. Zootaxa 1563:37–54
- Kakui K, Kajihara H, Mawatari SF (2010) A new species of *Nesotanaeis* Shiino, 1968 (Crustacea, Tanaidacea) from Japan, with a key to species and a note on male chelipeds. Zookeys 33:1–17
- Kakui K, Katoh T, Hiruta SF, Kobayashi N, Kajihara H (2011) Molecular systematics of Tanaidacea (Crustacea: Peracarida) based on 18S sequence data, with an amendment of suborder/superfamily-level classification. Zool Sci 28:749–757
- Kakui K, Kobayashi N, Kajihara H (2012) Phylogenetic position of *Arctotanais* in the suborder Tanaidomorpha (Peracarida: Tanaidacea). J Crustacean Biol 32:127–139
- Kakui K, Tomioka S, Yamasaki H (2014) Tanaidaceans from Rishiri Island (Crustacea: Peracarida). Rishiri Stud 33:7–12 [in Japanese with English abstract]
- Kikuchi S, Matsumasa M (1993) Two ultrastructurally distinct types of transporting

- tissues, the branchiostegal and the gill epithelia, in an estuarine tanaid, *Sinelobus stanfordi* (Crustacea, Peracarida). *Zoomorphology* 113:253–260
- Kito K (1975) Preliminary report on the phytal animals in the *Sargassum confusum* region in Oshoro Bay, Hokkaido. *J Fac Sci Hokkaido Univ Ser VI Zool* 20:141–158
- Kobayashi H, Hatada Y, Tsubouchi T, Nagahama T, Takami H (2012) The hadal amphipod *Hirondellea gigas* possessing a unique cellulase for digesting wooden debris buried in the deepest seafloor. *PLoS ONE* 7.
doi:10.1371/journal.pone.0042727
- Kudinova-Pasternak RK (1966a) On a new abyssal tanaidacean from the Pacific ultra-abyssals. *Crustaceana* 12:257–260
- Kudinova-Pasternak RK (1966b) Tanaidacea (Crustacea) of the Pacific ultra-abyssals. *Zool Zh* 45:518–535 [in Russian with English abstract]
- Kudinova-Pasternak RK (1970) [Tanaidacea of the Kurile–Kamchatka Trench]. *Tr Inst Okeanol* 86:341–381 [in Russian]
- Kudinova-Pasternak RK (1976) Additional notes of the fauna of Tanaidacea (Crustacea, Malacostraca) of the deep water trenches. *Tr Inst Okeanol* 99:115–125 [in Russian with English abstract]
- Kudinova-Pasternak RK (1978) Tanaidacea (Crustacea, Malacostraca) from the deep-sea trenches of the western part of the Pacific. *Tr Inst Okeanol* 108:115–135 [in Russian with English abstract]
- Kudinova-Pasternak RK (1984) The Tanaidacea (Crustacea, Malacostraca) of the Sea of Japan. *Zool Zh* 63:828–838 [in Russian with English abstract]
- Kussakin OG, Tzareva LA (1972) Tanaidacea from the coastal zones of the Middle Kurile Islands. *Crustaceana Suppl* 3:237–245
- Lang K (1953) *Apseudes hermaphroditicus* n. sp. A hermaphroditic tanaide from the Antarctic. *Ark Zool Ser* 2 4:341–350 + 4 pls
- Larsen K (2005) Deep-sea Tanaidacea (Peracarida) from the Gulf of Mexico. Brill, Leiden
- Larsen K (2006) Tanaidacea (Crustacea; Peracarida) from chemically reduced habitats—the hydrothermal vent system of the Juan de Fuca Ridge, Escabana Trough and Gorda Ridge, northeast Pacific. *Zootaxa* 1164:1–33
- Larsen K (2007) Family Agathotanaidae Lang, 1971a. In: Larsen K, Shimomura M (eds) Tanaidacea (Crustacea: Peracarida) from Japan III. The deep trenches; the Kurile–Kamchatka Trench and Japan Trench. *Zootaxa* 1599. Magnolia Press, Auckland, pp 41–60

- Larsen K (2012) A new family, genus and species of apseudomorphan Tanaidacea (Crustacea: Peracarida) from the Caribbean Sea. *J Mar Biol Assoc UK* 92:1089–1093
- Larsen K, Shimomura M (2006) Tanaidacea (Crustacea: Peracarida) from Japan. I. Apseudomorpha from the East China Sea, Seto Inland Sea, and Nansei Islands. *Zootaxa* 1341:29–48
- Larsen K, Shimomura M (2007a) Tanaidacea (Crustacea: Peracarida) from Japan. II. Tanaidomorpha from the East China Sea, the West Pacific Ocean and the Nansei Islands. *Zootaxa* 1464:1–43
- Larsen K, Shimomura M (eds) (2007b) Tanaidacea (Crustacea: Peracarida) from Japan III. The deep trenches; the Kurile–Kamchatka Trench and Japan Trench. *Zootaxa* 1599. Magnolia Press, Auckland
- Larsen K, Shimomura M (2008) Tanaidacea (Crustacea: Peracarida) from Japan. IV. Shallow-water species from Akajima with notes on the recolonization potential of tanaids. *Zootaxa* 1678:1–24
- Larsen K, Shimomura M (2009) Tanaidacea (Crustacea: Peracarida) from Japan. V. A new species of *Chauliopleona* from Nagasaki, Kyushu. *Spec Divers* 14:49–59
- Larsen K, Guțu M, Sieg J (2015) Order Tanaidacea Dana, 1849. In: Vaupel Klein JC von, Charmantier-Daures M, Schram FR (eds) *The Crustacea*. Revised and updated, as well as extended from the *Traité de Zoologie* 5. Brill, Leiden, pp 249–329
- Matsumasa M (1994) Effect of secondary substrate on associated small crustaceans in a brackish lagoon. *J Exp Mar Biol Ecol* 176:245–256
- Matsumasa M, Kurihara Y (1988) Distribution patterns of benthic small crustaceans and the environmental factors in a brackish shallow-water lagoon, Gamō lagoon. *Benth Res* 33/34:33–41 [in Japanese with English abstract]
- McClain C, Barry J (2014) Beta-diversity on deep-sea wood falls reflects gradients in energy availability. *Biol Lett* 10. doi:10.1098/rsbl.2014.0129
- McLelland JA (2007) Family Pseudotanaididae Sieg, 1976. In: Larsen K, Shimomura M (eds) Tanaidacea (Crustacea: Peracarida) from Japan III. The deep trenches; the Kurile–Kamchatka Trench and Japan Trench. *Zootaxa* 1599. Magnolia Press, Auckland, pp 87–99
- Mukai H (1971) The phytal animals on the thalli of *Sargassum serratifolium* in the *Sargassum* region, with reference to their seasonal fluctuations. *Mar Biol* 8:170–182
- Nakagami M, Takatsu T, Matsuda T, Takahashi T (2000) Feeding on harpacticoid

- copepods by marbled sole *Pleuronectes yokohamae* juveniles in the coastal areas of Tsugaru Strait, Hokkaido. *Nippon Suisan Gakk* 66:818–824 [in Japanese with English abstract]
- Nakamura Y, Sano M (2005) Comparison of invertebrate abundance in a seagrass bed and adjacent coral and sand areas at Amitori Bay, Iriomote Island, Japan. *Fish Sci* 71:543–550
- Nakamura Y, Horinouchi M, Nakai T, Sano M (2003) Food habits of fishes in a seagrass bed on a fringing coral reef at Iriomote Island, southern Japan. *Ichthyol Res* 50:15–22
- Nakamura Y, Hirota K, Shibuno T, Watanabe Y (2012) Variability in nursery function of tropical seagrass beds during fish ontogeny: timing of ontogenetic habitat shift. *Mar Biol* 159:1305–1315
- Nakano H, Kakui K, Kajihara H, Shimomura M, Jimi N, Tomioka S, Tanaka H, Yamasaki H, Tanaka M, Izumi T, Okanishi M, Yamada Y, Shinagawa H, Sato T, Tsuchiya Y, Omori A, Sekifuji M, Kohtsuka H (2015) JAMBIO Coastal Organism Joint Surveys reveals undiscovered biodiversity around Sagami Bay. *Reg Stud Mar Sci* 2 (Suppl):77–81
- Nakaoka M (2002) Predation on seeds of seagrasses *Zostera marina* and *Zostera caulescens* by a tanaid crustacean *Zeuxo* sp. *Aquat Bot* 72:99–106
- Nakaoka M, Toyohara T, Matsumasa M (2001) Seasonal and between-substrate variation in mobile epifaunal community in a multispecific seagrass bed of Otsuchi Bay, Japan. *Mar Ecol* 22:379–395
- Nunomura N (1985) Marine tanaid and isopod crustaceans off Kagawa Prefecture, Seto Inland Sea, Japan. *Spec Pub Mukaishima Mar Biol Stn* 1985:101–112
- Nunomura N (2005) A new species of genus *Apseudes* (Tanaidacea: Apseudidae) Okinawa, southern Japan. *Bull Toyama Sci Mus* 28:25–31
- Oliver JS, Oakden JM, Slaterry PN (1982) Phoxocephalid amphipod crustaceans as predators on larvae and juveniles in marine soft-bottom communities. *Mar Ecol Prog Ser* 7:179–184
- Oliver PG, Holmes AM, Killeen IJ, Turner JA (2010) Marine bivalve shells of the British Isles (Mollusca: Bivalvia). *Amgueddfa Cymru, National Museum Wales*. <http://naturalhistory.museumwales.ac.uk/britishbivalves/browse/record.php?-recid=276>. Accessed 6 Aug 2015
- Pennafirme S, Soares-Gomes A (2009) Population biology and reproduction of *Kalliapseudes schubartii* Mañé-Garzón, 1949 (Peracarida, Tanaidacea) in a tropical coastal lagoon, Itaipu, southeastern Brazil. *Crustaceana* 82:1509–1526

- Rumbold CE, Obenat SM, Leonardi MS, Spivak ED (2015) Intersex in the gonochoristic crustacean *Tanais dulongii* (Audouin, 1826) (Peracarida: Tanaidacea: Tanaididae): a comparison of external reproductive characteristics. *J Nat Hist* 49:775–788
- Saito N (1996) Notes on poorly known species of the crustaceans recorded from Kaminato Harbor, Hachijo Island, the Izu Islands. *I.O.P. Diving News* 7:2–5 [in Japanese with English abstract]
- Sánchez-García A, Penálver E, Fuente RP-de la, Delclós X (2015) A rich and diverse tanaidomorph (Crustacea: Tanaidacea) assemblage associated with Early Cretaceous resin-producing forests in North Iberia: palaeobiological implications. *J Syst Palaeontol* 13:645–676
- Sano M, Shimizu M, Nose Y (1984) Food habits of teleostean reef fishes in Okinawa Island, southern Japan. *Univ Mus Univ Tokyo Bull* 25:1–128
- Shaffer PL (1979) The feeding biology of *Podarke pugettensis* (Polychaeta: Hesionidae). *Biol Bull* 156:343–355
- Shiino SM (1937) On *Apseudes nipponicus* n. sp. (Crustacea, Tanaidacea). *Annot Zool Jpn* 16:53–62
- Shiino SM (1951a) Note on three species of Tanaidae from Japanese Coast. *Misc Rep Res Inst Nat Resour* 19–21:32–38 [in Japanese]
- Shiino SM (1951b) On two new species of the family Apseudidae found at Seto. *Rep Fac Fish Prefect Univ Mie* 1:12–25
- Shiino SM (1952) A new genus and two new species of the order Tanaidacea found at Seto. *Publ Seto Mar Biol Lab* 2:53–68
- Shiino SM (1963) Tanaidacea collected by Naga Expedition in the Bay of Nha-Trang, South Viet-Nam. *Rep Fac Fish Prefect Univ Mie* 4:437–507
- Shiino SM (1966) On *Kalliapseudes* (*Kalliapseudes*) *tomiokaensis* sp. nov. (Crustacea: Tanaidacea) from Japanese waters. *Rep Fac Fish Prefect Univ Mie* 5:473–488
- Sieg J (1980) Taxonomische Monographie der Tanaidae Dana 1849 (Crustacea: Tanaidacea). *Abh Senckenb Naturforsch Ges* 537:1–267
- Stephensen K (1936) A tanaid (*Tanais stanfordi* Richardson) found in fresh water in the Kurile Islands, with taxonomic remarks on the genus *Tanais* sensu lat. (*Tanais* Audouin et Milne-Edwards 1829, and *Anatanais* Nordenstam 1930). *Annot Zool Jpn* 15:361–373
- Suárez-Morales E, Londoño-Mesa M, Heard RW (2011) Discovery of a new genus of tanaidacean (Crustacea: Tanaidacea: Mirandotanaidae) found associated with a deep-sea terebellid polychaete. *Contr Zool* 80:157–167

- Takada Y, Abe O, Sibuno T (2012) Variations in cryptic assemblages in coral-rubble interstices at a reef slope in Ishigaki Island, Japan. *Fish Sci* 78:91–98
- Tsubaki K, Takeda M, Morii Y, Hashimoto J (2012) Feeding habits of the macrourid fish, *Caelorinchus anatirostris* from the upper continental slope southwest of Nagasaki, Japan. *Jpn J Benth* 67:9–14 [in Japanese with English abstract]
- Warén A, Carrozza F (1994) *Arculus sykesi* (Chaster), a leptonacean bivalve living on a tanaid Crustacea in the Gulf of Genova. *Boll Malacol* 29:303–306
- Wessel P, Smith WHF, Scharroo R, Luis J, Wobbe F (2013) Generic Mapping Tools: improved version released. *Eos, Trans Amer Geophys Union* 94:409–410
- Wirkner CS, Richter S (2008) Morphology of the haemolymph vascular system in Tanaidacea and Cumacea: —implications for the relationships of “core group” Peracarida (Malacostraca; Crustacea). *Arthropod Struct Dev* 37:141–154
- Wolff T (1956) Crustacea Tanaidacea from depths exceeding 6000 meters. *Galathea Rep* 2:187–241
- Yamada K, Hori M, Tanaka Y, Hasegawa N, Nakaoka M (2007) Temporal and spatial macrofaunal community changes along a salinity gradient in seagrass meadows of Akkeshi-ko estuary and Akkeshi Bay, northern Japan. *Hydrobiologia* 592:345–358
- Yamada K, Tanaka Y, Era T, Nakaoka M (2014) Environmental and spatial controls of macroinvertebrate functional assemblages in seagrass ecosystems along the Pacific coast of northern Japan. *Global Ecol Conserv* 2:47–61

Figure legends

Fig. 23.1 Examples of modes of life among tanaidaceans. **a** A sphyrapodid in its burrow, **b** a pagurapseudid inhabiting an empty gastropod shell, **c** a tanaidid in its self-woven, tube-like nest (*arrow*) on a seagrass leaf (tube-dweller)

Fig. 23.2 Map showing records of tanaidaceans from around Japan (only those identified to the species level). The bathymetric contour interval is 1000 m. The map, plots, and bathymetric contour lines were generated with GMT5 software (Wessel et al. 2013) using the ETOPO1 dataset (Amante and Eakins 2009). The source of all records is given in Appendix

Fig. 23.3 Two sympatric species in *Zeuxo* found around the Misaki Marine Biological Station; ethanol fixed specimens. **a** *Zeuxo* sp. 1, with the carapace having small white marks on a dark background, and a dark antennular article 3; this species is abundant on seagrass leaves and sandy bottoms, but rare in holdfasts, **b** *Zeuxo* sp. 2, with the carapace having large white marks and a thick white transverse line on a dark background, and a white antennular article 3; this species is common in holdfasts but rare on seagrass and sandy bottoms

Fig. 23.4 Map showing records of sphyrapodids from around Japan. The five bathymetric contour lines indicate 200, 400, 600, 800, and 1000 m. The map, plots, and bathymetric contour lines were generated with GMT5 software (Wessel et al. 2013) using the ETOPO1 dataset (Amante and Eakins 2009). Based on data from Kakui et al. (2007); Kakui and Kajihara (2011); Błażewicz-Paszkowycz et al. (2013); and Kakui (unpubl. data)

Fig. 23.5 Apseudomorphs new to Japan and/or undescribed, collected from waters in the vicinity of marine biological stations (**a–c**) or during research cruises (**d–f**); fresh specimens. **a d e** Apseudids, **b f** parapseudids, **c** a pagurapseudid in gastropod shell

Table 23.1 Tanaidacean species recorded around Japan

Species/classification	reference ^a	Species/classification	reference ^a
APSEUDOMORPHA			
Apseudidae			
<i>Apseudes nipponicus</i>	Shiino 1937	<i>Akanthophoreus imputatus</i>	LS 2007b
<i>Carpoapseudes spinigena</i>	LS 2007b	<i>Akanthophoreus lispopygmos</i>	BP 2013
<i>Carpoapseudes varindex</i>	LS 2007b	<i>Akanthophoreus longiremis</i> ^b	KP 1978
<i>Fageapseudes brachyomus</i>	LS 2007b	<i>Akanthophoreus undulatus</i>	LS 2007b
<i>Paradoxapseudes littoralis</i>	Shiino 1952	<i>Akanthophoreus verutus</i>	BP 2013
Kalliapseudidae		<i>Chauliopleona armata</i> ^b	KP 1984
<i>Phoxokalliapseudes tomiokaensis</i>	Shiino 1966	<i>Chauliopleona hansknehti</i>	LS 2007a
Metapseudidae		<i>Chauliopleona sinusa</i>	LS 2009
<i>Apseudomorpha albida</i>	Shiino 1951b	Anarthruridae	
<i>Hoplomachus toyoshious</i>	LS 2006	<i>Anarthruropsis langi</i>	KP 1976
<i>Synapseudes setoensis</i>	Shiino 1951b	<i>Anarthruropsis longa</i>	KP 1984
Pagurapseudidae		<i>Keska sei</i>	BP 2013
<i>Indoapseudes bamberi</i>	KaN 2015	<i>Siphonolabrum tenebrosus</i>	LS 2007b
Parapseudidae		Colletteidae	
<i>Longiflagrum nasutus</i>	Nunomura 2005	<i>Collettea cylindrata</i> ^b	KP 1970
<i>Longipedis fragilis</i>	LS 2006	<i>Collettea minima</i>	KP 1978
<i>Parapseudes algicola</i>	Shiino 1952	<i>Leptognathiopsis langi</i>	LS 2007b
<i>Parapseudes arenamans</i>	LS 2008	<i>Nippognathiopsis petila</i>	BP 2013
Sphyrapodidae		<i>Tumidochelia knighti</i>	LS 2007a
<i>Kudinopasternakia balanorostrata</i>	KaK 2011	Cryptocopidae	
<i>Pseudosphyrapus cuspidiger</i>	KaK 2011	<i>Cryptocopoides arcticus</i> ^b	KP 1978
<i>Pseudosphyrapus malyutinae</i>	BP 2013	<i>Cryptocopoides pacificus</i>	LS 2007b
<i>Pseudosphyrapus quintolongus</i>	Ka et al. 2007	Heterotanoididae	
<i>Pseudosphyrapus serratus</i> ^b	KP 1984	<i>Heterotanoides ornatus</i>	KP 1976
TANAIDOMORPHA		Leptocheliidae	
NEOTANAOIDEA		<i>Leptochelia itoi</i>	Ishimaru 1985
Neotanaidae		<i>Leptochelia savignyi</i>	Shiino 1951a
<i>Neotanais americanus</i>	KP 1978	<i>Makassaritanais modestus</i>	KT 1972
<i>Neotanais kuroshio</i>	LS 2007b	<i>Mesotanais birdi</i>	LS 2007a
<i>Neotanais oyashio</i>	LS 2007b	Leptognathiidae	
<i>Neotanais tuberculatus</i>	KP 1970	<i>Biarticulata greveae</i>	KP 1976
<i>Neotanais wolffi</i>	KP 1966b	<i>Biarticulata parelegans</i>	KP 1970
PARATANAOIDEA		<i>Forcipatia rotundicauda</i>	LS 2007b
Agathotanaidae		<i>Leptognathia aneristus</i>	LS 2007b
<i>Agathotanais hadalis</i>	LS 2007b	<i>Leptognathia bamberi</i>	LS 2007a
<i>Agathotanais ingolfi</i> ^b	KP 1970	Nototanaidae	
<i>Agathotanais misakiensis</i>	KaKo 2015	<i>Nesotanais ryukyuensis</i>	Ka et al. 2010
<i>Agathotanais toyoshioae</i>	KaKo 2015	<i>Nototanoides ohtsukai</i>	KaY 2013
<i>Paragathotanais abyssorum</i>	LS 2007b	<i>Paranesotanais longicephalus</i>	LS 2008
<i>Paranarthrura vitjazi</i>	LS 2007b	Paratanaidae	
Akanthophoreidae		<i>Paratanais impressus</i>	KT 1972
<i>Akanthophoreus crassicauda</i>	LS 2007b	Pseudotanaidae	
<i>Akanthophoreus gracilis</i> ^b	KP 1984	<i>Pseudotanais abathagastor</i>	BP 2013
		<i>Pseudotanais affinis</i> ^b	KP 1984
		<i>Pseudotanais intortus</i>	BP 2013
		<i>Pseudotanais nipponicus</i>	LS 2007b
		<i>Pseudotanais nordenskioldi</i>	KP 1978

Table 23.1 (continued)

Species/classification	reference ^a	Species/classification	reference ^a
<i>Pseudotanaïs soja</i>	BP 2013	<i>Typhlotanaïs magdalensis</i>	LS 2007a
<i>Pseudotanaïs vitjazi</i>	KP 1970	<i>Typhlotanaïs ohtsukae</i>	LS 2007a
Tanaellidae		<i>Typhlotanaïs rectus</i>	KP 1978
<i>Arthrura andriashevi</i>	KP 1966a	<i>Typhlotanaïs simplex</i>	BP 2013
<i>Arthrura longicephala</i>	KP 1978	Family incertae sedis	
<i>Tanaella forcifera</i> ^b	KP 1984	<i>Leptognathia microcephala</i>	LS 2007b
<i>Tanaella kummritzia</i>	LS 2007a	<i>Leptognathia vinogradovae</i>	KP 1970
Tanaopsididae		<i>Metatanaïs cylindricus</i>	Shiino 1952
<i>Tanaopsis curta</i>	KP 1984	<i>Robustochelia robusta</i>	KP 1978
<i>Tanaopsis rugaris</i>	BP 2013		
Typhlotanaidae		TANAIDOIDEA	
<i>Larsenotanaïs kamchatikus</i>	LS 2007b	Tanaididae	
<i>Paratyphlotanaïs japonicus</i>	BP 2013	<i>Arctotanaïs alascensis</i>	Ka et al. 2012
<i>Peraeospinosus magnificus</i>	LS 2007b	<i>Protanaïs birsteini</i>	BP 2015b
<i>Torquella angularis</i>	LS 2007b	<i>Sinelobus stanfordi</i>	AO 1990
<i>Torquella elegans</i>	KP 1978	<i>Tanaïs tinhauae</i>	Ka et al. 2011
<i>Torquella grandis</i>	KP 1970	<i>Tanaïs vanis</i> ^b	Sieg 1980
<i>Typhlamia mucronata</i>	KP 1970	<i>Zeuxo coralensis</i>	Sieg 1980
<i>Typhlotanaïs compactus</i>	LS 2007b	<i>Zeuxo maledivensis</i>	Sieg 1980
<i>Typhlotanaïs kussakini</i>	KP 1970	<i>Zeuxo normani</i>	Shiino 1951a
<i>Typhlotanaïs longicephala</i>	KP 1970		

AO, Ariyama and Ohtani; BP, Błażewicz-Paszkowycz et al.; Ka, Kakui; KaK, Kakui and Kajihara; KaKo, Kakui and Kohtsuka; KaN, Kakui and Naruse; KaY, Kakui and Yamasaki; KP, Kudinova-Pasternak; KT, Kussakin and Tzareva; LS, Larsen and Shimomura

^a only a selected source is presented, ^b the distribution around Japan is questionable

a

b

a

b

c

d

e

f

Appendix: List of literatures for Fig. 23.2

- Ariyama H, Ohtani M (1990) Record of tanaidacean Crustacea *Sinelobus stanfordi* (Richardson) from the Yodo River in Osaka. Nanki Seibutsu 32: 23–26. [in Japanese and English abstract and captions]
- Błażewicz-Paszkowycz M, Bamber RN, Jóźwiak P (2013) Tanaidaceans (Crustacea: Peracarida) from the SoJaBio joint expedition in slope and deeper waters in the Sea of Japan. Deep-Sea Res II 86–87: 181–213.
- Błażewicz-Paszkowycz M, Kobylecka E, Jennings RN (2015) Redescription of wood-associated tanaidacean *Protanais birsteini* (Kudinova-Pasternak, 1970) and its relationship within the Tanaididae. Deep-Sea Res II 111: 333–342.
- Edger GJ (2008) Shallow water Tanaidae (Crustacea: Tanaidacea) of Australia. Zootaxa 1836: 1–92.
- Ishimaru S (1985) A new species of *Leptochelia* (Crustacea, Tanaidacea) from Japan, with a redescription of *L. savignyi* (Krøyer, 1842). Publ Seto Mar Biol Lab 30: 241–267.
- Kakui K, Hiruta C (2014) Diverse pereopodal secretory systems implicated in thread production in an apseudomorph tanaidacean crustacean. J Morphol 275: 1041–1052.
- Kakui K, Kajihara H (2011) *Pseudosphyrapus cuspidiger* sp. nov. from the Nansei Islands, Japan, with a supplementary description of *Kudinopasternakia balanorostrata* Kakui, Kajihara and Mawatari, 2007 (Crustacea: Tanaidacea: Apseudomorpha: Sphyrapodidae). Bull Natl Nat Sci, Ser A Suppl 5: 53–70.
- Kakui K, Kohtsuka H (2015) Two new shallow-water species of *Agathotanaïs* (Crustacea: Tanaidacea) from Japan. Species Divers 20: 45–58.
- Kakui K, Naruse T (2015) *Indoapseudes bamberi* sp. nov. (Crustacea: Tanaidacea: Pagurapseudidae: Hodometricinae) from Iriomote Island, Ryukyu Islands, southwestern Japan. Zootaxa 3995: 138–148.
- Kakui K, Yamasaki H (2013) Nototanais (Crustacea: Tanaidacea) from Japan, with the description of a new species of *Nototanoides*. Species Divers 18: 245–254.
- Kakui K, Kajihara H, Mawatari SF (2007) Two new sphyrapodid species (Crustacea: Tanaidacea: Apseudomorpha) from southwestern Japan. Zootaxa 1563: 37–54.
- Kakui K, Kajihara H, Mawatari SF (2010) A new species of *Nesotanais* Shiino, 1968 (Crustacea, Tanaidacea) from Japan, with a key to species and a note on male chelipeds. Zookeys 33: 1–17.
- Kakui K, Katoh T, Hiruta SF, Kobayashi N, Kajihara H (2011) Molecular systematics of Tanaidacea (Crustacea: Peracarida) based on 18S sequence data, with an amendment of suborder/superfamily-level classification. Zool Sci 28: 749–757.
- Kakui K, Kobayashi N, Mawatari SF (2012) Phylogenetic position of *Arctotanais* in the suborder Tanaidomorpha (Peracarida: Tanaidacea). J Crustacean Biol 32: 127–139.
- Kakui K, Tomioka S, Yamasaki H (2014) Tanaidaceans from Rishiri Island (Crustacea: Peracarida). Rishiri Stud 33: 7–12. [in Japanese with English abstract]
- Kudinova-Pasternak RK (1966a) On a new abyssal tanaidacean from the Pacific ultra-abyssals. Crustaceana 12: 257–260.
- Kudinova-Pasternak RK (1966b) Tanaidacea (Crustacea) of the Pacific ultra-abyssals. Zool Zh 45: 518–535. [in Russian with English abstract]

- Kudinova-Pasternak RK (1970) [Tanaidacea of the Kurile-Kamchatka Trench]. Tr Inst Okeanol 86: 341–381. [In Russian]
- Kudinova-Pasternak RK (1976) Additional notes of the fauna of Tanaidacea (Crustacea, Malacostraca) of the deep water trenches. Tr Inst Okeanol 99: 115–125. [in Russian with English abstract]
- Kudinova-Pasternak RK (1978) Tanaidacea (Crustacea, Malacostraca) from the deep-sea trenches of the western part of the Pacific. Tr Inst Okeanol 108: 115–135. [in Russian with English abstract]
- Kudinova-Pasternak RK (1984) The Tanaidacea (Crustacea, Malacostraca) of the Sea of Japan. Zool Zh 63: 828–838. [in Russian with English abstract]
- Kussakin OG, Tzareva LA (1972) Tanaidacea from the coastal zones of the Middle Kurile Islands. Crustaceana Suppl 3: 237–245.
- Lang K (1966) Taxonomische und phylogenetische Untersuchungen über die Tanaidaceen 2. Die Gattung *Parapseudes* G. O. Sars. Ark Zool Ser 2 18: 549–566 + 1 pl.
- Larsen K, Shimomura M (2006) Tanaidacea (Crustacea: Peracarida) from Japan. I. Apseudomorpha from the East China Sea, Seto Inland Sea, and Nansei Islands. Zootaxa 1341: 29–48.
- Larsen K, Shimomura M (2007a) Tanaidacea (Crustacea: Peracarida) from Japan. II. Tanaidomorpha from the East China Sea, the West Pacific Ocean and the Nansei Islands. Zootaxa 1464: 1–43.
- Larsen K, Shimomura M (eds) (2007b) Tanaidacea (Crustacea: Peracarida) from Japan III. The deep trenches; the Kurile-Kamchatka Trench and Japan Trench. Zootaxa 1599. Magnolia Press, Auckland.
- Larsen K, Shimomura M (2008) Tanaidacea (Crustacea: Peracarida) from Japan. IV. Shallow-water species from Akajima with notes on the recolonization potential of tanaids. Zootaxa 1678: 1–24.
- Larsen K, Shimomura M (2009) Tanaidacea (Crustacea: Peracarida) from Japan. V. A new species of *Chauliopleona* from Nagasaki, Kyushu. Species Divers 14: 49–59.
- Matsumasa M (1994) Effect of secondary substrate on associated small crustaceans in a brackish lagoon. J Exp Mar Biol Ecol 176: 245–256.
- Nunomura N (1985) Marine tanaid and isopod crustaceans off Kagawa prefecture, Seto Inland Sea, Japan. Spec Pub Mukaishima Mar Biol Stn 1985: 101–112.
- Nunomura N (2005) A new species of genus *Apseudes* (Tanaidacea: Apseudidae) Okinawa, southern Japan. Bull Toyama Sci Mus 28: 25–31.
- Saito N (1996) Notes on poorly known species of the crustaceans recorded from Kaminato Harbor, Hachijo Island, the Izu Islands. I.O.P. Diving News 7: 2–5. [in Japanese with English abstract and captions]
- Shiino SM (1937) On *Apseudes nipponicus* n. sp. (Crustacea, Tanaidacea). Annot Zool Jpn 16: 53–62.
- Shiino SM (1951a) Note on three species of Tanaidae from Japanese Coast. Misc Rep Res Inst Nat Resour 19–21: 32–38. [in Japanese]
- Shiino SM (1951b) On two new species of the family Apseudidae found at Seto. Rep Fac Fish Prefect Univ Mie 1: 12–25.
- Shiino SM (1952) A new genus and two new species of the order Tanaidacea found at Seto. Publ Seto Mar Biol Lab 2: 53–68.
- Shiino SM (1966) On *Kalliapseudes (Kalliapseudes) tomiokaensis* sp. nov. (Crustacea: Tanaidacea)

- from Japanese waters. Rep Fac Fish Prefect Univ Mie 5: 473–488.
- Sieg J (1980) Taxonomische Monographie der Tanaidae Dana 1849 (Crustacea: Tanaidacea). Abh Senckenb Naturforsch Ges 537: 1–267.
- Stephensen K (1936) A tanaid (*Tanais stanfordi* Richardson) found in fresh water in the Kurile Islands, with taxonomic remarks on the genus *Tanais* sensu lat. (*Tanais* Audouin et Milne-Edward 1829, and *Anatanaïs* Nordenstam 1930). Annot Zool Jpn 15: 361–373.
- Uéno M (1936) Crustacea Malacostraca collected in the lakes of the Island of Kunasiri. Bull Biogeogr Soc Jpn 6: 247–252.