Title	Effects of N-acetylcysteine on root resorption after tooth replantation
Author(s)	Nishimi, Mitsuhiro; Nakamura, Koichi; Hisada, Akina; Endo, Kazuki; Ushimura, Shuya; Yoshimura, Yoshitaka; Yawaka, Yasutaka
Citation	Pediatric Dental Journal, 30(2), 72-79 https://doi.org/10.1016/j.pdj.2020.05.002
Issue Date	2020-08
Doc URL	http://hdl.handle.net/2115/82323
Rights	© 2020. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	PDJA-D-20-00001_R1.pdf

[Instructions for use](#)

Pediatric Dental Journal

Effects of N-acetylcysteine on root resorption after tooth replantation

--Manuscript Draft--

Manuscript Number:	PDJA-D-20-00001R1
Article Type:	Research Paper
Keywords:	dental trauma; NAC; root resorption; odontoclasts
Corresponding Author:	Koichi Nakamura Hokkaido university Sapporo, Japan
First Author:	Mitsuhiko Nishimi
Order of Authors:	Mitsuhiko Nishimi Koichi Nakamura Akina Hisada Kazuki Endo Shuya Ushimura Yoshitaka Yoshimura Yasutaka Yawaka
Abstract:	<p>In the case of complete luxation with dental trauma, delayed replantation leads to a higher probability of root resorption. We evaluated the efficacy of N-acetylcysteine (NAC) as a root canal medication for trauma-associated root resorption.</p> <p>A rat dental trauma model was used. In the test group, mesial root pulpectomy was carried out after tooth extraction. NAC was applied as a root canal medication followed by replantation. We euthanized the rats after two weeks, excised the maxilla, and prepared thin sections, followed by H-E and TRAP staining. Furthermore, RAW264.7 cells were treated with RANKL and NAC. The effects of NAC on osteoclast differentiation were investigated by determining the numbers of osteoclasts and nuclei and quantifying osteoclast-related mRNA expression levels by real-time PCR.</p> <p>The level of root resorption and number of odontoclasts tended to be lower in the NAC groups than in the control group. In terms of osteoclast differentiation, the total number of osteoclasts was lower in both the 5 and 10 mmol/L NAC groups than in the control group. Real-time PCR revealed that the expression levels of RANK , NFATc1 , DC-STAMP , CD47 , and cathepsin K were lower in the 10 mmol/L NAC group.</p> <p>NAC tended to inhibit root resorption and reduce the number of odontoclasts. NAC also inhibited the expression of osteoclast-related genes. Overall, NAC reduced the number of odontoclasts, inhibited bone resorption, and influenced the inhibition of osteoclast differentiation, fusion, and functions. Therefore, we suggest that NAC may be beneficial in the treatment of root resorption.</p>
Suggested Reviewers:	
Opposed Reviewers:	
Response to Reviewers:	

1 INTRODUCTION

Dental trauma takes on complete luxation, in case of the permanent teeth, which occurs in 1.5% to 16% of trauma cases [1]. The prognosis is relatively favorable if treatment is carried out within an hour after luxation, but a previous report indicates that replantation commonly occurs anywhere from 1 to 4 hours after injury [2]. Delaying replantation results in a poorer prognosis, and research by Guedes *et al.* showed that the incidence of inflammatory root resorption 1 to 8 years after trauma is 34.34% [3].

If considerable time has elapsed since luxation and the tissue surrounding the root canal is not expected to survive, a root canal treatment is necessary. Drying of the periodontal membrane is considered to be the principal cause of trauma-associated root resorption, but pulp necrosis is also involved [4]. Trauma guidelines published by the International Association of Dental Traumatology recommend pulp treatment outside the oral cavity of luxated teeth for which one or more hour(s) has passed since injury [5]. Root canal treatments, including the treatment of luxated teeth, require anti-inflammatory and antibacterial effects to fight inflammation occurring at the time of trauma and infections that develop after luxation, as well as inhibitory effects on root resorption when the prognosis is poor. An agent that is currently in widespread use is calcium hydroxide

(Ca(OH)₂). Calcium hydroxide is used as a root canal medication for traumatized teeth as it has the necessary antibacterial [6], anti-inflammatory [7], hard-tissue-forming [8], and root resorption inhibitory activities [9]. However, leakage of calcium hydroxide outside the apical foramen can lead to complications [10], and reports indicate that its long-term use weakens the root dentin [11], making long-term use problematic.

Therefore, the prospect of developing a new root canal medication remains desirable.

The amino acid N-acetylcysteine (NAC) is a glutathione precursor with antioxidant activities in the human body. It has been reported that NAC inhibit a bone resorption due to the decrease of osteoclasts [12]. If NAC shows inhibitory effects on root resorption, it will be possible to justify its use as a novel root canal medication.

Unfortunately, there are few reports of its inhibitory effects on root resorption.

A rat traumatized tooth model was prepared to verify the effects of NAC on root resorption in traumatized teeth. The model used the first maxillary molars, and NAC was used as a root canal medication for treating root resorption. In addition to the animal model, we treated RAW264.7 cells with receptor activator of nuclear factor κ -B ligand (RANKL), an osteoclast differentiation factor, and NAC. The effects of NAC on osteoclast differentiation were investigated.

2 MATERIALS AND METHODS

2.1 Preparation of a rat traumatized tooth model

The experimental animals were 6-week-old male SD rats (Sankyo Labo Service Corporation, Inc. Tokyo, Japan), and NAC (Sigma-Aldrich Corporation, Missouri, USA) was used as a root canal medication. The study was approved by Hokkaido University's Animal Committee (protocol approval no.: 14-0063) and was carried out in accordance with its instructions.

The right and left first maxillary molars were extracted under total anesthesia achieved using pentobarbital (Somnopentyl[®]; Kyoritsu Seiyaku Corporation, Tokyo, Japan) and isoflurane (Escain[®]; Pfizer, Inc., Tokyo, Japan) and allowed to dry for 1 hour. After removing the molar caps with a round bur and expanding to no. 25 with a K-file, the root canal interior was washed with sodium hypochlorite (Neocleaner[®]; Neo Dental Chemical Products Co., Ltd. Tokyo, Japan). In the control group, we replanted teeth after drying without other treatment. The three test groups included treatments with 1, 5, and 10 mmol/L NAC before replantation of the root canal (n=4). We sealed the apical end of the root with adhesive resin cement (Super-Bond[®]; Sun Medical Co., Ltd, Shiga, Japan) and the coronal end with glass ionomer cement (Fuji Ionomer Type II[®]; GC, Tokyo, Japan) before drying. Replantation was carried out, followed by grinding of the

opposing teeth.

The rats were euthanized two weeks later. Perfusion fixation was performed using 4% paraformaldehyde - maleic acid buffer (Wako Pure Chemical Industries, Ltd., Osaka, Japan), followed by excision and immersion fixation of the maxilla. After decalcification with EDTA, we embedded the sample in paraffin by standard reported methods. We took 5- μ m sections perpendicular to the tooth axis. Hematoxylin-eosin (H-E) staining was carried out, and root sections were photographed. The root section areas were measured using Image J (NIH, Maryland, USA), and the residual root section area (%) as a proportion of the mean root section area in untreated teeth was determined for each treatment group (n = 4). Staining was also carried out with tartrate-resistant acid phosphatase (TRAP) containing Fast Red Violet LB Salt (Sigma-Aldrich). We used light microscopy to observe odontoclasts in contact with the root surface.

2.2 Preparation of cells

RAW264.7 cells (ATCC no.: TIB-71TM; Virginia, USA), monocytic macrophages derived from murine ascites, were used as osteoclast precursor cells. RAW264.7 cells express RANK, and differentiation to osteoclasts can be achieved by stimulation with RANKL even without stimulation by a macrophage colony-stimulating factor [13]. The

cells were cultured in Dulbecco's modified Eagle's medium (D-MEM; Wako Pure Chemical) containing 10% bovine fetal serum (Invitrogen, California, USA) and 66.5 µg/L kanamycin sulfate (Meiji Seika Pharma Co., Ltd., Tokyo, Japan). The cells were cultured at 37 °C in a 5% carbon dioxide atmosphere in a 100-mm standard dish (BD Falcon, New Jersey, USA), after which they were washed with Dulbecco's phosphate-buffered saline containing no calcium and magnesium ions (PBS) and harvested using PBS containing 0.48 mmol/L EDTA.

2.3 Determining the number of viable cells

Cultured RAW264.7 cells were inoculated onto a 96-well plate to a cell density of 1.0×10^4 cells per well. The cells were treated with α -minimum essential medium (α -MEM; Wako Pure Chemical) containing 10% fetal bovine serum, 434.4 mg/L L-alanyl-L-glutamine (Wako Pure Chemical), 72.7 mg/L L-ascorbate-2-phosphate (Sigma-Aldrich), and 66.5 µg/L kanamycin sulfate, to which 100 mmol/L NAC stock solution (adjusted to pH 7.2) was added to final concentrations of 1, 5, and 10 mmol/L. The control group was the same but without NAC added. The suspensions were cultured at 37 °C in a 5% carbon dioxide atmosphere. We considered the inoculation of RAW264.7 cells onto the plate to be culture day 1. We replaced the culture medium on day 3 and

day 5. We measured the intracellular ATP and determined the viable cell number. The ViaLight™ plus kit (LONZA, Maryland, USA) was used for measurement, and a multi-label counter (Wallac 1420 ARVOSX; PerkinElmer, Inc., Massachusetts, USA) was used to measure fluorescence (n = 8).

2.4 Confirmation of osteoclast differentiation

Cultured RAW264.7 cells were inoculated onto a 24-well plate to a cell density of 1.0×10^4 cells per well. The only substance added was RANKL, to a final concentration of 50 ng/mL in α -MEM in the control group (as used for determining the number of viable cells). Three test groups were also established with NAC concentrations of 1, 5, and 10 mmol/L added at the same time as RANKL. Both the control and test groups were all cultured at 37 °C under a 5% carbon dioxide atmosphere. The date of inoculation was considered culture day 1, and the culture medium was replaced on day 3. On day 5, the cells were washed with PBS, fixed with 10% formalin, and then washed with distilled water. We stained cells with TRAP stain (pH 5.0) containing Fast Red Violet LB Salt. We determined osteoclast morphology by light microscopy.

2.5 Real-time polymerase chain reaction (PCR)

Cultured RAW264.7 cells were inoculated onto a six-well plate to a cell density of 5.0×10^4 cells per well. For the control group, the only substance added was RANKL, to a final concentration of 50 ng/mL in α -MEM (as used for determining the number of viable cells). Three test groups were also set up with NAC at concentrations of 1, 5, and 10 mmol/L added at the same time as RANKL. The control and test groups were cultured for 12 hours, 24 hours, and 5 days. After culturing, RNA was harvested from the cells using Trizol (Invitrogen). The reverse transcription reaction was carried out with Rever Tra Ace- α FSK-101 (Toyobo Co., Ltd., Osaka, Japan), using 1.0 μ g of RNA to synthesize cDNA. We carried out real-time PCR with an ABI7300 PCR system (Applied Biosystems; Thermo Fisher Scientific, Inc., Massachusetts, USA) using the following primers (Applied Biosystems; Thermo Fisher Scientific): TRAP (Mm00475698_m1), RANK (Mm00437132_m1), cathepsin-K (Cath-K; Mm00484039_m1), NFATc1 (Mm00479445_m1), DC-STAMP (Mm04209236_m1), OC-STAMP (Mm00512445_m1), and CD47 (Mm00495011_m1). The gene expression levels of the samples were standardized using the glyceraldehyde 3-phosphate dehydrogenase expression level and calculated by the $2^{-\Delta\Delta C_t}$ method (n = 4).

2.6 Statistical analyses

All data were expressed as means $\pm$ standard deviations. A Bonferroni's test was used to determine statistical significance, and differences were taken to be significant at $P < 0.05$.

3 RESULTS

3.1 Histological findings

In untreated teeth, we found calcified material around the root dentin. In the control group, TRAP staining confirmed existence of many odontoclasts (Fig. 1A). After replantation with 1 mmol/L NAC application, we found root resorption and odontoclasts, whereas with 5 and 10 mmol/L NAC, the numbers of both resorption lacunae and odontoclasts tended to decrease (Fig. 1A). Measurement of the residual root section area proportions showed a significant difference between the control and 10 mmol/L NAC groups (Fig. 1B).

3.2 Effects of NAC on osteoclast differentiation

We assessed the effects of NAC on osteoclast differentiation. To confirm the presence of viable cells, NAC was added to RAW264.7 cells at various concentrations, and we measured intracellular ATP levels after culture for 5 days. Treatments did not show a decrease in ATP, irrespective of the NAC concentration (Fig. 2A). The number of osteoclasts was also determined in the control group (RANKL only) and in the test groups (RANKL+NAC). In the control and 1 mmol/L NAC groups, large osteoclasts were found, whereas in the 5 and 10 mmol/L NAC groups, the osteoclasts tended to be

smaller (Fig. 2B). The numbers of osteoclasts with two or more nuclei decreased significantly in the 5 and 10 mmol/L NAC treatment groups compared with the control group (Fig. 2C). Comparing the osteoclasts by the number of nuclei, the numbers with two and three nuclei decreased significantly in the 10 mmol/L NAC group. Similarly, the numbers with four and five nuclei decreased significantly in the 5 mmol/L NAC group compared with the control group (Fig. 2D).

3.3 Osteoclast-related gene expression

We performed real-time PCR to quantify osteoclast-related gene mRNA expression. The group treated with only RANKL was considered the control group, and the groups treated with each NAC concentration, cultured for 12 hours, 24 hours, and 5 days, were taken to be the test groups. The mRNA expression after 12 hours decreased significantly for *DC-STAMP* with 5 and 10 mmol/L NAC and *CD47* with 1 and 5 mmol/L. However the mRNA expression increased significantly for *NFATc1* with 1 and 5 mmol/L. There was no significant difference in the others (Fig. 3A). After 24 hours, it decreased significantly for both *NFATc1* and *DC-STAMP* with 10 mmol/L NAC, but the mRNA expression increased significantly for *cathepsin K* and *CD47* with 1 mmol/L. There was no significant difference for *TRAP* and *RANK* (Fig. 3B). After 5 days, the mRNA

expressed decreased significantly for *RANK* with 5 and 10 mmol/L NAC, *NFATc1* with 5 mmol/L NAC, *DC-STAMP* with 5 and 10 mmol/L NAC, *CD47* with 5 and 10 mmol/L NAC, *OC-STAMP* with 5 mmol/L NAC, *cathepsin K* with 5 and 10 mmol/L NAC, and *TRAP* with 1 mmol/L NAC (Fig. 3C).

4 DISCUSSION

This study used a rat traumatized tooth model followed by replantation to evaluate the effects of NAC application on root resorption. This animal model was complemented by cell culture to address whether the addition of RANKL and NAC to RAW264.7 cells affected osteoclast differentiation. Luxated teeth have a favorable prognosis after replantation if drying time is kept short and/or they are kept in an appropriate storage solution [14, 15]. However, with prolonged drying time and delayed replantation, the periodontal membrane and dental pulp tissues undergo necrosis, and root resorption frequently occurs in cases with particularly poor prognoses. The International Association of Dental Traumatology's guidelines [5] define delayed replantation as occurring when the drying time is 1 hour or more. Therefore, in this study, the drying time was set at 1 hour for a traumatized tooth model. Lacunae with odontoclasts were found as histological signs of resorption in both the control and 1 mmol/L NAC groups.

The numbers of resorption lacunae and numbers of odontoclasts in contact with the root tended to decrease in the 5 and 10 mmol/L NAC groups (Fig. 1A). Furthermore, with respect to the root section area, root resorption was found to be inhibited in the 10 mmol/L NAC group (Fig. 1B). Bacteria and endotoxins derived from the necrotic pulp tissue may move from dentinal tubules into the gaps and damaged areas left by root resorption and periodontal membrane necrosis [4]. Therefore, luxation may have resulted in the periodontal membrane and pulp necrosis, probably resulting in root resorption in the control group. However, in the 1 mmol/L NAC group, approximately the same degree of root resorption was found in the control group, irrespective of pulpectomy and medication application. This result suggests that periodontal membrane and pulp necrosis are not the sole causes of root resorption and that odontoclast induction may occur due to mechanical stimulation at the time of luxation. Light microscopic examination showed a tendency toward inhibition of osteoclast differentiation in cell cultures with 5 and 10 mmol/L NAC (Fig. 2B), and the total osteoclast number also decreased in comparison with the control group. In examining osteoclasts with different numbers of nuclei, two and three nuclei significantly decreased with 10 mmol/L NAC and four and five nuclei significantly decreased with 5 mmol/L NAC when compared to the control group (Fig. 2C, D, Table 1). NAC treated

RAW264.7 cells showed no differences in the intracellular ATP levels between the control and any of the NAC concentrations (Fig. 2A). In other words, the decreases in total osteoclast numbers were not due solely to cell death; the inhibition of osteoclast differentiation is likely also involved. A decrease in the number of oligonucleated cells is likely due to inhibition at an early stage in the differentiation process, whereas a decrease in the number of polynucleate cells is due to inhibition of fusion between cells, which occurs later in the differentiation process.

Please insert Table I here.

The cells responsible for root resorption are osteoclasts (i.e., odontoclasts). For tissue resorption, it is important for monocyte/macrophage lineage precursor cells to differentiate into mononucleate osteoclasts and fuse to form polynucleate osteoclasts.

Osteoclast differentiation requires the master transcription factor NFATc1 [16] and RANK, RANKL-receptor [17]. DC-STAMP is important [18] for osteoclast fusion.

CD47 is involved in the fusion between oligonucleate osteoclasts [19], whereas fusion between polynucleate osteoclasts involves OC-STAMP [20]. In addition, cathepsin K, a cysteine protease that breaks down type-I collagen, is important for bone resorption by osteoclasts [21].

Please insert Table 2 here

In the present study, we found that 10 mmol/L NAC had inhibitory effects on trauma-associated root resorption (Fig. 3; Table 2). In addition, the real-time PCR showed a decreased expression of *RANK*, *NFATc1*, *DC-STAMP*, and *CD47* mRNA with 10 mmol/L NAC. Decreases in *RANK* and *NFATc1* affect osteoclast differentiation, whereas the decreases in *DC-STAMP* and *CD47* affect fusion between osteoclasts. This is consistent with the reductions in oligonucleated cells and numbers of osteoclasts in the present study. The number of osteoclasts with two or more nuclei decreased with 10 mmol/L NAC; the real-time PCR results, however, showed no difference from the control in the expression of *TRAP*, an osteoclast marker gene. *TRAP* is expressed even by mononucleated osteoclasts, but in the present study, two or more nuclei were the criterion for the classification of cells as osteoclasts. This classification may account for the discrepancy between these results. After 5 days, a *OC-STAMP* expression was significantly increased with 10 mmol/L NAC, whereas no significant difference from the control was found in the number of polynucleated cells. This finding is probably due to a decrease in the number of minority nuclei required for fusion of the polynucleated cells. Moreover, at day 5, *cathepsin K* mRNA expression also decreased. For these reasons, osteoclast functions may be inhibited along with osteoclast differentiation and fusion.

In this study, we identified calcified material around the root canal in rats. At many loci, the calcified material was incorporated into the root dentin without a border between the two. Therefore, investigation of the effects of NAC on the root included calcified material. The results of real-time PCR with 5 mmol/L NAC were not markedly different from those with 10 mmol/L NAC, and when comparing root resorption, the 5 mmol/L NAC group was not significantly different from the control group. In cell culture, effects were found without changes in NAC concentration. In the rat model, on applying the root canal medication and sealing both the apical and coronal sides, NAC in the tissues surrounding the root was likely released from only the dentinal tubules. Therefore, it is probable that the observed effects resulted from NAC levels that were, in reality, lower than those relating to the treatment concentrations. In future research, it will be necessary to investigate the appropriate concentrations for root canal medication. The results of this study suggest that in a rat traumatized tooth model, 10 mmol/L NAC can inhibit root resorption by reducing the levels of transcription factors and genes that control osteoclast differentiation and fusion. There have been reports of complications with calcium hydroxide treatment due to apical exudation. Because NAC is an antioxidant amino acid, it is highly probable that it would be absorbed even if apical exudation did occur. In addition, since calcium hydroxide has a pH of 12, it may

establish an alkaline environment for an extended time, which can result in degradation of root tissues and reduced dentin strength [22, 23]. By contrast, the NAC used in this study is adjusted to a neutral pH of approximately 7.2 and, therefore, may prevent a decrease in dentin strength. NAC has antibacterial activities [24, 25] and is reported to have anti-inflammatory activities due to inhibition of the inflammatory cytokine TNF- α [7]. Based on these reports, NAC could be used to treat inflammation due to bacterial infection of the dental pulp or trauma. For that reason, NAC offers one possible option as a root canal medication for traumatized teeth.

The periodontal tissues of rats and humans have various histological similarities [26] and, as a result, many dental studies use rats [27]. The results of the present study reveal a need for further research investigating the comparison with calcium hydroxide, which is currently widely used for root canal medication, the appropriate NAC concentrations for root canal medication and the effects of NAC on dentin strength in clinical application. These should expand into studies on human teeth.

5 CONCLUSIONS

The use of NAC in a rat traumatized tooth model resulted in the inhibition of root resorption and a tendency toward a decrease in the number of odontoclasts. NAC

application with cell culture resulted in the inhibition of osteoclast-related gene expression. Several studies report that NAC reduces the number of odontoclasts, inhibits root resorption, and is involved in inhibiting osteoclast differentiation and fusion and osteoclast functions. Therefore, there is a potential for the application of NAC in the treatment of root resorption.

CONFLICTS OF INTEREST

There are no conflicts of interest associated with this study.

ACKNOWLEDGMENTS

This work was supported in part by JSPS KAKENHI (No. 19K19256, 19K10374). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

REFERENCES

- [1] Andreasen JO. Etiology and pathogenesis of traumatic dental injuries. A clinical study of 1,298 cases. *Scand J Dent Res* 1970;78:329-42.
- [2] Zhang X, Gong Y. Characteristics of avulsed permanent teeth treated at Beijing Stomatological Hospital. *Dent Traumatol* 2011;27:379-84.
- [3] Yang M, Kahn AM. Insulin-stimulated NADH/NAD⁺ redox state increases NAD(P)H oxidase activity in cultured rat vascular smooth muscle cells. *Am J Hypertens* 2006;19:587-

92.

- [4] Andreasen JO. Relationship between surface and inflammatory resorption and changes in the pulp after replantation of permanent incisors in monkeys. *J Endod* 1981;7:294-301.
- [5] Flores MT, Andersson L, Andreasen JO, et al. Guidelines for the management of traumatic dental injuries. II. Avulsion of permanent teeth. *Dent Traumatol* 2007;23:130-6.
- [6] Silva LA, Romualdo PC, Silva RA, et al. Antibacterial Effect of Calcium Hydroxide With or Without Chlorhexidine as Intracanal Dressing in Primary Teeth With Apical Periodontitis. *Pediatr Dent* 2017;39:28-33.
- [7] Pinar Karapinar S, Ulum YZ, Ozcelik B, et al. The effect of N-acetylcysteine and calcium hydroxide on TNF- α and TGF- β 1 in lipopolysaccharide-activated macrophages. *Arch Oral Biol* 2016;68:48-54.
- [8] Wang S, Sasaki Y, Ogata Y. Calcium hydroxide regulates bone sialoprotein gene transcription in human osteoblast-like Saos2 cells. *J Oral Sci* 2011;53:77-86.
- [9] Lengheden A, Blomlöf L, Lindskog S. Effect of immediate calcium hydroxide treatment and permanent root-filling on periodontal healing in contaminated replanted teeth. *Scand J Dent Res* 1991;99:139-46.
- [10] Sharma S, Hackett R, Webb R, et al. Severe tissue necrosis following intra-arterial injection of endodontic calcium hydroxide: a case series. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008;105:666-9.
- [11] Andreasen JO, Farik B, Munksgaard EC. Long-term calcium hydroxide as a root canal dressing may increase risk of root fracture. *Dent Traumatol* 2002;18:134-7.
- [12] Jay J. Cao, Matthew J. Picklo. N-Acetylcysteine Supplementation Decreases Osteoclast Differentiation and Increases Bone Mass in Mice Fed a High-Fat Diet. *J. Nutr.* 2014;144:289-96.
- [13] Islam S, Hassan F, Tumurkhuu G, et al. Receptor activator of nuclear factor-kappa B ligand induces osteoclast formation in RAW 264.7 macrophage cells via augmented production of macrophage-colony-stimulating factor. *Microbiol Immunol* 2008;52:585-90.
- [14] Leite MC, Okamoto T. The influence of extra-oral time upon healing after tooth replantation. A histological study in rat incisors. *J Nihon Univ Sch Dent* 1984;26:316-30.
- [15] Pettiette M, Hupp J, Mesaros S, et al. Periodontal healing of extracted dogs' teeth air-dried for extended periods and soaked in various media. *Endod Dent Traumatol* 1997;13:113-8.
- [16] Takayanagi H, Kim S, Koga T, et al. Induction and activation of the transcription factor NFATc1 (NFAT2) integrate RANKL signaling in terminal differentiation of osteoclasts. *Dev Cell* 2002;3:889-901.
- [17] Dougall WC, Glaccum M, Charrier K, et al. RANK is essential for osteoclast and lymph

node development. *Genes Dev* 1999;13:2412-24.

[18] Xing L, Xiu Y, Boyce BF. Osteoclast fusion and regulation by RANKL-dependent and independent factors. *World J Orthop* 2012;3:212-22.

[19] Lundberg P, Koskinen C, Baldock PA, et al. Osteoclast formation is strongly reduced both in vivo and in vitro in the absence of CD47/SIRPalpha-interaction. *Biochem Biophys Res Commun* 2007;352:444-8.

[20] Witwicka H, Hwang SY, Reyes-Gutierrez P, et al. Studies of OC-STAMP in Osteoclast Fusion: A New Knockout Mouse Model, Rescue of Cell Fusion, and Transmembrane Topology. *PLoS One* 2015;10:e0128275.

[21] Troen BR. The role of cathepsin K in normal bone resorption. *Drug News Perspect* 2004;17:19-28.

[22] Rosenberg B, Murray PE, Namerow K. The effect of calcium hydroxide root filling on dentin fracture strength. *Dent Traumatol* 2007;23:26-9.

[23] Batur YB, Erdemir U, Sancakli HS. The long-term effect of calcium hydroxide application on dentin fracture strength of endodontically treated teeth. *Dent Traumatol* 2013;29:461-4.

[24] Moon JH, Jang EY, Shim KS, et al. In vitro effects of N-acetyl cysteine alone and in combination with antibiotics on *Prevotella intermedia*. *J Microbiol* 2015;53:321-9.

[25] Amaral EP, Conceição EL, Costa DL, et al. N-acetyl-cysteine exhibits potent anti-mycobacterial activity in addition to its known anti-oxidative functions. *BMC Microbiol* 2016;16:251.

[26] Klausen B. Microbiological and immunological aspects of experimental periodontal disease in rats: a review article. *J Periodontol* 1991;62:59-73.

[27] Esteves JC, Marão HF, Silva PI, et al. Delayed tooth replantation following root canal filling with calcium hydroxide and MTA: Histomorphometric study in rats. *Arch Oral Biol* 2015;60:1254-62.

Figure legends

Fig. 1. Tissue images with different NAC concentrations applied to root canal

The first maxillary molars were extracted, pulpectomy was carried out from the coronal

side, and NAC was applied at various concentrations. After drying for 1 hour, replantation was carried out. A) Calcified material was found around the root canal in untreated teeth. Resorption lacunae are indicated in the control and 1 mmol/L NAC groups, with fewer in the 5 and 10 mmol/L NAC groups compared with the control group. Scale bar = 100 μ m. B) Root resorption was inhibited with 10 mmol/L NAC compared with the control group. Residual root section area (%) = root section area in samples from each test group/mean root section area in untreated tooth $\times$ 100. Error bars represent mean $\pm$ SD; n = 4; p < 0.01 (**), p < 0.05 (*).

Fig. 2. Effects of NAC on osteoclast differentiation

NAC was added to RAW264.7 cells at various concentrations. After 5 days of culture, intracellular ATP levels were measured using the ViaLight™ Plus kit, and osteoclasts were observed with TRAP staining. A) ATP levels decreased in comparison with the control group (no added NAC) with any added concentration of NAC. Mean $\pm$ SD; n = 8. B) Large osteoclasts were seen in the control and 1 mmol/L NAC groups, whereas cells trended smaller in the 5 and 10 mmol/L NAC groups ($\rightarrow$: large osteoclasts). Scale bar=250 μ m. C) The total number of osteoclasts (i.e., two or more nuclei) decreased significantly in the 5 and 10 mmol/L NAC groups. Mean $\pm$ SD; n = 4; p < 0.05 (*). D) Classifying osteoclasts by the number of nuclei, osteoclasts with two and three nuclei

decreased significantly in the 10 mmol/L NAC treatment group. The number with four and five nuclei decreased significantly in the 5 mmol/L NAC group. Mean $\pm$ SD; n = 4; $p < 0.01$ (**); $p < 0.05$ (*).

Fig. 3. Changes in osteoclast-related gene expression after culture for 12 hours, 24 hours and 5 days.

RANKL and different concentrations of NAC to RAW264.7 were added to cells, and after 12 hours, 24 hours and 5 days. we compared the mRNA expression levels. A) After 12 hours, compared to control, significant decreases in *DC-STAMP* expression were found in the 5 and 10 mmol/L NAC groups, and *CD47* expression in the 1 and 5 mmol/L NAC group. However, significant increase in *NFATc1* expression were found in the 1 and 5 mmol/L NAC group. There was no significant difference in the others. B) After 24 hours, compared to control, significant decreases were found in *NFATc1* and *DC-STAMP* expression in the 10 mmol/L NAC group. However, significant increase in *Cath-K* and *CD47* expression were found in the 1 mmol/L NAC group. There was no significant difference for *TRAP*, *RANK* and *OC-STAMP*. C) After 5 days, compared to control, significant decreases in expression were found for *TRAP*, *RANK*, *Cath-K*, *DC-STAMP* and *CD47*, and for *RANK*, *Cath-K*, *NFATc1*, *DC-STAMP*, *OC-STAMP* and *CD47* in the 5 mmol/L NAC group, and for *RANK*, *Cath-K*, *DC-STAMP*, *OC-STAMP* and *CD47* in the

10 mmol/L NAC group.

Mean $\pm$ SD; n = 4; p < 0.01 (**); p < 0.05 (*)

control

1 mmol/L NAC

5 mmol/L NAC

10 mmol/L NAC

Table 1. Effects of NAC on osteoclast numbers

	1 mmol/L NAC	5 mmol/L NAC	10 mmol/L NAC
Number of osteoclasts (two or more nuclei)	—	↓	↓
2 or 3 nuclei	—	—	↓
4 or 5 nuclei	—	↓	—

Table 2. Effects of NAC on osteoclast-related gene expression

		1 mmol/L NAC	5 mmol/L NAC	10 mmol/L NAC
12 h	<i>TRAP</i>	—	—	—
	<i>RANK</i>	—	—	—
	<i>Cath-K</i>	—	—	—
	<i>NFATc1</i>	↑	↑	—
	<i>DC-STAMP</i>	—	↓	↓
	<i>OC-STAMP</i>	—	—	—
	<i>CD47</i>	↓	↓	—
24 h	<i>TRAP</i>	—	—	—
	<i>RANK</i>	—	—	—
	<i>Cath-K</i>	↑	—	—
	<i>NFATc1</i>	—	—	↓
	<i>DC-STAMP</i>	—	—	↓
	<i>OC-STAMP</i>	—	—	—
	<i>CD47</i>	↑	—	—
Day 5	<i>TRAP</i>	↓	—	—
	<i>RANK</i>	↓	↓	↓
	<i>Cath-K</i>	↓	↓	↓
	<i>NFATc1</i>	—	↓	—
	<i>DC-STAMP</i>	↓	↓	↓
	<i>OC-STAMP</i>	—	↓	↑
	<i>CD47</i>	↓	↓	↓