



Title	Double coverings of arrangement complements and 2-torsion in Milnor fiber homology
Author(s)	Yoshinaga, Masahiko
Citation	European Journal of Mathematics, 6(3), 1097-1109 <a href="https://doi.org/10.1007/s40879-019-00387-8">https://doi.org/10.1007/s40879-019-00387-8</a>
Issue Date	2020-09
Doc URL	<a href="http://hdl.handle.net/2115/82566">http://hdl.handle.net/2115/82566</a>
Rights	The final publication is available at <a href="http://link.springer.com">link.springer.com</a>
Type	article (author version)
File Information	Eur. J. Math.6-3_1097-1109.pdf



[Instructions for use](#)

# DOUBLE COVERINGS OF ARRANGEMENT COMPLEMENTS AND 2-TORSION IN MILNOR FIBER HOMOLOGY

MASAHIKO YOSHINAGA

*Dedicated to the memory of Stefan Papadima*

ABSTRACT. We prove that the mod 2 Betti numbers of double coverings of a complex hyperplane arrangement complement are combinatorially determined. The proof is based on a relation between the mod 2 Aomoto complex and the transfer long exact sequence.

Applying the above result to the icosidodecahedral arrangement (16 planes in the three dimensional space related to the icosidodecahedron), we conclude that the first homology of the Milnor fiber has non-trivial 2-torsion.

## CONTENTS

1. Introduction	2
2. Generalities on hyperplane arrangements	3
3. Finite coverings and combinatorial structures	5
3.1. Finite abelian covering	5
3.2. Double covering and transfer exact sequence	6
3.3. Mod 2 Betti number of $\mathbb{Z}_4$ -liftable double coverings	6
3.4. Double coverings of arrangement complements	7
4. 2-torsions in Milnor fiber homology	8
4.1. Icosidodecahedral arrangement $\mathcal{A}_{\mathcal{ID}}$	8
4.2. Mod 2 Aomoto complex of icosidodecahedral arrangement	10
4.3. Milnor fiber of icosidodecahedral arrangement	11
References	13

---

*Date:* November 14, 2019.

*2010 Mathematics Subject Classification.* Primary 52C35, Secondary 20F55.

*Key words and phrases.* Hyperplane arrangements, Milnor fiber, monodromy, Aomoto complex, transfer map, Icosidodecahedron.

The author thanks Professor Toshiyuki Akita for helpful discussion on mod 2 Betti numbers. The author also deeply thanks the referee for careful reading of the manuscript and for many suggestions. This work was partially supported by JSPS KAKENHI Grant Numbers JP18H01115, JP15KK0144.

## 1. INTRODUCTION

An arrangement of hyperplanes is a finite collection of hyperplanes in a finite dimensional vector (affine, or projective) space. For a complex arrangement, we can associate several topological spaces: the complement, Milnor fiber, their covering spaces, boundary manifolds and so on. These provides important spaces in topology such as the classifying space of the pure braid (Artin) group and their subgroups.

Another important aspect of arrangement is the combinatorial structure. An arrangement defines the poset of subspaces expressed as intersections of hyperplanes. These subspaces can be also considered as (the closure of) strata of a stratification of the total space. As is naturally expected, the poset of intersections has a lot of information on the associated topological spaces. Indeed, some of topological invariant (e.g., cohomology ring of the complement [22]. See also §2 for more details) is determined by the poset of intersections, while some other can not be determined by the poset (e.g., the fundamental group [27]). Furthermore, there are many invariant whose relations to the poset of intersections are yet unclear.

Recently, Milnor fibers, and more generally, covering spaces of the arrangement complements received considerable amount of attention. See for [33] for survey on the topic. Among other results, we just recall some of recent results on the Milnor fiber of an arrangement:

- Explicit computations for interesting examples [2, 11, 20, 36].
- Upper/lower bounds of monodromy eigenspaces [3, 5, 25, 35, 36].
- Vanishing of non-trivial monodromy eigenspaces [4, 29, 30].
- Examples of arrangements with multiplicities whose first homology group of the Milnor fiber has torsion [6].
- Examples of arrangements whose higher degree homology of the Milnor fiber has torsion [9].
- Purely combinatorial description of monodromy eigenspaces for line arrangements which have only double and triple intersections. [10, 18, 26].

The purpose of this paper is to study the mod 2 homology and 2-torsion in the integral homology of covering spaces of arrangement complements. The main results of this paper are as follows. The first result is concerning mod 2 homology of the double covering.

**Theorem 1.1.** *(See Theorem 3.7 for more precise statement.) The mod 2 Betti numbers of double covers of arrangement complements are combinatorially determined.*

More precisely, we will obtain a combinatorial expression of the rank of mod 2 homology in terms of the mod 2 Aomoto complex.

The second result is about the Milnor fiber of the icosidodecahedral arrangement which is the arrangement of 16 planes in  $\mathbb{R}^3$  associated to the icosidodecahedron. (See §4.1 for definition).

**Theorem 1.2.** (See Theorem 4.3 (3).) *Let  $F_{\mathcal{A}_{\mathcal{ID}}}$  be the Milnor fiber of the icosidodecahedral arrangement  $\mathcal{A}_{\mathcal{ID}}$ . Then  $H_1(F_{\mathcal{A}_{\mathcal{ID}}}, \mathbb{Z})$  has 2-torsion.*

The organization of this paper is as follows. In §2, we review several results concerning Milnor fiber of arrangements, intersection poset, Orlik-Solomon algebra, and Aomoto complex, which are necessary for stating results and proofs.

§3 summarizes results on abelian covering spaces. After recalling the transfer long exact sequence of a double covering in §3.1-§3.2, we prove, in §3.3, a key lemma (Lemma 3.2) which guarantees the first mod 2 Betti number does not decrease by taking certain (“ $\mathbb{Z}_4$ -liftable”) double coverings. In §3.4, we prove the first main result, that is a combinatorial formula for the mod 2 Betti numbers of double coverings of an arrangement complement (Theorem 3.7).

In §4.1, we introduce the icosidodecahedral arrangement  $\mathcal{A}_{\mathcal{ID}}$ , and prove in §4.2 and §4.3 that the first homology of the Milnor fiber has 2-torsion.

**Notation and Convention.** We will use the following convention throughout the paper.

- In this paper,  $\mathbb{Z}_n$  denotes the cyclic group (also a finite ring)  $\mathbb{Z}/n\mathbb{Z}$ .
- For a space  $X$  with the homotopy type of a finite CW complex,  $b_i(X) = \text{rank}_{\mathbb{Z}} H^i(X, \mathbb{Z})$  denotes the Betti number, and  $\bar{b}_i(X) = \text{rank}_{\mathbb{Z}_2} H^i(X, \mathbb{Z}_2)$  denotes the mod 2 Betti number.
- A covering  $Y \rightarrow X$  always means an unbranched covering. Unless otherwise stated, we assume  $X$  and  $Y$  are connected.
- We will assume that the base point  $x_0 \in X$  is fixed when we consider the fundamental group  $\pi_1(X) = \pi_1(X, x_0)$ .
- For an element  $\gamma \in \pi_1(X)$ , denote by  $[\gamma]$  its homology class. If  $(Y, y_0) \rightarrow (X, x_0)$  is a covering,  $\tilde{\gamma}$  denotes the lifting of  $\gamma$  starting from the base point  $y_0$ . Note that  $\tilde{\gamma}$  is not necessarily a closed curve.

## 2. GENERALITIES ON HYPERPLANE ARRANGEMENTS

In this section, we summarize notation and several results on hyperplane arrangements. See [23, 24] for more details.

Let  $\mathcal{A} = \{H_1, H_2, \dots, H_n\}$  be a collection of affine hyperplanes in  $\mathbb{C}^\ell$ . We denote by  $M(\mathcal{A}) = \mathbb{C}^\ell \setminus \bigcup_{i=1}^n H_i$  the complement. Let  $\overline{H_\infty}$  be a projective hyperplane in  $\mathbb{C}\mathbb{P}^\ell$ . Using the identification  $\mathbb{C}\mathbb{P}^\ell = \mathbb{C}^\ell \sqcup \overline{H_\infty}$ ,  $\mathcal{A}$  naturally induces an arrangement of  $n + 1$  projective hyperplanes

$\overline{\mathcal{A}} = \{\overline{H_1}, \dots, \overline{H_n}, \overline{H_\infty}\}$ . A projective hyperplane  $\overline{H_i}$  defines a linear hyperplane  $\widetilde{H_i}$  in  $\mathbb{C}^{\ell+1}$ . The collection of these linear hyperplanes  $\widetilde{\mathcal{A}} = \{\widetilde{H_1}, \dots, \widetilde{H_n}, \widetilde{H_\infty}\}$  is called the coning of  $\mathcal{A}$  (which is also denoted by  $c\mathcal{A}$ ). We can also define the opposite operation, which is called the deconing of  $\widetilde{\mathcal{A}}$  with respect to  $\widetilde{H_\infty}$  and denoted by  $\mathcal{A} = d_{\widetilde{H_\infty}} \widetilde{\mathcal{A}}$ .

There is a natural projection  $p : M(c\mathcal{A}) \rightarrow M(\mathcal{A})$ . It is easily seen that  $M(c\mathcal{A}) \simeq M(\mathcal{A}) \times \mathbb{C}^\times$ . Let  $\alpha_i : \mathbb{C}^{\ell+1} \rightarrow \mathbb{C}$  be a non-zero linear form such that  $\widetilde{H_i} = \alpha_i^{-1}(0)$  ( $i = 1, \dots, n, \infty$ ). Then  $Q = \prod_i \alpha_i$  is a homogeneous polynomial of degree  $n+1$ .  $F := \{x \in \mathbb{C}^{\ell+1} \mid Q(x) = 1\} \subset \mathbb{C}^{\ell+1}$  is called the Milnor fiber of  $c\mathcal{A}$ . Since  $Q$  is homogeneous, the cyclic group  $\mathbb{Z}/(n+1)\mathbb{Z} \simeq \{\zeta \in \mathbb{C} \mid \zeta^{n+1} = 1\}$  acts on  $F$  defined by the map  $\mu : F \rightarrow F, x \mapsto \zeta \cdot x$ . This action is nothing but the monodromy action of the fibration  $Q : M(c\mathcal{A}) \rightarrow \mathbb{C}^\times$ . The monodromy map  $\mu : F \rightarrow F$  induces a linear map on homology  $\mu_* : H_k(F, \mathbb{Z}) \rightarrow H_k(F, \mathbb{Z})$ . Since the map has finite order, the homology with coefficients in  $\mathbb{C}$  is decomposed into the direct sum of eigenspaces,

$$H_k(F, \mathbb{C}) = \bigoplus_{\zeta^{n+1}=1} H_k(F, \mathbb{C})_\zeta,$$

where  $H_k(F, \mathbb{C})_\zeta$  is the  $\zeta$ -eigenspace of  $\mu_*$ . Since  $M(\mathcal{A})$  can be identified with the quotient  $F/\langle \mu \rangle$ , we have  $H_k(F, \mathbb{C})_1 \simeq H_k(M(\mathcal{A}), \mathbb{C})$ .

Given an affine arrangement  $\mathcal{A} = \{H_1, \dots, H_n\}$ , non-empty intersections  $X = \bigcap_{H \in \mathcal{S}} H$  ( $\mathcal{S} \subset \mathcal{A}$ ) form a poset with respect to reverse inclusion, which is denoted by  $L(\mathcal{A})$  and called the intersection poset. We say that  $\mathcal{S} \subset \mathcal{A}$  does not intersect if  $\bigcap_{H \in \mathcal{S}} H = \emptyset$ . A subset  $\mathcal{S} \subset \mathcal{A}$  is called dependent if  $\bigcap_{H \in \mathcal{S}} H \neq \emptyset$  and  $\text{codim} \bigcap_{H \in \mathcal{S}} H < \#\mathcal{S}$ . For  $X \in L(\mathcal{A})$ , we denote by  $\mathcal{A}_X := \{H \in \mathcal{A} \mid H \supset X\}$  the localization of  $\mathcal{A}$  at  $X$ .

From the intersection poset, the Orlik-Solomon algebra  $A_{\mathbb{Z}}^*(\mathcal{A})$  is defined as follows. Let  $E = \bigoplus_{i=1}^n \mathbb{Z}e_i$  be the free abelian group generated by the symbols  $e_i$  corresponding to the hyperplanes  $H_i$ . Let  $\wedge E = \mathbb{Z} \oplus E \oplus E^{\wedge 2} \oplus \dots \oplus E^{\wedge n}$  be the exterior algebra on  $E$ . For given  $\mathcal{S} = \{i_1, i_2, \dots, i_k\} \subset \mathcal{A}$  with  $i_1 < \dots < i_k$ , let  $e_{\mathcal{S}} := e_{i_1} \wedge \dots \wedge e_{i_k}$ , and define  $\partial e_{\mathcal{S}} \in E^{\wedge(k-1)}$  by

$$\partial e_{\mathcal{S}} = \sum_{p=1}^k (-1)^{p-1} e_{i_1} \wedge \dots \widehat{e_{i_p}} \wedge \dots \wedge e_{i_k}.$$

Define  $I(\mathcal{A})$  to be the ideal of  $\wedge E$  generated by the following elements.

$$\{\partial e_{\mathcal{S}} \mid \mathcal{S} \text{ is dependent}\} \cup \{e_{\mathcal{S}} \mid \mathcal{S} \text{ does not intersect}\}.$$

The quotient ring  $A_{\mathbb{Z}}^*(\mathcal{A}) := \wedge E / I(\mathcal{A})$  is called the Orlik-Solomon algebra. For any abelian group  $R$ , denote  $A_R^*(\mathcal{A}) := A_{\mathbb{Z}}^*(\mathcal{A}) \otimes_{\mathbb{Z}} R$ . Note that when  $R$  is a commutative ring,  $A_R^*(\mathcal{A})$  becomes a graded commutative  $R$ -algebra.

Brieskorn's Lemma [23, Corollary 3.73] shows that the degree  $k$  component of the Orlik-Solomon algebra is

$$A_R^k(\mathcal{A}) \simeq \bigoplus_{\substack{X \in L(\mathcal{A}) \\ \text{codim } X = k}} A_R^k(\mathcal{A}_X).$$

Orlik and Solomon [22] proved that each homology group  $H_k(M(\mathcal{A}), \mathbb{Z})$  is torsion free and the cohomology ring is isomorphic to the Orlik-Solomon algebra. Namely, for any commutative ring  $R$ , we have an isomorphism

$$H^*(M(\mathcal{A}), R) \simeq A_R^*(\mathcal{A})$$

as graded commutative  $R$ -algebras.

The following will be used later.

**Lemma 2.1.** *Let  $\mathcal{A} = \{H_1, \dots, H_n\}$  be an arrangement of  $n$  lines ( $n \geq 3$ ) in  $\mathbb{C}^2$  with unique intersection (Figure 1). Let  $R$  be an integral domain. Let  $\eta = \sum_{i=1}^n a_i e_i$  and  $\omega = \sum_{i=1}^n b_i e_i \in A_R^1(\mathcal{A})$ . Then the following are equivalent.*

- (a)  $\eta \wedge \omega = 0$ .
- (b)  $\sum_{i=1}^n a_i = \sum_{i=1}^n b_i = 0$  or  $\omega, \eta$  are linearly dependent (i. e., there are  $c_1, c_2 \in R$ , not both zero, such that  $c_1 \eta + c_2 \omega = 0$ ).

*Proof.* See [38, Proposition 2.1] (or [34, Lemma 3.1]). □




FIGURE 1.  $n$  lines with one intersection ( $n \geq 3$ )

For a given  $\omega \in A_R^1(\mathcal{A})$ , Orlik-Solomon algebra determines the cochain complex  $(\mathcal{A}_R^\bullet(\mathcal{A}), \omega \wedge -)$ , which is called the Aomoto complex. This complex plays crucial role in the computation of twisted cohomology groups [1, 12, 13, 19, 25, 31, 34, 38].

### 3. FINITE COVERINGS AND COMBINATORIAL STRUCTURES

**3.1. Finite abelian covering.** Let  $G$  be a finite abelian group. Recall that a group homomorphism  $w : \pi_1(X) \rightarrow G$  determines a finite abelian covering  $p(w) : X^w \rightarrow X$ . Since  $G$  is abelian, we have

$$\text{Hom}(\pi_1(X), G) = \text{Hom}(H_1(X, \mathbb{Z}), G) \simeq H^1(X, G).$$

The element  $w \in H^1(X, G)$  is called the characteristic class of the covering. Note that  $\text{Ker}(w : \pi_1(X) \rightarrow G)$  is isomorphic to  $\pi_1(X^w)$  and  $\pi_1(X)/\pi_1(X^w) \simeq \text{Im}(w : \pi_1(X) \rightarrow G)$ .

**3.2. Double covering and transfer exact sequence.** (See [15, §4.3] for details.) Now we consider the double covering  $p : Y \rightarrow X$  with the unique nontrivial deck transformation  $\sigma : Y \rightarrow Y$ . Assume that  $Y$  is connected. Denote the characteristic class by  $w \in H^1(X, \mathbb{Z}_2) = \text{Hom}(\pi_1(X), \mathbb{Z}_2)$ . Recall that the transfer map

$$\tau_* : H_k(X, \mathbb{Z}_2) \longrightarrow H_k(Y, \mathbb{Z}_2),$$

is given by  $[\gamma] \mapsto [p^{-1}(\gamma)]$ . Denote by  $\tau^* : H^k(Y, \mathbb{Z}_2) \rightarrow H^k(X, \mathbb{Z}_2)$  the induced map between cohomology groups.

**Example 3.1.** Let us closely look at the transfer map  $\tau_*$  in degree one. Let  $\gamma \in \pi_1(X)$ . Then either  $w(\gamma) = 0$  or  $\neq 0$ .

- (1) Suppose  $w(\gamma) \neq 0$ , equivalently,  $\gamma \notin \pi_1(Y) (\subset \pi_1(X))$ . Then the lift  $\tilde{\gamma}$  is no longer a cycle. Note that  $p^{-1}(\gamma)$  is the union of  $\tilde{\gamma}$  and  $\sigma(\tilde{\gamma})$ , which is the lift of  $\gamma^2$ . Since  $\gamma^2 \in \pi_1(Y) \subset \pi_1(X)$ , we have  $\tau_*([\gamma]) = [\gamma^2]$ .
- (2) Suppose  $w(\gamma) = 0$ . Then the lift  $\tilde{\gamma}$  is a cycle on  $Y$ . Hence  $\tau_*([\gamma]) = [\tilde{\gamma}] + [\sigma(\tilde{\gamma})]$ .

The transfer map fits into the following long exact sequence.

$$(3.1) \quad \cdots H^{k-1}(X, \mathbb{Z}_2) \xrightarrow{w \cup} H^k(X, \mathbb{Z}_2) \xrightarrow{p^*} H^k(Y, \mathbb{Z}_2) \xrightarrow{\tau^*} H^k(X, \mathbb{Z}_2) \xrightarrow{w \cup} H^{k+1}(X, \mathbb{Z}_2) \cdots$$

**3.3. Mod 2 Betti number of  $\mathbb{Z}_4$ -liftable double coverings.** In this section, we prove an inequality between the mod 2 first Betti numbers of a double covering  $Y \rightarrow X$ , which will play a crucial role later in §4.3.

**Lemma 3.2.** *Let  $w : \pi_1(X) \rightarrow \mathbb{Z}_4$  be a surjective homomorphism. By composing the canonical surjective homomorphism  $\mathbb{Z}_4 \rightarrow \mathbb{Z}_2$ , we obtain an epimorphism  $\bar{w} : \pi_1(X) \rightarrow \mathbb{Z}_2$ . Consider the associated double coverings  $X^w \rightarrow X^{\bar{w}}$  and  $X^{\bar{w}} \rightarrow X$ .*

- (1)  $\bar{w} \in H^1(X, \mathbb{Z}_2)$  satisfies  $\bar{w} \cup \bar{w} = 0$ .
- (2)  $\bar{b}_1(X^{\bar{w}}) \geq \bar{b}_1(X)$ .

*Proof.* (1) Recall that the exact sequence of abelian groups  $0 \rightarrow \mathbb{Z}_2 \rightarrow \mathbb{Z}_4 \rightarrow \mathbb{Z}_2 \rightarrow 0$  induces the exact sequence

$$H^1(X, \mathbb{Z}_4) \xrightarrow{\varphi} H^1(X, \mathbb{Z}_2) \xrightarrow{\beta} H^2(X, \mathbb{Z}_2).$$

The first map  $\varphi$  sends  $w$  to  $\bar{w}$ . The second map  $\beta : H^1(X, \mathbb{Z}_2) \rightarrow H^2(X, \mathbb{Z}_2)$  is the so-called Bockstein homomorphism, which is  $\beta(x) = x \cup x$  [14, Section 4.L]. Since  $\beta \circ \varphi = 0$ , we have  $\bar{w} \cup \bar{w} = 0$ .

(2) Since  $p^* : H^0(X, \mathbb{Z}_2) \simeq H^0(X^{\bar{w}}, \mathbb{Z}_2)$  is an isomorphism, the beginning of the transfer exact sequence is as follows.

$$0 \rightarrow H^0(X, \mathbb{Z}_2) \xrightarrow{\bar{w} \cup} H^1(X, \mathbb{Z}_2) \xrightarrow{p^*} H^1(X^{\bar{w}}, \mathbb{Z}_2) \xrightarrow{\tau^*} H^1(X, \mathbb{Z}_2) \xrightarrow{\bar{w} \cup} H^2(X, \mathbb{Z}_2).$$

We have

$$\text{rank}_{\mathbb{Z}_2} H^1(X^{\bar{w}}, \mathbb{Z}_2) = \text{rank}_{\mathbb{Z}_2} H^1(X, \mathbb{Z}_2) - 1 + \text{rank}_{\mathbb{Z}_2} \text{Im}(\tau^*).$$

Since  $\bar{w} \in \text{Ker}(\bar{w} \cup) = \text{Im}(\tau^*)$ ,  $\text{rank}_{\mathbb{Z}_2} \text{Im}(\tau^*) \geq 1$ . This completes the proof.  $\square$

*Remark 3.3.* Without the assumption of  $\mathbb{Z}_4$ -liftability in Lemma 3.2 (2), the inequality between mod 2 Betti numbers does not hold in general. For example, the double covering  $S^m \rightarrow \mathbb{R}\mathbb{P}^m$  does not satisfy the inequality for  $m \geq 2$ . Indeed  $\text{rank}_{\mathbb{Z}_2} H^1(S^m, \mathbb{Z}_2) = 0$  and  $\text{rank}_{\mathbb{Z}_2} H^1(\mathbb{R}\mathbb{P}^m, \mathbb{Z}_2) = 1$ .

**Corollary 3.4.** (1) *Let  $X_k \rightarrow X_{k-1} \rightarrow \cdots \rightarrow X_0$  ( $k \geq 2$ ) be a tower of double coverings of connected spaces (i.e., each  $X_i \rightarrow X_{i-1}$  is a double covering) such that  $X_k \rightarrow X_0$  is a cyclic  $\mathbb{Z}_{2^k}$ -covering. Then*

$$\bar{b}_1(X_{k-1}) \geq \bar{b}_1(X_{k-2}) \geq \cdots \geq \bar{b}_1(X_1) \geq \bar{b}_1(X_0).$$

(2) *Let  $w : \pi_1(X) \rightarrow \mathbb{Z}$  be a surjective homomorphism, and  $\bar{w} : \pi_1(X) \rightarrow \mathbb{Z}_{2^k}$  be the induced surjection ( $k \geq 1$ ). Then*

$$\bar{b}_1(X^{\bar{w}}) \geq \bar{b}_1(X).$$

*Proof.* (1) is proved by induction on  $k$  using Lemma 3.2. (2) follows immediately from (1).  $\square$

**3.4. Double coverings of arrangement complements.** Now let us formulate a problem asking whether the Betti numbers of finite coverings of arrangement complements are combinatorially determined.

**Problem 3.5.** *Let  $\mathcal{A}$  be an arrangement in  $\mathbb{C}^\ell$ . Let  $G$  be a finite abelian group, and  $R$  be an abelian group. Let  $w \in A_G^1(\mathcal{A}) \simeq \text{Hom}(\pi_1(M(\mathcal{A})), G)$ . Describe the cohomology groups  $H^k(M(\mathcal{A})^w, R)$  (or their ranks) in terms of  $L(\mathcal{A})$  and  $w \in A_G^1(\mathcal{A})$ .*

**Example 3.6.** Let  $\mathcal{A} = \{H_1, \dots, H_n\}$  be an arrangement of  $n$  hyperplanes and  $G = \mathbb{Z}_{n+1}$  be the cyclic group of order  $(n+1)$ . Let  $w := e_1 + e_2 + \cdots + e_n \in A_{\mathbb{Z}_{n+1}}^1(\mathcal{A})$ . Then the corresponding covering  $M(\mathcal{A})^w$  is homeomorphic to the Milnor fiber  $F$  of the coning  $c\mathcal{A}$ .

Problem 3.5 is widely open. Indeed, many research problems are related to Problem 3.5 [8, 16, 17, 28]. For example, the notion of characteristic variety is deeply related to the computation of  $\text{rank}_{\mathbb{Z}} H^k(M(\mathcal{A})^w, \mathbb{Z})$ . See [32, 33] for surveys of the topic.



As a special case of Problem 3.5, we now prove that the mod 2 Betti numbers of the double covering  $M(\mathcal{A})^w$  of an arrangement complement are combinatorially determined.

**Theorem 3.7.** *Let  $w \in A_{\mathbb{Z}_2}^1(\mathcal{A})$ ,  $w \neq 0$ . Then the  $k$ -th mod 2 Betti number ( $k \geq 0$ ) of the double covering  $M(\mathcal{A})^w$  is expressed as follows.*

$$(3.2) \quad \overline{b}_k(M(\mathcal{A})^w) = b_k(M(\mathcal{A})) + \text{rank}_{\mathbb{Z}_2} H^k(A_{\mathbb{Z}_2}^\bullet(\mathcal{A}), w \wedge -).$$

*Proof.* Denote  $\mathcal{Z}^k := \text{Ker}(w \wedge : A_{\mathbb{Z}_2}^k(\mathcal{A}) \rightarrow A_{\mathbb{Z}_2}^{k+1}(\mathcal{A}))$  and  $\mathcal{B}^k := \text{Im}(w \wedge : A_{\mathbb{Z}_2}^{k-1}(\mathcal{A}) \rightarrow A_{\mathbb{Z}_2}^k(\mathcal{A}))$ . From the transfer long exact sequence, we have the following exact sequence.

$$0 \longrightarrow \mathcal{B}^k \longrightarrow A_{\mathbb{Z}_2}^k(\mathcal{A}) \longrightarrow H^k(M(\mathcal{A})^w, \mathbb{Z}_2) \longrightarrow \mathcal{Z}^k \longrightarrow 0.$$

Since  $\mathcal{Z}^k / \mathcal{B}^k \simeq H^k(A_{\mathbb{Z}_2}^\bullet(\mathcal{A}), w \wedge -)$ ,

$$\begin{aligned} \text{rank}_{\mathbb{Z}_2} H^k(M(\mathcal{A})^w, \mathbb{Z}_2) &= \text{rank}_{\mathbb{Z}_2} A_{\mathbb{Z}_2}^k(\mathcal{A}) + \text{rank}_{\mathbb{Z}_2} \mathcal{Z}^k - \text{rank}_{\mathbb{Z}_2} \mathcal{B}^k \\ &= b_k(M(\mathcal{A})) + \text{rank}_{\mathbb{Z}_2} H^k(A_{\mathbb{Z}_2}^\bullet(\mathcal{A}), w \wedge -). \end{aligned}$$

□

*Remark 3.8.* Note that Theorem 3.7 gives only mod 2 Betti numbers. It is not clear whether we can combinatorially describe Betti numbers or (mod 2) cohomology ring structure of the double covering.

## 4. 2-TORSIONS IN MILNOR FIBER HOMOLOGY

### 4.1. Icosidodecahedral arrangement $\mathcal{A}_{\mathcal{ID}}$ .

**Definition 4.1.** The icosidodecahedral arrangement  $\mathcal{A}_{\mathcal{ID}}$  is the coning of the 15 affine lines in Figure 2. Namely,  $\mathcal{A}_{\mathcal{ID}}$  consists of 16 planes in  $\mathbb{R}^3$ . (Another deconing of  $\mathcal{A}_{\mathcal{ID}}$  is depicted in Figure 3.)

Let us briefly comment on the naming ‘‘Icosidodecahedral arrangement’’. Actually,  $\mathcal{A}_{\mathcal{ID}}$  can be constructed from the icosidodecahedron as follows, which seems to be the most symmetric realization of  $\mathcal{A}_{\mathcal{ID}}$ .

The icosidodecahedron (Figure 4) is a polyhedron which is commonly obtained as vertex truncations of the icosahedron and the dodecahedron (by truncating mid points of edges). An icosidodecahedron has 32 faces (20 triangles and 12 pentagons), 60 edges and 30 vertices (Figure 4). We can choose 10 edges to form the equator of the polyhedron, which are lying on a plane in  $\mathbb{R}^3$ . Similarly, we obtain 6 planes in total (they correspond to the blue lines  $H_{11}, \dots, H_{15}, H_{16}$  in Figures 2 and 3).

Each pentagonal face of the icosidodecahedron has five diagonals. There are 60 such diagonals in all. If we choose appropriately consecutive six of them, they are lying on a plane (the red diagonals in Figure 4). We obtain




FIGURE 2. A decoupling  $d_{H_{16}}\mathcal{A}_{\mathcal{ID}}$  of the icosidodecahedral arrangement (with respect to a blue line at infinity  $H_{16}$ ). The coloring expresses the nontrivial cocycle in the mod 2 Ao-moto complex (§4.2).




FIGURE 3. Another decoupling  $d_{H_1}\mathcal{A}_{\mathcal{ID}}$  of the icosidodecahedral arrangement (with respect to a red line  $H_1$  at infinity)

in this manner 10 planes consisting of diagonals (Figure 5). As a result, we have an arrangement of 16 planes in  $\mathbb{R}^3$ , which is isomorphic to  $\mathcal{A}_{\mathcal{ID}}$ .

For computations, we use mainly the deconing from Figure 2.




FIGURE 4. Icosidodecahedron (blue) and a diagonal plane (red)




FIGURE 5. Icosidodecahedral arrangement

**4.2. Mod 2 Aomoto complex of icosidodecahedral arrangement.** Let  $d\mathcal{A}_{\mathcal{ID}} := d_{H_{16}}\mathcal{A}_{\mathcal{ID}}$  be the affine arrangement in Figure 2. Let  $w_2 := e_1 + \cdots + e_{15} \in A_{\mathbb{Z}_2}^1(d\mathcal{A}_{\mathcal{ID}})$ . For a subset  $\mathcal{S} \subset d\mathcal{A}_{\mathcal{ID}}$ , let  $e_{\mathcal{S}} := \sum_{H_i \in \mathcal{S}} e_i$ . Obviously, every element in  $A_{\mathbb{Z}_2}^1(d\mathcal{A}_{\mathcal{ID}})$  can be expressed as  $e_{\mathcal{S}}$ , where  $\mathcal{S}$  is a subset of  $d\mathcal{A}_{\mathcal{ID}}$ .

**Proposition 4.2.**  $\text{rank}_{\mathbb{Z}_2} H^1(A_{\mathbb{Z}_2}^\bullet(d\mathcal{A}_{\mathcal{ID}}), w_2 \wedge -) = 1$ .

*Proof.* Let  $\mathcal{S}_0 := \{1, 2, \dots, 10\}$  (red in Figure 2) and  $\mathcal{S}_1 := \{11, 12, 13, 14, 15\}$  (blue in Figure 2). Note that at each intersection  $p$ , the localization  $(d\mathcal{A}_{\mathcal{ID}})_p$  consists of either

- two lines from  $\mathcal{S}_0$ , or
- two from  $\mathcal{S}_0$  and two from  $\mathcal{S}_1$ .

This property, together with Lemma 2.1, enables us to conclude

$$w_2 \wedge e_{\mathcal{S}_0} = w_2 \wedge e_{\mathcal{S}_1} = 0.$$

Therefore,  $\text{rank}_{\mathbb{Z}_2} H^1(A_{\mathbb{Z}_2}^\bullet(d\mathcal{A}_{TD}), w_2 \wedge -) \geq 1$ .

Next we show that  $[e_{\mathcal{S}_0}] = [e_{\mathcal{S}_1}]$  is the unique nonzero cohomology class. Suppose  $w_2 \wedge e_{\mathcal{S}} = 0$  for some  $\mathcal{S} \subset d\mathcal{A}_{TD}$ . If  $\mathcal{S} \cap \mathcal{S}_0 \neq \emptyset$ , choose an  $i \in \mathcal{S} \cap \mathcal{S}_0$ , then by Lemma 2.1, all  $j$  such that  $i$  and  $j$  intersect at a double point must be contained in  $\mathcal{S}$ . Thus, if  $\mathcal{S} \cap \mathcal{S}_0 \neq \emptyset$ ,  $\mathcal{S} \supset \mathcal{S}_0$ . By replacing  $e_{\mathcal{S}}$  by  $e_{\mathcal{S}} + w_2 = e_{d\mathcal{A}_{TD} \setminus \mathcal{S}}$ , we may assume  $\mathcal{S} \cap \mathcal{S}_0 = \emptyset$ , in other words,  $\mathcal{S} \subset \mathcal{S}_1$ . Again by Lemma 2.1,  $\mathcal{S}$  must be either  $\mathcal{S}_1$  or  $\emptyset$ . This completes the proof.  $\square$

### 4.3. Milnor fiber of icosidodecahedral arrangement.

**Theorem 4.3.** *Let  $F_{\mathcal{A}_{TD}}$  be the Milnor fiber of the icosidodecahedral arrangement  $\mathcal{A}_{TD}$ . Then,*

- (1)  $\text{rank}_{\mathbb{Z}} H^1(F_{\mathcal{A}_{TD}}, \mathbb{Z}) = 15$ .
- (2)  $\text{rank}_{\mathbb{Z}_2} H^1(F_{\mathcal{A}_{TD}}, \mathbb{Z}_2) \geq 16$ .
- (3) *The integral first homology group  $H_1(F_{\mathcal{A}_{TD}}, \mathbb{Z})$  has 2-torsion.*

*Proof.* (1) Recall that  $H_1(F_{\mathcal{A}_{TD}}, \mathbb{Z})$  admits  $\mathbb{Z}_{16}$  action by monodromy. Then the homology with complex coefficients has eigenspace decomposition

$$H^1(F_{\mathcal{A}_{TD}}, \mathbb{C}) = \bigoplus_{\lambda^{16}=1} H^1(F_{\mathcal{A}_{TD}}, \mathbb{C})_{\lambda}.$$

Each eigenspace  $H^1(F_{\mathcal{A}_{TD}}, \mathbb{C})_{\lambda}$  is known to be isomorphic to  $H^1(M(d\mathcal{A}_{TD}), \mathcal{L}_{\lambda})$  [7], where  $\mathcal{L}_{\lambda}$  is the complex rank one local system on  $M(d\mathcal{A}_{TD})$  which has monodromy  $\lambda$  along the meridian of each line  $H \in d\mathcal{A}_{TD}$  in  $\mathbb{C}^2$ . In particular, the 1-eigen space is  $H^1(F_{\mathcal{A}_{TD}}, \mathbb{C})_1 \simeq H^1(M(d\mathcal{A}_{TD}), \mathbb{C}) \simeq \mathbb{C}^{15}$ . For  $\lambda \neq 1$ , there are several practical ways to check that  $H^1(F_{\mathcal{A}_{TD}}, \mathbb{C})_{\lambda} = 0$ . One of the methods is to apply the result by Esnault-Schechtman-Viehweg [12] and Schechtman-Varchenko-Terao [31]. Let  $a_1, \dots, a_{16} \in \mathbb{C}$ . Assume the following three conditions.

- (P1)  $\exp(2\pi\sqrt{-1} \cdot a_i) = \lambda$ ,
- (P2)  $\sum_{i=1}^{16} a_i = 0$ .
- (P3) Let  $\overline{\mathcal{A}_{TD}}$  be the induced projective arrangement of 16 lines. For each quadruple intersection  $p \in \mathbb{CP}^2$  of  $\overline{\mathcal{A}_{TD}}$ , the sum  $\sum_{H_i \ni p} a_i$  is not contained in  $\mathbb{Z}_{>0}$ .

(In some literature, such a local system is called admissible [21].) Then [12, 31] asserts that

$$(4.1) \quad H^k(M(d\mathcal{A}_{TD}), \mathcal{L}_{\lambda}) \simeq H^k(A_{\mathbb{C}}^\bullet(d\mathcal{A}_{TD}), \eta \wedge -),$$

for  $k \geq 0$ , where  $\eta = \sum_{i=1}^{15} a_i e_i \in \mathcal{A}_{\mathbb{C}}^1(d\mathcal{A}_{\mathcal{ID}})$ .

Here let us illustrate how to prove  $H^1(M(d\mathcal{A}_{\mathcal{ID}}), \mathcal{L}_\lambda) = 0$  for  $\lambda = -1$ . We can choose  $a_1, \dots, a_{16}$  as

$$\begin{aligned} a_1 &= a_2 = \cdots = a_9 = a_{10} = \frac{1}{2}, \\ a_{11} &= a_{12} = \cdots = a_{15} = -\frac{1}{2} \\ a_{16} &= -\frac{5}{2}. \end{aligned}$$

Then at each quadruple point, the sum of  $a_i$ 's is either 0 or  $-2$ . Thus the conditions (P1), (P2) and (P3) are verified. Set  $\eta := \sum_{i=1}^{15} a_i e_i \in \mathcal{A}_{\mathbb{C}}^1(d\mathcal{A}_{\mathcal{ID}})$ . We can check  $H^k(A_{\mathbb{C}}^\bullet(d\mathcal{A}_{\mathcal{ID}}), \eta \wedge -) = 0$  by arguments similar to the proof of Proposition 4.2. The vanishing of the eigenspaces for other eigenvalues  $\lambda$  can be proved in a similar way.

We can obtain the same result also by using resonant band algorithms formulated in [36, 37].

(2) follows from Theorem 3.7, Proposition 4.2 and Corollary 3.4.

(3) By universal coefficient theorem and (1), if  $H_1(F_{\mathcal{A}_{\mathcal{ID}}}, \mathbb{Z})$  does not have 2-torsion,  $H_1(F_{\mathcal{A}_{\mathcal{ID}}}, \mathbb{Z}_2) \simeq H_1(F_{\mathcal{A}_{\mathcal{ID}}}, \mathbb{Z}) \otimes \mathbb{Z}_2$  has rank 15 over  $\mathbb{Z}_2$ . This contradicts (2).  $\square$

*Remark 4.4.* The proof of Theorem 4.3 works more generally. Let  $\mathcal{A}$  be an arrangement in  $\mathbb{C}^3$ . Suppose

- $\#\mathcal{A}$  is a power of 2.
- The first cohomology of the mod 2 Aomoto complex of the deconing of  $\mathcal{A}$  does not vanish.
- The first cohomology of the Milnor fiber  $H^1(F_{\mathcal{A}}, \mathbb{C})$  does not have nontrivial monodromy eigenspaces.

Then we can conclude  $H_1(F_{\mathcal{A}}, \mathbb{Z})$  has 2-torsion.

*Remark 4.5.* Proposition 4.2 and Theorem 4.3 (1) show that  $\mathcal{A}_{\mathcal{ID}}$  is a counterexample to a conjectures in [26, Conjecture 1.9]. More precisely, the equality  $\beta_2 = e_2$  in [26] does not hold for  $\mathcal{A}_{\mathcal{ID}}$ .

*Remark 4.6.* Enrique Artal-Bartolo communicated to us that he checked by computer that  $H_1(F_{\mathcal{A}_{\mathcal{ID}}}, \mathbb{Z}) \simeq \mathbb{Z}^{15} \oplus \mathbb{Z}_2$ . It would be a challenging problem to develop a method which can check the result theoretically. The following are also interesting problems.

- (1) Describe the mod 2 Betti numbers of the Milnor fibers of arrangements.
- (2) Describe the mod 2 cohomology rings of double covers of arrangement complements.

## REFERENCES

- [1] K. Aomoto, M. Kita, Theory of hypergeometric functions. With an appendix by Toshitake Kohno. Translated from the Japanese by Kenji Iohara. Springer Monographs in Mathematics. Springer-Verlag, Tokyo, 2011. xvi+317 pp
- [2] P. Bailet, A. Dimca, M. Yoshinaga, A vanishing result for the first twisted cohomology of affine varieties and applications to line arrangements. *Manuscripta Math.* **157** (2018), no. 3-4, 497-511.
- [3] P. Bailet, S. Settepanella, Homology graph of real arrangements and monodromy of Milnor fiber. *Adv. in Appl. Math.* **90** (2017), 46-85.
- [4] P. Bailet, M. Yoshinaga, Degeneration of Orlik-Solomon algebras and Milnor fibers of complex line arrangements. *Geom. Dedicata* **175** (2015), 49-56.
- [5] N. Budur, A. Dimca, M. Saito, First Milnor cohomology of hyperplane arrangements. *Topology of algebraic varieties and singularities*, 279-292, Contemp. Math., 538, Amer. Math. Soc., Providence, RI, 2011.
- [6] D. C. Cohen, G. Denham, A. Suciu, Torsion in Milnor fiber homology. *Algebr. Geom. Topol.* **3** (2003), 511-535.
- [7] D. C. Cohen, A. Suciu, On Milnor fibrations of arrangements. *J. London Math. Soc.* **51** (1995), no. 2, 105-119.
- [8] D. C. Cohen, A. Suciu, Characteristic varieties of arrangements. *Math. Proc. Cambridge Philos. Soc.* **127** (1999), no. 1, 33-53.
- [9] G. Denham, A. Suciu, Multinets, parallel connections, and Milnor fibrations of arrangements. *Proceedings of the London Mathematical Society* **108** (2014) no. 6, 1435-1470.
- [10] A. Dimca, Monodromy of triple point line arrangements. *Singularities in geometry and topology 2011*, 7180, Adv. Stud. Pure Math., 66, Math. Soc. Japan, Tokyo, 2015.
- [11] A. Dimca, G. Sticlaru, On the Milnor monodromy of the exceptional reflection arrangement of type  $G_{31}$ . *Doc. Math.* **23** (2018), 1-14.
- [12] Esnault, Schechtman, Viehweg, H. Esnault, V. Schechtman, and E. Viehweg, Cohomology of local systems of the complement of hyperplanes. *Invent. Math.* **109** (1992), 557-561
- [13] M. Falk, Arrangements and cohomology. *Ann. Comb.* **1** (1997), no. 2, 135-157.
- [14] A. Hatcher, Algebraic topology. Cambridge University Press, Cambridge, 2002. xii+544 pp.
- [15] J. -C. Hausmann, Mod two homology and cohomology. Universitext. Springer, Cham, 2014. x+535 pp.
- [16] E. Hironaka, Abelian coverings of the complex projective plane branched along configurations of real lines. *Mem. Amer. Math. Soc.* **105** (1993), no. 502, vi+85 pp.
- [17] A. Libgober, On the homology of finite abelian coverings, *Topology Appl.* **43** (1992) 157-166.
- [18] A. Libgober, On combinatorial invariance of the cohomology of the Milnor fiber of arrangements and the Catalan equation over function fields. *Arrangements of hyperplanes-Sapporo 2009*, 175-187, Adv. Stud. Pure Math., 62, Math. Soc. Japan, Tokyo, 2012.
- [19] A. Libgober, S. Yuzvinsky, Cohomology of the Orlik-Solomon algebras and local systems. *Compositio Math.* **121** (2000), no. 3, 337-361.
- [20] A. D. Măcinic, S. Papadima, On the monodromy action on Milnor fibers of graphic arrangements. *Topology Appl.* **156** (2009), no. 4, 761-774.

- [21] S. Nazir, Z. Raza, Admissible local systems for a class of line arrangements. *Proc. Amer. Math. Soc.* **137** (2009), no. 4, 1307-1313.
- [22] P. Orlik, L. Solomon, Combinatorics and topology of complements of hyperplanes. *Invent. Math.* **56** (1980), 167-189.
- [23] P. Orlik, H. Terao, Arrangements of hyperplanes. Grundlehren der Mathematischen Wissenschaften, 300. Springer-Verlag, Berlin, 1992. xviii+325 pp.
- [24] P. Orlik, H. Terao, Arrangements and hypergeometric integrals. Second edition. MSJ Memoirs, 9. Mathematical Society of Japan, Tokyo, 2007. x+112 pp.
- [25] S. Papadima, A. Suciu, The spectral sequence of an equivariant chain complex and homology with local coefficients. *Trans. A. M. S.*, **362** (2010), no. 5, 2685-2721.
- [26] S. Papadima, A. Suciu, The Milnor fibration of a hyperplane arrangement: from modular resonance to algebraic monodromy, *Proc. London Math. Soc.* **114** (2017), no. 6, 961-1004.
- [27] G. L. Rybnikov, On the fundamental group of the complement of a complex hyperplane arrangement. *Funktsional. Anal. i Prilozhen.*, **45** (2):71-85 (2011).
- [28] M. Sakuma, Homology of abelian coverings of links and spatial graphs, *Canad. J. Math.* **47** (1) (1995) 201-224.
- [29] M. Salvetti, M. Serventi, Twisted cohomology of arrangements of lines and Milnor fibers. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **17** (2017), no. 4, 1461-1489. 5
- [30] M. Salvetti, M. Serventi, Arrangements of lines and monodromy of associated Milnor fibers. *J. Knot Theory Ramifications* **25** (2016), no. 12, 1642014, 17 pp.
- [31] V. Schechtman, H. Terao, and A. Varchenko, Local systems over complements of hyperplanes and the Kac-Kazhdan conditions for singular vectors. *J. Pure Appl. Algebra* **100** (1995), 93-102.
- [32] A. Suciu, Fundamental groups of line arrangements: Enumerative aspects *Advances in algebraic geometry motivated by physics (Lowell, MA, 2000)*, 43-79, Contemporary Mathematics, vol. 276, Amer. Math. Soc., Providence, RI, 2001.
- [33] A. Suciu, Hyperplane arrangements and Milnor fibrations, *Ann. Fac. Sci. Toulouse Math. (6)* **23** (2014), no. 2, 417-481.
- [34] M. Torielli, M. Yoshinaga, Resonant bands, Aomoto complex, and real 4-nets. *Journal of Singularities*, **11** (2015) 33-51.
- [35] K. Williams, The homology groups of the Milnor fiber associated to a central arrangement of hyperplanes in  $\mathbb{C}^3$ . *Topology Appl.* **160** (2013), no. 10, 1129-1143.
- [36] M. Yoshinaga, Milnor fibers of real line arrangements. *Journal of Singularities*, **7** (2013), 220-237.
- [37] M. Yoshinaga, Resonant bands and local system cohomology groups for real line arrangements. *Vietnam J. Math.* **42** (2014), no. 3, 377-392.
- [38] S. Yuzvinsky, Cohomology of the Brieskorn-Orlik-Solomon algebras. *Comm. Algebra* **23** (1995), no. 14, 5339-5354.

MASAHIKO YOSHINAGA, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE,  
 HOKKAIDO UNIVERSITY, KITA 10, NISHI 8, KITA-KU, SAPPORO 060-0810, JAPAN.  
*E-mail address:* yoshinaga@math.sci.hokudai.ac.jp