

Tube-constructing paratanaoidean tanaidaceans (Crustacea: Peracarida): a brief review

Keiichi Kakui

Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan.
e-mail: kakui@eis.hokudai.ac.jp, Tel: +81-11-706-2750.

Abstract

This study summarizes previous reports of tubes constructed with thread or mucus by paratanaoidean tanaidaceans. A literature survey found 34 genera in 14 extant families to contain species with for which information exists on tubes, whereas five families (Akanthophoreidae, Heterotanoididae, Paranarthrurellidae, Pseudozeuxidae, and Teleotanaidae) lacked any records of tube-use.

Key words: benthos; Paratanaoidea; Tanaidacea; Tanaidomorpha; thread; tube dweller

Tanaidacea is an order of benthic crustaceans that contains about 1500 described species worldwide (Anderson 2020) and comprises five superfamilies (Apseudoidea, Cretitanaoidea, Neotanaoidea, Paratanaoidea, and Tanaidoidea) in three suborders (Anthracocaridomorpha, Apseudomorpha, and Tanaidomorpha) (Kakui et al. 2011; Heard et al. 2020). Many tanaidaceans construct tubes with thread or mucus in bottom sediments and on biotic or abiotic substrata (Larsen 2005; Kakui 2016; hereafter ‘tube-dwellers’); in some groups, one or both ends of tubes can be sealed (Hassack and Holdich 1987). Although a few tube-dwelling species are known in Apseudoidea (Kakui and Hiruta 2014, 2017), most known tube dwellers belong to Paratanaoidea or Tanaidoidea and use a thoracic gland system (Kaji et al. 2016) to construct their tubes.

Paratanaoidea comprises more than 170 genera in one extinct and 19 extant families (Anderson 2020; Heard et al. 2020). Paratanaoideans are almost exclusively tube dwellers (Larsen et al.

2015), but actual observations of their tubes have been restricted to a few groups. Hassack and Holdich (1987), in a review that included previous records of tube construction in Tanaidacea as well as new observations from fixed samples, found that twelve paratanaoidean genera included tube-constructing species. Several papers subsequently reported additional paratanaoideans having tubes.

My literature survey detected 34 genera in 14 extant families that contain species for which information exists on tubes (Table 1); Fig. 1 shows two examples of paratanaoidean tubes. Several previous studies have included photos or illustrations of paratanaoidean tubes or threads (e.g., Bückle Ramírez 1965; Greve 1967; Lang 1973; Heard 1982; Bird and Holdich 1985; Hassack and Holdich 1987; Sieg and Zibrowius 1988; Krasnow and Taghon 1997; Larsen and Cunha 2006; Blazewicz-Paszkowycz et al. 2012; Gellert and Błażewicz 2019). There are no records of tube use in the remaining five extant families in Paratanaoidea (Akanthophoreidae, Heterotanoidi-



Fig. 1. Two examples of paratanaoidean tubes. A, tube of *Tanaella kommitzia* Larsen and Shimomura, 2007 (Tanaellidae), with an individual inside. B, tube of *Nesotanaeis* cf. *ryukyensis* Kakui et al., 2010 (Nototanaidae).

dae, Paranarthrurellidae, Pseudozeuxidae, and Teleotanaidae).

Several previous studies found no tubes in limited groups; for example, Bird (2004: p. 1381) noted, “No tube has been definitively associated with any of the *Paratyphlotanaeis* species reported here.” Failure to observe tubes is often interpreted to mean that tubes were broken or lost during sample processing and fixation (Bird 2004; Larsen 2005; Błażewicz-Paszkowycz 2007). There may be validity to this, as tubes of *Paranarthrura*, which had not been reported by 1989 (Bird and Holdich 1989a), were reported later (Bird 2010), and Błażewicz-Paszkowycz (2007) observed that the tubes of *Nototanaeis antarcticus* disintegrated during sample processing. Or, it may imply the possibility that there may be groups constructing

tubes only for limited periods in their life (e.g., as a brood-nursery). On the other hand, several researchers have suggested that tube construction occurs in relatively few paratanaoidean groups (see Larsen 2005). Uncertainty remains because observations of both tubes and thread-producing glands are very limited for paratanaoideans. To understand the taxonomic distribution of tube building members in Paratanaoidea, further reports are badly needed.

Acknowledgments

I thank Matthew H. Dick for reviewing the manuscript and editing the English. This study was funded in part by KAKENHI grant JP19K06800 from the Japan Society for the Promotion of Science (JSPS).

Table 1. List of paratanaoidean groups in which tubes have been observed.

Family	Genus	Species	Source	
Agathotanaidae	<i>Agathotana</i>	<i>A. ingolfi</i>	Bird (2010)	
	<i>Paranarthrura</i>	<i>P. insignis</i>	Bird (2010)	
Anarthruridae	<i>Anarthrura</i>	<i>A. simplex</i>	Hassack and Holdich (1987)	
	<i>Macilenta</i>	<i>M. ewae</i>	Gellert and Błażewicz (2019)	
	<i>Anarthruopsis</i>	<i>A. sp.</i>	Reidenauer and Thistle (1985)	
	<i>Waki</i>	<i>W. australiensis</i>	Gellert and Błażewicz (2019)	
Colletteidae	<i>Collettea</i>	<i>C. cylindrata</i>	Hassack and Holdich (1987)	
	<i>Haplocope</i>	<i>H. angusta</i>	Greve (1967)	
	<i>Nematotana</i>	<i>N. mirabilis</i>	Bird and Holdich (1985)	
Cryptocopidae	<i>Cryptocopoides</i>	<i>C. arcticus</i>	Hassack and Holdich (1987)	
Leptocheliidae	<i>Chondrochelia</i>	<i>C. dubia</i> , <i>C. savignyi</i>	Lang (1973); Holdich and Jones (1983); Krasnow and Taghon (1997)	
		<i>Hargeria</i>	<i>H. rapax</i>	Heard (1982)
	<i>Heterotana</i>	<i>H. oerstedii</i> , <i>H. sp.</i>	Bückle Ramírez (1965)	
	<i>Konarus</i>	<i>K. straddi</i>	Bamber (2013)	
	<i>Parakonarus</i>	<i>P. kopure</i>	Bird (2011)	
	<i>Poorea</i>	<i>P. nobbi</i>	Bamber (2005)	
	Leptognathiidae	<i>Leptognathia</i>	<i>L. breviremis</i> , <i>L. tenella</i>	Greve (1967); Hassack and Holdich (1987)
	Mirandotanaidae	<i>Mirandotana</i>	<i>M. vorax</i>	Sieg (1986)
	Nototanaidae	<i>Nesotana</i>	<i>N. lacustris</i> , <i>N. cf. ryukyuensis</i>	Shiino (1968); this study
<i>Nototana</i>			<i>N. antarcticus</i> , <i>N. dimorphus</i>	Oliver and Slattery (1985); Błażewicz-Paszkowycz (2007)
<i>Bathytana</i>		<i>B. culteriformis</i>	Larsen and Wilson (1998)	
Paratanaidae	<i>Paratana</i>	<i>P. hamulus</i>	Bird and Bamber (2013)	
	<i>Pseudobathytana</i>	<i>P. gibberosus</i>	Larsen and Heard (2001)	
Pseudotanaidae	(not mentioned)	(not mentioned)	Bird and Holdich (1989b)	
Tanaellidae	<i>Tanaella</i>	<i>T. sp.</i> , <i>T. kommitzia</i>	Larsen and Heard (2004); this study	
Tanaissuidae	<i>Tanaissus</i>	<i>T. lilljeborgi</i> , <i>T. psammophilus</i>	Siewing (1953); Heard et al. (2004)	
Tanaopsiidae	<i>Tanaopsis</i>	<i>T. graciloides</i>	Bamber (1986)	
Typhlotanaidae	<i>Obesutana</i>	<i>O. sigridi</i>	Larsen and Cunha (2006); Larsen et al. (2006)	
		<i>Peraeospinosus</i>	<i>P. magnificus</i>	Kudinova-Pasternak (1969)
		<i>Pulcherella</i>	<i>P. pulcher</i>	Hassack and Holdich (1987)
	<i>Typhlotana</i>	<i>T. aequiremis</i> , <i>T. assimilis</i> , <i>T. brevicornis</i> , <i>T. solidus</i> , <i>T. spinicauda</i> , <i>T. trispinosus</i>	Greve (1967); Hassack and Holdich (1987)	
		<i>Typhlotanoides</i>	<i>T. spp.</i>	Hassack and Holdich (1987)
		<i>Bifidia</i>	<i>B. scleractinicola</i>	Sieg and Zibrowius (1988)
		<i>Exspina</i>	<i>E. typica</i>	Hassack and Holdich (1987)

References

- Anderson, G. (2020). Tanaidacea—Forty Years of Scholarship, Version 3.0. <https://aquila.usm.edu/tanaids30/5/> (accessed 12 December 2020).
- Bamber, R. N. (1986). The marine fauna of the Cullercoats District 20 Tanaidacea. Rep. Dove Mar. Lab. 33: 1–14.
- Bamber, R. N. (2005). The tanaidaceans (Arthropoda: Crustacea: Peracarida: Tanaidacea) of Esperance, Western Australia, Australia. In: F. E. Wells, D. I. Walker, G. A. Kendrick (Eds.) The Marine Flora and Fauna of Esperance, Western Australia. Western Australian Museum, Perth, p. 613–727.
- Bamber, R. N. (2013). A re-assessment of *Konarus* Bamber, 2006 and sympatric leptocheliids from Australasia, and of *Pseudoleptochelia* Lang, 1973 (Crustacea: Peracarida: Tanaidacea). *Zootaxa* 3694: 1–39.
- Bird, G. J., Holdich, D. M. (1985). A remarkable tubicolous tanaid (Crustacea: Tanaidacea) from the Rockall Trough. *J. Mar. Biol. Assoc. UK* 65: 563–572.
- Bird, G. J., Holdich, D. M. (1989a). Deep-sea Tanaidacea (Crustacea) of the North-east Atlantic: the genus *Paranarthrura* Hansen. *J. Nat. Hist.* 23: 137–167.
- Bird, G. J., Holdich, D. M. (1989b). Tanaidacea (Crustacea) of the north-east Atlantic: the subfamily Pseudotanainae (Pseudotanaidae) and the family Nototanaidae. *Zool. J. Linnean Soc.* 97: 233–298.
- Bird, G. J. (2004). The Tanaidacea (Crustacea, Peracarida) of the North-East Atlantic: the shelf and bathyal *Paratyphlotanais* of the ‘Atlantic Margin’. *J. Nat. Hist.* 38: 1359–1384.
- Bird, G. J. (2010). Tanaidacea (Crustacea, Peracarida) of the North-east Atlantic: the Agathotanaidae of the AFEN, BIOFAR and BIOICE projects, with a description of a new species of *Paragathotanais* Lang. *Zootaxa* 2730: 1–22.
- Bird, G. J. (2011). Paratanaoidean tanaidaceans (Crustacea: Peracarida) from littoral and shallow sublittoral habitats in New Zealand, with descriptions of three new genera and seven new species. *Zootaxa* 2891: 1–62.
- Bird, G. J., Bamber, R. N. (2013). New littoral, shelf, and bathyal Paratanaidae (Crustacea: Peracarida: Tanaidacea) from New Zealand, with descriptions of three new genera. *Zootaxa* 3676: 1–71.
- Błażewicz-Paszkowycz, M. (2007). A revision of the family Typhlotanaidae Sieg 1984 (Crustacea: Tanaidacea) with the remarks on the Nototanaidae Sieg, 1976. *Zootaxa* 1598: 1–141.
- Błażewicz-Paszkowycz, M., Bamber, R., Anderson, G. (2012). Diversity of Tanaidacea (Crustacea: Peracarida) in the world’s oceans—how far have we come? *PLoS ONE* 7: e33068.
- Bückle Ramírez, L. F. (1965). Untersuchungen über die Biologie von *Heterotanais oerstedii* Kröyer (Crustacea, Tanaidacea). *Z. Morphol. Ökol. Tiere* 55: 714–782.
- Gellert, M., Błażewicz, M. (2019). New species of Anarthruridae (Tanaidacea: Crustacea) of the western Australian slope. *Mar. Biodivers.* 49: 583–601.
- Greve, L. (1967). On the tube building of some Tanaidacea. *Sarsia* 29: 295–298.
- Hassack, E., Holdich, D. M. (1987). The tubicolous habit amongst the Tanaidacea (Crustacea, Peracarida) with particular reference to deep-sea species. *Zool. Scr.* 16: 223–233.
- Heard, R. W. (1982). Guide to Common Tidal Marsh Invertebrates of the Northeastern Gulf of Mexico. Mississippi-Alabama Sea Grant Consortium, Mississippi.
- Heard, R. W., Hansknecht, T., Larsen, K. (2004). An illustrated identification guide to Florida Tanaidacea (Crustacea: Peracarida) occurring in depths of less than 200 m. Florida Department of Environmental Protection, Florida.
- Heard, R. W., Morales-Núñez, A. G., Serrano-Sánchez, M. L., Coutiño, M. A., Barragán, R., Vega, V. J. (2020). A new family, genus and species of Tanaidacea (Crustacea; Apseudomorpha) from the Lower Cretaceous (Aptian) of Chiapas, Mexico: Systematic revisions, including designation of two new Paleozoic families, and paleoenvironmental observations. *J. S. Am. Earth Sci.* 102: 102609.
- Holdich, D. M., Jones, J. A. (1983). Tanaids. Cambridge University Press, Cambridge.
- Kaji, T., Kakui, K., Miyazaki, N., Murata, K., Palmer, A. R. (2016). Mesoscale morphology at nanoscale resolution: serial block-face scanning electron microscopy reveals fine 3D detail of a novel silk spinneret system in a tube-building tanaid crustacean. *Front. Zool.* 13: 14.
- Kakui, K., Katoh, T., Hiruta, S. F., Kobayashi, N., Kajihara, H. (2011). Molecular systematics of Tanaidacea (Crustacea: Peracarida) based on 18S sequence data, with an amendment of

- suborder/superfamily-level classification. *Zool. Sci.* 28: 749–757.
- Kakui, K., Hiruta, C. (2014). Diverse pereopodal secretory systems implicated in thread production in an apseudomorph tanaidacean crustacean. *J. Morphol.* 275: 1041–1052.
- Kakui, K. (2016). Review of the taxonomy, diversity, ecology, and other biological aspects of order Tanaidacea from Japan and surrounding waters. In: M. Motokawa, H. Kajihara (Eds.) *Species diversity of animals in Japan*. Springer, Berlin, p. 603–627.
- Kakui, K., Hiruta, C. (2017). Tube construction by a tanaidacean crustacean using a novel mucus secretion system involving the anal opening. *Zool. Lett.* 3: 20.
- Krasnow, L. D., Taghon, G. L. (1997). Rate of tube building and sediment particle size selection during tube construction by the tanaid Crustacea, *Leptochelia dubia*. *Estuaries* 20: 534–546.
- Kudinova-Pasternak, R. K. (1969). A case of extramarsupial development of eggs in *Typhlotanais magnificus* (Crustacea, Tanaidacea) living in the tube. *Tr. Inst. Okeanol.* 48: 1737–1738. [in Russian with English abstract]
- Lang, K. (1973). Taxonomische und phylogenetische Untersuchungen über die Tanaidaceen (Crustacea) 8. Die Gattungen *Leptochelia* Dana, *Paratanais* Dana, *Heterotanais* G. O. Sars und *Nototanais* Richardson. Dazu einige Bemerkungen über die Monokonophora und ein Nachtrag. *Zool. Scr.* 2: 197–229.
- Larsen, K., Wilson, D. F. (1998). Tanaidomorphan systematics—is it obsolete? *J. Crustacean Biol.* 18: 346–362.
- Larsen, K., Heard, R. W. (2001). A new tanaidacean subfamily, Bathytanaidinae (Crustacea: Paratanaididae), from the Australian continental shelf and slope. *Zootaxa* 19: 1–22.
- Larsen, K., Heard, R. (2004). Revision of the tanaidomorphan deep-sea genus *Tanaella* (Crustacea: Tanaidacea). *J. Nat. Hist.* 38: 549–579.
- Larsen, K. (2005). Deep-sea Tanaidacea (Peracarida) from the Gulf of Mexico. Brill, Leiden.
- Larsen, K., Błażewicz-Paszkowycz, M., Cunha, M. R. (2006). Tanaidacean (Crustacea: Peracarida) fauna from chemically reduced habitats—the Lucky Strike hydrothermal vent system, Mid-Atlantic Ridge. *Zootaxa* 1187: 1–36.
- Larsen, K., Cunha, M. R. (2006). *Obesutanais sigridi* Larsen, Błażewicz-Paszkowycz & Cunha, in press. *Danisia* 18: 376.
- Larsen, K., Guțu, M., Sieg, J. (2015). Order Tanaidacea Dana, 1849. In: J. C. von Vaupel Klein, M. Charmantier-Daures, F. R. Schram (Eds.) *The Crustacea. Revised and Updated, as well as Extended from the Traité de Zoologie* 5. Brill, Leiden, p. 249–329.
- Oliver, J. S., Slattery, P. N. (1986). Effects of crustacean predators on species composition and population structure of soft-bodied infauna from McMurdo Sound, Antarctica. *Ophelia* 24: 155–175.
- Reidenauer, J. A., Thistle, D. (1985). The tanaid fauna from a region of the deep North Atlantic where near-bottom current velocities are high. *Oceanol. Acta* 8: 355–360.
- Shiino, S. M. (1968). A tanaid crustacean, *Nesotanais lacustris* gen. et sp. nov., from Lake Tegano, Rennell Island. *Nat. Hist. Rennell Island, British Solomon Islands* 5: 153–168.
- Sieg, J. (1986). Crustacea Tanaidacea of the Antarctic and the Subantarctic. 1. On Material Collected at Tierra del Fuego, Isla de los Estados, and the west coast of the Antarctic Peninsula. American Geophysical Union, Washington.
- Sieg, J., Zibrowius, H. (1988). Association of a tube inhabiting tanaidacean, *Bifidia scleractinicola* gen. nov., sp. nov., with bathyal scleractinians off New Caledonia (Crustacea Tanaidacea - Cnidaria Scleractinia). *Mésogée* 48: 189–199.
- Siewing, R. (1953). Morphologische Untersuchungen an Tanaidaceen und Lophogastriden. *Z. Wiss. Zool.* 157: 333–426.

Received: 13 December 2020 | Accepted: 11 February 2021 | Published: 12 February 2021