



Title	The Gauss maps of Demoulin surfaces with conformal coordinates In Memory of Professor Zhengguo Bai(1916-2015)
Author(s)	Inoguchi, Jun-ichi; Kobayashi, Shimpei
Citation	Science China-mathematics, 64(7), 1479-1492 https://doi.org/10.1007/s11425-020-1738-0
Issue Date	2021-09-28
Doc URL	http://hdl.handle.net/2115/82779
Rights	This is a post-peer-review, pre-copyedit version of an article published in Science China-mathematics. The final authenticated version is available online at: http://dx.doi.org/10.1007/s11425-020-1738-0 .
Type	article (author version)
File Information	Sci. China-Math.64-7_1479-1492.pdf



[Instructions for use](#)

THE GAUSS MAPS OF DEMOULIN SURFACES WITH CONFORMAL COORDINATES

JUN-ICHI INOYUCHI AND SHIMPEI KOBAYASHI

In Memory of Professor Zhengguo Bai (1916-2015)

ABSTRACT. Demoulin surfaces in real projective 3-space are investigated. Our result enable us to establish a generalized Weierstrass type representation for definite Demoulin surfaces by virtue of primitive maps into a certain semi-Riemannian 6-symmetric space.

INTRODUCTION

Professor Zhengguo Bai have done great contributions in projective differential geometry. For example, he solved the so-called Fubini's problem [22] (*cf* [28]).

Projective differential geometry of surfaces is a treasure box of infinite dimensional integrable systems. For instance, harmonic maps of Riemann surfaces into complex projective space $\mathbb{C}\mathbb{P}^n$ (the $\mathbb{C}\mathbb{P}^n$ -sigma models in particle physics) are typical examples of 2-dimensional integrable systems. One of the key clue of the study of harmonic maps into complex projective space is the use of *harmonic sequences* introduced by Chern and Wolfson [8]. It should be emphasized that the basic idea of harmonic sequence goes back to *Laplace sequence* in classical projective differential geometry, see [3].

From modern point of view, the Laplace sequence produces 2-dimensional Toda field equation of type A_∞ , see [9, 25]. In particular, the periodic Laplace sequence produces 2-dimensional periodic Toda field equations. For example, Laplace sequences of period 2 produce sinh-Gordon equation. *Tițeica equation* is obtained as Laplace sequence of period 3, and it is a structure equation of affine spheres [11]. Laplace sequences of period 4 were studied by Su [26, 27]. Hu gave a Darboux matrix, that is, the simple type dressing for such a sequence [14].

This article addresses Laplace sequences of period 6. The Toda field equation derived from those sequences is a structure equation of *Demoulin surfaces* in real projective 3-space $\mathbb{R}\mathbb{P}^3$.

Godeaux gave a method for studying projective surfaces through their Plücker images in real projective 5-space $\mathbb{R}\mathbb{P}^5$. His method relies on the consideration of the Laplace sequence associated with the Plücker image, called the *Godeaux sequence*. For a characterization of

2010 *Mathematics Subject Classification*. Primary 53A20, Secondary 53C43, 37K10.

Key words and phrases. Demoulin surface, Wilczynski frame, Gauss map.

The first named author is partially supported by JSPS Kakenhi Grant Number JP19K03461.

The second named author is partially supported by JSPS Kakenhi Grant Number JP18K03265.

Demoulin surfaces in terms of Godeaux sequences, see [25, §4.8]. Bai [21] studied Godeaux sequences of quadrics.

In [15], the second named author considered two Gauss maps of surfaces in $\mathbb{R}\mathbb{P}^3$ with indefinite *projective metric* and characterized projective minimal surfaces and Demoulin surfaces in terms of harmonicities of the Gauss maps. In this paper, we consider those surfaces with *positive definite* projective metric. This paper is organized as follows: After preparing prerequisite knowledge on projective surface theory in Sections 1-3, we parametrize the space of all conformal 2-spheres in $\mathbb{R}\mathbb{P}^3$ in Section 4. We will show that the space of all conformal 2-spheres is realized as a semi-Riemannian symmetric space. The Gauss maps introduced in this paper take values in this symmetric space. In Section 5, we introduce the first-order Gauss map for a surface in $\mathbb{R}\mathbb{P}^3$ as a congruence of conformal 2-spheres which has the first-order contact to the surface. Definite Demoulin surfaces are characterized as surfaces with conformal first order Gauss map. In addition definite Demoulin surfaces and definite projective minimal coincidence surfaces are characterized by harmonicity of first order Gauss map. In the final section we will show that every definite Demoulin surface can be constructed by a primitive map into certain semi-Riemannian 6-symmetric space fibered over the semi-Riemannian symmetric space of all conformal 2-spheres.

Throughout this paper, we use the following abbreviation:

$$\text{diag}(a_1, a_2, \dots, a_n) = \begin{pmatrix} a_1 & & & \\ & a_2 & & \\ & & \ddots & \\ & & & a_n \end{pmatrix}, \quad \text{offdiag}(a_1, a_2, \dots, a_n) = \begin{pmatrix} & & & a_1 \\ & & & \\ & & a_2 & \\ & & & \ddots \\ a_n & & & \end{pmatrix}.$$

1. PROJECTIVE SURFACE THEORY

Let $\mathfrak{f}: M \rightarrow \mathbb{R}\mathbb{P}^3$ be an immersed surface in the real projective 3-space $\mathbb{R}\mathbb{P}^3$. Take a simply connected region $\mathbb{D} \subset M$ and homogeneous coordinate vector field $f = (f^0, f^1, f^2, f^3): \mathbb{D} \rightarrow \mathbb{R}^4 \setminus \{\mathbf{0}\}$. Let D be the natural affine connection on \mathbb{R}^4 and Ω a volume element so that $D\Omega = 0$. Thus $(\mathbb{R}^4, D, \Omega)$ is an *equiaffine 4-space*. One can take a vector field ξ transversal to both f and the radial vector field $\zeta = \sum_{i=0}^3 x^i \partial / \partial x^i$. Then ξ induces an affine connection ∇ on \mathbb{D} and symmetric tensor fields h and T via the *Gauss formula*:

$$D_X f_* Y = f_*(\nabla_X Y) + h(X, Y)\xi + T(X, Y)\zeta, \quad X, Y \in \Gamma(T\mathbb{D}).$$

Moreover we have the following *Weingarten formula*:

$$D_X \xi = -f_*(SX) + \tau(X)\xi + \rho(X)\zeta.$$

The triplet (\mathbb{D}, f, ξ) is a *centroaffine surface* (of codimension 2) in \mathbb{R}^4 in the sense of [19, 20]. We introduce an area element ϑ on \mathbb{D} by

$$\vartheta(X, Y) = \Omega(f_* X, f_* Y, \xi, \zeta).$$

The *cubic form* C is defined by

$$C = \nabla h + \tau \otimes h.$$

The non-degeneracy of h is independent of the choice of ξ . In addition, the conformal class $[h]$ of h is independent of ξ . Thus the property “ h is *positive definite*” is well defined for f . Throughout this article, we assume that h is positive definite.

When we take ξ so that $\tau = 0$, then (\mathbb{D}, f, ξ) is said to be *equiaffine*. An equiaffine centroaffine immersion f is said to be *Blaschke* if ϑ coincides with the area element of the metric h .

On the other hand Nomizu and Sasaki [19] showed that there exists a transversal vector field ξ such that

$$(1.1) \quad \operatorname{tr}_h T + \operatorname{tr} S = 0.$$

Such a vector field is called a *pre-normalized transversal vector field*. In particular, pre-normalized transversal vector field ξ so that (\mathbb{D}, f, ξ) is a Blaschke immersion is unique up to sign. In such a choice, the pair surface (f, ξ) is called a *pre-normalized Blaschke immersion*.

Let us take another homogeneous coordinate vector field $\tilde{f} = \phi f$. Here ϕ is a smooth (nonzero) function. Then the connection $\tilde{\nabla}$ induced from \tilde{f} is projectively equivalent to ∇ . The equiaffine property is preserved under the change f by ϕf .

Let us denote by ∇^h the Levi-Civita connection of h . Then the scalar field $J = h(K, K)/2$ is called the *Fubini-Pick invariant* of f . Here $K = \nabla - \nabla^h$. The Riemannian metric Jh is projectively invariant and called the *projective metric* of \mathfrak{f} . Although C itself is *not* projective invariant, its conformal class is projective invariant (see [18]). When (f, ξ) is pre-normalized Blaschke, the projective metric is given by $h(\nabla h, \nabla h)h/8$.

For more details on centroaffine immersions and projective immersions, we refer to [19, 20].

2. WILCZYNSKI FRAMES

Let $\mathfrak{f} : M \rightarrow \mathbb{R}\mathbb{P}^3$ be an immersed surface with *positive definite projective metric*. We regard M as a Riemann surface with respect to the conformal structure $[Jh]$ determined by the projective metric Jh .

We take a simply connected complex coordinate region \mathbb{D} with coordinate $z = x + iy$ on \mathbb{D} and a lift $f = (f^0, f^1, f^2, f^3) : \mathbb{D} \rightarrow \mathbb{R}^4 \setminus \{\mathbf{0}\}$. Then the *canonical system* of \mathfrak{f} is given by

$$(2.1) \quad f_{zz} = bf_{\bar{z}} + pf, \quad f_{\bar{z}\bar{z}} = \bar{b}f_z + \bar{p}f$$

for some smooth functions b and p , see [25, p. 121]. Note that the subscripts z and \bar{z} denote the partial derivative of z and \bar{z} , respectively:

$$\frac{\partial}{\partial z} := \frac{1}{2} \left(\frac{\partial}{\partial x} - \sqrt{-1} \frac{\partial}{\partial y} \right), \quad \frac{\partial}{\partial \bar{z}} := \frac{1}{2} \left(\frac{\partial}{\partial x} + \sqrt{-1} \frac{\partial}{\partial y} \right).$$

Assume that $f^0 \neq 0$, then \mathfrak{f} is given by the inhomogeneous coordinate $\mathfrak{f} = (f^1, f^2, f^3)/f^0$. The canonical system is rewritten as

$$(2.2) \quad \mathfrak{f}_{zz} = b\mathfrak{f}_{\bar{z}} - 2(\log f^0)_z \mathfrak{f}_z, \quad \mathfrak{f}_{\bar{z}\bar{z}} = \bar{b}\mathfrak{f}_z - 2(\log f^0)_{\bar{z}} \mathfrak{f}_{\bar{z}}.$$

The integrability condition of the canonical system is (cf. [25, §2.3]):

$$p_{\bar{z}} = b\bar{b}_z + \frac{1}{2}b_z\bar{b} - \frac{1}{2}b_{z\bar{z}},$$

$$\text{Im}(b_{z\bar{z}\bar{z}} - b\bar{b}_{z\bar{z}} - 2b\bar{p}_{\bar{z}} - 2b_z\bar{b}_z - 4b_z\bar{p}) = 0.$$

The Fubini-Pick invariant is given by $J = 8|b|^2$ and hence the projective metric is $8|b|^2 dzd\bar{z}$. The cubic form of \mathfrak{f} is given by $C = -2(b dz^3 + \bar{b}d\bar{z}^3)$ (see [24, p. 54, Definition, §4.8]). Note that when f is pre-normalized Blaschke, then the projective metric is expressed as $2|b|^2 dzd\bar{z}$.

Hereafter we assume that $b \neq 0$. Note that when $C = 0$, \mathfrak{f} is a part of a quadratic surface (Wilczynski [30], Pick [23]. See also [24, Theorem 4.4].)

The *Wilczynski frame* F of \mathfrak{f} is defined by

$$F = (f, f_1, f_2, \eta),$$

where

$$f_1 := f_z - \frac{\bar{b}_z}{2\bar{b}}f, \quad f_2 := f_{\bar{z}} - \frac{b_{\bar{z}}}{2b}f, \quad \eta = f_{z\bar{z}} - \frac{\bar{b}_z}{2\bar{b}}f_{\bar{z}} - \frac{b_{\bar{z}}}{2b}f_z + \left(\frac{|b_z|^2}{4|b|^2} - \frac{|b|^2}{2}\right)f.$$

Then a straightforward computation shows that the Wilczynski frame F satisfies the following equations:

$$(2.3) \quad F_z = FU \quad \text{and} \quad F_{\bar{z}} = FV,$$

where

$$U = \begin{pmatrix} \bar{b}_z/(2\bar{b}) & P & k & b\bar{P} \\ 1 & -\bar{b}_z/(2\bar{b}) & 0 & k \\ 0 & b & \bar{b}_z/(2\bar{b}) & P \\ 0 & 0 & 1 & -\bar{b}_z/(2\bar{b}) \end{pmatrix},$$

$$V = \begin{pmatrix} b_{\bar{z}}/(2b) & \bar{k} & \bar{P} & \bar{b}P \\ 0 & b_{\bar{z}}/(2b) & \bar{b} & \bar{P} \\ 1 & 0 & -b_{\bar{z}}/(2b) & \bar{k} \\ 0 & 1 & 0 & -b_{\bar{z}}/(2b) \end{pmatrix}.$$

Here we introduced functions k and P and Q of as follows:

$$(2.4) \quad k = \frac{|b|^2 - (\log b)_{z\bar{z}}}{2},$$

$$(2.5) \quad P = p + \frac{b_{\bar{z}}}{2} - \frac{\bar{b}_{z\bar{z}}}{2\bar{b}} + \frac{\bar{b}_z^2}{4\bar{b}^2}.$$

The compatibility conditions of (2.3) are

$$(2.6) \quad \bar{P}_z = k_{\bar{z}} + k \frac{b_{\bar{z}}}{b},$$

$$(2.7) \quad \text{Im}(\bar{b}P_z + 2\bar{b}_z P) = 0.$$

These equations are nothing but the *projective Gauss-Codazzi equations* of a surface \mathfrak{f} . One can see that Pdz^2 and $2b^2Pdz^4$ are globally defined on M and projectively invariant [13].

Since both U and V are trace free, the Wilczynski frame F takes values in $\text{SL}_4\mathbb{C}$ up to initial condition. Moreover, if we choose at some base point $z_* \in \mathbb{D}$ and $F(z_*) = \text{id}$, then the frame F takes values in $\text{SL}_4\mathbb{R}$ by conjugation of a simple complex matrix:

$$(2.8) \quad \text{Ad}(L)F \in \text{SL}_4\mathbb{R}, \quad L = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2} & 0 & 0 & 0 \\ 0 & \sqrt{-1} & -\sqrt{-1} & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & \sqrt{2} \end{pmatrix}.$$

3. PROJECTIVE MINIMAL SURFACES AND DEFINITE DEMOULIN SURFACES

A surface $\mathfrak{f} : M \rightarrow \mathbb{RP}^3$ with positive definite projective metric is said to be a *projective minimal surface* if it is a critical point of the area functional of the projective metric (called the *projective area functional*): Then the projective minimality can be computed as in [29]:

$$(3.1) \quad \bar{b}P_z + 2\bar{b}_z P = 0.$$

where the functions P is defined in (2.5). It should be remarked that the projective minimality (3.1) implies the second equation (2.7) of the projective Gauss-Codazzi equations. There is a particular class of projective minimal surfaces with positive definite projective metric.

A surface with positive definite projective metric is said to be a *definite Demoulin surface* if it satisfies $P = 0$. The Demoulin property is originated from Demoulin transformations of surfaces in \mathbb{RP}^3 . For more details, we refer to [25].

4. THE PLÜCKER QUADRIC AND THE SPACE OF CONFORMAL SPHERES

4.1. The Plücker quadric. Take a volume element Ω on \mathbb{R}^4 parallel with respect to the natural affine connection D . Then we can introduce a scalar product $\langle \cdot, \cdot \rangle$ on $\wedge^2 \mathbb{R}^4$ by

$$\langle \alpha, \beta \rangle = \Omega(\alpha \wedge \beta), \quad \alpha, \beta \in \wedge^2 \mathbb{R}^4.$$

One can check that $\langle \cdot, \cdot \rangle$ is of signature $(3, 3)$. In fact, let $\{e_0, e_1, e_2, e_3\}$ be the natural basis of \mathbb{R}^4 . Denote by $\{e^0, e^1, e^2, e^3\}$ the dual basis of $\{e_0, e_1, e_2, e_3\}$. Then with respect to the volume element $\Omega = e^0 \wedge e^1 \wedge e^2 \wedge e^3$, the basis $\{e_0 \wedge e_1, e_0 \wedge e_2, e_0 \wedge e_3, e_1 \wedge e_2, e_3 \wedge e_1, e_2 \wedge e_3\}$ of $\wedge^2 \mathbb{R}^4$, the scalar product $\langle \cdot, \cdot \rangle$ is determined by the matrix $\text{offdiag}(1, 1, 1, 1, 1, 1)$. The special linear group $\text{SL}_4\mathbb{R}$ acts on $\wedge^2 \mathbb{R}^4$ via the action:

$$\text{SL}_4\mathbb{R} \times \wedge^2 \mathbb{R}^4 \rightarrow \wedge^2 \mathbb{R}^4; \quad (g, v \wedge w) \mapsto gv \wedge gw.$$

One can see that this action is isometric with respect to $\langle \cdot, \cdot \rangle$. This fact implies the Lie group isomorphism $\mathrm{PSL}_4\mathbb{R} \cong \mathrm{SO}_{3,3}^+$. Here $\mathrm{SO}_{3,3}^+$ denotes the identity component of the semi-orthogonal group $\mathrm{O}_{3,3}$.

Next we consider the Plücker embedding of the Grassmannian manifold $\mathrm{Gr}_2(\mathbb{R}^4)$ of 2-planes in \mathbb{R}^4 into the projective 5-space $\mathbb{RP}^5 = \mathbb{P}(\wedge^2\mathbb{R}^4)$. The Plücker coordinates of the 2-plane spanned by (a^0, a^1, a^2, a^3) and (b^0, b^1, b^2, b^3) is $[p_{01} : p_{02} : p_{03} : p_{23} : p_{31} : p_{12}]$, where

$$(4.1) \quad p_{ij} = \det \begin{pmatrix} a^i & b^i \\ a^j & b^j \end{pmatrix}.$$

The Plücker coordinates $[p_{01} : p_{02} : p_{03} : p_{23} : p_{31} : p_{12}]$ of $a \wedge b$ satisfies the *quadratic Plücker relation*:

$$(4.2) \quad p_{01}p_{23} + p_{02}p_{31} + p_{03}p_{12} = 0.$$

Thus the Plücker image of $\mathrm{Gr}_2(\mathbb{R}^4)$ is a projective variety (called the *Plücker quadric*) of \mathbb{RP}^5 determined by the equation (4.2). Moreover the Plücker relation means that $(p_{01}, p_{02}, p_{03}, p_{23}, p_{31}, p_{12})$ is null with respect to $\langle \cdot, \cdot \rangle$. Namely the Plücker image of $\mathrm{Gr}_2(\mathbb{R}^4)$ is the projective light cone $\mathbb{P}(\mathcal{L})$ of $\wedge^2\mathbb{R}^4 = \mathbb{R}^{3,3}$.

Now let us consider a line ℓ in \mathbb{RP}^3 connecting two points $a = [a^0 : a^1 : a^2 : a^3]$ and $b = [b^0 : b^1 : b^2 : b^3]$. The Plücker image of ℓ in $\mathbb{RP}^5 = \mathbb{P}(\wedge^2\mathbb{R}^4)$ is

$$a \wedge b = [p_{01} : p_{02} : p_{03} : p_{23} : p_{31} : p_{12}]$$

with (4.1). Hence the space \mathcal{P} of lines in \mathbb{RP}^3 is identified with the Plücker quadric. This identification is called the *Klein correspondence*.

Remark 4.1. The conformal compactification of semi-Euclidean 4-space $\mathbb{R}^{2,2}$ of neutral signature is obtained as the projective light cone $\mathbb{P}(\mathcal{L}) \subset \mathbb{RP}^5$ equipped with the conformal structure induced from $\mathbb{R}^{3,3}$. The action of $\mathrm{PSL}_4\mathbb{R} \cong \mathrm{SO}_{3,3}^+$ on $\mathbb{P}(\mathcal{L})$ is conformal. One can see that the Plücker quadric $\mathcal{P} = \mathbb{P}(\mathcal{L})$ is isomorphic to $\mathrm{Gr}_2(\mathbb{R}^4) \cong (\mathbb{S}^2 \times \mathbb{S}^2)/\mathbb{Z}_2$ (equipped with the standard conformal structure of neutral signature) as a conformal manifold. Note that on $\mathbb{P}(\mathcal{L})$, there exists a complex structure compatible to the standard neutral metric. The standard neutral metric is neutral Kähler with respect to the complex structure. In particular, the Kähler form is regarded as a standard symplectic form on $\mathbb{P}(\mathcal{L})$. For more information on conformal geometry of $\mathbb{P}(\mathcal{L})$, see [17].

4.2. The space of conformal spheres. A quadric in \mathbb{RP}^3 is a surface of the form $\{[v] \in \mathbb{RP}^3 \mid q(v, v) = 0\}$, where q is a scalar product of \mathbb{R}^4 . For our purpose we choose a Lorentzian scalar product $q = \langle \cdot, \cdot \rangle$ on \mathbb{R}^4 and regard it as a Minkowski 4-space $\mathbb{R}^{1,3}$. Then the quadric is nothing but the conformal 2-sphere (Riemann sphere) in \mathbb{RP}^3 . The space of conformal 2-spheres in \mathbb{RP}^3 is parametrized as the space \mathcal{Q} of 3×3 symmetric matrices with determinant one and signature $(1, 3)$. In fact, the conformal 2-sphere is given by the Lorentzian scalar product $q(u, v) = uQv^T$ with $Q \in \mathcal{Q}$.

The special linear group $\mathrm{SL}_4\mathbb{R}$ acts transitively on \mathcal{Q} via the action $(g, Q) \mapsto gQg^T$ with $g \in \mathrm{SL}_4\mathbb{R}$ and $Q \in \mathcal{Q}$. The stabilizer at

$$(4.3) \quad \hat{J}_1 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix},$$

is given by $\hat{K}_1 = \{a \in \mathrm{SL}_4\mathbb{R} \mid a\hat{J}_1a^T = \hat{J}_1\}$, which is isomorphic to the identity component $\mathrm{SO}_{1,3}^+$ of the semi-orthogonal group $\mathrm{O}_{1,3}^+$ of signature $(1, 3)$. Thus \mathcal{Q} is isomorphic to the homogeneous space $\mathrm{SL}_4\mathbb{R}/\mathrm{SO}_{1,3}^+ \cong \mathrm{SO}_{3,3}^+/\mathrm{SO}_{1,3}^+$.

We introduce a scalar product $\langle \cdot, \cdot \rangle$ at $Q \in \mathcal{Q}$ by

$$\langle X, Y \rangle_Q = \mathrm{Tr}(Q^{-1}XQ^{-1}Y), \quad X, Y \in T_Q\mathcal{Q}.$$

Note that at the origin of $\mathrm{SO}_{3,3}^+/\mathrm{SO}_{1,3}^+$, and $8\langle \cdot, \cdot \rangle$ is the Killing form of $\mathfrak{sl}_4\mathbb{R}$. This scalar product is invariant under the action of $\mathrm{SL}_4\mathbb{R}$. In fact,

$$\langle gXg^T, gYg^T \rangle_{gQg^T} = \mathrm{Tr}((gQg^T)^{-1}gXg^T(gQg^T)^{-1}gYg^T) = \langle X, Y \rangle_Q.$$

Thus $\mathcal{Q} = \mathrm{SL}_4\mathbb{R}/\hat{K}_1$ is a semi-Riemannian symmetric space corresponding to the outer involution

$$\hat{\tau}_1(X) = \hat{J}_1(X^T)^{-1}\hat{J}_1.$$

Remark 4.2. The space of lines in the Plücker quadric \mathcal{P} is identified with the Grassmannian manifold of all null 2-planes in $\mathbb{R}^{3,3}$:

$$\mathcal{Z} = \{W \in \mathrm{Gr}_2(\mathbb{R}^{3,3}) \mid W \text{ is a null 2-plane in } \mathbb{R}^{3,3}\} \cong \mathrm{SO}_{3,3}^+/\mathrm{SO}_{2,2}^+.$$

For surfaces in \mathbb{RP}^3 with *indefinite* projective metric, two kinds of Gauss maps are considered in our previous work [15]. Those Gauss maps take value in the space of quadrics determined by scalar products of signature $(2, 2)$ of \mathbb{R}^4 . The space of all quadrics derived from such scalar products are identified with the semi-Riemannian symmetric space $\mathrm{SO}_{3,3}^+/\mathrm{SO}_{2,2}^+$.

5. DEMOULIN SURFACES AND THE FIRST ORDER GAUSS MAPS

In this section, we define the first-order Gauss map for a surface in \mathbb{RP}^3 .

5.1. First-order Gauss map. Let $f : M \rightarrow \mathbb{RP}^3$ be a surface and F the corresponding Wilczynski frame defined in (2.3) with a base point $z_* \in \mathbb{D}$ and $F(z_*) = \mathrm{id}$. Let L be the matrix defined in (2.8) and \hat{F} be the $\mathrm{SL}_4\mathbb{R}$ matrix such that

$$\mathrm{Ad}(L)F = \hat{F}.$$

We now define the *first order Gauss map* g_1 as follows:

$$(5.1) \quad g_1 = \hat{F}\hat{J}_1\hat{F}^T = \mathrm{Ad}(L)(FJ_1F^T),$$

where the matrix \hat{J}_1 is the one given by (4.3) and $J_1 = \mathrm{offdiag}(1, 1, 1, 1)$. Note that $\mathrm{Ad}(L)J_1 = \hat{J}_1$. Therefore the map g_1 takes values in the space of conformal 2-spheres:

$$(5.2) \quad g_1 : M \rightarrow \mathcal{Q} \cong \mathrm{SL}_4\mathbb{R}/\hat{K}_1 = \mathrm{SL}_4\mathbb{R}/\mathrm{SO}_{1,3}^+.$$

This map g_1 is known to be a quadric which has the first order contact to the surface and it does not have the second order contact, see [16, Section 22].

We now characterize the Demoulin surface by the first-order Gauss map.

Proposition 5.1. *The first-order Gauss map g_1 of a surface \mathfrak{f} in \mathbb{RP}^3 with positive definite projective metric is conformal if and only if \mathfrak{f} is a definite Demoulin surface.*

Proof. A direct computation shows that

$$\partial_z g_1 = 2(LF) \begin{pmatrix} b\bar{P} & k & P & 0 \\ k & 0 & 0 & 1 \\ P & 0 & b & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} (LF)^T, \quad \partial_{\bar{z}} g_1 = 2(LF) \begin{pmatrix} \bar{b}P & \bar{P} & \bar{k} & 0 \\ \bar{P} & \bar{b} & 0 & 0 \\ \bar{k} & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} (LF)^T.$$

Thus

$$\langle \partial_z g_1, \partial_z g_1 \rangle = 16P, \quad \langle \partial_{\bar{z}} g_1, \partial_{\bar{z}} g_1 \rangle = 16\bar{P} \quad \text{and} \quad \langle \partial_z g_1, \partial_{\bar{z}} g_1 \rangle = \langle \partial_{\bar{z}} g_1, \partial_z g_1 \rangle = 8(k + \bar{k}) + 4|b|^2.$$

Since the coordinates (z, \bar{z}) are null for the conformal structure induced by \mathfrak{f} , the first-order Gauss map g_1 is conformal if and only if $P = 0$. \square

5.2. Demoulin surfaces and projective minimal coincidence surfaces. We set

$$G = \text{Ad}(L^{-1})\text{SL}_4\mathbb{R} = \{L^{-1}XL \mid X \in \text{SL}_4\mathbb{R}\} \subset \text{SL}_4\mathbb{C},$$

where L is defined in (2.8). The closed subgroup G is a real form of $\text{SL}_4\mathbb{C}$ which is isomorphic to $\text{SL}_4\mathbb{R}$. The space \mathcal{Q} of conformal 2-spheres is isomorphic to G/K_1 , where K_1 is

$$K_1 = \{a \in G \mid aJ_1a^T = J_1\}.$$

Let τ_1 be the outer involution on the G associated to G/K_1 given by

$$\tau_1(a) = J_1 (a^T)^{-1} J_1, \quad a \in G.$$

By abuse of notation, we denote the differential of τ_1 by the same letter τ_1 :

$$(5.3) \quad \tau_1(X) = -J_1 X^T J_1, \quad X \in \mathfrak{g}.$$

Let us consider the eigenspace decomposition of \mathfrak{g} with respect to τ_1 , that is, $\mathfrak{g} = \mathfrak{k}_1 \oplus \mathfrak{p}_1$, where \mathfrak{k}_1 is the (+1)-eigenspace and \mathfrak{p}_1 is the (-1)-eigenspace as follows:

$$\mathfrak{k}_1 = \left\{ \begin{pmatrix} a_{11} & a_{12} & a_{13} & 0 \\ a_{21} & a_{22} & 0 & -a_{13} \\ a_{31} & 0 & -a_{22} & -a_{12} \\ 0 & -a_{31} & -a_{21} & -a_{11} \end{pmatrix} \in \mathfrak{g} \right\}, \quad \mathfrak{p}_1 = \left\{ \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & -a_{11} & a_{23} & a_{13} \\ a_{31} & a_{32} & -a_{11} & a_{12} \\ a_{41} & a_{31} & a_{21} & a_{11} \end{pmatrix} \in \mathfrak{g} \right\}.$$

We decompose the Maurer-Cartan form according to this decomposition $\alpha = F^{-1}dF = Udz + Vd\bar{z}$ along the Lie algebra decomposition $\mathfrak{g} = \mathfrak{k}_1 \oplus \mathfrak{p}_1$. First we decompose U and V as

$$U = U_{\mathfrak{k}_1} + U_{\mathfrak{p}_1}, \quad V = V_{\mathfrak{k}_1} + V_{\mathfrak{p}_1}, \quad U_{\mathfrak{k}_1}, V_{\mathfrak{k}_1} \in \mathfrak{k}_1, \quad U_{\mathfrak{p}_1}, V_{\mathfrak{p}_1} \in \mathfrak{p}_1.$$

Next, set $\alpha_{\mathfrak{k}_1} = U_{\mathfrak{k}_1}dz + V_{\mathfrak{k}_1}d\bar{z}$ and $\alpha_{\mathfrak{p}_1} = U_{\mathfrak{p}_1}dz + V_{\mathfrak{p}_1}d\bar{z}$, then we obtain the expression

$$\alpha = \alpha_{\mathfrak{t}_1} + \alpha_{\mathfrak{p}_1} = U_{\mathfrak{t}_1} dz + V_{\mathfrak{t}_1} d\bar{z} + U_{\mathfrak{p}_1} dz + V_{\mathfrak{p}_1} d\bar{z},$$

where $U = U_{\mathfrak{t}_1} + U_{\mathfrak{p}_1}$ and $V = V_{\mathfrak{t}_1} + V_{\mathfrak{p}_1}$. Let us insert the spectral parameter $\lambda \in \mathbb{S}^1$ into U and V as follows:

$$U^\lambda = U_{\mathfrak{t}_1} + \lambda^{-1}U_{\mathfrak{p}_1} \quad \text{and} \quad V^\lambda = V_{\mathfrak{t}_1} + \lambda V_{\mathfrak{p}_1}.$$

Then a \mathbb{S}^1 -family of 1-forms α_λ is defined as follows:

$$(5.4) \quad \alpha^\lambda = \alpha_{\mathfrak{t}_1} + \lambda^{-1}\alpha'_{\mathfrak{p}_1} + \lambda\alpha''_{\mathfrak{p}_1} = U^\lambda dz + V^\lambda d\bar{z}.$$

Using the matrices U^λ and V^λ , they are explicitly given as follows:

$$U^\lambda = \begin{pmatrix} \bar{b}_z/(2\bar{b}) & \lambda^{-1}P & \lambda^{-1}k & \lambda^{-1}b\bar{P} \\ \lambda^{-1} & -\bar{b}_z/(2\bar{b}) & 0 & \lambda^{-1}k \\ 0 & \lambda^{-1}b & \bar{b}_z/(2\bar{b}) & \lambda^{-1}P \\ 0 & 0 & \lambda^{-1} & -\bar{b}_z/(2\bar{b}) \end{pmatrix}, \quad V^\lambda = \begin{pmatrix} b_z/(2b) & \lambda\bar{k} & \lambda\bar{P} & \lambda\bar{b}\bar{P} \\ 0 & b_z/(2b) & \lambda c & \lambda Q \\ \lambda & 0 & -b_z/(2b) & \lambda\bar{k} \\ 0 & \lambda & 0 & -b_z/(2b) \end{pmatrix}.$$

After these preparation, we obtain the following theorem.

Theorem 5.2. *Let \mathfrak{f} be a surface in \mathbb{RP}^3 with positive definite projective metric and g_1 the first-order Gauss map defined in (5.2). Moreover, let $\{\alpha^\lambda\}_{\lambda \in \mathbb{S}^1}$ be a family of 1-forms defined in (5.4). Then the following three properties are mutually equivalent:*

1. *The surface \mathfrak{f} is a definite Demoulin surface or a projective minimal coincidence surface.*
2. *The first-order Gauss map g_1 is a harmonic map into \mathcal{Q} .*
3. *$\{d + \alpha^\lambda\}_{\lambda \in \mathbb{S}^1}$ is a family of flat connections on $\mathbb{D} \times G$.*

Proof. Let us first compute the flatness

$$d\alpha^\lambda + \frac{1}{2}[\alpha^\lambda \wedge \alpha^\lambda] = 0, \quad \lambda \in \mathbb{S}^1$$

for the connection $d + \alpha^\lambda$ on $\mathbb{D} \times G$. A straightforward computation shows that $d\alpha^\lambda + \frac{1}{2}[\alpha^\lambda \wedge \alpha^\lambda] = 0$ holds for all $\lambda \in \mathbb{S}^1$ if and only if

$$(5.5) \quad P_{\bar{z}} = 0, \quad k_{\bar{z}} + k \frac{b_{\bar{z}}}{b} = 0, \quad \bar{b}P_z + 2\bar{b}_z P = 0.$$

One can see that this system implies the projective Gauss-Codazzi equations (2.6)–(2.7). In particular, the third equation is nothing but the projective minimality equation (3.1).

Every definite Demoulin surface clearly satisfies the above flatness condition (zero curvature equations) since $P = 0$.

Assume that $P \neq 0$. The first equation of (5.5) means that Pdz^2 is a holomorphic differential. From the third equation together with the holomorphicity of P , one can deduce that $(\log \bar{b})_z$ is holomorphic. Hence $(\log b/\bar{b})_{z\bar{z}} = 0$. Via the holomorphic coordinate change of z preserving the form of canonical system, we can assume that $b = \bar{b}$, *i.e.*, b is real¹.

¹The transformation rule of b under the conformal change of coordinates $w(z)$ is given by $\tilde{b} = (\bar{w}_z/w_z^2)b$, and thus $\tilde{b} = \bar{\tilde{b}}$ can be achieved by a suitable choice of the function $w(z)$ under the condition $(\log b/\bar{b})_{z\bar{z}} = 0$, see [13, Section 3].

Then (2.4) implies that $2k = b^2$. By using the second equation of (5.5), b is constant and k is a real constant. By using the third equation again, we get P is constant. This implies that $P = p$ is a non-zero constant. After these reparametrization, the canonical system is rewritten as

$$f_{zz} = bf_{\bar{z}} + pf, \quad f_{\bar{z}\bar{z}} = bf_z + pf.$$

A surface satisfying the above equation is a special case of the *coincidence surface*, [25, Example 2.19]. In fact, it is easy to see that the surface is a projective minimal coincidence surface. Thus the equivalence of the claim 1 and claim 3 follows.

The equivalence of the claims 2 and 3 follows from Proposition A.2, since the \mathbb{S}^1 -family of 1-forms α^λ is given by the involution τ_1 and it defines the semi-Riemannian symmetric space $\mathcal{Q} = \mathrm{SL}_4\mathbb{R}/K_1$. \square

Corollary 5.3. *Retaining the assumptions in Theorem 5.2, the following are equivalent:*

1. *The surface \mathfrak{f} is a definite Demoulin surface.*
2. *The first-order Gauss map g_1 is a conformal harmonic map into \mathcal{Q} .*

Proof. From Proposition 5.1, it is easy to see that the first-order Gauss map is conformal if and only if it satisfies that $P = 0$, that is, the surface is a definite Demoulin surface. Moreover, from Theorem 5.2 the Gauss map of the Demoulin surface is harmonic. \square

This corollary implies that if \mathfrak{f} is a definite Demoulin surface or a projective minimal coincidence surface, then there exists a \mathbb{S}^1 -parameter family of smooth map $F_\lambda : \mathbb{D} \times \mathbb{S}^1 \rightarrow G$ which is a solution to

$$(F_\lambda)^{-1}dF_\lambda = \alpha^\lambda$$

under initial condition $F_\lambda(z_*) = \mathrm{id}$. One can see that F_λ is regarded as a smooth map of \mathbb{D} into the following twisted loop group

$$\Lambda G_{\tau_1} = \{g : \mathbb{S}^1 \rightarrow G \mid \tau_1 g(\lambda) = g(-\lambda)\}$$

of G . The ΛG_{τ_1} -valued map F_λ is referred as to the *extended Wilczynski frame* of a definite Demoulin surface.

Precisely speaking, the extended Wilczynski frame F_λ is not the Wilczynski frame of a Demoulin surface or a projective minimal coincidence surface except for $\lambda = 1$. By conjugating F_λ by $DF_\lambda D^{-1}$ with $D = \mathrm{diag}(1, \lambda, \lambda^{-1}, 1)$, the frames $DF_\lambda D^{-1}$ give a family of Wilczynski frames for Demoulin surfaces or projective minimal coincidence surfaces. The corresponding Demoulin surfaces or projective minimal coincidence surfaces have the same projective metric $8|b|^2 dzd\bar{z}$ but the different conformal classes of cubic forms $\lambda^{-3}bdz^3$. Moreover, the functions P changes as $\lambda^{-2}P$.

6. PRIMITIVE LIFTS

We now show that the extended Wilczynski frame for a Demoulin surface has an additional order three cyclic symmetry. Let σ be an order three automorphism on the complexification $\mathrm{SL}_4\mathbb{C}$ of G as follows:

$$\sigma X = \mathrm{Ad}(E)X, \quad X \in \mathrm{SL}_4\mathbb{C},$$

where $E = \text{diag}(1, \epsilon^2, \epsilon, 1)$ with $\epsilon = e^{2\pi\sqrt{-1}/3}$. It should be emphasized that σ preserves the real form G . Thus σ is regarded as an automorphisms of G .

Next, one can check that F_λ satisfies the symmetry $\sigma(F_\lambda) = F_{\epsilon\lambda}$, since U^λ and V^λ satisfy the same symmetry. It is also easy to see that τ_1 and σ commute, and $\kappa = \tau_1 \circ \sigma$ defines an automorphism of order six. We obtain a regular semi-Riemannian 6-symmetric space G/K (see Appendix A.1), where

$$(6.1) \quad K = \{\text{diag}(k_1, k_2, k_2^{-1}, k_1^{-1}) \mid k_1 \in \mathbb{R}^\times, k_2 \in \mathbb{S}^1\} \cong \text{SO}_{1,1} \times \text{SO}_2.$$

Note that G/K is identified with $\{gJg^T \mid g \in G\}$, where $J = EJ_1$. There is a homogeneous projection

$$\pi : G/K \rightarrow G/K_1; \quad gK \longmapsto gK_1.$$

The extended Wilczynski frame F_λ satisfies the symmetry

$$\kappa(F_\lambda) = F_{-\epsilon\lambda}.$$

Note that $-\epsilon$ is the 6th root of unity. From the above argument, it is easy to see that the extended Wilczynski frame $F_\lambda = F(\lambda)$ for a Demoulin surface is an element of the twisted loop group of G :

$$\Lambda G_\kappa = \{g : \mathbb{S}^1 \rightarrow G \mid \kappa g(\lambda) = g(-\epsilon\lambda)\}.$$

Theorem 6.1. *The first-order Gauss map of a Demoulin surface, which is conformal harmonic into $\mathcal{Q} = G/K_1$, can be obtained by the homogeneous projection of a primitive map into the regular semi-Riemannian 6-symmetric space $G/K \cong \text{SL}_4\mathbb{R}/\text{SO}_{1,1} \times \text{SO}_2$.*

Proof. The 0th-eigenspace $\mathfrak{g}_0^{\mathbb{C}}$ and ± 1 st-eigenspaces $\mathfrak{g}_{\pm 1}^{\mathbb{C}}$ of the derivative of the order six automorphism $\kappa = \tau_1 \circ \sigma$ are described as follows:

$$\mathfrak{g}_0^{\mathbb{C}} = \{\text{diag}(a_{11}, a_{22}, -a_{22}, -a_{11}) \mid a_{11} \in \mathbb{R}, a_{22} \in \mathbb{C}\},$$

and

$$\mathfrak{g}_{-1}^{\mathbb{C}} = \left\{ \left(\begin{array}{cccc} 0 & 0 & a_{13} & 0 \\ a_{21} & 0 & 0 & a_{13} \\ 0 & a_{32} & 0 & 0 \\ 0 & 0 & a_{21} & 0 \end{array} \right) \middle| a_{ij} \in \mathbb{C} \right\}, \quad \mathfrak{g}_1^{\mathbb{C}} = \left\{ \left(\begin{array}{cccc} 0 & a_{12} & 0 & 0 \\ 0 & 0 & a_{23} & 0 \\ a_{31} & 0 & 0 & a_{12} \\ 0 & a_{31} & 0 & 0 \end{array} \right) \middle| a_{ij} \in \mathbb{C} \right\}.$$

From the matrices U^λ and V^λ in (5.4) with $P = 0$, we see that the condition in Definition A.1 of primitive map is satisfied. The stabilizer of κ is the closed subgroup K given by (6.1). Therefore there is a primitive map $g = FJF^T$ $J = EJ_1$ into the 6-symmetric space G/K such that $\pi \circ g = \text{Ad}(L^{-1})g_1$. Since $\text{Ad}(L^{-1}) : \text{SL}_4\mathbb{R}/\hat{K}_1 \rightarrow G/K_1$ is an isometry, $g_1 = \text{Ad}(L)(\pi \circ g)$ is harmonic. \square

This theorem enable us to establish a generalized Weierstrass type representation for definite Demoulin surfaces by virtue of primitive maps into the semi-Riemannian 6-symmetric space G/K , see [12].

Remark 6.2. Corresponding result theorem for indefinite Demoulin surfaces was obtained by the second named author in the preprint version of [15].

APPENDIX A. PRIMITIVE HARMONIC MAPS

A.1. Homogeneous geometry. Let G be a semi-simple real Lie group with automorphism τ of order $k \geq 2$. We consider a reductive homogenous space G/K equipped with a G -invariant semi-Riemannian metric satisfying the following three conditions:

- The closed subgroup H satisfies $G_\tau^\circ \subset K \subset G_\tau$. Here G_τ is the Lie subgroup of all fixed points of τ and G_τ° the identity component of it.
- The G -invariant semi-Riemannian metric is derived from (a constant multiple of) the Killing form of G .
- The Lie algebra \mathfrak{k} of K is *non-degenerate* with respect to the induced scalar product.

The resulting homogeneous semi-Riemannian space G/K is called a regular *semi-Riemannian k -symmetric space*. Note that a regular semi-Riemannian 2-symmetric spaces is just a semi-Riemannian symmetric space. Since \mathfrak{k} is non-degenerate, the orthogonal complement \mathfrak{p} of \mathfrak{k} is non-degenerate and can be identified with the tangent space of G/K at the origin $o = K$. The Lie algebra \mathfrak{g} is decomposed into the direct sum:

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}.$$

of linear subspaces.

We denote the induced Lie algebra automorphism of \mathfrak{g} by the same letter τ . Now we have the eigenspace decomposition of the complexified Lie algebra $\mathfrak{g}^\mathbb{C}$;

$$\mathfrak{g}^\mathbb{C} = \sum_{j \in \mathbb{Z}_k} \mathfrak{g}_j^\mathbb{C},$$

where $\mathfrak{g}_j^\mathbb{C}$ is the eigenspace of τ with eigenvalue ω^j . Here ω is the (primitive) k -th root of unity. In particular, $\mathfrak{g}_0^\mathbb{C} = \mathfrak{k}^\mathbb{C}$ and $\mathfrak{g}_{-1}^\mathbb{C} = \overline{\mathfrak{g}_1^\mathbb{C}}$. Let us define a subbundle $[\mathfrak{g}_j^\mathbb{C}]$ of $G/K \times \mathfrak{g}$ by

$$[\mathfrak{g}_j^\mathbb{C}]_{g \cdot o} = \text{Ad}(g)\mathfrak{g}_j^\mathbb{C}.$$

Then the complexified tangent bundle $T^\mathbb{C}G/K$ is expressed as

$$T^\mathbb{C}G/K = \sum_{j \in \mathbb{Z}_k, j \neq 0} [\mathfrak{g}_j^\mathbb{C}].$$

A.2. Primitive maps. A smooth map $\psi : \Sigma \rightarrow N$ of a Riemann surface Σ into a semi-Riemannian manifold N is said to be a *harmonic map* if its *tension field* $\text{tr}(\nabla d\psi)$ vanishes.

For smooth maps into regular semi-Riemannian k -symmetric spaces with $k > 2$, the notion of primitive map was introduced by Burstall-Pedit [5] (see also Bolton-Pedit-Woodward [2]).

Definition A.1. Let $\psi : \Sigma \rightarrow G/K$ be a smooth map of a Riemann surface Σ into a regular semi-Riemannian k -symmetric space with $k > 2$. Then ψ is said to be a *primitive map* if $d\psi(T'\Sigma) \subset [\mathfrak{g}_{-1}^\mathbb{C}]$. Here $T'\Sigma$ denotes the $(1,0)$ -tangent bundle of Σ .

Black [1] showed that primitive maps are *equi-harmonic*, that is, harmonic with respect to suitable invariant metrics on G/K (see also [5]). In addition primitive maps well behave with respect to homogeneous projections [5, Theorem 3.7].

Theorem A.1. *Let H be a closed subgroup of G satisfying*

- $K \subset H$.
- The Lie algebra \mathfrak{h} of H is non-degenerate.
- The decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}$ is reductive and stable under τ . Here \mathfrak{q} is the orthogonal complement of \mathfrak{h} .

Denote by $\pi_H : G/K \rightarrow G/H$ be the homogenous projection. Then for any primitive map ψ , $\pi_H \circ \psi$ is a harmonic map into G/H .

Note that when $k = 2$, $[\mathfrak{g}_{-1}^{\mathbb{C}}] = T^{\mathbb{C}}G/K$ and the primitivity condition is vacuous. On the other hand when $k > 2$, every primitive map is harmonic with respect to the Killing metric. To provide a unified description, we recall the following terminology from [6].

Definition A.2. A smooth map $\psi : \Sigma \rightarrow G/K$ into a regular semi-Riemannian k -symmetric space is said to be a *primitive harmonic map* if it is primitive for $k > 2$ and harmonic if $k = 2$.

Now let $\psi : \mathbb{D} \rightarrow G/H$ be a smooth map from a simply connected Riemann surface \mathbb{D} into a regular semi-Riemannian k -symmetric space G/K with $k \geq 2$. Take a frame $\Psi : \mathbb{D} \rightarrow G$ of ψ and put $\alpha := \Psi^{-1}d\Psi$. Then we have the identity (*Maurer-Cartan equation*):

$$d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0.$$

Decompose α along the Lie algebra decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ as

$$\alpha = \alpha_{\mathfrak{k}} + \alpha_{\mathfrak{p}}, \quad \alpha_{\mathfrak{k}} \in \mathfrak{k}, \quad \alpha_{\mathfrak{p}} \in \mathfrak{p}.$$

We decompose $\alpha_{\mathfrak{p}}$ with respect to the conformal structure of \mathbb{D} as

$$\alpha_{\mathfrak{p}} = \alpha'_{\mathfrak{p}} + \alpha''_{\mathfrak{p}}.$$

Here $\alpha'_{\mathfrak{p}}$ and $\alpha''_{\mathfrak{p}}$ are the $(1, 0)$ and $(0, 1)$ -part of $\alpha_{\mathfrak{p}}$, respectively. Since G is a real Lie group, $\alpha''_{\mathfrak{p}}$ is the conjugate of $\alpha'_{\mathfrak{p}}$.

Now let us assume that ψ is a primitive harmonic map, then $\alpha'_{\mathfrak{p}}$ is $[\mathfrak{g}_{-1}^{\mathbb{C}}]$ -valued and $\alpha''_{\mathfrak{p}}$ is $[\mathfrak{g}_1^{\mathbb{C}}]$ -valued, respectively. Hence the decomposition of α is rewritten as

$$\alpha = \alpha'_{-1} + \alpha_0 + \alpha'_1.$$

Now let us introduce a spectral parameter $\lambda \in \mathbb{S}^1$ into α as

$$\alpha^\lambda := \alpha_0 + \lambda^{-1}\alpha'_{-1} + \lambda\alpha''_1.$$

We arrive at the *zero curvature representation* of primitive harmonic maps:

Proposition A.2. *Let \mathbb{D} be a connected open subset of \mathbb{C} . Let $\psi : \mathbb{D} \rightarrow G/K$ be a primitive harmonic map. Then the loop of connections $d + \alpha^\lambda$ is flat for all λ , that is,*

$$d\alpha^\lambda + \frac{1}{2}[\alpha^\lambda \wedge \alpha^\lambda] = 0$$

for all λ .

Conversely assume that \mathbb{D} is simply connected. Let $\alpha^\lambda = \alpha_0 + \lambda^{-1}\alpha'_{-1} + \lambda\alpha''_1$ be an \mathbb{S}^1 -family of \mathfrak{g} -valued one-forms which satisfies

$$d\alpha^\lambda + \frac{1}{2}[\alpha^\lambda \wedge \alpha^\lambda] = 0$$

for all $\lambda \in \mathbb{S}^1$. Then there exists a one-parameter family of maps $\Psi_\lambda : \mathbb{D} \rightarrow G$ such that

$$\Psi_\lambda^{-1}d\Psi_\lambda = \alpha^\lambda,$$

and

$$\psi_\lambda = \Psi_\lambda \bmod K : \mathbb{D} \rightarrow G/K$$

is primitive harmonic for all λ .

APPENDIX B. PROJECTIVE MINIMAL SURFACES AND THE CONFORMAL GAUSS MAPS

B.1. Conformal Gauss map. Let $\mathfrak{f} : M \rightarrow \mathbb{RP}^3$ be a surface with Wilczynski frame F as in subsection 5.1. We define a map g_2 by

$$g_2 = \hat{F}\hat{J}_2\hat{F}^T = -\text{Ad}(L)(FJ_2F^T), \quad \hat{J}_2 = -LJ_2L^T,$$

where $J_2 = \text{offdiag}(1, -1, -1, 1)$ (cf. [24, §4.1]). Analogous to the first-order Gauss map g_1 , g_2 takes value in the space \mathcal{Q} of conformal 2-spheres in \mathbb{RP}^3 . More precisely, since the matrix \hat{J}_2 is of signature $(1, 3)$, it is a point of \mathcal{Q} . Thus \mathcal{Q} is realized as a homogeneous space $\text{SL}_4\mathbb{R}/\hat{K}_2$, where \hat{K}_2 is the stabilizer at $\hat{J}_2 \in \mathcal{Q}$ explicitly given by $\hat{K}_2 = \{X \in \text{SL}_4\mathbb{R} \mid XJ_2X^T = J_2\}$, which is also isomorphic to $\text{SO}_{1,3}^+$. Thus the map g_2 takes value in $\text{SL}_4\mathbb{R}/\hat{K}_2$:

$$(B.1) \quad g_2 : M \rightarrow \mathcal{Q} \cong \text{SL}_4\mathbb{R}/\hat{K}_2 = \text{SL}_4\mathbb{R}/\text{SO}_{1,3}^+.$$

This map g_2 is known to be a *Lie quadric* which has the second order contact to the surface, see [16, Section 18]. The map g_2 has been called the *conformal Gauss map* for a surface \mathfrak{f} in \mathbb{RP}^3 , see [29, 4]. In [18], the conformal Gauss map g_2 was called the *projective Gauss map*. In classical literature, g_2 was called the *congruence of Lie quadrics*.

Proposition B.1 (Theorem 3 in [4]). *The conformal Gauss map g_2 is conformal map.*

Proof. As in the proof of Proposition 5.1, a direct computation shows that

$$\partial_z g_2 = -2(LF) \text{diag}(b\bar{P}, 0, -b, 0)(LF)^T \quad \text{and} \quad \partial_{\bar{z}} g_2 = -2(LF) \text{diag}(\bar{b}P, -\bar{b}, 0, 0)(LF)^T.$$

Thus

$$\langle \partial_z g_2, \partial_z g_2 \rangle = \langle \partial_{\bar{z}} g_2, \partial_{\bar{z}} g_2 \rangle = 0 \quad \text{and} \quad \langle \partial_z g_2, \partial_{\bar{z}} g_2 \rangle = 4|b|^2 \neq 0.$$

Since the coordinates (z, \bar{z}) are null for the conformal structure induced by \mathfrak{f} , the conformal Gauss map g_2 is conformal. \square

Remark B.2. *The Hodge star operator \star on $\wedge^2\mathbb{R}^{1,3}$ is introduced by*

$$\langle a, b \rangle = \Omega(a \wedge \star b).$$

Since $\langle \cdot, \cdot \rangle$ is Lorentzian, \star satisfies $\star^2 = -1$. Thus the complexification $(\wedge^2 \mathbb{R}^{1,3})^{\mathbb{C}} \cong \wedge^2 \mathbb{C}^{1,3}$ has the eigenspace decomposition

$$(\wedge^2 \mathbb{R}^4)^{\mathbb{C}} = \mathbb{S} \oplus \bar{\mathbb{S}},$$

where \mathbb{S} is the $\sqrt{-1}$ -eigenspace of \star . In this way, a quadric $Q \in \mathcal{Q}$ corresponds to a complex linear subspace \mathbb{S} of $(\wedge^2 \mathbb{R}^4)^{\mathbb{C}}$. The correspondence $Q \mapsto \mathbb{S}$ defines a smooth bijection from the space \mathcal{Q} of conformal 2-spheres in $\mathbb{R}\mathbb{P}^3$ to the space

$$\mathcal{G}_{2,0}^{3,3} = \{\mathbb{S} \subset (\mathbb{R}^{3,3})^{\mathbb{C}} \mid \mathbb{S} \cap \mathbb{S}^{\perp} = \{0\}, \bar{\mathbb{S}} = \mathbb{S}^{\perp}\}.$$

Under this identification, g_2 is regarded as a smooth map into $\mathcal{G}_{2,0}^{3,3}$ in [4, p. 183], [7, p. 30].

B.2. Projective minimal surfaces and the conformal Gauss maps. The space \mathcal{Q} of conformal 2-spheres in $\mathbb{R}\mathbb{P}^3$ is isomorphic to the semi-Riemannian symmetric space G/K_2 , where

$$K_2 = \{a \in G \mid aJ_2a^T = J_2\}.$$

Let τ_2 be the outer involution on G associated to the symmetric space G/K_2 defined by:

$$\tau_2(a) = J_2 (a^T)^{-1} J_2,$$

where $a \in G$. By abuse of notation, we denote the differential of τ_2 by the same letter τ_2 which is an outer involution on \mathfrak{g} :

$$(B.2) \quad \tau_2(X) = -J_2 X^T J_2,$$

where $X \in \mathfrak{g}$. Let us consider the eigenspace decomposition of \mathfrak{g} with respect to τ_2 , that is, $\mathfrak{g} = \mathfrak{k}_2 \oplus \mathfrak{p}_2$, where \mathfrak{k}_2 is the $(+1)$ -eigenspace and \mathfrak{p}_2 is the (-1) -eigenspace as follows:

$$\mathfrak{k}_2 = \left\{ \begin{pmatrix} a_{11} & a_{12} & a_{13} & 0 \\ a_{21} & a_{22} & 0 & a_{13} \\ a_{31} & 0 & -a_{22} & a_{12} \\ 0 & a_{31} & a_{21} & -a_{11} \end{pmatrix} \in \mathfrak{g} \right\}, \quad \mathfrak{p}_2 = \left\{ \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & -a_{11} & a_{23} & -a_{13} \\ a_{31} & a_{32} & -a_{11} & -a_{12} \\ a_{41} & -a_{31} & -a_{21} & a_{11} \end{pmatrix} \in \mathfrak{g} \right\}.$$

According to this decomposition $\mathfrak{g} = \mathfrak{k}_2 \oplus \mathfrak{p}_2$, the Maurer-Cartan form $\alpha = F^{-1}dF = Udz + Vd\bar{z}$ can be decomposed into

$$\alpha = \alpha_{\mathfrak{k}_2} + \alpha_{\mathfrak{p}_2} = U_{\mathfrak{k}_2}dz + V_{\mathfrak{k}_2}d\bar{z} + U_{\mathfrak{p}_2}dz + V_{\mathfrak{p}_2}d\bar{z},$$

where $U = U_{\mathfrak{k}_2} + U_{\mathfrak{p}_2}$ and $V = V_{\mathfrak{k}_2} + V_{\mathfrak{p}_2}$. Let us insert the parameter $\lambda \in \mathbb{S}^1$ into U and V in a manner similar to section 5.2:

$$U^\lambda = U_{\mathfrak{k}_2} + \lambda^{-1}U_{\mathfrak{p}_2} \quad \text{and} \quad V^\lambda = V_{\mathfrak{k}_2} + \lambda V_{\mathfrak{p}_2}.$$

Then a family of 1-forms α^λ is defined as follows:

$$(B.3) \quad \alpha^\lambda = \alpha_{\mathfrak{k}_2} + \lambda^{-1}\alpha'_{\mathfrak{p}_2} + \lambda\alpha''_{\mathfrak{p}_2} = U^\lambda dz + V^\lambda d\bar{z}.$$

In fact the matrices U^λ and V^λ are explicitly given as follows:

$$(B.4) \quad U^\lambda = \begin{pmatrix} \frac{\bar{b}_z}{2b} & P & k & \lambda^{-1}b\bar{P} \\ 1 & -\frac{\bar{b}_z}{2b} & 0 & k \\ 0 & \lambda^{-1}b & \frac{\bar{b}_z}{2b} & P \\ 0 & 0 & 1 & -\frac{\bar{b}_z}{2b} \end{pmatrix}, \quad V^\lambda = \begin{pmatrix} \frac{b_z}{2b} & \bar{b} & \bar{P} & \lambda\bar{b}P \\ 0 & \frac{b_z}{2b} & \lambda\bar{b} & \bar{P} \\ 1 & 0 & -\frac{b_z}{2b} & \bar{k} \\ 0 & 1 & 0 & -\frac{b_z}{2b} \end{pmatrix}.$$

Then the projective minimal surface can be characterized by the harmonicity of the conformal Gauss map [29], [4, Theorem 7], and by a family of flat connections.

Theorem B.3 ([29], [4]). *Let \mathfrak{f} be a surface in \mathbb{RP}^3 and g_2 the conformal Gauss map defined in (B.1). Moreover, let $\{\alpha^\lambda\}_{\lambda \in \mathbb{S}^1}$ be a family of 1-forms defined in (B.3). Then the following are mutually equivalent:*

1. *The surface \mathfrak{f} is a projective minimal surface.*
2. *The conformal Gauss map g_2 is a conformal harmonic map into \mathcal{Q} .*
3. *$\{\alpha^\lambda\}_{\lambda \in \mathbb{S}^1}$ is a family of flat connections on $\mathbb{D} \times G$.*

Proof. Let us compute the flatness conditions of $d + \alpha^\lambda$, that is, the Maurer-Cartan equation $d\alpha^\lambda + \frac{1}{2}[\alpha^\lambda \wedge \alpha^\lambda] = 0$. It is easy to see that except for the (1, 4)-entry, the Maurer-Cartan equation is equivalent to (2.6). Moreover, the λ^{-1} -term and the λ -term of the (1, 4)-entry are equivalent to that the first equation and the second equation in (3.1), respectively. Thus the equivalence of (1) and (3) follows.

The equivalence of (2) and (3) follows from Proposition A.2, since the family of 1-forms α^λ is given by the involution τ_2 and it defines the semi-Riemannian symmetric space $\mathcal{Q} = \mathrm{SL}_4\mathbb{R}/K_2$. \square

Remark B.4. The above theorem implies that if \mathfrak{f} is a projective minimal surface, then there exists a family of projective minimal surface \mathfrak{f}^λ ($\lambda \in \mathbb{S}^1$) such that $\mathfrak{f}^\lambda|_{\lambda=1} = \mathfrak{f}$. Projective minimal surfaces of the family have the same projective metric $8|b|^2 dzd\bar{z}$ but the different conformal classes of cubic forms $\lambda^{-1}b dz^3$.

REFERENCES

- [1] M. Black, *Harmonic maps into Homogeneous Spaces*, Pitman Research Notes in Mathematics Series **255**, 1991.
- [2] J. Bolton, F. Pedit, L. M. Woodward, Minimal surfaces and the affine Toda field model, *J. Reine Angew. Math.* **459** (1995), 119–150.
- [3] J. Bolton, L. M. Woodward, Congruence theorems for harmonic maps from a Riemann surface into $\mathbb{C}P^n$ and S^n , *J. London Math. Soc.* (2) **45** (1992), 363–376.
- [4] F. E. Burstall, U. Hertrich-Jeromin, Harmonic maps in unfashionable geometries, *Manuscripta. Math.* **108** (2002), 171–189.
- [5] F. E. Burstall, F. Pedit, Harmonic maps via Adler-Kostant-Symes theory, in: *Harmonic Maps and Integrable Systems* (A. P. Fordy, J. C. Wood eds.), Aspects of Mathematics **E23** (1994), Vieweg, pp. 221–272.
- [6] F. E. Burstall, F. Pedit, Dressing orbits of harmonic maps, *Duke Math. J.* **80** (1985), 353–382.
- [7] D. J. Clarke, *Integrability in Submanifold Geometry*, Thesis, Univ. Bath., 2012.

- [8] S. S. Chern, J. G. Wolfson, Harmonic maps of the two-sphere into a complex Grassmannian manifolds II, *Ann. of Math. (2)* **125** (1987), 301–335.
- [9] G. Darboux, *Leçons sur la théorie générale des surfaces* I, 1914, II, 1915, second edition, Gauthier-Villars.
- [10] A. Demoulin, Sur deux transformations des surfaces dont les quadriques de Lie n'ont que deux ou trois points caractéristiques, *Bull de l'Acad Belgique* **19** (1933), 479–502, 579–592, 1352–1363.
- [11] J. F. Dorfmeister, W. Freyn, S.-P. Kobayashi, E. Wang, Survey on real forms of the complex $A_2^{(2)}$ -Toda equation and surface theory. *Complex Manifolds* **6** (2019), 194–227.
- [12] J. Dorfmeister, I. McIntosh, F. Pedit, H. Wu, On the meromorphic potential for a harmonic surface in a k -symmetric space, *Manuscripta Math.* **92** (1997) no. 2, 143–152.
- [13] F. E. Ferapontov, Integrable systems in projective differential geometry, *Kyushu J. Math.* **54** (2000), no. 1, 183–215.
- [14] H. S. Hu, Darboux transformations of Su-chain, in: *Differential Geometry. Proceedings of the Symposium in honor of Professor Su Buchin on his 90th Birthday* (C. H. Gu, H. S. Hu, Y. L. Xin eds.), World Scientific, 1993, pp. 325–380.
- [15] S.-P. Kobayashi, A loop group method for Demoulin surfaces in the 3-dimensional real projective space, *Differential Geom. Appl.*, **40** (2015), 57–66 ([arXiv:1301.6325v2\[math.DG\]](https://arxiv.org/abs/1301.6325v2))
- [16] E. P. Lane, *Projective Differential Geometry of Curves and Surfaces*, University of Chicago Press, 1932.
- [17] C. LeBrun, L. J. Mason, Nonlinear gravitons, null geodesics, and holomorphic disks, *Duke Math. J.* **136** (2007), 205–273.
- [18] E. Musso, L. Nicolodi, Tableaux over Lie algebras, integrable systems, and classical surface theory, *Comm. Anal. Geom.* **14** (2006), 475–496.
- [19] K. Nomizu, T. Sasaki, Centroaffine immersions of codimension two and projective hypersurface theory, *Nagoya Math. J.* **132** (1993), 63–90.
- [20] K. Nomizu, T. Sasaki, *Affine Differential Geometry. Geometry of Affine Immersions*, Cambridge Tracts in Mathematics **11** (1994), Cambridge University Press.
- [21] C. Pa (Zhengguo Bai), A new definition of the Godeaux sequence of quadrics, *Amer. J. Math.* **69** (1947), 117–120.
- [22] C. Pa (Zhengguo Bai), On the surfaces whose asymptotic curves of one system are projectively equivalent, *Univ. Nac. Tucumán Revista* **A3** (1942), 341–349.
- [23] G. Pick, Über affine Geometrie IV. Differentialinvarianten der Flächen gegenüber affinen Transformationen, *Berichte Verh Ges. Wiss. Leipzig* **69** (1917), 107–136.
- [24] T. Sasaki, *Projective Differential Geometry and Linear Homogenous Differential Equations*, Rokko Lectures in Math. **5** (1999), Kobe Univ.
- [25] T. Sasaki, Line congruence and transformation of projective surfaces, *Kyushu J. Math.* **60** (2006), 101–243.
- [26] B. Su, On certain periodic sequence of Laplace of period four in ordinary space, *Sci. Rep. Tohoku Imperial Univ.* **25** (1936), 227–256.
- [27] B. Su, On certain couples of closed Laplace sequences of period four in ordinary space, *Chinese Math.* **5** (1964), 151–174.
- [28] B. Su, The growth and development of differential geometry in China. (Japanese) *Sūgaku* **35** (1983), no. 3, 221–228.
- [29] G. Thomsen, Sulle superficie minime proiettive, *Annali Mat.* **5** (1928), 169–184
- [30] E. J. Wilczynski, *Projective Differential Geometry of Curves and Ruled Surfaces*, Teubner, 1906.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF TSUKUBA, TSUKUBA, 305-8571, JAPAN
E-mail address: `inoguchi@math.tsukuba.ac.jp`

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO, 060-0810, JAPAN
E-mail address: `shimpei@math.sci.hokudai.ac.jp`