
| Title | Serotonergic control in initiating defensive responses to unexpected tactile stimuli in the trap-jaw ant Odontomachus kuroiwae |
|------------------------|--|
| Author(s) | Aonuma, Hitoshi |
| Citation | Journal of experimental biology, 223(19), jeb228874 https://doi.org/10.1242/jeb.228874 |
| Issue Date | 2020-10 |
| Doc URL | http://hdl.handle.net/2115/82905 |
| Туре | article (author version) |
| Additional Information | There are other files related to this item in HUSCAP. Check the above URL. |
| File Information | S4_20200709.pdf (Supplementary Information (Fig S2)) |





S4. Amount of precursor and catabolites of biogenic amines in the brain. Box-and-whisker graphs indicate minimum, median, maximum, 25% percentile and 75% percentile. **A:** Amount of 5HTP in the brain. There was no significant difference between turn and dart (Unpaired t test with Welch's correction: p = 0.21). **B:** Amount of Nac-5HT in the brain. There was no significantly difference between turn and dart (Unpaired t test with Welch's correction: p = 0.18). **C:** Amount of Nac-DA in the brain. There was no significant difference between turn and dart (Unpaired t test with Welch's correction: p = 0.80). **D:** Amount of Nac-TA in the brain. There was no significant difference between turn and dart (Unpaired t test with Welch's correction: p = 0.75).