Title	Rapid detection of rifampicin-resistant Mycobacterium tuberculosis, based on isothermal DNA amplification and DNA chromatography
Author(s)	Takarada, Yutaka; Koder, Takuya; Kobayashi, Kumi; Nakajima, Chie; Kawase, Mitsuo; Suzuki, Yasuhiko
Citation	Journal of microbiological methods, 177, 106062 https://doi.org/10.1016/j.mimet.2020.106062
Issue Date	2020-10
Doc URL	http://hdl.handle.net/2115/83154
Rights	© 2020. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0
Type	article (author version)
File Information	MS-Takarada-1_Main Text_2020.09.14R2.pdf

[Instructions for use](#)

Title: Rapid detection of rifampicin-resistant *Mycobacterium tuberculosis*, based on isothermal DNA amplification and DNA chromatography

Authors;

Yutaka Takarada ^{1),*,§}, Takuya Kodera ^{1),§}, Kumi Kobayashi ¹⁾, Chie Nakajima^{2),3)}, Mitsuo Kawase¹⁾, Yasuhiko Suzuki ^{2),3),*}

1) TBA Co., LTD, T-Biz 307, 6-6-40 Aramaki-Aza-Aoba, Aoba-ku, Sendai, 980-8579 Japan

2) Hokkaido University Research Center for Zoonosis Control, Sapporo, Japan

3) Hokkaido University, GI-CoRE Global Station for Zoonosis Control, Sapporo, 001-0020 Japan.

E-mail addresses:

y-takara@t-bioarray.com , t-kodera@ecei.tohoku.ac.jp , k-kobaya@t-bioarray.com , cnakajim@czc.hokudai.ac.jp, m-kawase@ecei.tohoku.ac.jp, suzuki@czc.hokudai.ac.jp

[§]YT and TK equally contributed to this work.

*Corresponding author:

Yutaka Takarada

TBA Co.,LTD, T-Biz 307, 6-6-40 Aramaki-Aza-Aoba, Aoba-ku, Sendai, 980-8579 Japan

E-mail: y-takara@t-bioarray.com

Telephone: +81-22-795-7204; FAX: +81-22-795-7204

27 Co-corresponding author:
28 Yasuhiko Suzuki
29 Kita 20, Nishi 10, Kita-ku, Sapporo, 001-0020, Japan
30 E-mail: suzuki@czc.hokudai.ac.jp
31 Telephone: +81-11-706-9503; FAX: +81-11-706-7310
32
33

Abstract

Rapid and easy detection of nucleotide point mutations in bacterial pathogens associated with drug resistance is essential for the proper use of antimicrobials. Here, we developed a rapid and simple method for the detection of mutations using Loop-mediated isothermal amplification (LAMP) combined with the single-tag hybridization (STH) chromatographic printed array strips (PAS) method. This procedure is able to detect four mutations (C1349T, A1295C, G1303T, A1304T) in Rifampicin Resistance Determining Region (RRDR) of rifampicin-resistant *Mycobacterium tuberculosis* (RR-TB), simultaneously.

LAMP reactions contained a LAMP primer and eight allele-specific primers for each mutation. The allele-specific primers products were detected by nucleic acid chromatography using PAS. Four detection lines were detected there, one of which was detected at different positions depend on the wild type and the mutant type. We carried out the four mutations detection using 31 genomic DNA (2 A1295T, 1 G1303T, 6 A1304T, 22 C1349T) from clinical isolate. The mutations have been confirmed by sequence analysis. The detection results were completely consistent with the sequence analysis. In the present study, four mutations could be detected, but only 60% of RR-TB could be detected with these four. It is expected that the detection rate will increase by adding more mutant primers.

The combined LAMP and STH chromatographic PAS method is a simple and rapid method for detecting point mutations in clinical isolates as a point-of-care testing (POCT) technique. In addition, it does not require special equipment and can meet the demand in areas where drug-resistant bacteria are endemic, such as developing countries.

Keywords

- 60 *Mycobacterium tuberculosis*, Rapid detection, Loop-mediated isothermal amplification,
61 DNA-Chromatography, Mutation detection
62

Introduction

Tuberculosis (TB) caused by *Mycobacterium tuberculosis* (MTB) is an infectious disease that affects more than 10 million people worldwide every year, mainly in developing countries (WHO, 2019). In addition, MTB is still a global leading cause of death (WHO, 2019). Although in general TB cases have gradually decreased annually, the incidence of MTB resistant to rifampicin and multi-drug treatments (RR-TB/MDR-TB) have increased five-fold in the past decade (WHO, 2019). Therefore, measures are taken to prevent drug-resistant tuberculosis from spreading. Of special importance is that diagnosis and treatments be properly conducted in developing countries.

For the treatment of MTB, rifampicin is the first-line drug in a combination supported by 3-4 other drugs. Therefore, the presence or absence of rifampicin resistance is a very important factor in appropriate TB treatment. Approximately 78-100% of rifampicin-resistant MTB (RR-MTB) has mutations in the 81 bp region (Rifampicin Resistance Determining Region: RRDR) of the RNA polymerase subunit β (*rpoB*) gene (Telenti et al., 1993; Rahim et al., 2012; Aye et al., 2016).

Typically, diagnosis of mycobacterial infections, including drug-resistance MTB, is carried out by culturing, but these bacteria are slow growing and thus culturing is time-consuming. Therefore, a method to quickly identify drug resistance is needed.

Molecular diagnostic methods can quickly identify drug resistance, which can help expedite the development of new drugs for the treatment of drug-resistant MTB.

The World Health Organization (WHO) recommended Xpert MTB/RIF® assay as rapid diagnostic test for the detection of TB and RR-TB (WHO, 2017). This assay is a single-use sample-processing cartridge system with integrated multicolor real-time PCR to detect majority of RR associated mutations. WHO states that the system requires no additional experimental equipment and is suitable for use at all levels of the health system, but because of its sophisticated system, it requires a stable, uninterrupted

electrical supply. And, this system cannot be used to monitor treatment and requires traditional cultures and drug susceptibility test to detect resistance to anti-tuberculosis drugs other than rifampicin. In addition, this system has low throughput, it was unable to meet the demands from high TB endemic regions (Tam et al., 2017).

Loop-mediated isothermal amplification (LAMP) is an isothermal amplification method that uses four primers and a DNA polymerase with strand displacement activity (Notomi et al., 2000; Mori et al., 2001; Nagamine et al., 2001). The four primers recognize six regions and therefore show very high specificity compared to conventional PCR. The amplification product has a structure with several loops and an inverted repeat sequence. Additional primers designed to the loop position, the efficiency and specificity can be further increased. Since reaction can be carried out at a fixed temperature, the reaction products can be detected by the naked eye, no expensive instruments are needed, LAMP is recognized as a powerful tool for point-of-care genetic testing. Detection of mutations using LAMP has been reported, which was either allele-specific LAMP (Badolo et al., 2012; Badolo et al., 2015) or PNA clamp method using a PNA probe (Itonaga et al., 2016; Sakai et al., 2017). Nonetheless, multi-mutations were difficult to detect simultaneously by these reported methods.

The single-tag hybridization (STH) chromatographic printed array strip (PAS) method is also an innovative multi-detection method. The PCR products using primer-attached tag sequences and primers with biotin are mixed with streptavidin-bound colored beads. The resulting mixture then binds chromatographically to the anti-tag probe on the strip. Different anti-tag probes are located at different sites on the strip, so that multi-detection is possible. In this method, individual amplicons of multiplex PCR products can be easily detected with high sensitivity within 10-15 min, with the naked eye (Monden et al., 2014; Tian et al., 2014; Tian et al., 2016; Saito et al., 2018). As for the detection of drug resistant genes using this method, there are cases in which the

mecA and *blaZ* of the *Staphylococcus aureus* were detected in blood culture samples (Ohshiro et al., 2016) and the carbapenemase gene of carbapenem-resistant bacterium was detected in fecal samples (Shanmugakani et al., 2017). More recently, the gene for pork in beef meatballs (Riztyan et al., 2018) and the gene for genetically modified crops in soybeans (Takabatake et al., 2018) were detected.

In the present study, we developed a rapid and simple method for the detection of mutations using LAMP combined with the STH chromatographic PAS method. We were able to detect four mutations (C1349T, A1295C, G1303T, A1304T) for RRDR of RR-TB, simultaneously.

1. Materials and Methods

1.1 DNA preparation

For evaluation of LAMP primers, extract DNA were used from T702 (*M. tuberculosis*), KPM1001 (*M. kansasii*), KPM3011 (*M. avium*) and KPM2783 (*M. intracellulare*).

To verify the detection of mutations, 31 DNAs (2 of A1295C, 1 of G1303T, 6 of A1304T and 22 of C1349T) extracted from clinical isolates obtained in Bangladesh were used. The drug resistance has been confirmed by drug susceptibility test, and their mutations have been confirmed by a sequence analysis. (Rahim et al., 2012).

1.2 LAMP primer and allele-specific primer

The MTB-specific LAMP primer sets for Rifampicin Resistance Determining Region (RRDR) of *rpoB* gene were designed using online software (PrimerExplorerV4, http://primerexplorer.jp/v4_manual/index.html). The F1 region of forward inner primer (FIP) and B1 region of backward inner primer (BIP) were designed on the outer side of RRDR. The allele-specific primers to detect mutation were designed between the F1

and B1 region (Fig. 1). The primers for individual mutations or wild types were tagged with different tags. The LAMP primer set and allele specific primers are listed in Table 1 and the assigned positions are shown in Fig. 2.

1.3 LAMP reaction

LAMP reactions were carried out in a 10- μ l reaction mixture consisting of 0.8 μ M biotinylated FIP primer, 0.8 μ M biotinylated BIP primer, 0.2 μ M of each outer primer (F3 and B3), 0.8 μ M of each loop primer (FLP and BLP), 1.4 mM dNTP, 1 μ l 10 $\times$ reaction buffer (Nippon gene KK), 4.8 U of Bst DNA polymerase (Nippon gene KK), 1 μ l of 3,000-fold diluted SYBR Green I and 1 μ l of extracted DNA. Fluorescence of SYBR Green I was measured every minute for 50 min at 68 $^{\circ}$ C during the ongoing reaction, using a thermal cycler Dice real time system (TAKARA Bio Inc.). To terminate the reaction, the mixture was incubated for 2 min at 80 $^{\circ}$ C.

For multiple mutant detection, allele-specific primers for detecting mutations were added to the reaction mixture at a concentration of 0.2 μ M each. In individual mutation detection, the reaction was carried out with two primers (wild and mutant) at the mutation position. Eight primers were used in the reaction for simultaneous detection of four positions. The LAMP reaction condition was same as above.

1.4 Detection design of STH chromatographic PAS

Fig. 2 shows the detection design of PAS for mutations. On the individual detection line of C-PAS, 8, 8 oligonucleotide complementary tags (named c-tags 1 - 8) with distinct nucleotide sequences were previously immobilized from the lower end of PAS. In Table 1, c-tags attached to their respective allele-specific primer are shown.

1.5 Detection of mutations from LAMP products using STH chromatographic PAS

After the LAMP reaction, 10 µl of the products were diluted 100-fold with distilled water and mixed with 2 µl of streptavidin coated blue latex beads and 10 µl of a developing solution (TBA Co., Ltd.). Subsequently, a C-PAS8 membrane strip (TBA Co., Ltd.) was dipped into the mixture for 15-20 min. The appearing blue line was dependent on the type of the tag bound to the primer and hence, it determined what mutations were present. To prevent contamination, reagent preparation and detection sites were strictly isolated. In fact, it took place on the 1st and 4th floors of the building.

2. Results

2.1 LAMP reaction for RRDR region of TB

One FIP, three BIP, two FLP and four BLP primers were examined to amplify the 81bp of the RRDR region. To select the optimal primer combinations, the concentration, the reaction temperature and time were examined by real time amplification using SYB Green I. The selected primers and conditions were able to amplify the TB-specific target region. The detection limit was 200 fg/test using the extracted DNA from isolate strain. (Fig. 3)

2.2 Detection of the individual mutations using STH chromatographic PAS by LAMP

The detection reaction of individual mutations was conducted using the extracted genomic DNA (10 pg/test) . Sequences of the allele-specific primers are shown in Table 1. LAMP reactions were carried out containing LAMP primers and two allele-specific primers (wild and mutant-types) of each target sequence in RRDR. After the LAMP reaction, elongation products of each specific primer were detected by nucleic acid chromatography using PAS. The results are shown in Fig. 4. Genotypes of C1349 (wild type: C), 1349T (mutation: T), A1295 (wild type: A), 1295C (mutation: C), A1304

(wild type: A), 1304T (mutation: T), G1303 (wild type: G) and 1303T (mutation: T) were detected by tags 1, 2, 3, 4, 5, 6, 7 and 8, respectively. According to the analysis by Rahim et al., the mutation of nucleotide position 1349 was the most frequent, followed by 1333 to 1334 and 1303 to 1304, followed by 1295,1345,1355 with low frequency. Since there are many types of mutations in 1333 to 1334 and they are close to 1349, so in order to accurate detection of the 1349, we do not select mutations from 1333 to 1334, and the following 1303 to 1304 and 1295 was selected.

2.3 Detection of the multi-mutations using STH chromatographic PAS by LAMP

Reactions of the multi-mutation detection were carried out using 31 genomic DNA from analyzed mutations and 1 wild type DNA. Moreover, LAMP reactions contained a LAMP primer and eight allele-specific primers for each mutation. After the LAMP reactions, elongation products of each allele-specific primer were detected by nucleic acid chromatography using PAS. Four detection lines were detected there, one of which was detected at different positions depend on the wild type and the mutant type. The results are shown in Fig. 5 and summarized Table 2. In samples with a mutation, a positive line was shifted from the wild type to the mutant type corresponding to the specific mutation. This result indicated that each specific primer underwent a mutation-specific elongation reaction and could be accurately detected by nucleic acid chromatography using PAS.

3. Discussion

The isothermal amplification method is extremely useful for genetic point of care testing (POCT) because special equipment such as PCR is not needed. However, many of the known isothermal amplification methods (NASBA, SDA, RCA) are complicated and costly, using heat sensitive enzymes and requiring multiple enzymes. Primers are

complex, but once the primer design is completed, the LAMP method is comparable in sensitivity and reproducibility to PCR reactions. Moreover, only one type of thermostable enzyme is used, and hence the cost is reduced. By contrast, the LAMP method is not good for multi-detection. As a strand displacement enzyme is used, neither the Taqman probe method nor the Molecular Beacon probe can be applied to LAMP. Furthermore, since the LAMP product is very complicated, multi-detection by electrophoresis is difficult to achieve.

STH chromatographic PAS is a rapid and easy detection method for multi-amplification products. It has been reported that multi-LAMP reaction products can be easily detected (Takabatake et al., 2018). In this study, allele-specific primers were used to detect mutations using the LAMP amplification method. The principle of LAMP reaction using allele-specific primers is shown in Fig. 6a. First, LAMP primers were designed to squeeze in between the mutation site of RRDR. In LAMP amplification, a dumbbell structure is generated after the first few steps. An allele-specific primer with tag anneals to the target sequence in RRDR, and DNA strands are only synthesized when the primer sequences fully match with the target sequences. The product is then displaced by the loop primer elongation. The replacement product gets looped and extends, with the biotin-LAMP primer reacting at the loop part. Next, the allele-specific primer reacts and elongates from the single-stranded portion of the product, and the biotin-LAMP primer product is displaced. Subsequently, the allele-specific primer reacts and expands to the biotin-LAMP primer product, after which the reaction is terminated. At this point, the terminated product has biotin and a tag. Detection of the product with biotin and a tag is shown in Fig. 6b (Monden et al., 2014; Tian et al., 2014; Tian et al., 2016; Saito et al., 2018).

In the present study, it was possible to detect 60% or more of RR-MTB by detecting the aforementioned four mutations. With this new method, it was proved that it is

possible to simultaneously react multiple mutagenic primers with one LAMP reaction. To detect additional mutations, it is possible to add a mutation primers or perform another reaction with the same LAMP primer and different mutation primers.

Xpert MTB/RIF® assay is a useful method to detect the RR-TB (Helb et al., 2010). According to its principle, when a wild type sequence is present, the signal of the molecular beacon probe is detected and the signal disappears when a mutated sequence is present (El-Hajj et al., 2001). In other words, the detection rate would differ depending on the abundance ratio of the susceptible and the resistant bacteria in the specimen. If the majority of the samples were susceptible bacteria and only small amounts of resistant bacteria were present, there would be a possibility that no resistant bacteria could be detected.

Line probe assay is also a useful method to detect RR-TB (Beenhouwer et al., 1995). As this method positively detects the mutant sequence, it is less affected by the abundance ratio of the resistant bacteria. However, the line probe assay requires 3 hours or longer and 7 or more detection steps to detect mutant sequence after PCR, and it needs heating equipment. In contrast, mutant sequences can be detected by STH chromatographic PAS method at room temperature in 15-20 minutes and following only two steps after LAMP.

Since the combined LAMP and STH chromatographic PAS method does not require special equipment and complicated work after the Xpert assay or Line probe assay is completed, it can be implemented anywhere. For example, it can be conducted and tested on site in a remote location and or in areas with unstable power supply. However, there are some issues in the combined LAMP and the STH chromatographic PAS method that need to be addressed. There is a risk of contamination as the lid of LAMP reaction tube needs to be opened after LAMP is completed. Further improvement of the method could eliminate the risk of contamination. A simple device with LAMP reaction

chamber connected to the STH chromatographic PAS cassette enables chromatographic development without opening the lid after LAMP reaction. This will be reported in our future publication.

In conclusion, the combined LAMP and STH chromatographic PAS method does not require special equipment and readily detects several mutations in a quick manner, which meets the demand for it in areas with high prevalence of drug-resistant bacteria such as those in developing countries. In addition, it is suitable for POCT.

Acknowledgment

We thank Dr. Aki Tamaru for providing DNA extracted from clinically isolated *M. tuberculosis*.

Fund

This study is based on previous results obtained from a project subsidized by the New Energy and Industrial Technology Development Organization (NEDO). The research was supported in part by a grant from the Ministry of Education, Culture, Sports, Science and Technology, Japan(MEXT), the Joint Research Program of the Research Center for Zoonosis Control, Hokkaido University to YS, and in part by Japan Agency for Medical Research and Development (AMED) under Grant Number JP20jk0210005, JP20jm0110021 and JP20wm0125008 to YS.

References

1. Aye KS, Nakajima C, Yamaguchi T, Win MM, Shwe MM, Win AA, Lwin T, Nyunt WW, Ti T, Suzuki Y. Genotypic characterization of multi-drug-resistant

296 *Mycobacterium tuberculosis* isolates in Myanmar. J Infect Chemother.
 297 2016;22:174-9. DOI: [10.1016/j.jiac.2015.12.009](https://doi.org/10.1016/j.jiac.2015.12.009)

298 2. Badolo A, Okado K, Guelbeogo WM, Aonuma H, Bando H, Fukumoto S, Sagnon
 299 N, Kanuka H. Development of an allele-specific, loop-mediated, isothermal
 300 amplification method (AS-LAMP) to detect the L1014F *kdr-w* mutation in
 301 *Anopheles gambiae* s.l. Malar J. 2012;11:227. DOI: [10.1186/1475-2875-11-227](https://doi.org/10.1186/1475-2875-11-227)

302 3. Badolo A, Bando H, Traore A, Ko-ketsu M, Guelbeogo WM, Kanuka H, Ranson
 303 H, Sagnon N, Fukumoto S. Detection of G119S *ace-1^R* mutation in field-collected
 304 *Anopheles gambiae* mosquitoes using allele-specific loop-mediated isothermal
 305 amplification (AS-LAMP) method. Malar J. 2015;14:477. DOI: [10.1186/s12936-](https://doi.org/10.1186/s12936-015-0968-9)
 306 [015-0968-9](https://doi.org/10.1186/s12936-015-0968-9)

307 4. De Beenhouwer, H., Lhiang, Z., Jannes, G., Mijs, W., Machtelinckx, L., Rossau,
 308 R., Traore, H., Portaels, F. Rapid detection of rifampicin resistance in sputum and
 309 biopsy specimens from tuberculosis patients by PCR and line probe assay. Tuber
 310 Lung Dis. 1995;76:425-30. DOI: [10.1016/0962-8479\(95\)90009-8](https://doi.org/10.1016/0962-8479(95)90009-8)

311 5. Global Health Observatory (GHO) data repository. Geneva, Switzerland: World
 312 Health Organization (WHO).
 313 <http://apps.who.int/gho/data/node.main.1315?lang=en>

314 6. Itonaga M, Matsuzaki I, Warigaya K, Tamura T, Shimizu Y, Fujimoto M, Kojima
 315 F, Ichinose M, Murata S. Novel methodology for rapid detection of KRAS
 316 mutation using PNA-LNA mediated loop-mediated isothermal amplification. PLoS
 317 ONE. 2016;11:e0151654. DOI: [10.1371/journal.pone.0151654](https://doi.org/10.1371/journal.pone.0151654)

318 7. Monden Y, Takasaki K, Futo S, Niwa K, Kawase M, Akitake H, Tahara M. A
 319 rapid and enhanced DNA detection method for crop cultivar discrimination. J
 320 Biotechnol. 2014;185:57-62. DOI: [10.1016/j.jbiotec.2014.06.013](https://doi.org/10.1016/j.jbiotec.2014.06.013)

8. Mori Y, Nagamine K, Tomita N, Notomi T. Detection of loop mediated isothermal amplification reaction by turbidity derived from magnesium pyrophosphate formation. *Biochem Biophys Res Commun*. 2001;289:150-4.
DOI: [10.1006/bbrc.2001.5921](https://doi.org/10.1006/bbrc.2001.5921)
9. Nagamine K, Watanabe K, Ohtsuka K, Hase T, Notomi T. Loop-mediated isothermal amplification reaction using a nondenatured template. *Clin Chem*. 2001;47:1742–3. PMID: 11514425
10. Notomi T, Okayama H, Masubuchi H, Yonekawa T, Watanabe K, Amino N, Hase T. Loop-mediated isothermal amplification of DNA. *Nucleic Acids Res*. 2000;28:e63. DOI: [10.1093/nar/28.12.e63](https://doi.org/10.1093/nar/28.12.e63)
11. Ohshiro T, Miyagi C, Tamaki Y, Mizuno T, Ezaki T. Development of a rapid diagnostic method for identification of *Staphylococcus aureus* and antimicrobial resistance in positive blood culture bottles using a PCR-DNA-chromatography method. *J Infect Chemother*. 2016;22:372-6. DOI: [10.1016/j.jiac.2016.02.013](https://doi.org/10.1016/j.jiac.2016.02.013)
12. Rahim Z, Nakajima C, Raqib R, Zaman K, Endtz HP, van der Zanden AG, Suzuki Y. Molecular mechanism of rifampicin and isoniazid resistance in *Mycobacterium tuberculosis* from Bangladesh. *Tuberculosis* 2012;92:529-34.
DOI: [10.1016/j.tube.2012.07.005](https://doi.org/10.1016/j.tube.2012.07.005)
13. Takasaki, K., Yamakoshi, Y., Futo, S., 2018. Single-Laboratory Validation of Rapid and Easy DNA Strip for Porcine DNA Detection in Beef Meatballs. *J AOAC Int*. 2018;101:1653-6. DOI: [10.5740/jaoacint.17-0394](https://doi.org/10.5740/jaoacint.17-0394)
14. Saito T, Kikuchi A, Kaneko A, Isozumi R, Teramoto I, Kimura M, Hirasawa N, Hiratsuka M. Rapid and sensitive multiplex single-tube nested PCR for the identification of five human *Plasmodium* species. *Parasitol Int*. 2018;67:277-83.
DOI: [10.1016/j.parint.2018.01.005](https://doi.org/10.1016/j.parint.2018.01.005)

15. Sakai J, Maeda T, Tarumoto N, Misawa K, Tamura S, Imai K, Yamaguchi T, Iwata S, Murakami T, Maesaki S. A novel detection procedure for mutations in the 23S rRNA gene of *Mycoplasma pneumoniae* with peptide nucleic acid-mediated isothermal amplification assay. J Microbiol Methods. 2017;141:90-6. DOI: [10.1016/j.mimet.2017.08.009](https://doi.org/10.1016/j.mimet.2017.08.009)
16. Shanmugakani RK, Akeda Y, Yamamoto N, Sakamoto N, Hagiya H, Yoshida H, Takeuchi D, Sugawara Y, Kodera T, Kawase M, Laolerd W, Chaihongsan N, Santanirand P, Ishii Y, Hamada S, Tomono K. PCR-Dipstick Chromatography for Differential Detection of Carbapenemase Genes Directly in Stool Specimens. Antimicrob Agents Chemother. 2017;61:e00067-17. DOI: [10.1128/AAC.00067-17](https://doi.org/10.1128/AAC.00067-17)
17. Takabatake R, Kagiya Y, Minegishi Y, Futo S, Soga K, Nakamura K, Kondo K, Mano J, Kitta K. Rapid Screening Detection of Genetically Modified Crops by Loop-Mediated Isothermal Amplification with a Lateral Flow Dipstick. J Agric Food Chem. 2018;66:7839-45. DOI: [10.1021/acs.jafc.8b01765](https://doi.org/10.1021/acs.jafc.8b01765)
18. Tam KK-G, Leung KS-S, To SW-C, Siu GK-H, Lau TC-K, Shek VC-M, Tse CW-S, Wong SS-Y, Ho P-L, Yam W-C. Direct detection of *Mycobacterium tuberculosis* and drug resistance in respiratory specimen using Abbott Realtime MTB detection and RIF/INH resistance assay. Diagn Microbiol Infect Dis. 2017;89,:118-24. DOI: [10.1016/j.diagmicrobio.2017.06.018](https://doi.org/10.1016/j.diagmicrobio.2017.06.018)
19. Telenti A, Imboden P, Marchesi F, Lowrie D, Cole S, Colston MJ, Matter L, Schopfer K, Bodmer T, Detection of rifampicin-resistance mutations in *Mycobacterium tuberculosis*. Lancet 1993;341:647-50. DOI: [10.1016/0140-6736\(93\)90417-f](https://doi.org/10.1016/0140-6736(93)90417-f)

- 370 20. Tian L, Sato T, Niwa K, Kawase M, Tanner ACR, Takahashi N. Rapid and
371 sensitive PCR-dipstick DNA chromatography for multiplex analysis of oral
372 microbiota. Biomed Res Int. 2014: 180323. DOI: [10.1155/2014/180323](https://doi.org/10.1155/2014/180323)
- 373 21. Tian L, Sato T, Niwa K, Kawase M, Mayanagi G, Washio J, Takahashi N. PCR-
374 dipstick DNA chromatography for profiling of a subgroup of cariesassociated
375 bacterial species in plaque from healthy coronal surfaces and periodontal pockets.
376 Biomed Res Int. 2016;37:29-36. DOI: [10.2220/biomedres.37.29](https://doi.org/10.2220/biomedres.37.29)
- 377 22. World Health Organization (WHO). Global tuberculosis report 2019. Geneva,
378 Switzerland: World Health Organization; 2019.
379 [https://apps.who.int/iris/bitstream/handle/10665/329368/9789241565714-](https://apps.who.int/iris/bitstream/handle/10665/329368/9789241565714-eng.pdf?ua=1)
380 [eng.pdf?ua=1](https://apps.who.int/iris/bitstream/handle/10665/329368/9789241565714-eng.pdf?ua=1)
381

382 Table1. Nucleotide sequence of LAMP primers for RRDR and allele-specific primers for mutation
383 detection

Use	Name	Sequence
LAMP	bFIP	biotin-CTTGATCGCGGCGACCAACGGACGTGGAGGCGATCACA
	F3	AGCGGATGACCACCCAG
	FLP	GGATGTTGATCAACGTCTGCG
	bBIP	biotin-ACGTGCACCCGTCGCACTTGTTGGGCCCCCTCAGG
	B3	AGCGAGCCGATCAGACC
	BLP	CGGATGTGCCCCGATCGAAA
Allele-detection	C1349	tag1-CCACAAGCGCCGACTGTC
	1349T	tag2-CCACAAGCGCCGACTGcT
	A1295	tag4-GGCACCAGCCAGCTGAGCgA
	1295C	tag3-GGCACCAGCCAGCTGAGCgC
	A1304	tag5-ACAGCGGGTTGTTCTGGT
	1304T	tag6-ACAGCGGGTTGTTCTGGA
	G1303	tag7-CCAGCTGAGCCAATTCATGG
	1303T	tag8-CCAGCTGAGCCAATTCATGT

384 The small character shows artificial mutations.

385

Table2. Characteristics of samples used for this study and reactivities to tags

No.	Name	Mutation position	tag 1	tag 2	tag 3	tag 4	tag 5	tag 6	tag 7	tag 8
1	MTB	Wild	+	-	-	+	+	-	+	-
2	1-70	A1295C	+	-	+	-	+	-	+	-
3	8-11		+	-	+	-	+	-	+	-
4	2P-14	G1303T	+	-	-	+	+	-	-	+
5	11-72	A1304T	+	-	-	+	-	+	+	-
6	OM-4		+	-	-	+	-	+	+	-
7	OM-5		+	-	-	+	-	+	+	-
8	OM-10		+	-	-	+	-	+	+	-
9	OM-19		+	-	-	+	-	+	+	-
10	OM-95		+	-	-	+	-	+	+	-
11	9-8	C1349T	-	+	-	+	+	-	+	-
12	OM-2		-	+	-	+	+	-	+	-
13	OM-6		-	+	-	+	+	-	+	-
14	OM-7		-	+	-	+	+	-	+	-
15	OM-9		-	+	-	+	+	-	+	-
16	OM-12		-	+	-	+	+	-	+	-
17	OM-13		-	+	-	+	+	-	+	-
18	OM-14		-	+	-	+	+	-	+	-
19	OM-16		-	+	-	+	+	-	+	-
20	OM-18		-	+	-	+	+	-	+	-
21	OM-20		-	+	-	+	+	-	+	-
22	OM-21		-	+	-	+	+	-	+	-
23	OM-25		-	+	-	+	+	-	+	-
24	OM-26		-	+	-	+	+	-	+	-
25	OM-29		-	+	-	+	+	-	+	-
26	OM-31		-	+	-	+	+	-	+	-
27	OM-35		-	+	-	+	+	-	+	-
28	OM-37		-	+	-	+	+	-	+	-
29	OM-38		-	+	-	+	+	-	+	-
30	OM-42		-	+	-	+	+	-	+	-
31	OM-55		-	+	-	+	+	-	+	-
32	OM-57		-	+	-	+	+	-	+	-

Figure legends

Figure 1. Design of the LAMP primer and the allele-specific primer

DNA sequence of partial *rpoB* gene containing RRDR region and position of mutations, and LAMP and allele-specific primers designed for this study. The mutation nucleotide is shown in red and bolded. The nucleotides shown in blue are the region of RRDR. The LAMP primers were designed to amplify the region of RRDR. The 3' end nucleotide of the allele-specific primers were assigned to the base corresponding to the wild type and the mutant type, respectively.

Figure 2. Detection design of STH chromatographic PAS for four mutations of RRDR.

Prior to development, C-PAS8 had eight orange lines of immobilized c-tag and 1 flow control. The orange lines disappeared after development. During the development, each tag in the primer hybridized to the each c-tag independently and streptavidin coated blue latex beads bound to the biotin conjugate primer.

Figure 3. Real time detection for LAMP reaction of RRDR.

The sensitivity and specificity of the LAMP reaction was measured with a real-time PCR instrument (Dice real time system). The results showed a detection limit of 200fg/test for the RRDR gene from the MTB genome DNA and no cross reactivity for *Mycobacterium avium*, *Mycobacterium intracellulare* and *Mycobacterium kansasii*.

Figure 4. Detection results of individual mutation.

Individual mutation was detected using the allele-specific primer for each mutation. No.1 and 2 used 531 wild and mutation primers, respectively. The wild type genome used No.1 reaction and the mutation type genome used No. 2 reaction. The results showed No.1 detected only wild type primer signal (tag1) and No. 2 detected only

mutation type signal (tag2). No. 3 and 4 used 513 wild and mutation primers, respectively. The wild type genome used No. 3 reaction and the mutation type genome used No. 4 reaction. The results showed No. 3 detected only wild type primer signal (tag4) and No. 4 detected only mutation type signal (tag3). No. 5 and 6 used 516 wild and mutation primers, respectively. The wild type genome used No. 5 reaction and the mutation type genome used No. 6 reaction. The results showed No. 5 detected only wild type primer signal (tag7) and No. 6 detected only mutation type signal (tag8). No.7 and 8 used 516 wild and mutation primers, respectively. The wild type genome used No.7 reaction and the mutation type genome used No.8 reaction. The results showed the No.7 detected only wild type primer signal (tag5) and No. 8 detected only mutation type signal (tag6).

Figure 5. Detection results of multi-detection

Multi-mutations were detected using the eight allele-specific primers for each mutation. The extracted genome sample are summarized in Table 2. The results show that four mutations were precisely detected.

Figure 6. Detection principle for STH chromatographic PAS using LAMP

(A) To detect with the STH chromatographic PAS method, an amplification product with a tag and biotin will be synthesized by isothermal amplification. In this scheme, the extension product of tag-attached allele-specific primer forms a double strand with biotin-attached to the elongation product of FIP. (B) Biotin labeled LAMP products are labeled by avidin-coated blue beads. Blue beads labeled LAMP products are trapped by anti-tag oligonucleotides printed on the strip membrane via the tag at the end of LAMP products.

Takarada et al. Figure 1

Takarada et al. Figure 3

Takarada et al. Figure 4

A

B

Avidin-coated Blue Beads
with developing buffer