Title	Pumilio1 phosphorylation precedes translational activation of its target mRNA in zebrafish oocytes
Author(s)	Saitoh, Atsushi; Takada, Yuki; Horie, Mayu; Kotani, Tomoya
Citation	Zygote, 26(5), 372-380 https://doi.org/10.1017/S0967199418000369
Issue Date	2018-10
Doc URL	http://hdl.handle.net/2115/83283
Rights	This article has been published in a revised form in Zygote https://doi.org/10.1017/S0967199418000369 . This version is published under a Creative Commons CC-BY-NC-ND. No commercial re-distribution or re-use allowed. Derivative works cannot be distributed. © Cambridge University Press.
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	Zygote 26-5_372 - 380.pdf

[Instructions for use](#)

Pumilio1 phosphorylation precedes translational activation of its target mRNA in zebrafish oocytes

Atsushi Saitoh¹, Yuki Takada¹, Mayu Horie¹, and Tomoya Kotani^{1,2,*}

¹Biosystems Science Course, Graduate School of Life Science, Hokkaido University, Sapporo 060-0810, Japan

²Department of Biological Sciences, Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan

*Corresponding author.

Tomoya Kotani,

Department of Biological Sciences, Faculty of Science, Hokkaido University,

North 10 West 8, Sapporo, Hokkaido 060-0810, Japan

Tel.: +81-11-706-4455

Fax.: +81-11-706-4455

E-mail: tkotani@sci.hokudai.ac.jp

A running headline: Pumilio1 phosphorylation in oocytes

Summary

Translational regulation of mRNAs is crucial for promoting various cellular and developmental processes. Pumilio1 (Pum1) has been shown to play key roles in translational regulation of target mRNAs in many systems of diverse organisms. In zebrafish immature oocytes, Pum1 was shown to bind to *cyclin B1* mRNA and promote the formation of *cyclin B1* RNA granules. This Pum1-mediated RNA granule formation seemed critical to determine the timing of translational activation of *cyclin B1* mRNA during oocyte maturation, leading to activation of maturation/M-phase-promoting factor (MPF) at the appropriate timing. Despite its fundamental importance, the mechanisms of translational regulation by Pum1 remain elusive. In this study, we examined the phosphorylation of Pum1 as a first step to understand the mechanisms of Pum1-mediated translation. SDS-PAGE analyses and phosphatase treatments showed that Pum1 was phosphorylated at multiple sites during oocyte maturation. This phosphorylation began in an early period after induction of oocyte maturation, which preceded the polyadenylation of *cyclin B1* mRNA. Interestingly, depolymerization of actin filaments in immature oocytes caused phosphorylation of Pum1, disassembly of *cyclin B1* RNA granules, and polyadenylation of *cyclin B1* mRNA but not translational activation of the mRNA. Overexpression of the Pum1 N-terminus prevented the phosphorylation of Pum1, disassembly of *cyclin B1* RNA granules, and translational activation of the mRNA even after induction of oocyte maturation. These results suggest that Pum1 phosphorylation in the early period of oocyte maturation is one of key processes for promoting the disassembly of *cyclin B1* RNA granules and translational activation of target mRNA.

Key words: vertebrate, RNA granule, translation, oocyte maturation, cytoplasmic polyadenylation

Introduction

Timings and sites of protein synthesis are fundamentally important for promoting diverse biological processes including meiotic cell cycles, early development and neuronal plasticity. RNA-binding proteins have been shown to drive the localization and translational regulation of target mRNAs, by which protein synthesis is achieved at the right time and place (Kloc et al., 2002; Kotani et al., 2017; Martin and Ephrussi, 2009). Pumilio (Pum) is a sequence-specific RNA-binding protein that belongs to the PUF (Pumilio and Fem-3 mRNA-binding factor) family, which is highly conserved in eukaryotes from yeast to human (Spassov and Jurecic, 2003b; Wickens et al., 2002). Pum was identified in *Drosophila* as a protein that is essential for posterior patterning of embryos and development of germline cells (Asaoka-Taguchi et al., 1999; Forbes and Lehmann, 1998; Lehmann and Nussleinvolhard, 1987; Lin and Spradling, 1997). *Drosophila* Pum targets *hunchback* mRNA in the posterior patterning and *cyclin B* mRNA in the germline development (Asaoka-Taguchi et al., 1999; Murata and Wharton, 1995). In both cases, Pum represses the translation of target mRNAs in spatially and temporally regulated manners. FBF proteins, members of the PUF family in *C. elegans*, also function in germline development by regulating translation of their target mRNAs during embryogenesis (Crittenden et al., 2002; Zhang et al., 1997).

Two Pum proteins, Pumilio1 (Pum1) and Pumilio2 (Pum2), have been found in all vertebrates studied (Kurisaki et al., 2007; Lee et al., 2008; Nakahata et al., 2001; Padmanabhan and Richter, 2006; Spassov and Jurecic, 2002; Spassov and Jurecic, 2003a; White et al., 2001). Recent studies have shown that Pum1-deficient mice are viable but have decreased fertility due to an extensive increase in apoptosis of spermatocytes in males (Chen et al., 2012) and due to a decrease in diplotene-stage oocytes in females (Mak et al., 2016). Pum1 deficiency also causes neuronal degeneration in the brain (Gennarino et al., 2015). Mouse Pum1 has been shown to target thousands of mRNAs in the testis and brain (Chen et al., 2012; Zhang et al., 2017). In addition, the amount of protein products, but not the amount of mRNAs, was increased in Pum1-deficient mice, indicating that Pum1 represses translation of target mRNAs. All of the findings from various species show the importance of Pum-mediated translational regulation in embryogenesis, spermatogenesis and neurogenesis. However, the mechanisms of Pum-mediated translational regulation remain to be elucidated.

Temporal regulation of mRNA translation has been most extensively studied in

oocytes. In many vertebrates, oocytes are arrested at prophase I of the meiotic cell cycle and deposit tens of thousands of mRNAs for promoting oocyte maturation and development (Harvey et al., 2013; Masui and Clarke, 1979). Fully grown oocytes in this stage are unable to be fertilized and they are called immature oocytes. It has been estimated that more than one thousand mRNAs are translationally repressed in immature oocytes and begin to be translated after resumption of the meiotic cell cycle at timings specific to distinct mRNAs (Chen et al., 2011; Kotani et al., 2017). Oocytes are arrested again at metaphase II until fertilization. This process is called oocyte maturation, and oocytes arrested at metaphase II are called mature oocytes.

Translation of *cyclin B1* mRNA is repressed in immature oocytes and de-repressed after initiation of oocyte maturation. Synthesized Cyclin B1 protein from the stored mRNA functions as a regulatory subunit of maturation/M-phase-promoting factor (MPF), a master regulator in the progression of oocyte maturation. Extensive studies using *Xenopus* oocytes have shown that translational activation of dormant mRNAs including *cyclin B1* mRNA is mainly directed by cytoplasmic polyadenylation elements (CPEs)-mediated polyadenylation of the mRNAs (de Moor and Richter, 1999; McGrew et al., 1989; Sheets et al., 1994). The CPE-binding protein CPEB has been shown to drive translational repression and activation of target mRNAs including *cyclin B1* mRNA (de Moor and Richter, 1999; Hake and Richter, 1994; Tay et al., 2000). Pum1 targets *cyclin B1* mRNA in zebrafish, *Xenopus* and mouse oocytes (Kotani et al., 2013; Nakahata et al., 2001; Nakahata et al., 2003; Ota et al., 2011a; Pique et al., 2008). We previously showed that *cyclin B1* mRNA forms RNA granules in immature oocytes of zebrafish and mouse (Kotani et al., 2013). These RNA granules disassemble at the timing of translational activation during oocyte maturation (Horie and Kotani, 2016; Kotani et al., 2013; Nukada et al., 2015). By generating transgenic zebrafish expressing reporter mRNAs in oocytes, we demonstrated that binding of Pum1 to *cyclin B1* mRNA is essential for the formation of RNA granules. This Pum1-mediated granule formation seems to be important for regulating the timing of translational activation of *cyclin B1* mRNA after initiation of oocyte maturation (Kotani et al., 2013). However, the mechanism by which translational timing of *cyclin B1* mRNA is regulated by Pum1 remains unresolved.

In this study, we analyzed the phosphorylation of Pum1 in zebrafish oocytes. Pum1 was phosphorylated at multiple sites during oocyte maturation. This phosphorylation

was initiated in an early period after induction of oocyte maturation, preceding the translational activation of *cyclin B1* mRNA. In addition, Pum1 phosphorylation was found to be linked to disassembly of RNA granules and polyadenylation of *cyclin B1* mRNA. Our findings provide the basis for understanding the mechanisms of translational regulation of mRNAs by Pum1.

Materials and Methods

Preparation of oocytes

All animal experiments in this study were approved by the Committee on Animal Experimentation, Hokkaido University. Zebrafish ovaries were dissected from adult females in zebrafish Ringer's solution (116 mM NaCl, 2.9 mM KCl, 1.8 mM CaCl₂, and 5 mM HEPES; pH 7.2). Oocytes were manually isolated from ovaries with forceps under a dissecting microscope. Oocyte maturation was induced by stimulation with 1 μ g/ml of 17 α ,20 β -dihydroxy-4-pregnen-3-one, an MIH in fish. For *in situ* hybridization analysis, oocytes were fixed with 4% paraformaldehyde in PBS (4% PFA/PBS) overnight at 4°C. For poly(A) test (PAT) assay, oocytes were extracted with Trizol reagent (Invitrogen) and total RNA was used for RNA ligation-coupled RT-PCR. For immunoblotting and immunoprecipitation analyses, oocytes were homogenized with an equal volume of ice-cold extraction buffer (EB: 100 mM β -glycerophosphate, 20 mM HEPES, 15 mM MgCl₂, 5 mM EGTA, 1 mM dithiothreitol, 100 μ M (*p*-amidinophenyl)methanesulfonyl fluoride, and 3 μ g/ml leupeptin; pH 7.5). After centrifugation at 15,000 g for 10 min at 4°C, the supernatant was collected and used for SDS-PAGE analysis and immunoprecipitation.

Section *in situ* hybridization

Section *in situ* hybridization with the tyramide signal amplification (TSA) Plus DNP system (PerkinElmer) was performed according to the procedure reported previously (Takei et al., 2018). Briefly, fixed oocytes were dehydrated, embedded in paraffin, and cut into 12- μ m-thick sections. A digoxigenin (DIG)-labeled antisense RNA probe for the full length of *cyclin B1* was used for detection of the *cyclin B1* gene transcript. No signal was detected with sense probes. After hybridization and washing, samples were incubated with anti-DIG-horseradish peroxidase (HRP) antibody (Roche) (1:500 dilution) for 30 min. The reaction with tyramide-dinitrophenyl (DNP) was performed

according to the manufacturer's instructions. The samples were then incubated overnight with anti-DNP-Alexa 488 antibody (Molecular Probes) (1:500 dilution). To detect nuclei, the samples were incubated with 10 μ g/ml Hoechst 33258 for 10 min. After being mounted with a Prolong Antifade Kit (Molecular probes), the samples were observed under an LSM5LIVE confocal microscope (Carl Zeiss).

Poly(A) test (PAT) assay

RNA ligation-coupled RT-PCR was performed according to the procedure reported previously (Charlesworth et al., 2004; Yasuda et al., 2010). Two μ g of total RNA extracted from pools of 15 zebrafish oocytes was ligated to 0.4 μ g of P1 anchor primer (5'-P-GGT CAC CTT GAT CTG AAG C-NH₂-3') in a 10- μ l reaction using T4 RNA ligase (New England Biolabs) for 30 min at 37°C. The ligase was inactivated for 5 min at 92°C. Half of the RNA ligation reaction was used in a 25- μ l reverse transcription reaction using the Superscript III First Strand Synthesis System with a P1' primer (5'-GCT TCA GAT CAA GGT GAC CTT TTT-3'). Four μ l of the cDNA was used for the 1st PCR with the P1' primer and a *zcyclin B1*-PAT-f1 primer (5'-GAG GGC CTT TCT AAG CAT CTG GCT GTG-3') for 17 cycles. Two μ l of the 1st PCR reaction was used for the 2nd PCR with the P1' primer and a *zcyclin B1*-3'UTR-f primer (5'-TAC GGA TTT CTT CAC TGC CAT G-3') for 35 cycles. The PCR product was resolved on a 2% agarose gel in TAE buffer. We confirmed that the increase in PCR product length was due to elongation of the poly(A) tails by cloning the 2nd PCR products and sequencing them.

Immunoblotting and immunoprecipitation

Ten μ l of oocyte extracts was incubated with an anti-*Xenopus* Pum1 rabbit antibody, which was affinity-purified with zebrafish Pum1 (Kotani et al., 2013), for 1 h at 4°C. The extracts were then incubated with protein A-Sepharose beads (GE Healthcare) for 3 h at 4°C and washed three times with EB containing 1% Tween 20. The crude oocyte extracts and the immunoprecipitates of anti-Pum1 rabbit antibody were separated by SDS-PAGE, blotted onto an Immobilon membrane, and probed with an anti-*Xenopus* Pum1 monoclonal antibody (Pum2A5; Nakahata et al., 2001). CPEB was detected by immunoblotting the crude extracts with anti-CPEB monoclonal antibody (9G3; Takahashi et al., 2014). The Cyclin B1-Cdc2 complex was detected by immunoblotting

Suc1 precipitates from crude oocyte extracts with anti-Cdc2 (MC2-21; Tanaka and Yamashita, 1995) and anti-goldfish Cyclin B1 (B112; Katsu et al., 1993) monoclonal antibodies.

Phosphatase treatments

After immunoprecipitation of oocyte extracts with anti-Pum1 rabbit antibody, the immunoprecipitates were treated with calf intestinal alkaline phosphatase (CIAP; New England BioLabs) or λ -protein phosphatase (λ -PP; New England BioLabs) for 1 h at 30°C. The samples were then washed three times with EB and analyzed by immunoblotting.

Cytochalasin B treatment

To depolymerize actin filaments, oocytes were treated with 1 μ g/ml cytochalasin B (Sigma) for 3 h. Cytochalasin B was dissolved in dimethyl sulfoxide (DMSO) as stocks and diluted in Ringer's solution before use. As a control, oocytes were treated with 0.01% DMSO. After treatment with cytochalasin B or DMSO for 3 h, the oocytes were stimulated with MIH for induction of oocyte maturation.

RT-PCR analysis after IP (IP/RT-PCR)

One hundred μ l oocyte extracts was incubated with affinity-purified anti-Pum1 rabbit antibody or control rabbit IgG for 1 h at 4°C. The extracts were then incubated with protein A-Sepharose beads for 3 h at 4°C and washed five times with EB containing 1% Tween 20. After extraction of mRNAs from the beads with Trizol reagent, RT-PCR was performed using a *zcyclin B1*-f2 primer (5'-GAG GGC CTT TCT AAG CAT CTG GCT GTG-3') and a *zcyclin B1*-r2 primer (5'-TTA TTT GAA TTC AAA TGT ACA AAC TTG C-3').

Overexpression of Pum1 N terminus and immunostaining

mRNAs encoding GFP-Pum1N (Pum1N; 1-790 amino acids) (Kotani et al., 2013) and GFP were synthesized with an mMESSAGE mMACHINE SP6 kit (Life Technologies). Two ng of the mRNAs was injected into fully grown oocytes using a microinjector (CellTram vario). The oocytes were incubated for 3 h and stimulated with MIH. Before or at 90 min after MIH stimulation, the oocytes were fixed with 4% PFA/PBS for *in situ*

hybridization analysis or extracted with ice-cold EB for detection of Pum1 and Cyclin B1. To detect GFP-Pum1N and GFP, the oocytes were immunostained with anti-GFP mouse antibody (1:200 dilution; Roche) followed by anti-mouse IgG-Alexa Fluor 488 antibody (1:200 dilution; Molecular Probes), after hybridization and washing of the *cyclin B1* RNA probe in *in situ* hybridization analysis.

Results and Discussion

Phosphorylation of Pum1 during zebrafish oocyte maturation

As reported previously, *in situ* hybridization of zebrafish ovaries showed the distribution of *cyclin B1* mRNA as granular structures in the animal polar cytoplasm of fully grown immature oocytes beneath the micropyle (Fig. 1A) (Horie and Kotani, 2016; Kotani et al., 2013). The micropyle is a structure located at the animal pole of the fish chorion, through which a sperm enters into the oocyte cytoplasm. The *cyclin B1* RNA granules disassembled during oocyte maturation (Fig. 1B) (Horie and Kotani, 2016; Kotani et al., 2013). The PAT assay showed that poly(A) tails of *cyclin B1* mRNA were short in immature oocytes and became elongated during oocyte maturation (Fig. 1C) (Yasuda et al., 2010), accompanied by translational activation of mRNA (Kotani et al., 2013). Pum1 has been shown to bind to Pumilio-binding element (PBE) located in the 3' untranslated region (UTR) of *cyclin B1* mRNA (Kotani et al., 2013). Since binding of Pum1 to *cyclin B1* mRNA persists during oocyte maturation (Ota et al., 2011a; Takahashi et al., 2014) even after the mRNA is translated, we predicted that modification of Pum1 may be involved in the change of translational regulation of target *cyclin B1* mRNA from repression to activation states. One of the candidates is phosphorylation as observed in *Xenopus* Pum1 (Ota et al., 2011a). To assess this possibility, we examined whether zebrafish Pum1 is phosphorylated during oocyte maturation.

Pum1 was detected as an approximately 120-kDa protein in crude extracts of zebrafish immature oocytes by SDS-PAGE analysis followed by immunostaining with anti-Pum1 antibody (Fig. 1D). The electrophoretic mobility of Pum1 appeared to be reduced in mature oocytes, resulting in an apparent increase in the molecular weight of Pum1 (Fig. 1D). This reduction in Pum1 mobility in mature oocytes was confirmed by immunoprecipitation of Pum1 (Fig. 1E). The electrophoretic mobility of Pum1 in mature oocytes was partially but not completely recovered by treatment of the

precipitations with calf intestinal alkaline phosphatase (CIAP) (Fig. 1F). Treatment of the precipitations with λ -protein phosphatase (λ -PP) resulted in almost complete recovery of the mobility of Pum1 as a single band (Fig. 1G). These results indicate that Pum1 is phosphorylated at multiple sites and suggest that serine and threonine residues of Pum1 are mainly phosphorylated because λ -PP specifically dephosphorylates these residues (Barik, 1993). Notably, partial reduction in Pum1 mobility was observed in immature oocytes of some, but not all, batches (4 out of 18 batches) (Fig. 1G, compare with Fig. 1D-F), suggesting that a kinase responsible for Pum1 phosphorylation is at least partially activated and that the state of phosphorylation is equilibrated with that of dephosphorylation in immature oocytes.

Timing of Pum1 phosphorylation after induction of oocyte maturation

To assess whether Pum1 phosphorylation is involved in the translational regulation of target *cyclin B1* mRNA, we examined the timing of Pum1 phosphorylation after induction of oocyte maturation. Polyadenylation of *cyclin B1* mRNA began 60 min after induction of oocyte maturation (Fig. 2A), at which no oocyte underwent GVBD (Fig. 2B). As reported previously, the timing of *cyclin B1* mRNA poly(A) tail elongation (60 min) was approximately half of the time point at which 50% of the oocytes have undergone GVBD (120 min in this batch; Fig. 2B) (Horie and Kotani, 2016; Kotani et al., 2013). Time course analysis of Pum1 phosphorylation showed that partial but apparent phosphorylation had occurred at 30 min and then significant phosphorylation had occurred at 120 min after induction of oocyte maturation (Fig. 2C). We then focused on the initiation of Pum1 phosphorylation. Interestingly, slight phosphorylation of Pum1 was detected at 10 min and then apparent phosphorylation was detected at 30 min after induction of oocyte maturation (Fig. 2D). Since crude extracts were unable to clearly show the changes in Pum1 electrophoretic mobility, we performed immunoprecipitation of Pum1 and confirmed that Pum1 was slightly phosphorylated at 10 min and apparently phosphorylated at 30 min after induction of oocyte maturation (Fig. 2E). Significant phosphorylation of Pum1 was also observed 150 min after induction of oocyte maturation in this assay (Fig. 2E). The results showing a gradual increase in the molecular weight of Pum1 support our conclusion that Pum1 is phosphorylated at multiple sites during oocyte maturation (Fig.1). Taken together, these results indicate that Pum1 phosphorylation precedes the polyadenylation of target *cyclin*

B1 mRNA during normal oocyte maturation (Fig. 2F, compare with Fig. 2B).

Phosphorylation of Pum1 was previously reported in *Xenopus* oocytes (Ota et al., 2011a). However, a shift up of the Pum1 protein band, which was caused by phosphorylation as in the case of zebrafish Pum1, was shown to begin at a timing similar to that of Cyclin B1 protein synthesis. Therefore, whether Pum1 phosphorylation was the cause or consequence of translational activation of *cyclin B1* mRNA remains unclear. Our results show for the first time that Pum1 is phosphorylated prior to translational activation of its target mRNA, implying that this Pum1 modification may be a cause of translational activation of target mRNAs.

Effect of actin filament depolymerization on Pum1 phosphorylation

The granule formation of *cyclin B1* mRNA has been shown to be dependent on the binding of Pum1 and on the actin filaments assembled in the oocyte cytoplasm (Kotani et al., 2013; Nukada et al., 2015). However, the relationship between Pum1 modification and actin filament depolymerization has been completely unknown. We then examined the effects of depolymerization of actin filaments on Pum1 phosphorylation. Treatment of immature oocytes with cytochalasin B caused depolymerization of actin filaments (Nukada et al., 2015) and disassembly of *cyclin B1* RNA granules (Fig. 3A and B) (Kotani et al., 2013). In the oocytes treated with cytochalasin B for 3 h, Pum1 was phosphorylated at a level similar to that in oocytes stimulated with MIH for 3 h (Fig. 3C). RT-PCR analysis after immunoprecipitation with anti-Pum1 antibody of the oocyte extracts showed that the binding of Pum1 to *cyclin B1* mRNA was not affected by actin filament depolymerization (Fig. 3E).

As reported previously, Cyclin B1 protein was not detected in oocytes treated with cytochalasin B for 3 h (Fig. 3F, see the lane 0 min after stimulation with MIH) (Kotani et al., 2013), indicating that translation of *cyclin B1* mRNA is still repressed. However, interestingly, the PAT assay showed that poly(A) tails of *cyclin B1* mRNAs were elongated in oocytes treated with cytochalasin B (Fig. 3C). We then examined modifications of CPEB in the cytochalasin B-treated oocytes. CPEB was shown to be phosphorylated and degraded after induction of oocyte maturation in zebrafish (Takahashi et al., 2014) as in the cases of *Xenopus* (Mendez et al., 2002), mouse (Tay et al., 2000; Chen et al., 2011) and porcine (Komrskova et al., 2014) (see Fig. 3F control oocytes stimulated with MIH after DMSO treatment). In *Xenopus* oocytes, the CPEB

degradation has been shown to control the translational activation of *cyclin B1* mRNA (Mendez et al., 2002). In the cytochalasin B-treated oocytes, CPEB was degraded at a level similar to that in the MIH-stimulated oocytes (Fig. 3C). We confirmed that CPEB was phosphorylated shortly after cytochalasin B treatment, followed by gradual degradation (Fig. 3D). Collectively, our results indicate that neither Pum1 phosphorylation, CPEB degradation, nor polyadenylation of *cyclin B1* mRNA is sufficient for translational activation of the mRNA in oocytes. Instead, these events seemed to be linked to rapid activation of *cyclin B1* mRNA translation after the response to MIH, because synthesis of Cyclin B1 protein was detected immediately (30 min) after MIH stimulation in these oocytes (Fig. 3F), compared with the synthesis of Cyclin B1 in a later period (90 min) in control oocytes. Consistent with the rapid synthesis of Cyclin B1, appearance of active form of Cdc2 kinase (Fig. 3F, arrowheads) and occurrence of GVBD (Fig. 3G) were significantly accelerated in oocytes treated with cytochalasin B. Our results also indicate the existence of an unknown event(s) or factor(s) downstream of the MIH receptor that is required for the final trigger of translational activation of *cyclin B1* mRNA after Pum1 phosphorylation, CPEB degradation, and polyadenylation of the mRNA.

Effect of overexpression of GFP-Pum1N on Pum1 phosphorylation

Pum1 contains the highly conserved PUF domain at the C-terminal region that is responsible for binding to mRNAs (Fig. 4A) (Zamore et al., 1997). In the central region, Pum1 contains a Q/N-rich region that is thought to promote ordered aggregation of the proteins (Salazar et al., 2010). Overexpression of Pum1N, which lacks the PUF domain, affected the assembly of *cyclin B1* RNA granules and the translational activation of mRNA (Kotani et al., 2013). This suggested that overexpression of Pum1N affects Pum1 phosphorylation. We therefore examined the effects of overexpression of GFP-Pum1N on Pum1 phosphorylation.

As reported previously, GFP-Pum1N formed large aggregates surrounding *cyclin B1* RNA granules in the animal polar cytoplasm of immature oocytes, while GFP was diffused throughout the cytoplasm (Fig. 4B) (Kotani et al., 2013). These GFP-Pum1N aggregates prevented the disassembly of *cyclin B1* RNA granules. The *cyclin B1* RNA granules had almost completely disassembled in control GFP-expressing oocytes 90 min after induction of oocyte maturation, while the granules were maintained in GFP-

Pum1N-expressing oocytes (Fig. 4B) (Kotani et al., 2013). In response to MIH, endogenous Pum1 was phosphorylated in the control oocytes, while endogenous Pum1 phosphorylation was prevented in the oocytes expressing GFP-Pum1N (Fig. 4C). Consistent with our previous study, oocytes expressing GFP-Pum1N showed delay in the occurrence of GVBD and 70% of the oocytes did not undergo GVBD (Fig. 4D), accompanied by failure in Cyclin B1 synthesis (Kotani et al., 2013). An explanation for the effects of overexpression of GFP-Pum1N is that Pum1N proteins surrounding *cyclin B1* RNA granules prevented endogenous Pum1 phosphorylation by sequestering phosphorylation sites of Pum1 and therefore attenuated granule disassembly and translational activation of the mRNA. Alternatively, GFP-Pum1N affected endogenous Pum1 phosphorylation independently of the effects on granule disassembly and translational activation of the mRNA. Nevertheless, these results suggest a relation between Pum1 phosphorylation and disassembly of *cyclin B1* RNA granules, which may promote translational activation of the mRNA.

Pum1 phosphorylation and translational activation of *cyclin B1* mRNA

Pum1 has been shown to play fundamental roles in diverse biological processes by regulating translation of target mRNAs. However, the mechanisms of Pum1-mediated translation remain to be elucidated. In this study, we demonstrated that a) Pum1 was phosphorylated at multiple sites during oocyte maturation (Fig. 1), b) this phosphorylation preceded the polyadenylation and translational activation of its target mRNA, *cyclin B1* mRNA (Fig. 2), c) depolymerization of actin filaments resulted in Pum1 phosphorylation, *cyclin B1* RNA granule disassembly and polyadenylation of the mRNA without induction of oocyte maturation (Fig. 3), and d) overexpression of Pum1N prevented Pum1 phosphorylation, *cyclin B1* RNA granule disassembly and translational activation of the mRNA even after induction of oocyte maturation (Fig. 4). Since binding of Pum1 to *cyclin B1* mRNA is required for RNA granule formation (Kotani et al., 2013) and persists during oocyte maturation (Ota et al., 2011a; Takahashi et al., 2014), our findings suggest that Pum1 phosphorylation is an initial key step to promote the disassembly of *cyclin B1* RNA granules and polyadenylation of the mRNA, which accelerates the sequential activation of mRNA translation.

A signaling pathway upstream of Pum1 phosphorylation in zebrafish oocytes remains unknown. One of the candidates is Nemo-like kinase (NLK), a typical mitogen-

activated protein kinase. NLK is present in *Xenopus* immature oocytes and is activated in an early phase of oocyte maturation (Ota et al., 2011b). This kinase can directly phosphorylate Pum1 as well as CPEB and Pum2 *in vitro*. Whether NLK is expressed in zebrafish oocytes and is involved in Pum1 phosphorylation will be an interesting issue. Another possibility is that a kinase responsible for Pum1 phosphorylation is at least partially active in zebrafish immature oocytes, because our immunoprecipitation analysis showed that Pum1 was partially phosphorylated in immature oocytes of some batches (Fig. 1G and data not shown) and that Pum1 was immediately phosphorylated in response to MIH in most batches (Fig. 2). If this is the case, a phosphatase responsible for dephosphorylation of Pum1 would be present in immature oocytes, which would maintain the non-phosphorylated state of Pum1. In this scenario, MIH stimulation immediately inhibits the phosphatase in addition to fully activating the kinase responsible for Pum1 phosphorylation. As a next step to understand the mechanism of Pum1-mediated translation, identification of a kinase responsible for Pum1 phosphorylation will be necessary. Our time course analysis also showed significant phosphorylation of Pum1 after the occurrence of GVBD (Fig. 2). This would be caused by activated MPF as in the case of hyperphosphorylation of CPEB in *Xenopus*, porcine and mouse oocytes (Han et al., 2017; Komrskova et al., 2014; Mendez et al., 2002). The kinase responsible for significant phosphorylation of Pum1 and the importance of this second phase of phosphorylation remain to be elucidated.

An intriguing finding in this study is that the cytoplasmic polyadenylation is not sufficient for translational activation of *cyclin B1* mRNA (Fig. 3). Extensive studies using *Xenopus* and mouse oocytes have demonstrated that the cytoplasmic polyadenylation of dormant mRNAs stored in oocytes direct translational activation of the mRNAs (McGrew et al., 1989; Sheets et al., 1994; Stutz et al., 1998; Vassalli et al., 1989). Polyadenylation of *cyclin B1* mRNA has been well shown using *Xenopus* oocytes (Radford et al., 2008; Weill et al., 2012). In immature oocytes, both the deadenylase PARN and the poly(A) polymerase GLD2 associate with *cyclin B1* mRNA. The high activity of PARN sequesters GLD2 activity, shortening the poly(A) tails of mRNA. Multiple CPEBs bind to the CPEs of *cyclin B1* mRNA (deMoor and Richter, 1999). Pum1 binds CPEB, PARN and GLD2 in immature oocytes (Nakahata et al., 2003; Ota et al., 2011a). After initiation of oocyte maturation, several CPEBs are degraded (Mendez et al., 2002), and remaining CPEB recruits the cleavage and

polyadenylation specificity factor CPSF on the poly(A) signal AAUAAA in the terminal end of mRNA and excludes PARN from the mRNA. This remodeling promotes the polyadenylation of mRNA by GLD2. Subsequent binding of the poly(A)-binding protein PABP on the elongated poly(A) tail circularizes the mRNA by the interaction between PABP and the cap-binding factor eIF4E, which is mediated by a scaffold protein eIF4G, resulting in induction of the translational activation of mRNA (Cao and Richter, 2002). Although we could not rule out the possibility that the polyadenylation caused by actin filament depolymerization is partially different from that caused by MIH stimulation, our findings suggest that the induction of translational activation of *cyclin B1* mRNA requires another factor(s) or additional remodeling of the complex even after the polyadenylation of mRNA.

In *Xenopus* oocytes, Pum2 has been shown to function in oocyte maturation by regulating translation of mRNAs other than *cyclin B1* mRNA (Ota et al., 2011a; Padmanabhan and Richter, 2006). In addition, *Xenopus* Pum2 was shown to be phosphorylated during oocyte maturation (Ota et al., 2011a; Ota et al., 2011b), implying that phosphorylation is one of the important processes in Pum2-mediated translational regulation. The expression and involvement of Pum2 in zebrafish oocyte maturation remain to be elucidated. Our results in this study and future studies on the phosphorylation of Pum1 including the identification of phosphorylation sites will contribute to an understanding of the molecular mechanisms of Pum-mediated translational control in diverse systems and organisms.

Acknowledgements

We thank Dr. Yamashita for providing anti-Pum1 antibodies and Dr. Katsu for providing anti-CPEB antibody. This work was supported by a Grant-in-Aid for Scientific Research (16K07242 to T.K.) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

Conflicts of Interest

The authors have no conflicts of interest to declare.

References

Asaoka-Taguchi, M., Yamada, M., Nakamura, A., Hanyu, K. & Kobayashi, S. (1999).

- Maternal Pumilio acts together with Nanos in germline development in *Drosophila* embryos. *Nature Cell Biol.* 1, 431-437.
- Barik, S. (1993). Expression and biochemical properties of a protein serine/threonine phosphatase encoded by bacteriophage lambda. *PNAS* 90, 10633-10637.
- Cao, Q. & Richter, J.D. (2002). Dissolution of the maskin-eIF4E complex by cytoplasmic polyadenylation and poly(A)-binding protein controls cyclin B1 mRNA translation and oocyte maturation. *EMBO J.* 21, 3852-3862.
- Charlesworth, A., Cox, L.L. & MacNicol, A.M. (2004). Cytoplasmic polyadenylation element (CPE)- and CPE-binding protein (CPEB)-independent mechanisms regulate early class maternal mRNA translational activation in *Xenopus* oocytes. *J. Biol. Chem.* 279, 17650-17659.
- Chen, D., Zheng, W., Lin, A., Uyhazi, K., Zhao, H. & Lin, H. (2012). Pumilio 1 suppresses multiple activators of p53 to safeguard spermatogenesis. *Curr. Biol.* 22, 420-425.
- Chen, J., Melton, C., Suh, N., Oh, J.S., Horner, K., Xie, F., Sette, C., Belloch, R. & Conti, M. (2011). Genome-wide analysis of translation reveals a critical role for deleted in azoospermia-like (Dazl) at the oocyte-to-zygote transition. *Genes Dev.* 25, 755-766.
- Crittenden, S.L., Bernstein, D.S., Bachorik, J.L., Thompson, B.E., Gallegos, M., Petcherski, A.G., Moulder, G., Barstead, R., Wickens, M. & Kimble, J. (2002). A conserved RNA-binding protein controls germline stem cells in *Caenorhabditis elegans*. *Nature* 417, 660-663.
- de Moor, C.H. & Richter, J.D. (1999). Cytoplasmic polyadenylation elements mediate masking and unmasking of cyclin B1 mRNA. *EMBO J.* 18, 2294-2303.
- Forbes, A. & Lehmann, R. (1998). Nanos and Pumilio have critical roles in the development and function of *Drosophila* germline stem cells. *Development* 125, 679-690.
- Gennarino, V.A., Singh, R.K., White, J.J., De Maio, A., Han, K., Kim, J.Y., Jafar-Nejad, P., di Ronza, A., Kang, H., Sayegh, L.S., Cooper, T.A., Orr, H.T., Sillitoe, R.V. & Zoghbi, H.Y. (2015). Pumilio1 haploinsufficiency leads to SCA1-like neurodegeneration by increasing wild-type Ataxin1 levels. *Cell* 160, 1087-1098.
- Hake, L.E. & Richter, J.D. (1994). CPEB is a specificity factor that mediates cytoplasmic polyadenylation during *Xenopus* oocyte maturation. *Cell* 79, 617-

- Han, S.J., Martins, J.P.S., Yang, Y., Kang, M.K., Daldello, E.M. & Conti, M. (2017). The Translation of Cyclin B1 and B2 is Differentially Regulated during Mouse Oocyte Reentry into the Meiotic Cell Cycle. *Sci. Reports* 7, 14077.
- Harvey, S.A., Sealy, I., Kettleborough, R., Fenyes, F., White, R., Stemple, D. & Smith, J.C. (2013). Identification of the zebrafish maternal and paternal transcriptomes. *Development* 140, 2703-2710.
- Horie, M. & Kotani, T. (2016). Formation of mos RNA granules in the zebrafish oocyte that differ from cyclin B1 RNA granules in distribution, density and regulation. *Eur. J. Cell Biol.* 95, 563-573.
- Katsu, Y., Yamashita, M., Kajiura, H. & Nagahama, Y. (1993). Behavior of the components of maturation-promoting factor, cdc2 kinase and cyclin B, during oocyte maturation of goldfish. *Dev. Biol.* 160, 99-107.
- Kloc, M., Zearfoss, N.R. & Etkin, L.D. (2002). Mechanisms of subcellular mRNA localization. *Cell* 108, 533-544.
- Komrskova, P., Susor, A., Malik, R., Prochazkova, B., Liskova, L., Supolikova, J., Hladky, S. & Kubelka, M. (2014). Aurora kinase A is not involved in CPEB1 phosphorylation and cyclin B1 mRNA polyadenylation during meiotic maturation of porcine oocytes. *PLOS One* 9, 7.
- Kotani, T., Maehata, K. & Takei, N. (2017). Regulation of Translationally Repressed mRNAs in Zebrafish and Mouse Oocytes. *Results Probl. Cell Differ.* 63, 297-324.
- Kotani, T., Yasuda, K., Ota, R. & Yamashita, M. (2013). Cyclin B1 mRNA translation is temporally controlled through formation and disassembly of RNA granules. *J. Cell Biol.* 202, 1041-1055.
- Kurisaki, I., Iwai, T., Yamashita, M., Kobayashi, M., Ito, E. & Matsuoka, I. (2007). Identification and expression analysis of rainbow trout pumilio-1 and pumilio-2. *Cell Tissue Res.* 327, 33-42.
- Lee, J.Y., Lim, J.M., Kim, D.K., Zheng, Y.H., Moon, S., Han, B.K., Song, K.D., Kim, H. & Han, J.Y. (2008). Identification and gene expression profiling of the Pum1 and Pum2 members of the Pumilio family in the chicken. *Mol. Reprod. Dev.* 75, 184-190.
- Lehmann, R. & Nussleinvohard, C. (1987). Involvement of the Pumilio Gene in the

- Transport of an Abdominal Signal in the *Drosophila* Embryo. *Nature* 329, 167-170.
- Lin, H. & Spradling, A.C. (1997). A novel group of pumilio mutations affects the asymmetric division of germline stem cells in the *Drosophila* ovary. *Development* 124, 2463-2476.
- Mak, W., Fang, C., Holden, T., Dratver, M.B. & Lin, H. (2016). An Important Role of Pumilio 1 in Regulating the Development of the Mammalian Female Germline. *Biol. Reprod.* 94, 134.
- Martin, K.C. & Ephrussi, A. (2009). mRNA localization: gene expression in the spatial dimension. *Cell* 136, 719-730.
- Masui, Y. & Clarke, H.J. (1979). Oocyte maturation. *Int. Rev. Cytol.* 57, 185-282.
- McGrew, L.L., Dworkin-Rastl, E., Dworkin, M.B. & Richter, J.D. (1989). Poly(A) elongation during *Xenopus* oocyte maturation is required for translational recruitment and is mediated by a short sequence element. *Genes Dev.* 3, 803-815.
- Mendez, R., Barnard, D. & Richter, J.D. (2002). Differential mRNA translation and meiotic progression require Cdc2-mediated CPEB destruction. *EMBO J.* 21, 1833-1844.
- Murata, Y. & Wharton, R.P. (1995). Binding of pumilio to maternal hunchback mRNA is required for posterior patterning in *Drosophila* embryos. *Cell* 80, 747-756.
- Nakahata, S., Katsu, Y., Mita, K., Inoue, K., Nagahama, Y. & Yamashita, M. (2001). Biochemical identification of *Xenopus* Pumilio as a sequence-specific cyclin B1 mRNA-binding protein that physically interacts with a Nanos homolog, Xcat-2, and a cytoplasmic polyadenylation element-binding protein. *J. Biol. Chem.* 276, 20945-20953.
- Nakahata, S., Kotani, T., Mita, K., Kawasaki, T., Katsu, Y., Nagahama, Y. & Yamashita, M. (2003). Involvement of *Xenopus* Pumilio in the translational regulation that is specific to cyclin B1 mRNA during oocyte maturation. *Mech. Dev.* 120, 865-880.
- Nukada, Y., Horie, M., Fukui, A., Kotani, T. & Yamashita, M. (2015). Real-time imaging of actin filaments in the zebrafish oocyte and embryo. *Cytoskeleton* 72, 491-501.
- Ota, R., Kotani, T. & Yamashita, M. (2011a). Biochemical characterization of Pumilio1

- and Pumilio2 in *Xenopus* oocytes. *J. Biol. Chem.* 286, 2853-2863.
- Ota, R., Kotani, T. & Yamashita, M. (2011b). Possible involvement of Nemo-like kinase 1 in *Xenopus* oocyte maturation as a kinase responsible for Pumilio1, Pumilio2, and CPEB phosphorylation. *Biochemistry* 50, 5648-5659.
- Padmanabhan, K. & Richter, J.D. (2006). Regulated Pumilio-2 binding controls RINGO/Spy mRNA translation and CPEB activation. *Genes Dev.* 20, 199-209.
- Pique, M., Lopez, J.M., Foissac, S., Guigo, R. & Mendez, R. (2008). A combinatorial code for CPE-mediated translational control. *Cell* 132, 434-448.
- Radford, H.E., Meijer, H.A. & de Moor, C.H. (2008). Translational control by cytoplasmic polyadenylation in *Xenopus* oocytes. *Biochim. Biophys. Acta* 1779, 217-229.
- Salazar, A.M., Silverman, E.J., Menon, K.P. & Zinn, K. (2010). Regulation of synaptic Pumilio function by an aggregation-prone domain. *J. Neurosci.* 30, 515-522.
- Sheets, M.D., Fox, C.A., Hunt, T., Vande Woude, G. & Wickens, M. (1994). The 3'-untranslated regions of c-mos and cyclin mRNAs stimulate translation by regulating cytoplasmic polyadenylation. *Genes Dev.* 8, 926-938.
- Spassov, D.S. & Jurecic, R. (2002). Cloning and comparative sequence analysis of PUM1 and PUM2 genes, human members of the Pumilio family of RNA-binding proteins. *Gene* 299, 195-204.
- Spassov, D.S. & Jurecic, R. (2003a). Mouse Pum1 and Pum2 genes, members of the Pumilio family of RNA-binding proteins, show differential expression in fetal and adult hematopoietic stem cells and progenitors. *Blood Cells Mol. Dis.* 30, 55-69.
- Spassov, D.S. & Jurecic, R. (2003b). The PUF family of RNA-binding proteins: does evolutionarily conserved structure equal conserved function? *IUBMB life* 55, 359-366.
- Stutz, A., Conne, B., Huarte, J., Gubler, P., Volkel, V., Flandin, P. & Vassalli, J.D. (1998). Masking, unmasking, and regulated polyadenylation cooperate in the translational control of a dormant mRNA in mouse oocytes. *Genes Dev.* 12, 2535-2548.
- Takahashi, K., Kotani, T., Katsu, Y. & Yamashita, M. (2014). Possible involvement of insulin-like growth factor 2 mRNA-binding protein 3 in zebrafish oocyte maturation as a novel cyclin B1 mRNA-binding protein that represses the

- translation in immature oocytes. *Biochem. Biophys. Res. Commun.* 448, 22-27.
- Takei, N., Nakamura, T., Kawamura, S., Takada, Y., Satoh, Y., Kimura, A.P. & Kotani, T. (2018). High-Sensitivity and High-Resolution In Situ Hybridization of Coding and Long Non-coding RNAs in Vertebrate Ovaries and Testes. *Biol. Proced. Online* 20, 6.
- Tanaka, T. & Yamashita, M. (1995). Pre-MPF is absent in immature oocytes of fishes and amphibians except *Xenopus*. *Dev. Growth Differ.* 37, 387-393.
- Tay, J., Hodgman, R. & Richter, J.D. (2000). The control of cyclin B1 mRNA translation during mouse oocyte maturation. *Dev. Biol.* 221, 1-9.
- Vassalli, J.D., Huarte, J., Belin, D., Gubler, P., Vassalli, A., O'Connell, M.L., Parton, L.A., Rickles, R.J. & Strickland, S. (1989). Regulated polyadenylation controls mRNA translation during meiotic maturation of mouse oocytes. *Genes Dev.* 3, 2163-2171.
- Weill, L., Belloc, E., Bava, F.A. & Mendez, R. (2012). Translational control by changes in poly(A) tail length: recycling mRNAs. *Nature Struct. Mol. Biol.* 19, 577-585.
- White, E.K., Moore-Jarrett, T. & Ruley, H.E. (2001). PUM2, a novel murine puf protein, and its consensus RNA-binding site. *RNA* 7, 1855-1866.
- Wickens, M., Bernstein, D.S., Kimble, J. & Parker, R. (2002). A PUF family portrait: 3'UTR regulation as a way of life. *Trends Genet.* 18, 150-157.
- Yasuda, K., Kotani, T., Ota, R. & Yamashita, M. (2010). Transgenic zebrafish reveals novel mechanisms of translational control of cyclin B1 mRNA in oocytes. *Dev. Biol.* 348, 76-86.
- Zamore, P.D., Williamson, J.R. & Lehmann, R. (1997). The Pumilio protein binds RNA through a conserved domain that defines a new class of RNA-binding proteins. *RNA* 3, 1421-1433.
- Zhang, B., Gallegos, M., Puoti, A., Durkin, E., Fields, S., Kimble, J. & Wickens, M.P. (1997). A conserved RNA-binding protein that regulates sexual fates in the *C. elegans* hermaphrodite germ line. *Nature* 390, 477-484.
- Zhang, M., Chen, D., Xia, J., Han, W., Cui, X., Neuenkirchen, N., Hermes, G., Sestan, N. & Lin, H. (2017). Post-transcriptional regulation of mouse neurogenesis by Pumilio proteins. *Genes Dev.* 31, 1-16.

Figure Legends

Fig. 1. Phosphorylation of Pum1 during oocyte maturation in zebrafish. (A and B) *In situ* hybridization analysis of *cyclin B1* mRNA (green) in immature (A) and mature (B) oocytes. DNA is shown in blue. The chorion is outlined by broken lines. Dormant *cyclin B1* mRNA is stored as RNA granules localized in the animal polar cytoplasm of immature oocytes. These granules disappear during oocyte maturation. m, micropyle; f, follicle cells; c, chorion. Bars: 10 μ m. (C) PAT assay of *cyclin B1* mRNA in immature (Im) and mature (M) oocytes. The poly(A) tails of *cyclin B1* mRNA are short in immature oocytes and become elongated during oocyte maturation (arrowhead). (D and E) SDS-PAGE analyses of crude extracts (D) and immunoprecipitations with anti-Pum1 antibody (E) from immature (Im) and mature (M) oocytes followed by immunostaining with anti-Pum1 antibody. Asterisk shows background signals of yolk protein. The electrophoretic mobility of Pum1 is reduced in mature oocytes. (F and G) SDS-PAGE analyses of immunoprecipitations with anti-Pum1 antibody from immature (Im) and mature (M) oocytes stimulated with (+) or not stimulated with (-) CIAP (F) and λ -PP (G). The reduction in electrophoretic mobility of Pum1 in mature oocytes is recovered partially by CIAP and almost completely by λ -PP. Similar results were obtained from two independent experiments.

Fig. 2. Time course analyses of Pum1 phosphorylation after initiation of oocyte maturation. (A and B) Time course of PAT assay for *cyclin B1* mRNA (A) and GVBD (B) after MIH stimulation. Arrowheads indicate signals of poly(A) tail elongation. (C and D) Time course of Pum1 phosphorylation after MIH stimulation using crude oocyte extracts. A red arrow indicates an initial signal of phosphorylation. Black arrowheads indicate apparent phosphorylation. White arrowheads indicate significant phosphorylation. Dotted lines indicate the basal size of Pum1. Similar results were obtained from two independent experiments. (E and F) Time course of Pum1 phosphorylation using immunoprecipitations with anti-Pum1 antibody (E) and GVBD (F) after MIH stimulation. A red arrow indicates an initial signal of phosphorylation. A black arrowhead indicates apparent phosphorylation. A white arrowhead indicates significant phosphorylation. Pum1 phosphorylation began from 10 to 30 min after MIH stimulation. Similar results were obtained from five independent experiments.

Fig. 3. Effects of actin filament depolymerization on Pum1 phosphorylation and translational regulation of *cyclin B1* mRNA. (A and B) *In situ* hybridization analysis of *cyclin B1* mRNA (green) in oocytes treated with DMSO (Control) and cytochalasin B (CytoB). DNA is shown in blue. The chorion is outlined by broken lines. *cyclin B1* RNA granules are maintained in control oocytes (A), while they are dispersed throughout the cytoplasm in CytoB-treated oocytes (B). m, micropyle; f, follicle cells; c, chorion. Bars: 10 μ m. (C) Immunoblotting of Pum1 (upper) and CPEB (middle) and PAT assay for *cyclin B1* mRNA (lower) in oocytes treated with (+) or not treated with (-) CytoB or MIH. Arrowheads indicate Pum1 phosphorylation (upper) and poly(A) tail elongation of *cyclin B1* mRNA (lower). Similar results were obtained from four independent experiments. (D) Time course of CPEB phosphorylation after CytoB treatment. An arrowhead indicates a phosphorylated form of CPEB. (E) RT-PCR analysis of immunoprecipitations with control IgG (IgG) or anti-Pum1 (α -Pum1) antibody in control and CytoB-treated oocytes. (F) Time course of Cyclin B1 synthesis after stimulation with MIH in control and CytoB-treated oocytes. Arrows indicate initial signals of Cyclin B1 synthesis. Arrowheads indicate active forms of the Cdc2 kinase. Asterisks show nonspecific bands. Time course of CPEB phosphorylation and degradation is shown in control oocytes. (G) Time course of GVBD after stimulation with MIH in control and CytoB-treated oocytes. Similar results were obtained from three independent experiments.

Fig. 4. Effects of overexpression of Pum1N on endogenous Pum1 phosphorylation and translational regulation of *cyclin B1* mRNA. (A) Schematic diagrams of zebrafish Pum1 and Pum1N. The PUF domain is responsible for binding to PBE in target mRNAs. The Q/N-rich region is thought to promote protein aggregation. (B) *In situ* hybridization of *cyclin B1* mRNA (red) and immunostaining of GFP (green) in oocytes expressing GFP or GFP-Pum1N before (0 min) or at 90 min after stimulation with MIH. Bars: 5 μ m. (C) Pum1 phosphorylation in control or GFP-Pum1N-expressing oocytes stimulated with (+) or not stimulated with (-) MIH. Pum1 is phosphorylated in control oocytes but not in GFP-Pum1N-expressing oocytes after stimulation with MIH. Similar results were obtained from three independent experiments. (D) Time course of GVBD after stimulation with MIH in control and GFP-Pum1N-expressing oocytes. Similar results were obtained from three independent experiments.

Fig. 1

Fig. 2

Fig. 3

Fig. 4