Title	Evaluation of sensory loss obtained by modified-thoracoabdominal nerves block through perichondrial approach in patients undergoing gynecological laparoscopic surgery : a prospective observational study
Author(s)	Aikawa, Katsuhiro; Yokota, Isao; Maeda, Yosuke; Morimoto, Yuji
Citation	Regional Anesthesia & Pain Medicine, 47(2), 134-135 https://doi.org/10.1136/rapm-2021-102870
Issue Date	2022-02
Doc URL	http://hdl.handle.net/2115/83430
Rights	This article has been accepted for publication in Regional Anesthesia & Pain Medicine, 2022 following peer review, and the Version of Record can be accessed online at http://dx.doi.org/10.1136/rapm-2021-102870 . " © Authors (or their employer(s))OR " © BMJ Publishing Group Ltd " (for assignments of BMJ Case Reports) " 2022 " [Add where a funder mandates: " Reuse of this manuscript version (excluding any databases, tables, diagrams, photographs and other images or illustrative material included where a another copyright owner is identified)is permitted strictly pursuant to the terms of theCreative Commons Attribution-Non Commercial 4.0 International (CC-BY-NC 4.0) http://creativecommons.org https://creativecommons.org/licenses/by-nc/4.0/ "]
Rights(URL)	https://creativecommons.org/licenses/by-nc/4.0/
Type	article (author version)
File Information	Reg Anesth Pain Med rapm-2021-102870.pdf

Instructions for use

**Evaluation of Sensory Loss Obtained by Modified-Thoracoabdominal Nerves Block
Through Perichondrial Approach in Patients Undergoing Gynecological Laparoscopic
Surgery: A Prospective Observational Study**

Katsuhiro Aikawa, MD, PhD,[†] Isao Yokota, PhD, MPH,[‡] Yosuke Maeda, MD,[†] and Yuji

Morimoto, MD, PhD[†]

From the

[†]Department of Anesthesiology and Critical Care Medicine, Faculty of Medicine and Graduate School of Medicine, Sapporo, Japan

[‡]Department of Biostatistics, Faculty of Medicine and Graduate School of Medicine, Sapporo, Japan

Address correspondence to:

Katsuhiro Aikawa, MD, PhD, Department of Anesthesiology and Critical Care Medicine, Faculty of Medicine and Graduate School of Medicine, Hokkaido University, Kita-15, Nishi-7, Kita-ku, Sapporo 060-8638, Japan

Phone/Fax: +81-11-706-7861 (Email: katsuhiro.aikawa@med.hokudai.ac.jp)

ORCID iD

Katsuhiro Aikawa <http://orcid.org/0000-0003-0871-3956>

Running Head:

Evaluation of Sensory Loss of M-TAPA

Funding:

The authors have no sources of funding to declare for this research.

Conflicts of Interest:

The authors declare no conflicts of interest for this research.

Key words

modified-thoracoabdominal nerves block through perichondrial approach (M-TAPA)

thoracoabdominal nerves

Word count:

Trial registry number

UMIN000040296

Abbreviations

M-TAPA: modified-thoracoabdominal nerves block through perichondrial approach

ASA: American Society of Anesthesiologists

IQR: interquartile range

LA: local anesthetic

TAP: transversus abdominis plane

TAPB: transversus abdominis plane block

1 INTRODUCTION

2 Recently, Tulgar et al. reported modified-thoracoabdominal nerves block through perichondrial
3 approach (M-TAPA).¹ However, studies showing the area blocked by M-TAPA are lacking.

METHODS

In this prospective observational study, 30 patients undergoing gynecological laparoscopic surgery were recruited after obtaining ethical approval and written informed consent. Exclusion criteria were: ASA ≥ 3 , age < 20 or > 70 years, pregnancy, local anesthetics allergy, weight < 45 kg, body mass index > 35 kg/m², history of thoracotomy or laparotomy except for lower abdominal laparoscopic surgery, communication disorders, neuropathy, or operation time of > 6 hours.

M-TAPA technique

M-TAPA was performed bilaterally by a single anesthesiologist (KA) after general anesthesia induction (see Figure 1A–D). The needle tip was repositioned if intramuscular spread was observed.

Sensory level assessment

An investigator (KA or YM) assessed sensory levels T4–L1 using a pinprick test 2 hours postoperatively. The anterior and lateral cutaneous branches were evaluated at a vertical line 3–5 cm from the midline and midaxillary lines, respectively. A vertical force (1 newton) was visualized using a tension gauge. A 3-point numerical scale (0 = no pain, 1 = decreased pain, 2 = normal pain) was used: values of 0 or 1 were defined as effective. Normal sensation on the

22 shoulder was used for comparison.

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40 **RESULTS**

41 The median (IQR) highest sensory level was T7 (T5–8) in the anterior and T9 (T7–10) in the
42 lateral area. Sensory loss was not observed in the lateral area in five patients. The median (IQR)
43 number of blocked dermatomes was 5 (7–4) in the anterior and 4 (5–2) in the lateral area (Figure
44 2).

DISCUSSION

This study evaluated the sensory levels of anterolateral thoracoabdominal wall after performing M-TAPA using 25 mL of 0.25% ropivacaine and demonstrated its inconsistency. Because 10th costal cartilage usually contacts the transversus abdominis muscle, the injected local anesthetic (LA) spreads along the transversus abdominis plane (TAP). However, our results indicated that the LA pathway differed between TAP block (TAPB) and M-TAPA. Borglum et al.² reported that LA injected to the TAP does not overcome the linea semilunaris. The anterior abdominal wall is innervated by the upper (T6–9) and lower TAP plexus (T10–12) divided by the linea semilunaris; therefore, the hydrodissection technique³ or dual TAPB⁴ is required to simultaneously anesthetize both areas. Considering the LA was injected outside the linea semilunaris, and most patients had some sensory loss in the lateral area, the LA might spread laterocranially, resulting in a sensory block. Zinboonyahgoon et al.⁵ reported that a single LA injection to the endothoracic fascial plane, positioned immediately beneath the ribs, could achieve multi-level intercostal nerve block. Thus, that compartment could be the responsible pathway of M-TAPA. LA spread of fascial plane blocks is influenced by multiple factors.⁶ Therefore, the features of the individual fascia allowing cranial and lateral spread may affect the sensory area. The lower rate of dermatomal coverage of the T12 and L1 in the anterior area may be because of cranial needle direction.

This study has several limitations. Sensory areas were assessed postoperatively, and a previous

study demonstrated sensory ranges regressing over time.⁴ Hence, our results may be less than the maximum height and range. Furthermore, we cannot deny the existence of performance bias and the potential residual effect of opioids. Because of several incisions and dressings, we could not evaluate the blocked area as a plane, but only on the representative longitudinal lines.

81 **CONCLUSION**

82 The area blocked by M-TAPA varies and provides limited dermatomal coverage than
83 previously reported. Anterior sensory coverage was marginally better than lateral coverage.
84 Future studies should investigate the corresponding fascial plane, and perioperative analgesic
85 efficiency of M-TAPA.

86 **ACKNOWLEDGEMENTS**

87 We would like to thank the nurses in the gynecologic surgical ward for supporting our pinprick
88 tests, and Editage (www.editage.com) for English language editing.

REFERENCES

- 89 1 Tulgar S, Selvi O, Thomas DT, *et al.* Modified thoracoabdominal nerves block through
90 perichondrial approach (M-TAPA) provides effective analgesia in abdominal surgery and
91 is a choice for opioid sparing anesthesia. *J Clin Anesth* 2019;55:109.
- 92 2 Børglum J, Jensen K, Christensen AF, *et al.* Distribution patterns, dermatomal anesthesia,
93 and ropivacaine serum concentrations after bilateral dual transversus abdominis plane
94 block. *Reg Anesth Pain Med* 2012;37:294-301.
- 95 3 Hebbard PD, Barrington MJ, Vasey C. Ultrasound-guided continuous oblique subcostal
96 transversus abdominis plane blockade: description of anatomy and clinical technique. *Reg*
97 *Anesth Pain Med* 2010;35:436-41.
- 98 4 Lee TH, Barrington MJ, Tran TM, *et al.* Comparison of extent of sensory block following
99 posterior and subcostal approaches to ultrasound-guided transversus abdominis plane
100 block. *Anaesth Intensive Care* 2010;38:452-60.
- 101 5 Zinboonyahgoon N, Luksanapruksa P, Piyaselakul S, *et al.* The ultrasound-guided
102 proximal intercostal block: anatomical study and clinical correlation to analgesia for breast
103 surgery. *BMC Anesthesiol* 2019;19:94.
- 104 6 Black ND, Stecco C, Chan VWS. Fascial Plane Blocks: More Questions Than Answers?
105 *Anesth Analg* 2021;132:899-905.

LEGENDS for figures

Figure 1: M-TAPA technique

A: A high-frequency linear probe (6–13 MHz) is placed in the sagittal direction at the 10th costal margin, and transversus abdominis, internal oblique, and external oblique muscles are identified.

B: An 8-cm 18G Tuohy needle is inserted in the cranial direction using an in-plane technique. C: The needle tip is moved to the posterior aspect of the 10th costal cartilage, following 1 mL of saline injection to confirm the correct spread. D: Subsequently, 25 mL of 0.25% ropivacaine is administered along with an intermittent aspiration test. It should be noted that the needle tip never proceeded over the cranial edge of the 10th costal cartilage.

Abbreviations: C, costal cartilage. EOM, external oblique muscle. IOM, internal oblique muscle.

TAM, transversus abdominis muscle. LA: local anesthetics.

The white triangle symbolizes the 18G Touhy needle.

Figure 2: Percentages of successful blocks in each dermatome.

Anterior Area

Lateral Area