Title	Changes in subunit composition of NMDA receptors in animal models of schizophrenia by repeated administration of methamphetamine
Author(s)	Oka, Matsuhiko; Ito, Koki; Koga, Minori; Kusumi, Ichiro
Citation	Progress in neuro-psychopharmacology & biological psychiatry, 103, 109984 https://doi.org/10.1016/j.pnpbp.2020.109984
Issue Date	2020-12-20
Doc URL	http://hdl.handle.net/2115/83602
Rights	© 2020. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	Prog Neuropsychopharmacol Biol Psychiatry 103_109984.pdf

[Instructions for use](#)

Changes in subunit composition of NMDA receptors in animal models of schizophrenia by repeated administration of methamphetamine

Matsuhiko Oka^{a)}, Koki Ito^{a)}, Minoru Koga^{b)}, Ichiro Kusumi^{a)}

^{a)} Department of Psychiatry, Hokkaido University Graduate School of Medicine
West 7, North 15, Kita Ward, Sapporo City, Hokkaido, 060-8638, JAPAN

^{b)} Department of Psychiatry, National Defense Medical College
3-2, Namiki, Tokorozawa Saitama, 359-8513, Japan

Koki Ito: itokoki502@gmail.com

Minoru Koga: mkoga3@ndmc.ac.jp

Ichiro Kusumi: ikusumi@med.hokudai.ac.jp

Corresponding author:

Matsuhiko Oka

E-mail: m_oka@pop.med.hokudai.ac.jp

Department of Psychiatry, Hokkaido University Graduate School of Medicine
West 7, North 15, Kita Ward, Sapporo City, Hokkaido, 060-8638, JAPAN

Abstract

The dopamine and glutamate hypotheses reflect only some of the pathophysiological changes associated with schizophrenia. We have proposed a new “comprehensive progressive pathophysiology model” based on the “dopamine to glutamate hypothesis.” Repeated administration of methamphetamine (METH) at a dose of 2.5 mg/kg in rats has been used to assess dynamic changes in the pathophysiology of schizophrenia. Previous use of this model suggested *N*-methyl-D-aspartate receptor (NMDA-R) dysfunction, but the mechanism could only be inferred from limited, indirect observations. In the present study, we used this model to investigate changes in the expression of NMDA-R subunits. Repeated administration of METH significantly decreased the gene expression levels of glutamate ionotropic receptor NMDA type subunit (*Grin*) subtypes *Grin1* and *Grin2c* in the prefrontal cortex (PFC), *Grin1* and *Grin2a* in the hippocampus (HPC), and *Grin1*, *Grin2b*, and *Grin2d* in the striatum (ST). We observed a significant difference in *Grin1* expression between the PFC and ST. Furthermore, repeated administration of METH significantly decreased the protein expression of GluN1 in both cytosolic and synaptosomal fractions isolated from the PFC, and significantly decreased the protein expression of GluN1 in the cytosolic fraction, but not the synaptosomal fraction from the ST. These regional differences may be due to variations in the synthesis of GluN1 or

intracellular trafficking events in each area of the brain. Considering that knockdown of *Grin1* in mice affects vulnerability to develop schizophrenia, these results suggest that this model reflects some of the pathophysiological changes of schizophrenia, combining both the dopamine and glutamate hypotheses.

Keywords

Schizophrenia, NMDA receptor, GluN1, Animal model, Methamphetamine

Abbreviations

CNS, Central nervous system; HPC, Hippocampus; METH, Methamphetamine; NAc, Nucleus accumbens; NMDA-R, *N*-methyl-D-aspartate receptor; PBS, Phosphate-buffered saline; PFC, Prefrontal cortex; PPI, Prepulse inhibition; qRT-PCR, quantitative reverse transcription polymerase chain reaction; Sal, Saline; s.c., Subcutaneous injection; ST, Striatum; TBST, Tris-buffered saline containing 0.5 % Tween 20

1. Introduction

Schizophrenia is a mental illness that affects approximately 1% of the general population, regardless of sex, race, or nationality. The onset of schizophrenia typically occurs in the late teens to thirties, manifesting as a variety of symptoms, most of which are chronic in nature (Carlsson *et al.*, 1997; Lewis and Lieberman, 2000; Van Os. *et al.*, 2010). The precise etiology of schizophrenia remains controversial, but there is consensus that it is a multifactorial neurodevelopmental disorder that is influenced by both genetic and environmental factors. The pathology of schizophrenia progresses from hyperactivity of dopaminergic systems, which was originally posited by the dopamine hypothesis, to *N*-methyl-D-aspartate receptor (NMDA-R) dysfunction due to changes in expression or composition of NMDA-R subunits (Olney and Farber, 1995; Goff and Coyle, 2001).

NMDA-Rs, which are widely distributed throughout the central nervous system (CNS), are essential mediators of synaptic transmission and neuronal plasticity. NMDA-Rs are tetrameric receptors composed of two essential GluN1 subunits along with two GluN2 or GluN3 subunits, which have four (GluN2A - GluN2D) and two subtypes

(GluN3A and GluN3B), respectively. These NMDA-R subtypes differ in their molecular (subunit) composition, which is plastic and changes during development and in response to alterations in neuronal activity (Cull-Candy and Leszkiewicz, 2004; Traynelis *et al.*, 2010; Paoletti *et al.*, 2013). The protein name mirrors the gene name, with just the two-letter code difference (i.e., Grin1 translates to GluN1, Grin2a translates to GluN2A).

Postmortem analyses of brains taken from patients with schizophrenia have reported reduced expression of glutamate ionotropic receptor NMDA type subunit (Grin) subtypes GRIN1 and GRIN2C in the prefrontal cortex (PFC) (Akbarian *et al.*, 1996; Weickert *et al.*, 2013; Catts *et al.*, 2016), and GRIN1 and GRIN2A in the hippocampus (HPC) (Gao *et al.*, 2000; Law and Deakin, 2001); these genes encode three specific subunits of NMDA-Rs, GluN1, GluN2C, and GluN2A respectively. A meta-analysis has determined effect sizes for changes in mRNA and protein expression levels of the essential GluN1 subunit in the PFC in schizophrenia. In schizophrenic patients, compared to unaffected controls, the pooled effect size was -0.64 (95% confidence interval: -1.08 to -0.20) and -0.44 (95% confidence interval: -0.80 to -0.07) for reductions in GluN1 mRNA and protein expression, respectively (Catts *et al.*, 2016).

We previously reported that repeated administration of methamphetamine (METH), which increases dopamine levels in the nucleus accumbens (NAc) at a dose of 2.5 mg/kg, but not at 1.0 mg/kg, also increased glutamate levels in the medial prefrontal cortex (mPFC) and the NAc (Ito *et al.*, 2006a). At this dose, repeated administration of METH results in the following: (1) development of behavioral cross-sensitization to MK -801 (a non-competitive NMDA-R antagonist) (Ito *et al.*, 2006a); (2) prepulse inhibition (PPI) deficit (Abekawa *et al.*, 2008), which is an indicator of cognitive dysfunction; and (3) induction of apoptosis in the PFC (Abekawa *et al.*, 2008), indicating brain atrophy. In addition, administration of atypical antipsychotics and mood stabilizers attenuates some or all of these changes (Ito *et al.*, 2006b; Abekawa *et al.*, 2008; Nakato *et al.*, 2010; Abekawa *et al.*, 2011; Nakato *et al.*, 2011).

Based on these pathophysiological changes induced by repeated METH administration, we proposed a new “comprehensive progressive pathophysiology model” based on the “dopamine to glutamate hypothesis” (Abekawa *et al.*, 2012). This model mimics the dysfunction of NMDA-Rs in schizophrenia. However, in our previous study, the precise molecular mechanisms could not be determined because changes could only be inferred from indirect evidence (Table 1).

Table 1. Electrophysiological, molecular, and behavioral changes caused by repeated administration of METH.

	1.0 mg/kg	2.5 mg/kg
Development of behavioral sensitization (Ito <i>et al.</i> , 2006a)	+	+
Delayed increases in glutamate levels in the NAc and PFC (Ito <i>et al.</i> , 2006b; Abekawa <i>et al.</i> , 2008)	–	+
Development of behavioral cross-sensitization to MK-801 (Ito <i>et al.</i> , 2006b; Abekawa <i>et al.</i> , 2008)	–	+
PPI deficit (Abekawa <i>et al.</i> , 2008)	–	+
Apoptosis in the PFC (Abekawa <i>et al.</i> , 2008)	–	+

METH, Methamphetamine; NAc, Nucleus accumbens; PFC, Prefrontal cortex; PPI, Prepulse inhibition

116

117 Using this model, we aimed to assess changes in both the gene and protein expression
118 levels of NMDA-R subunits, including *Grin1*, *Grin2a*, *Grin2b*, *Grin2c*, and *Grin2d* in the
119 PFC, the HPC, and the striatum (ST) to more precisely elucidate the molecular
120 mechanisms driving changes in receptor functionality that mediate behavior.

121

122

123 2. Materials and methods

124

125 2.1. Animals

126 Seven-week-old male Sprague–Dawley rats (Sankyo Labo Service Corporation,
127 Inc., Japan), weighing 210–230 g at the start of the experiment, were housed in plastic
128 cages with dimensions of 30 × 25 × 18 cm, with a wire mesh top and sawdust bedding
129 (two rats / cage). The colony room was under controlled lighting (lights on from 7:00
130 A.M. to 7:00 P.M.), temperature (23 ± 1 °C), and humidity ($50 \pm 10\%$). Animals were
131 allowed free access to standard laboratory chow and tap water. Animals were handled
132 daily for at least four days before the start of the experiment and were tested only once in
133 each experiment. All experiments were ethically approved by the Animal Research

Committee of Hokkaido University (permission number 13–0137, 17–0086) and performed in accordance with ARRIVE guidelines, the Guide for the Care and Use of Laboratory Animals of Hokkaido University, and the guidelines established by the US National Institutes of Health guide for the care and use of laboratory animals (NIH Publications No. 8023, revised 1978).

2.2. Drugs and administration schedules

For this study, METH (Dainippon Sumitomo Pharma Co., Ltd., Japan) was dissolved in sterile physiological saline and injected subcutaneously at a volume of 1.0 mL/kg at a dose of 2.5 mg/kg. The control group was subcutaneously administered 1.0 mL/kg of saline. Injections were repeated five times on alternating days (treatment period). To avoid acute, confounding effects of having the drug on-board during testing, we allowed time for a sufficient withdrawal (wash-out) period (6–9 days), then rats were anesthetized by intraperitoneal injection of pentobarbital (30 mg/kg), decapitated, and the PFC, HPC, and ST were collected (the experimental timeline is shown in Fig. 1). The METH dose (2.5 mg/kg) was selected based on the optimal manifestation of electrophysiological, molecular, and behavioral changes we observed in our previous study (Table 1).

Fig. 1 Schema of the experimental timeline.

Rats were injected with saline or METH five times, followed by a 6–9 days withdrawal period before tissue collection.

METH, methamphetamine; PFC, prefrontal cortex; HPC, hippocampus; ST, striatum; s.c., subcutaneous injection.

2.3. Quantitative reverse transcription polymerase chain reaction (qRT-PCR)

Immediately after tissue collection, the right hemisphere of each brain was rapidly immersed in RNAlater™ Stabilization Solution (Thermo Fisher Scientific, USA), maintained at 4 °C for two days, then stored at –80°C until use.

Total RNA was extracted from the PFC, HPC, and ST of the right hemisphere using an SV Total RNA Isolation System (Z3105, Promega, USA). The cDNA was obtained by reverse transcription using ReverTra Ace® qPCR RT Master Mix (Toyobo Co., Ltd., Japan). PCR was performed using Thunderbird SYBR qPCR mix (Toyobo Co.,

Ltd., Japan); each sample was measured in triplicate. The amplification parameters were: 40 cycles of denaturing at 95 °C for 15 s and annealing and extension at 60 °C for 30 s. For each cycle, the fluorescent emission of SYBR green was quantified for each sample and used to calculate the threshold cycle numbers (Ct). The reaction conditions for each primer set for *Grin1*, *Grin2a*, *Grin2b*, *Grin2c*, *Grin2d*, and *Actb* genes were selected based on previous studies, which are shown in Supplementary Table 1.

The expression level of each gene was determined using the $\Delta\Delta\text{CT}$ method for qRT-PCR. *Actb*, the gene encoding β -actin levels were measured as the internal loading control. Relative gene expression was calculated using the 2- ddCt method.

2.4. Western blotting

Immediately after tissue collection, the left hemisphere of each brain was rapidly homogenized with a glass homogenizer using an ice-cold synaptosome isolation reagent (Syn-PER: Thermo Fisher Scientific, USA) containing 1 $\times$ protease inhibitor cocktail (cOmplete: Sigma-Aldrich, USA). The homogenate was centrifuged at 1200 $\times g$ for 10 min at 4 °C. The supernatant was transferred to a new tube and centrifuged at 15,000 $\times g$ for 20 min at 4 °C. After centrifugation, the supernatant was stored as the cytosolic fraction at -80 °C until protein analysis. The remaining pellet was resuspended in 1 $\times$

186 phosphate-buffered saline (PBS; pH 7.4, catalog number: 048–29,805, FUJIFILM Wako
187 Pure Chemical Corporation, Japan) and stored as the synaptosomal fraction at –80 °C
188 until protein analysis.

189 To quantify protein expression in the cellular fractions from each brain region,
190 10 µg of the cytoplasmic fraction and 2 µg of the synaptosome fraction were subjected to
191 gel electrophoresis at a constant voltage of 200 V for 30 min (4–15% Mini-PROTEAN®
192 TGX™: Bio-Rad), then transferred at a constant voltage of 100 V for 1 h to Amersham
193 Hybond polyvinylidene fluoride (PVDF) membranes (10,600,057, GE Healthcare Life
194 Science, USA) with a pore size of 0.2 µm in a Tris-glycine transfer buffer (25 mM tris
195 base, 192 mM glycine, 20% methanol). The samples were blocked for 1 h with 2% skim
196 milk in TBST (Tris-buffered saline containing 0.5% Tween 20) at room temperature, then
197 incubated overnight at 4 °C with the following primary antibodies: β-actin (1:1000
198 dilution; mouse monoclonal, 010–27,841, FUJIFILM Wako Pure Chemical Corporation,
199 Japan) and GluN1 (1:500 dilution; rabbit monoclonal, ab109182, Abcam, USA). After
200 washes with TBST (3 × 10 min), the blots were incubated for 1 h at room temperature
201 with horseradish peroxidase (HRP)-conjugated secondary antibodies (goat anti-rabbit
202 IgG or goat anti-mouse IgG; 1: 5000, Amersham Biosciences, UK), followed by washes
203 with TBST (3 × 10 min). The samples were incubated with ImmunoStar LD (FUJIFILM

Wako Pure Chemical Industries, Japan), a luminescent substrate, and images were captured using ImageQuant LAS 4000 (GE Healthcare Life Science, USA). The NIH ImageJ software was used to quantify the protein bands by densitometry. β -actin was used as the loading control.

2.5. Statistical analyses

JMP[®] Pro 14.0.0 (SAS Institute Inc., Cary, North Carolina, USA) software was used for statistical analyses. Group means of data quantified from qRT-PCR and Western blotting to assess mRNA and protein expression changes were compared using the Wilcoxon rank-sum test. Significance level was set as 0.05. Data are presented as mean $\pm$ standard error of the mean (*SEM*).

3. Results

The RNA expression levels of *Grin1*, *Grin2a*, *Grin2b*, *Grin2c*, and *Grin2d* in the PFC, HPC, and ST were quantified by qRT-PCR. Repeated administration of METH significantly decreased gene expression levels of *Grin1* ($Z = 3.17$, $p = .0015$) and *Grin2c* ($Z = 2.31$, $p = .0207$) in the PFC (Fig.2A), *Grin1* ($Z = 2.15$, $p = .0315$) and *Grin2a* ($Z =$

2.67, $p = .0076$) in the HPC (Fig.2B), and *Grin1* ($Z = 2.95$, $p = .0032$), *Grin2b* ($Z = 2.55$, $p = .0108$), and *Grin2d* ($Z = 2.09$, $p = .0364$) in the ST (Fig.2C). The expression of *Grin1* was significantly different in the PFC compared to the ST.

Fig. 2 Gene expression levels of NMDA-R subunits in various brain regions following repeated METH administration.

Repeated administration of 2.5 mg/kg METH significantly decreased expression levels of *Grin1* and *Grin2c* in the PFC (A), *Grin1* and *Grin2a* in the HPC (B), *Grin1*, *Grin2b*, and *Grin2d* in the ST (C).

Data are shown as mean $\pm$ SEM. * $p < .05$, ** $p < .0033$. NMDA-R, *N*-methyl-D-aspartate receptor; METH, methamphetamine; PFC, medial prefrontal cortex; HPC, hippocampus; ST, striatum; qRT-PCR, quantitative reverse transcription polymerase chain reaction.

($N = 30$ rats/group. Data were compared by Wilcoxon rank-sum tests with a

Bonferroni correction).

In order to validate this finding, we further analyzed the protein expression levels of GluN1 by western blotting in the cytosolic and synaptosomal fractions of the PFC and the ST, which were significantly decreased even after Bonferroni correction.

Repeated administration of METH significantly decreased the protein expression levels of GluN1 in both the cytosolic ($Z = 2.11$, $p = .0351$) and synaptosomal ($Z = 1.97$, $p = .0488$) fractions of tissue from the PFC (Fig.3A, C). Similarly, protein expression levels of GluN1 levels were also decreased in the cytosolic fractions ($Z = 2.57$, $p = .0102$) but not the synaptosomal fractions ($Z = 0.78$, $p = .4357$) of tissue from the ST (Fig.3B, D).

Fig. 3 Changes in protein expression levels of NMDA-R subunits in cellular fractions of tissue from various brain regions following repeated METH administration.

Repeated administration of METH significantly decreased the protein expression level of GluN1 in the cytosolic and synaptosomal fractions of the PFC (A, C), and the cytosolic fraction of the ST (B, D).

Data are expressed as the mean $\pm$ SEM. $*p < .05$. NMDA-R, *N*-methyl-D-aspartate receptor; METH, methamphetamine; PFC, prefrontal cortex; ST, striatum.

($N = 11-12$ rats/group; statistical comparisons were performed using the Wilcoxon rank-sum test).

4. Discussion

The results of this study suggest a decrease in population of NMDA-R subtype GluN1/2C or GluN1/2A/2C in the PFC, GluN1/2A in the HPC, and GluN1/2B, GluN1/2D, and GluN1/2B/2D in the ST. In accordance with our previous studies, we have determined the mechanism through which METH induces changes in NMDA-R expression and subunit composition. Dopamine is released following administration of METH, followed by glutamate release, which reduces levels of the GluN1 subunit, ultimately leading to decreased expression of functional NMDA-Rs. These alterations in gene expression of NMDA-R subunits are associated with the development of behavioral cross-sensitization to MK-801, PPI deficit, and apoptosis within the PFC (Table 1).

In the model used in this study, there was a decrease in the expression of GluN1 and decreased intracellular trafficking events that regulate NMDA-R expression (Fig.3A, B). After NMDA-Rs are synthesized in and exported from the endoplasmic reticulum, the receptors travel through the Golgi apparatus before being inserted into vesicles and directly trafficked to the plasma membrane or into dendrites. Alternatively, vesicles may be trafficked to dendritic Golgi outposts before reaching the cell membrane or to the

synapse (Horak *et al.*, 2014). Two molecules genetically linked to schizophrenia, neuregulin and serine–threonine phosphatase PP2B (also known as calcineurin) regulate these NMDA-R trafficking events. In a hypothetical model, activation of the ErbB4 receptor by neuregulin suppresses tyrosine phosphorylation of the GluN2A subunit, which promotes NMDA-R internalization. In another model, PP2B dephosphorylates and activates striatal-enriched tyrosine phosphatase (STEP), which, in turn, dephosphorylates tyrosine residues on GluN1 and/or GluN2 subunits to promote NMDA-R internalization (Lau *et al.*, 2007).

In this study, in the PFC, the protein expression level of GluN1 decreased in both the cytosolic and synaptosomal fractions following repeated METH administration, while in the ST, the protein expression level of GluN1 was significantly decreased in the cytosolic fraction, but there was no significant difference in expression in the synaptosomal fraction. These regional differences may be due to variable expression of functional NMDA-Rs, which is regulated by intracellular receptor trafficking (Ladépêche *et al.*, 2014). Depending on the number of NMDA-R subunits containing GluN1 and differences in intracellular trafficking mechanisms, the changes in NMDA-R expression may be differentially delayed across brain regions. It will be necessary to further assess

NMDA-R expression and associated intracellular trafficking in each region of the brain.

The METH models are considered limited, as they do not fully reflect the negative symptoms and cognitive impairment associated with the development of schizophrenia (Jones *et al.*, 2011; Marcotte *et al.*, 2001). There is also a theory that purely dopaminergic models may be unlikely to lead to a great improvement in efficacy or safety of antipsychotic drugs (Steeds *et al.*, 2015). In addition, pharmacological models using NMDA-R antagonists have been widely used but these typically produce only transient changes in behavior and brain function (Featherstone *et al.*, 2015). Because NMDA-R antagonists induce hypo-functionality of the receptor throughout the brain, these animal models cannot selectively target NMDA-Rs in specific neural circuits; thus, these models may fail to precisely elucidate the mechanisms resulting in the pathophysiology of schizophrenia (Olney *et al.*, 1999). At present, models mimicking the dysfunction of both presynaptic dopamine release and NMDA-R functionality may provide the best tools to explore the molecular, cellular, and behavioral aspects of schizophrenia (Howes *et al.*, 2015). While pharmacological models may never be able to accurately mimic all aspects of such a complex condition as schizophrenia, they may still be able to provide valuable insight into the neurobiological mechanisms underlying specific symptom domains

(Curran *et al.*, 2009). The METH model is a classical pharmacological model, but there are many points that need to be re-evaluated because multiple hypotheses are included depending on the protocol, as in this study.

Grin1 knockdown mice or GluN2A and GluN2B mutant mice are also commonly used animal models of schizophrenia (Lee *et al.*, 2019). GluN1-mediated deficits in either pyramidal or GABAergic neurons could cause an imbalance in neuronal excitation and inhibition in cortical neural circuitry, leading to development of behavioral phenotypes that mimic symptoms of schizophrenia. Due to overlapping roles of GluN2A and GluN2B subunits in learning and memory (Sakimura *et al.*, 1995; Kiyama *et al.*, 1998; Moriya *et al.*, 2000), GluN2A and GluN2B mutant mice would serve as great models to study the pathophysiology of cognitive changes associated with the development of schizophrenia. Heterozygous (GluN1 +/-) mice exhibit a 30% reduction in GluN1 receptor expression, and the current study suggests that these mice may be among the most sensitive models of increased vulnerability to schizophrenia (Featherstone *et al.*, 2015).

The model used in this study differs from the typical amphetamine and METH

models, which are based solely on the dopamine hypothesis; it is instead a model that reflects other aspects of the pathology of schizophrenia, combining both the dopamine and glutamate hypotheses. In studies with different protocols using METH, *Grin1* and GluN1 were not necessarily decreased (Simões *et al.*, 2007; González *et al.*, 2018), and in contrast to the results of the present study, GluN2B in the mPFC was decreased (Lominac *et al.*, 2016). The effect on NMDA-R varies with the dose of METH, and a protocol similar to that used in this study may be desirable to replicate both the dopaminergic overactivity and NMDA-R malfunction seen in schizophrenia.

In our model, we have identified four steps in the pathophysiological mechanisms driving changes in NMDA-R functionality. First, dopamine is released in the PFC and the NAc during psychotic episodes, with increased glutamate release in severe cases (Ito *et al.*, 2006b; Abekawa *et al.*, 2008). Second, increased glutamate release reduces GluN1 expression and the number of functional NMDA-Rs at the plasma membrane (Fig. 3). As NMDA-Rs become desensitized and dysfunctional, glutamate release is eventually reduced (Abekawa *et al.*, 2012). Third, these changes result in a disease state with increased susceptibility to NMDA-R antagonism (possibly unresponsive to D₂ receptor antagonists) (Ito *et al.*, 2006b; Abekawa *et al.*, 2008), besides

the recognized dysfunction (Abekawa *et al.*, 2008; Abekawa *et al.*, 2012). Finally, cerebral atrophy occurs when AMPA receptors (α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid receptor) are repeatedly stimulated during the phase of increased synaptic glutamate release (Abekawa *et al.*, 2008). Peripheral blood levels of D-serine binding to GluN1 may be the greatest biological marker for diagnosis and treatment of patients with schizophrenia (Ohnuma and Arai, 2011).

Although schizophrenia is considered a neurodevelopmental disorder with deficits occurring during early brain development, the majority of animal studies have been focused on METH or NMDA-R antagonist-induced changes in adulthood (Harrison *et al.*, 2005; Fatemi and Folsom, 2009; Powell, 2010; Rapoport *et al.*, 2012). Further studies should address schizophrenia as a neurodevelopmental disorder by administering METH during pregnancy and in periods leading up to adulthood to examine its impact on behavior and neural circuitry during development (Lee *et al.*, 2019). Some studies further point to how re-expression or overexpression of NMDA-R subunits can rescue behavioral deficits associated with symptoms of schizophrenia, suggesting that enhancing levels of certain NMDA-R subunits may ameliorate hypo-functionality of the receptor.

5. Conclusion

In conclusion, repeated METH administration induces changes in not only dopaminergic systems but also glutamatergic systems, and it alters NMDA receptor function and subunit expression. This suggests that our model reflects some of the pathophysiological changes of schizophrenia, and may be useful for identifying new therapeutic agents for the treatment of schizophrenia.

Contributors

MO, KI, and MK designed the study; MO and MK performed the experiments; MO analyzed the data and wrote the first draft of the manuscript; all authors contributed to the interpretation of the data and commented on the manuscript. All authors have approved the final manuscript.

Funding source

The present study was supported by Novartis Research Grants 2018.

Ethical statement

393 All experiments were ethically approved by the Animal Research Committee of
394 Hokkaido University (permission number 13–0137, 17–0086), and performed in
395 accordance with ARRIVE guidelines, the Guide for the Care and Use of Laboratory
396 Animals of Hokkaido University, and the guidelines established by the US National
397 Institutes of Health guide for the care and use of laboratory animals (NIH Publications
398 No. 8023, revised 1978).

399

400 **Declaration of Competing Interest**

401 The authors declare that there is no conflict of interest.

402

403 **Acknowledgements**

404 The authors thank Mariko Tonosaki, Akiko Kato, and Keisuke Makihara for
405 technical assistance and Yuki Omiya for comments and suggestions. We would like to
406 thank Editage (www.editage.com) for English language editing.

407

408 **Appendix A. Supplementary data**

409 Supplementary data to this article can be found online at
410 <https://doi.org/10.1016/j.pnpbp.2020.109984>.

References

- Abekawa, T., Ito, K., Nakagawa, S., Nakato, Y., Koyama, T., 2008. Olanzapine and risperidone block a high dose of methamphetamine-induced schizophrenia-like behavioral abnormalities and accompanied apoptosis in the medial prefrontal cortex. *Schizophr. Res.* 101, 84–94.
- Abekawa, T., Ito, K., Nakagawa, S., Nakato, Y., Koyama, T., 2011. Effects of aripiprazole and haloperidol on progression to schizophrenia-like behavioural abnormalities and apoptosis in rodents. *Schizophr. Res.* 125, 77–87.
- Abekawa, T., Ito, K., Nakato, Y., Koyama, T., 2012. Innovation of an animal model for the pathophysiology of schizophrenia. *Seishin Shinkeigaku Zasshi* 114, 81–98.
- Akbarian, S., Sucher, N.J., Bradley, D., Tafazzoli, A., Trinh, D., Hetrick, W.P., Potkin, S.G., Sandman, C.A., Bunney, W.E., Jones, E.G., 1996. Selective alterations in gene expression for NMDA receptor subunits in prefrontal cortex of schizophrenics. *J. Neurosci.* 16, 19–30.
- Carlsson, A., Hansson, L.O., Waters, N., Carlsson, M.L., 1997. Neurotransmitter aberrations in schizophrenia: new perspectives and therapeutic implications. *Life Sci.* 61, 75–94.
- Catts, V.S., Lai, Y.L., Weickert, C.S., Weickert, T.W., Catts, S.V., 2016. A quantitative review of the postmortem evidence for decreased cortical *N*-methyl-D-aspartate receptor expression levels in schizophrenia: how can we link molecular abnormalities to mismatch negativity deficits? *Biol. Psychol.* 116, 57–67.
- Cull-Candy, S.G., Leszkiewicz, D.N., 2004. Role of distinct NMDA receptor subtypes at central synapses. *Sci. STKE* 2004.
- Curran, H.V., D’Souza, D.C., Robbins, T.W., Fletcher, P., 2009. Editorial: modelling psychosis. *Psychopharmacology* 206, 513–514.
- Fatemi, S.H., Folsom, T.D., 2009. The neurodevelopmental hypothesis of schizophrenia, revisited. *Schizophr. Bull.* 35, 528–548.
- Featherstone, R.E., Shin, R., Kogan, J.H., Liang, Y., Matsumoto, M., Siegel, S.J., 2015. Mice with subtle reduction of NMDA NR1 receptor subunit expression have a selective decrease in mismatch negativity: implications for schizophrenia prodromal population. *Neurobiol. Dis.* 73, 289–295.

444 Gao, X.M., Sakai, K., Roberts, R.C., Conley, R.R., Dean, B., Tamminga, C.A., 2000.
445 Iontropic glutamate receptors and expression of *N*-methyl-D-aspartate receptor
446 subunits in subregions of human hippocampus: effects of schizophrenia. *Am. J.*
447 *Psychiatry* 157, 1141–1149.

448 Goff, D.C., Coyle, J.T., 2001. The emerging role of glutamate in the pathophysiology
449 and treatment of schizophrenia. *Am. J. Psychiatry* 158, 1367–1377

450 González, B., Jayanthi, S., Gomez, N., Torres, O.V., Sosa, M.H., Bernardi, A., Urbano,
451 F.J., García-Rill, E., Cadet, J.-L., Bisagno, V., 2018. Repeated methamphetamine
452 and modafinil induce differential cognitive effects and specific histone acetylation
453 and DNA methylation profiles in the mouse medial prefrontal cortex. *Prog.*
454 *Neuropsychopharmacol. Biol. Psychiatry* 82, 1–11.

455 Harrison, P.J., Weinberger, D.R., 2005. Schizophrenia genes, gene expression, and
456 neuropathology: on the matter of their convergence. *Mol. Psychiatry* 10, 40–68.

457 Horak, M., Petralia, R.S., Kaniakova, M., Sans, N., 2014. ER to synapse trafficking of
458 NMDA receptors. *Front. Cell. Neurosci.* 8.

459 Howes, O., McCutcheon, R., Stone, J., 2015. Glutamate and dopamine in schizophrenia:
460 an update for the 21st century. *J. Psychopharmacol.* 29, 97–115.

461 Ito, K., Abekawa, T., Koyama, T., 2006a. Relationship between development of cross-
462 sensitization to MK-801 and delayed increases in glutamate levels in the nucleus
463 accumbens induced by a high dose of methamphetamine. *Psychopharmacology*
464 (Berl). 187, 293–302.

465 Ito, K., Abekawa, T., Koyama, T., 2006b. Valproate blocks high-dose
466 methamphetamine-induced behavioral cross-sensitization to locomotion-inducing
467 effect of dizocilpine (MK-801), but not methamphetamine. *Psychopharmacology*
468 (Berl). 186, 525–533.

469 Jones, C., Watson, D., Fone, K., 2011. Animal models of schizophrenia. *Br. J.*
470 *Pharmacol.* 164, 1162–1194.

471 Kiyama, Y., Manabe, T., Sakimura, K., Kawakami, F., Mori, H., Mishina, M., 1998.
472 Increased thresholds for long-term potentiation and contextual learning in mice
473 lacking the NMDA-type glutamate receptor epsilon1 subunit. *J. Neurosci.* 18,
474 6704–6712.

475 Ladépêche, L., Dupuis, J.P., Groc, L., 2014. Surface trafficking of NMDA receptors:
476 gathering from a partner to another. *Semin. Cell Dev. Biol.* 27, 3–13.

477 Lau, C.G., Zukin, R.S., 2007. NMDA receptor trafficking in synaptic plasticity and
478 neuropsychiatric disorders. *Nat. Rev. Neurosci.* 8, 413–426.

479 Law, A.J., Deakin, J.F.W., 2001. Asymmetrical reductions of hippocampal NMDARI

glutamate receptor mRNA in the psychoses. *Neuroreport* 12, 2971–2974.

Lee, G., Zhou, Y., 2019. NMDAR hypofunction animal models of schizophrenia. *Front. Mol. Neurosci.* 12.

Lewis, D.A., Lieberman, J.A., 2000. Catching up on schizophrenia: natural history and neurobiology. *Neuron* 28, 325–334.

Lominac, K.D., Quadir, S.G., Barrett, H.M., McKenna, C.L., Schwartz, L.M., Ruiz, P.N., Wroten, M.G., Campbell, R.R., Miller, B.W., Holloway, J.J., Travis, K.O., Rajasekar, G., Maliniak, D., Thompson, A.B., Urman, L.E., Kippin, T.E., Phillips, T.J., Szumlinski, K.K., 2016. Prefrontal glutamate correlates of methamphetamine sensitization and preference. *Eur. J. Neurosci.* 43, 689–702.

Marcotte, E.R., Pearson, D.M., Srivastava, L.K., 2001. Animal models of schizophrenia: a critical review. *J. Psychiatry Neurosci.* 26, 395–410.

Moriya, T., Kouzu, Y., Shibata, S., Kadotani, H., Fukunaga, K., Miyamoto, E., Yoshioka, T., 2000. Close linkage between calcium/calmodulin kinase II α/β and NMDA-2A receptors in the lateral amygdala and significance for retrieval of auditory fear conditioning. *Eur. J. Neurosci.* 12, 3307–3314.

Nakato, Y., Abekawa, T., Ito, K., Inoue, T., Koyama, T., 2010. Lamotrigine blocks the initiation and expression of repeated high-dose methamphetamine-induced prepulse inhibition deficit in rats. *Neurosci. Lett.* 481, 183–187.

Nakato, Y., Abekawa, T., Ito, K., Inoue, T., Koyama, T., 2011. Lamotrigine blocks apoptosis induced by repeated administration of high-dose methamphetamine in the medial prefrontal cortex of rats. *Neurosci. Lett.* 490, 161–164.

Ohnuma, T., Arai, H., 2011. Significance of NMDA receptor-related glutamatergic amino acid levels in peripheral blood of patients with schizophrenia. *Prog. Neuro-Psychopharmacology Biol. Psychiatry* 35, 29–39.

Olney, J.W., Farber, N.B., 1995. Glutamate receptor dysfunction and schizophrenia. *Arch. Gen. Psychiatry* 52, 998–1007.

Olney, J.W., Newcomer, J.W., Farber, N.B., 1999. NMDA receptor hypofunction model of schizophrenia. *J. Psychiatr. Res.* 33, 523–533.

Paoletti, P., Bellone, C., Zhou, Q., 2013. NMDA receptor subunit diversity: impact on receptor properties, synaptic plasticity and disease. *Nat. Rev. Neurosci.* 14, 383–400.

Powell, S.B., 2010. Models of neurodevelopmental abnormalities in schizophrenia. *Curr. Top. Behav. Neurosci.* 4, 435–481.

Rapoport, J.L., Giedd, J.N., Gogtay, N., 2012. Neurodevelopmental model of schizophrenia: update 2012. *Mol. Psychiatry* 17, 1228–1238.

- Sakimura, K., Kutsuwada, T., Ito, I., Manabe, T., Takayama, C., Kushiya, E., Yagi, T., Shinichi, A., Inoue, Y., Sugiyama, H., Mishina, M., 1995. Reduced hippocampal LTP and spatial learning in mice lacking NMDA receptor $\epsilon 1$ subunit. *Nature* 373, 151-155.
- Simões, P.F., Silva, A.P., Pereira, F.C., Marques, E., Grade, S., Milhazes, N., Borges, F., Ribeiro, C.F., Macedo, T.R., 2007. Methamphetamine induces alterations on hippocampal NMDA and AMPA receptor subunit levels and impairs spatial working memory. *Neuroscience* 150, 433–441.
- Steeds, H., Carhart-Harris, R.L., Stone, J.M., 2015. Drug models of schizophrenia. *Ther. Adv. Psychopharmacol.* 5, 43-58.
- Traynelis, S.F., Wollmuth, L.P., McBain, C.J., Menniti, F.S., Vance, K.M., Ogden, K.K., Hansen, K.B., Yuan, H., Myers, S.J., Dingledine, R., 2010. Glutamate receptor ion channels: structure, regulation, and function. *Pharmacol. Rev.* 62, 405-496
- Van Os, J., Kenis, G., Rutten, B.P.F., 2010. The environment and schizophrenia. *Nature* 468, 203–212.
- Weickert, C.S., Fung, S.J., Catts, V.S., Schofield, P.R., Allen, K.M., Moore, L.T., Newell, K.A., Pellen, D., Huang, X.F., Catts, S.V., Weickert, T.W., 2013. Molecular evidence of *N*-methyl-D-aspartate receptor hypofunction in schizophrenia. *Mol. Psychiatry* 18, 1185–1192.