Title	alpha-aminoisobutyric acid mimics the effect of 1-aminocyclopropane-1-carboxylic acid to promote sexual reproduction in the marine red alga <i>Pyropia yezoensis</i> (Rhodophyta)
Author(s)	Endo, Harune; Mizuta, Hiroyuki; Uji, Toshiki
Citation	Journal of Applied Phycology, 33(2), 1081-1087 https://doi.org/10.1007/s10811-020-02326-7
Issue Date	2021-01-02
Doc URL	http://hdl.handle.net/2115/83824
Rights	This is a post-peer-review, pre-copyedit version of an article published in [insert journal title]. The final authenticated version is available online at: https://doi.org/10.1007/s10811-020-02326-7
Type	article (author version)
File Information	article.pdf

[Instructions for use](#)

1 α -aminoisobutyric acid mimics the effect of 1-aminocyclopropane-1-carboxylic acid to
2 promote sexual reproduction in the marine red alga *Pyropia yezoensis* (Rhodophyta)

3
4 Harune Endo

5 Laboratory of Aquaculture Genetics and Genomics, Division of Marine Life Science,
6 Faculty of Fisheries Sciences, Hokkaido University, Hakodate 041-8611, Japan

7
8 Hiroyuki Mizuta

9 Laboratory of Aquaculture Genetics and Genomics, Division of Marine Life Science,
10 Faculty of Fisheries Sciences, Hokkaido University, Hakodate 041-8611, Japan

11
12 Toshiki Uji¹

13 Laboratory of Aquaculture Genetics and Genomics, Division of Marine Life Science,
14 Faculty of Fisheries Sciences, Hokkaido University, Hakodate 041-8611, Japan

15
16 Running title: AIB promotes reproduction in *Pyropia*

¹ Tel/Fax: +81-138-40-8864

E-mail: t-uji@fish.hokudai.ac.jp

Abstract

1-aminocyclopropane-1-carboxylic acid (ACC), a precursor for ethylene, stimulates the switch from a vegetative to a sexual reproductive phase in the marine red alga *Pyropia* species. This study explored the effects of ethylene biosynthesis inhibitors on the sexual reproduction of gametophytes of the red alga *Pyropia yezoensis* to gain a functional understanding of the role of ACC as a plant hormone in red algae. Here we show that two inhibitors of ACC synthesis in higher plants, 2-aminoethoxyvinyl glycine (AVG) and aminooxyacetic acid (AOA), had no effect on the growth and gametogenesis of *P. yezoensis*. In contrast, exogenous application of α -aminoisobutyric acid (AIB), a structural analog of ACC that blocks the conversion of ACC to ethylene in higher plants, induced the formation of spermatangia and carpospores, in a similar manner as ACC, without endogenous ACC accumulation. The treatment of AIB failed to inhibit ethylene production in the gametophytes. The present results suggest that AIB mimics the effect of ACC to induce the sexual reproduction and support our previous study that ACC has a role in the regulation of the sexual reproduction independent from ethylene.

Keywords: 1-aminocyclopropane-1-carboxylic acid, ethylene, plant hormone, *Pyropia*, red algae, sexual reproduction

35 Abbreviations: ACC, 1-aminocyclopropane-1-carboxylic acid; ACS, ACC synthase;
36 ACO, ACC oxidase; AIB, α -aminoisobutyric acid; AOA, aminooxyacetic acid; AVG,
37 2-aminoethoxyvinyl glycine
38
39

Introduction

Plants produce hormones (also known as phytohormones or plant growth regulators) that regulate plant growth, development, and environmental responses (Vanstraelen and Benkova 2012; Xia et al. 2015; Verma et al. 2016). In plant hormone research, the use of a wide variety of small molecules such as agonists, antagonists, and inhibitors of the biosynthetic pathway and transports have significantly enhanced our understanding of the molecular basis of hormone actions in higher plants (Fonseca et al. 2014; Rigal et al. 2014). In macroalgae, treatment with auxin efflux inhibitors, such as naphthylphthalamic acid (NPA), elevates indole-3-acetic acid (IAA) accumulation and reduces environmental polarization in response to gravity and light vectors in embryos of *Fucus distichus*, a brown alga (Basu et al. 2002; Sun et al. 2004). Jasmonate (JA) is related to cystocarp development in the red macroalga *Grateloupia imbricate*, and the application of phenidone, a specific inhibitor of lipoxygenases (LOX) that catalyzes the first step in the biosynthesis of JA, decreases the number of the cystocarps concomitant with the repression of the JA release (Garcia-Jimenez et al. 2016). In addition, callus induction from leaf explants of the brown alga *Sargassum horneri* was achieved when grown in medium supplemented with uniconazole, a triazole-type inhibitor of cytochrome P450 enzymes for gibberellin biosynthesis, trans-zeatin biosynthesis, and

abscisic acid catabolism (Uji et al. 2016b). However, in contrast to higher plants, few reports are available on the use of small molecules involved in plant hormones in macroalgal research, which hinders the elucidation of the functional roles of plant hormones.

Ethylene, a simple gaseous plant hormone, is derived from methionine (Met), which is first converted to S-adenosyl-methionine (SAM) by SAM synthetase. SAM is then converted into 1-aminocyclopropane-1-carboxylic acid (ACC), an ethylene precursor (Yang and Hoffman 1984). This step is catalyzed by ACC synthase (ACS), an aminotransferase that requires pyridoxal 5'-phosphate (PLP) as a cofactor. In the second step, ACC is oxidized by an O₂-activating non-heme iron enzyme, ACC oxidase (ACO), giving rise to ethylene, carbon dioxide (CO₂), and hydrogen cyanide. Ethylene production can be manipulated by the inhibition of ACS and ACO activity (Fig. 1). For instance, 2-aminoethoxyvinyl glycine (AVG) and aminoxyacetic acid (AOA) inhibit the activity of ACS and consequently decrease ethylene production by preventing the production of ACC (Amrhein and Wenker 1979; Boller et al. 1979; Yang and Hoffman 1984). Additionally, α -aminoisobutyric acid (AIB) is a structural analog of ACC that blocks ACO activity by acting as a competitive inhibitor of the ACC substrate (Satoh and Esashi 1980; 1982; Serrano et al. 1990). The pharmacological manipulation of

ethylene biosynthesis using these inhibitors has increased our understanding of the roles of ethylene in higher plants (Tamimi and Timko 2003; Tian et al. 2014).

The marine red alga *Pyropia* (formerly *Porphyra*), a genus of Bangiophyceae, is a significant marine crop that is harvested to produce the food “nori”. The heteromorphic life history of *Pyropia* is comprised of a blade gametophyte and filamentous sporophyte (Fig. 2). This genus forms male (spermatia) and female (carpogonia) gametes during sexual reproduction. After fertilization, successive cell divisions occur to produce clones of the zygote called carpospores that will grow into sporophytes. Our previous research revealed that the exogenous application of ACC promoted gametogenesis and enhanced the antioxidant capacity in accompany with ethylene emission in the monoecious *P. yezoensis* (Uji et al. 2016a). Research on the dioecious species *P. pseudolinearis* revealed that ACC can modulate the expression of genes involved in the regulation of cell division and cell wall organization, which leads to the formation of spermatangia and parthenosporangia in male and female gametophytes, respectively (Yanagisawa et al. 2019). Our recent study suggests that ACC acts as a signaling molecule independent from ethylene in the regulation of sexual reproduction through alterations to the redox state in *P. yezoensis* (Uji et al. 2020).

In the present study, we investigated the effects of three ethylene biosynthetic

pathway inhibitors on the sexual reproduction of *P. yezoensis* gametophytes to gain a better understanding of ACC plant hormone action in *Pyropia* species. The present results suggest that AIB mimics the effect of ACC to induce the sexual reproduction in *P. yezoensis*.

Materials and methods

Algal cultures

Standard cultivation of gametophytes of *P. yezoensis* strain TU-1 was conducted at 15 °C in glass flasks (150 mL volume) with 100 mL of medium. The medium consisted of autoclaved seawater enriched with sterile vitamin-free Provasoli solution (PES; Provasoli, 1968). Light was supplied by cool-white fluorescent lamps at 60 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ irradiance with a photoperiod regime of 10 h light:14 h dark (short day condition) or 14 h light:10 h dark (long day condition).

Chemical treatments

The gametophytes of *P. yezoensis* were treated with the chemical compounds generally used as ethylene biosynthesis inhibitors in higher plants. Five individual immature gametophytes of ca. 20 mm blade length that were microscopically determined to bear

only vegetative cells were cultured in glass flasks (150 mL volume) with 100 mL PES medium with either three inhibitors prepared each concentration: 0, 5, 50 μ M aminoethoxyvinylglycine (AVG) (Cayman Chemical Company, MI, USA); 50, 500 μ M α -aminoisobutyric acid (AIB) (Tokyo Chemical Industry, Tokyo, Japan); 50, 500 μ M aminooxyacetic acid (AOA) (FUJIFILM Wako Pure Chemical Corporation, Osaka, Japan); or 50, 500 μ M ACC (Tokyo Chemical Industry). Two pieces of glass (20 mm $\times$ 25 mm) were placed on the bottom of the culture flask to capture the formation of carpogonia, the female gametes. The gametophytes were cultured with the chemical reagents for 7 d, then the number of gametophytes that had formed clusters of spermatangia were counted under a Leica DM 5000 B microscope (Leica Microsystems, Tokyo, Japan). In this study, a mature gametophyte was defined as a thallus bearing at least five clusters of spermatangia because carpogonium from *P. yezoensis* are almost indistinguishable from surrounding vegetative cells. In addition, after 14 d, the numbers of discharged carpospores attached to the pieces of glass were counted under a microscope as an index of formation of carpogonia. The gametophyte blade lengths were measured after 10 d and growth rate was calculated as the mean percentage of length increase per day using the following formula: Growth rate = $[100(BL_t - BL_0)/BL_0]/t$, where BL_0 = initial blade length, BL_t = blade length at days of culture, t =

culture time. The experiments under the long day or the short day culture were conducted at three or eight replicates with 5 thalli each condition, respectively. In these experiments, thalli treated without inhibitors were used as a control.

Measurement of ethylene production

To examine whether AIB has an inhibitory effect on ethylene production in gametophytes, we measured ethylene levels after 10 d treatments with AIB. Vegetative gametophytes (fresh weight: FW ca. 0.1 g) were cultured in 20 mL aluminum sealed vials with butyl-gum septums (GL Science, Tokyo, Japan) with 10 mL of medium containing 0, 500 μ M ACC, or 500 μ M ACC and AIB, at 15 °C under a 10 h light:14 h dark photoperiod regime using cool-white fluorescent lamps at 60 μ mol photons $\text{m}^{-2} \text{s}^{-1}$. After 10 d, the samples were analyzed by GC analysis of the volatile compounds. For the static headspace GC analysis, the sample vial was transferred into the HS-20 headspace auto-sampler (Shimadzu Corporation Kyoto, Japan) of the GC apparatus. The headspace gas in the vial was automatically pressurized at 60 °C for 2 min and then immediately injected through a loop into a GC-2014AFSC (Shimadzu) equipped with an HP-1 capillary column (50 m length, 0.32 mm i.d. and 1.05 μ m film thickness; Agilent Technologies, CA, USA) and a flame ionization detector. An initial oven

temperature of 40 °C for 5 min was followed by heating at 3 °C/min to 70°C, then 20 °C/min to 200 °C, and finally, the temperature was held at 200 °C for 4 min. Both the injection port and the flame ionization detector were set at 250 °C. The content of ethylene was expressed in nmol day⁻¹ g⁻¹ FW. These experiments were repeated in triplicate and thalli treated with ACC were used as a control.

Measurement of ACC content

ACC concentrations in the gametophytes treated with AIB were determined following the method of Wächter et al. (1999) with minor modifications. The algal materials were harvested, immediately frozen with liquid nitrogen, and stored at -80 °C until the extraction of ACC. A 1.5 g sample (FW) was ground in liquid nitrogen with a mortar and pestle, followed by homogenization in 7.5 mL 80% methanol with 2 mg/L dibutylhydroxytoluene. The extract was stored at room temperature for 45 min, then centrifuged for 15 min at 2000 × g at 20 °C. The extraction was repeated with 6 mL 80% methanol and the supernatants were evaporated to dryness with a Rotary Evaporator N-N/NVC-1100 (EYELA, Osaka, Japan) at 40 °C. The dry residue was dissolved in 2 mL distilled water and 4 mL dichloromethane, then centrifuged at 2000 ×

g at 20 °C for 5 min. The conversion of ACC to ethylene was performed following the procedure described by Lizada and Yang (1979). A 1.6 mL aliquot of the upper aqueous phase was mixed with 0.1 mL HgCl₂ (20 mM) in a 20 mL aluminum sealed vial with a butyl-gum septum. NaOCl-solution (5% NaOCl and 50% NaOH (2:1, v/v)) was injected through the septum with a syringe. The vials were shaken, incubated for 5 min and then the sample was assayed for ethylene by GC analysis according to the method described above. The ACC content was expressed as nmol g⁻¹ FW. These experiments were repeated in triplicate and thalli treated without ACC/AIB were used as a control.

Statistical analysis

Data are expressed as mean ± standard deviation (SD). The results of treatments with and without inhibitors of the ethylene biosynthetic pathway were analyzed using Mann-Whitney's U test. For all analyses, $p < 0.05$ or 0.01 were considered statistically significant.

Results

Effects of ethylene inhibitors on growth and gametogenesis

When the gametophytes were cultured with 5, 50 µM AVG or 5, 50 µM AOA under the

long day culture condition that promotes the sexual reproduction, there were no significant differences in spermatangia formation and growth rates between thalli treated with and without the inhibitors (Table 1). These results imply that ACS inhibitors generally used in higher plants are ineffective in the gametophytes. Contrary to expectations, gametophytic thalli treated with 50 μ M or 500 μ M AIB formed significantly more spermatangia clusters and gametophytes cultured with 500 μ M AIB exhibited slight growth repression compared to the thalli in the absence of chemical compounds (control) (Table 1).

Comparison of the effects of AIB and ACC on induction of sexual reproduction

The results in the effects of AIB prompted us to compare with the induction level of AIB and ACC on the sexual reproduction in the gametophytes. The gametophytes were cultured with 50, 500 μ M ACC or 50, 500 μ M AIB under the short day culture condition that represses the sexual reproduction. Treatment with 500 μ M AIB promoted the formation of spermatangia on the upper parts of the thalli, whereas 500 μ M ACC promoted spermatangia formation on the upper to middle sections (Fig. 3). All of the gametophyte thalli that were cultured with 50, 500 μ M ACC or 50, 500 μ M AIB formed spermatangia, whereas only 22.5% of thalli formed spermatangia when the

gametophytes were cultured without chemical compounds (control) (Fig. 4a). The growth rate of gametophytes cultured in medium containing 50 and 500 μ M ACC exhibited 14.6% and 13.7%, respectively, whereas that of gametophytes grown with 50 and 500 μ M AIB exhibited 20.2% and 12.2%, respectively (Fig. 4b). The number of discharged carpospores from the gametophytes treated with ACC or AIB was significantly higher than the control, but AIB has less effective than ACC (Fig. 4c). These results indicate that AIB can induce the sexual reproduction in a similar manner as ACC.

Evaluation of AIB as an inhibitor of ACC to ethylene conversion

The ACC contents of thalli treated with AIB were investigated to consider the possibility that AIB promoted the sexual reproduction through endogenous ACC accumulation by blocking the conversion of ACC to ethylene in the gametophytes. As shown in Fig.5, exogenous application of 50 and 500 μ M ACC increased the contents of ACC *in vivo*, whereas no accumulation of ACC was confirmed in thalli supplemented with 500 μ M AIB. These results indicated that AIB can induce the sexual reproduction without endogenous ACC accumulation and raised the possibility that AIB is ineffective inhibitor of ACO. To confirm this, the ethylene production was examined in thalli

220 treated with AIB. Since our previous study revealed small amount of ethylene release in
221 the gametophytes under normal conditions (Uji et al. 2016a), we used thalli increased
222 ACC/ethylene *in vivo* by supplement with 500 μ M ACC as a control. The results
223 showed that there were no significant differences in ethylene production between the
224 thalli treated without AIB and with 0.5 mM AIB (Fig.6). In addition, no inhibition of
225 ethylene production in the gametophytes treated with the even high doses of AIB (10
226 mM AIB) (Fig.6) which the concentration strongly inhibits the conversion of ACC to
227 ethylene in higher plants (Sato and Esashi 1982; Serrano et al. 1990).

228 229 Discussion

230 Our recent study of *P. yezoensis* suggests that ACC acts as a signaling molecule
231 independent of ethylene signaling in the regulation of sexual reproduction through
232 alterations to the redox state (Uji et al. 2020). Interestingly, previous studies also
233 indicate that ACC regulates plant development in the higher plant *Arabidopsis thaliana*,
234 independent of its role as an ethylene precursor (Vanderstraeten et al. 2019; Polko and
235 Kieber 2019). Based on these findings, Vanderstraeten et al. (2019) suggest that AIB
236 may have the capacity to interact with ACC binding proteins (e.g., putative ACC
237 receptors), because AIB is structurally similar to ACC. Supporting this possibility, the

present study revealed that AIB promoted the formation of spermatangia and carpospores in gametophytes without endogenous ACC accumulation in a similar manner as ACC. However, the comparative experiment between ACC and AIB treatments revealed that AIB promoted the formation of spermatangia on the upper parts of the thalli, whereas ACC promoted spermatangia formation on the upper to middle sections (Fig. 3). In addition, the number of discharged carpospores in the gametophytes treated with AIB was smaller than that of ACC (Fig. 4). These results suggest that AIB binds on the same target of ACC such as a putative ACC receptor to act ACC signaling, but the binding capacity of AIB may be weaker than that of ACC in *P. yezoensis*. Future work is needed to identify ACC receptor(s) and ACC signaling pathway(s) to uncover the mechanism regulating the sexual reproduction in *Pyropia*.

Recent genetic analyses of mutants that affect biosynthesis or plant hormone responses have revealed a number of signaling pathways in higher plants (Browse 2009; Zhao and Li 2012). To elucidate the role of ACC, the analysis of ACC response to ethylene-insensitive mutant contributed to the understanding of the ACC dependent pathway (Tsuchisaka et al. 2009; Tsang et al. 2011). However, a rapid and simple method of obtaining stable mutants has not yet been established for red macroalgae. Thus, knowledge about the effects of small molecules that act as agonists, antagonists,

and inhibitors of biosynthetic pathways for plant hormones is indispensable to advance macroalgal plant hormones research. In the previous study, exogenous application of ACC induced ethylene emission in *P. yezoensis*, indicating that the effect of ACC treatment is indistinguishable between the ACC- and ethylene-dependent pathways (Uji et al. 2016). In contrast, AIB mimics the effect of ACC in *P. yezoensis* gametophytes and thereby discriminates between ACC- and ethylene-responses, indicating that AIB could provide a useful approach to elucidate the ACC-dependent, and ethylene-independent pathways in red algae.

Genome sequence analyses have revealed that the homolog of the ACO gene is absent in the genomes of red algae, including *P. yezoensis*. Consistent with this, AIB application failed to inhibit ethylene production in the gametophytes (Fig. 6). In addition to the absent ACO gene, there is no homolog for the receptors and components of the ethylene signaling pathway in the red algal genome. Consistent with these findings, our recent study revealed that exogenous application of ACC promoted the formation of spermatia and carporspores in the gametophytes, whereas ethephon, an ethylene-releasing compound, did not stimulate sexual reproduction (Uji et al. 2020). In addition to the red algal genome, putative homologs for the perception and signaling of ethylene were apparently absent in chlorophyte green algae. This finding is supported

274 by a lack of ethylene-binding activity in two chlorophytes, *Chlamydomonas reinhardtii*
275 and *Acetabularia acetabulum* (Wang et al. 2006). In contrast, the charophyte green alga
276 *Spirogyra pratensis* induced cell elongation in response to ethylene and possesses
277 functionally conserved homologs of a complete set of ethylene-signaling genes (Ju et al.
278 2015). These findings suggest that the origin of ethylene response and signaling in
279 plants occurred during the evolution of the charophyte lineage, prior to the colonization
280 of land (Ju et al. 2015). However, exogenous application of ethylene promoted
281 tetrasporogenesis in the red marine alga *Pterocladia capillacea* (Garcia-Jimenez and
282 Robaina 2012) and regulated gene expression during carposporogenesis in the red
283 marine alga *Grateloupia imbricate* (Garcia-Jimenez et al. 2018). Hence, the
284 physiological roles of ethylene in red algae require further investigation.

285 In the present study, treatment with ACS inhibitors generally used in higher
286 plants, such as AVG and AOA did not have a significant effect on sexual reproduction in
287 *P. yezoensis*. On the other hand, exogenous application of AIB induced the formation of
288 spermatangia and carpospores without endogenous ACC accumulation, suggesting that
289 AIB have the capacity to interact with ACC binding proteins to induce the sexual
290 reproduction. This study supports that ACC acts as a plant hormone independent of
291 ethylene in *Pyropia*. Further studies on the identification of ACC receptor(s) and ACC

292 signaling pathway(s) could contribute to elucidate the role of ACC beyond the precursor
293 of ethylene in red algae.

294

295 Acknowledgments

296

297 We are grateful to Drs. Kazuo Miyashita, Masashi Hosokawa and Fumiaki Beppu
298 (Hokkaido University, Japan) for kindly providing GC-2014AFSC. This study was
299 supported by a grant-in-aid for Young Scientists (B) (16K18740 to TU) and Young
300 Scientists (19K15907 to TU) from the Japan Society for the Promotion of Science.
301 The authors declare no conflict of interest.

302 References

- 303 Adams DO, Yang SF (1979) Ethylene biosynthesis: identification of
304 1-aminocyclopropane-1-carboxylic acid as an intermediate in the conversion of
305 methionine to ethylene. *Proc Natl Acad Sci USA*. 76:170–174
- 306 Amrhein N, Wenker D (1979) Novel inhibitors of ethylene production in higher
307 plants. *Plant Cell Physiol* 20:1635–1642
- 308 Basu S, Sun HG, Brian L, Quatrano RL, Muday GK (2002) Early embryo development
309 in *Fucus distichus* is auxin sensitive. *Plant Physiol* 130:292–302
- 310 Boller T, Herner RC, Kende H (1979) Assay for and enzymatic formation of an
311 ethylene precursor, 1-aminocyclopropane-1-carboxylic acid. *Planta* 145:293–303
- 312 Bradford KJ, Hsiao TC, Yang SF (1982) Inhibition of ethylene synthesis in tomato
313 plants subjected to anaerobic root stress. *Plant Physiol* 70: 1503–1507
- 314 Browse J (2009) The power of mutants for investigating jasmonate biosynthesis and
315 signaling. *Phytochemistry* 70:1539–1546
- 316 Fonseca S, Rosado A, Vaughan-Hirsch J, Bishopp A, Chini A (2014) Molecular locks
317 and keys: the role of small molecules in phytohormone research. *Front Plant Sci*.
318 <https://doi.org/10.3389/fpls.2014.00709>
- 319 Garcia-Jimenez P, Brito-Romano O, Robaina RR (2016) Occurrence of jasmonates
320 during cystocarp development in the red alga *Grateloupia imbricata*. *J Phycol* 52:1085–
321 1093

322

323 Garcia-Jimenez P, Montero-Fernández M, Robaina RR (2018) Analysis of
324 ethylene-induced gene regulation during carposporogenesis in the red seaweed
325 *Grateloupia imbricata* (Rhodophyta). J Phycol 54:681–689

326 Garcia-Jimenez P, Robaina RR (2012) Effects of ethylene on tetrasporogenesis in
327 *Pterocliadiella capillacea* (Rhodophyta). J Phycol 48:710–715

328 Ju C, Van de Poel B, Cooper ED, Thierer JH, Gibbons TR, Delwiche CF, Chang C
329 (2015) Conservation of ethylene as a plant hormone over 450 million years of
330 evolution. *Nature Plants* 1:14004

331 Katoh Y, Hasegawa T, Suzuki T, Fujii T (1987) Changes in
332 1-aminocyclopropane-1-carboxylic acid content and ethylene production in hiproly
333 barley callus during differentiation. Agric biol chem 51:2185–2190

334 Lizada MC, Yang SF (1979) A simple and sensitive assay for
335 1-aminocyclopropane-1-carboxylic acid. Anal Biochem 100:140–145

336 Plettner I, Steinke M, Malin G (2005) Ethene (ethylene) production in the marine
337 macroalga *Ulva (Enteromorpha) intestinalis* L. (Chlorophyta, Ulvophyceae): effect of
338 light-stress and co-production with dimethyl sulphide. Plant Cell Environ 28:1136–
339 1145

340 Polko JK, Kieber JJ (2019) 1-aminocyclopropane 1-carboxylic acid and its emerging
341 role as an ethylene-independent growth regulator. Front Plant Sci.
342 <https://doi.org/10.3389/fpls.2019.01602>

343 Provasoli L (1968) Media and prospects for the cultivation of marine algae. In:

344 Watanabe A, Hattori A (eds) Culture and collections of algae, Proc U S-Japan Conf,
 345 Hakone, September 1966. Jpn Soc Plant Physiol Tokyo, 63–75

346 Rigal A, Ma Q, Robert S (2014) Unraveling plant hormone signaling through the use of
 347 small molecules. Front. Plant Sci. <https://doi.org/10.3389/fpls.2014.00373>

348 Satoh S, Esashi Y (1980) α -aminoisobutyric-acid - a probable competitive inhibitor of
 349 conversion of 1-aminocyclopropane-1-carboxylic acid to ethylene. Plant Cell
 350 Physiol 21:939–949

351 Satoh S, Esashi Y (1982) Effects of α -aminoisobutyric acid and D-amino and L-amino
 352 acids on ethylene production and content of 1-aminocyclopropane-1-carboxylic acid in
 353 cotyledonary segments of cocklebur seeds. Physiologia Plantarum 54:147–152

354 Serrano M, Romojaro F, Casas JL, Delrio JA, Acosta M (1990) Action and mechanism
 355 of α -aminoisobutyric acid as a retardant of cut carnation senescence. Sci
 356 Hortic 44:127–134

357 Sun HG, Basu S, Brady SR, Luciano RL, Muday GK (2004) Interactions between auxin
 358 transport and the actin cytoskeleton in developmental polarity of *Fucus distichus*
 359 embryos in response to light and gravity. Plant Physiol 135:266–278

360 Tamimi SM, Timko MP (2003) Effects of ethylene and inhibitors of ethylene synthesis
 361 and action on nodulation in common bean (*Phaseolus vulgaris* L.). Plant and
 362 Soil 257:125–131

363 Tian QY, Zhang XX, Ramesh S, Gilliam M, Tyerman SD, Zhang WH (2014) Ethylene
 364 negatively regulates aluminium-induced malate efflux from wheat roots and tobacco
 365 cells transformed with TaALMT1. *J Exp Bot* 65:2415–2426

366 Tsang DL, Edmond C, Harrington JL, Nuehse TS (2011) Cell wall integrity controls
 367 root elongation via a general 1-aminocyclopropane-1-carboxylic acid-dependent,
 368 ethylene-independent pathway. *Plant Physiol* 156:596–604

369 Tsuchisaka A, Yu G, Jin H, Alonso JM, Ecker JR, Zhang X, Gao S, Theologis A (2009)
 370 A combinatorial interplay among the 1-aminocyclopropane-1-carboxylate isoforms
 371 regulates ethylene biosynthesis in *Arabidopsis thaliana*. *Genetics* 183:979–1003

372 Uji T, Endo H, Mizuta H (2020) Sexual reproduction via a
 373 1-aminocyclopropane-1-carboxylic acid-dependent pathway through redox modulation
 374 in the marine red alga *Pyropia yezoensis* (Rhodophyta). *Front. Plant Sci.*
 375 <https://doi.org/10.3389/fpls.2020.00060>

376 Uji T, Matsuda R, Takechi K, Takano H, Mizuta H, Takio S (2016a) Ethylene
 377 regulation of sexual reproduction in the marine red alga *Pyropia yezoensis*
 378 (Rhodophyta). *J Appl Phycol* 28:3501–3509

379 Uji T, Nanaumi D, Kawagoe C, Saga N, Miyashita K (2016b) Factors influencing the
 380 induction of adventitious bud and callus in the brown alga *Sargassum horneri* (Turner)
 381 C. Agardh. *J Appl Phycol* 28:2435–2443

382 Vanderstraeten L, Depaepe T, Bertrand S, Van Der Straeten D (2019) The ethylene
 383 precursor ACC affects early vegetative development independently of ethylene
 384 signaling. *Front Plant Sci.* <https://doi.org/10.3389/fpls.2019.01591>

385 Vanstraelen M, Benkova E (2012) Hormonal interactions in the regulation of plant
 386 development. *Ann Rev Cell Dev Biology* 28:463–487

387 Verma V, Ravindran P, Kumar PP (2016) Plant hormone-mediated regulation of stress
 388 responses. *BMC Plant Biology.* <https://doi.org/10.1186/s12870-016-0771-y>

389 Wächter R, Fischer K, Gäbler R, Kühnemann F, Urban W, Bögemann GM, Voesenek L
 390 ACJ, Blom CWPM, Ullrich CI (1999) Ethylene production and ACC-accumulation in
 391 *Agrobacterium tumefaciens*-induced plant tumours and their impact on tumour and
 392 host stem structure and function. *Plant Cell and Environ* 22:1263–1273

393 Wang W, Esch JJ, Shiu SH, Agula H, Binder BM, Chang C, Patterson SE, Bleecker AB
 394 (2006) Identification of important regions for ethylene binding and signaling in the
 395 transmembrane domain of the ETR1 ethylene receptor of *Arabidopsis*. *Plant Cell*
 396 18:3429–3442

397 Xia XJ, Zhou YH, Shi K, Zhou J, Foyer CH, Yu JQ (2015) Interplay between reactive
 398 oxygen species and hormones in the control of plant development and stress
 399 tolerance. *J Exp Bot* 66:2839–2856

400 Yanagisawa R, Sekine N, Mizuta H, Uji T (2019) Transcriptomic analysis under
 401 ethylene precursor treatment uncovers the regulation of gene expression linked to

402 sexual reproduction in the dioecious red alga *Pyropia pseudolinearis*. J Appl Phycol
 403 31:3317–3329

404 Yang SF, Hoffman NE (1984) Ethylene biosynthesis and its regulation in higher plants.
 405 Annu Rev Plant Physiol Plant Mol Biol 35:155–189

406 Yu YB, Yang SF (1979) Auxin-induced ethylene production and its inhibition by
 407 aminoethoxyvinylglycine and cobalt ion. Plant Physiol 64:1074–1077

408 Zhao BL, Li J (2012) Regulation of brassinosteroid biosynthesis and inactivation. J
 409 Integr Plant Biol 54:746–759

410
 411
 412
 413
 414

415 Figure legends

416 Figure 1. The ethylene biosynthetic pathway and the inhibitors used in this study.

417 Met, Methionine; SAM, S-adenosyl-methionine; ACC,
 418 1-aminocyclopropane-1-carboxylic acid; AVG, 2-aminoethoxyvinyl glycine; AOA,
 419 aminooxyacetic acid; AIB, α -aminoisobutyric acid

420

421 Figure 2. Life cycle of *Pyropia* species

422 The macroscopic leafy gametophytes of *Pyropia*, which are harvested to produce the
 423 food for nori or laver, bear non-motile male (spermatia) and female (carpogonia)

gametes on the thallus during sexual reproduction. Fertilization occurs when the female gametes are still retained on the gametophytes and successive cell divisions produce clones of the zygote (carpospores), which develop into microscopic filamentous sporophytes referred as the conchocelis. The mature sporophytes produce conchospores that are released from conchosporangia and settle on the substratum, where they germinate to form new gametophytes. Some *Pyropia* such as *P. yezoensis* can also reproduce asexually in the gametophytes by asexual spores called archeospores.

Figure 3. Effect of α -aminoisobutyric acid (AIB) on sexual reproduction in *Pyropia yezoensis* gametophytes under the short day culture condition. Gametophytes were cultured in media containing 0 or 500 μ M ACC, or 500 μ M AIB. The thalli treated with ACC and AIB formed spermatangia, which were clear or discolored. Scale bar = 10 mm.

Figure 4 α -aminoisobutyric acid (AIB) mimics the effect of ACC on the sexual reproduction in *Pyropia yezoensis* gametophytes under the short day culture condition. (a) Formation of spermatangia clusters on gametophytes after 7 d treatment with ACC or AIB. (b) Growth rate of gametophytes after 10 d treatment with ACC or AIB. (c) The

number of carpospores released from gametophytes after 14 d treatment with ACC or AIB. Data are expressed as mean $\pm$ SD. of eight independent experiments with five thalli for each condition. Asterisks and double asterisks indicate significant differences at $p < 0.05$ and $p < 0.01$, respectively, between control and treatments.

Figure 5. Effect of α -aminoisobutyric acid (AIB) on ACC accumulation in *Pyropia yezoensis* gametophytes. Gametophytes were cultured in media containing 0 (control) or 50, 500 μ M ACC, or 500 μ M AIB. The ACC production is expressed as mean $\pm$ SD (nmol g⁻¹ fresh wt.).

Figure 6. Effect of α -aminoisobutyric acid (AIB) on ethylene production in *Pyropia yezoensis* gametophytes treated with ACC. The ethylene production is expressed as mean $\pm$ SD (nmol day⁻¹ g⁻¹ fresh wt.).

Table 1 Effect of ethylene biosynthetic pathway inhibitors on spermatogenesis and growth rate in *Pyropia yezoensis* gametophytes under the long day culture condition

Treatments (μM)	% of thallus with cluster of spermatangia	Elongation rate (mean $\pm$ SD% d^{-1})
Control (No treatment)	33.3 $\pm$ 21.6	17.3 $\pm$ 1.2
AVG 5	40.0 $\pm$ 37.4	12.5 $\pm$ 5.2
AVG 50	33.3 $\pm$ 29.4	11.9 $\pm$ 1.5
AOA 5	33.3 $\pm$ 16.3	15.2 $\pm$ 3.0
AOA 50	66.7 $\pm$ 8.2	13.5 $\pm$ 0.7
AIB 50	100.0 $\pm$ 0.0 *	12.8 $\pm$ 2.1
AIB 500	100.0 $\pm$ 0.0 *	8.4 $\pm$ 5.7

Data are expressed as means $\pm$ SD of three independent experiments with 5 thalli each condition

Asterisks indicate significant differences at $p < 0.05$ between control and treatments.

Elongation rate = $[100(\text{BL}_t - \text{BL}_0)/\text{BL}_0]/t$, BL_0 = initial blade length, BL_t = blade length at 10 days culture, t = culture time.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6