Title	Modified thoracoabdominal nerves block through perichondrial approach (M-TAPA) provides a sufficient postoperative analgesia for laparoscopic sleeve gastrectomy
Author(s)	Aikawa, Katsuhiro; Tanaka, Nobuhiro; Morimoto, Yuji
Citation	Journal of Clinical Anesthesia, 59, 44-45 https://doi.org/10.1016/j.jclinane.2019.06.020
Issue Date	2020-02
Doc URL	http://hdl.handle.net/2115/84181
Rights	© 2019. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	letter to the editor.pdf

[Instructions for use](#)

Modified thoracoabdominal nerves block through perichondrial approach (M-TAPA) provides a sufficient postoperative analgesia for laparoscopic sleeve gastrectomy.

Katsuhiro Aikawa^a, MD, PhD katsuhiro.aikawa@gmail.com,

Nobuhiro Tanaka^a, MD dorami1105@hotmail.com,

Yuji Morimoto^a, MD, PhD morim2@med.hokudai.ac.jp,

^aDepartment of Anesthesiology and Critical Care Medicine, Faculty of Medicine and Graduate School of Medicine Hokkaido University, Kita-15, Nishi-7, Kita-ku, Sapporo 060-8638, Japan

Phone: +81-11-706-7861

Corresponding author: Katsuhiro Aikawa

Kita-15, Nishi-7, Kita-ku, Sapporo 060-8638, Japan

Phone: +81-11-706-7861

Running Title: M-TAPA for laparoscopic sleeve gastrectomy.

Disclosures: The authors received no funding for this study.

19 **Key words**

20 Obesity, bariatric surgery, modified thoracoabdominal nerves block, postoperative
21 analgesia.

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37 Dear editor,

38 Achieving sufficient postoperative analgesia in patients who undergo laparoscopic
39 sleeve gastrectomy (LSG) is challenging. In fact, epidural anesthesia is technically
40 difficult given the excessive subcutaneous fat, which increases the risk of serious
41 complications. Moreover, patients with this condition often have comorbidities that
42 require anticoagulation therapy. Although ultrasound-guided transversus abdominis plane
43 (TAP) block may be beneficial, it is still a matter of debate [1].

44 Recently, modified thoracoabdominal nerves block through perichondrial approach
45 (M-TAPA) has been reported as a novel and promising technique that provides effective
46 analgesia in the anterior and lateral thoracoabdominal wall [2]. Herein, we present a
47 successful case of LSG managed with M-TAPA.

48
49 A 46-year old female patient (156 cm, 99kg) with diabetes, sleep apnea syndrome,
50 and hypertension was scheduled for LSG. Epidural anesthesia was avoided considering
51 that she had undergone thoracic spine surgery. A bilateral M-TAPA was selected for
52 opioid-sparing postoperative analgesia. Following an uneventful induction and
53 intubation, a linear transducer was placed on the costochondral angle in the sagittal
54 plane [3] (Fig. 1A). A total of 60 mL of 0.25% ropivacaine (30 mL for each side) was

bilaterally injected into the layer between the transversus abdominis muscle and the lower aspect of the costal cartilage (Fig. 1B). Anesthesia was maintained with 0.8 MAC desflurane and 0.15-0.35 $\mu\text{g/kg/min}$ remifentanyl infusion. Hemodynamic stability was maintained throughout the anesthesia. The operation time was 104 min, and the total dose of intraoperative fentanyl was 400 μg . She had no pain at discharge from the operation theater.

Although the patient was administered a continuous infusion of fentanyl to control visceral pain in the ward, the postoperative pain was adequately controlled. By performing repetitive pinprick tests, we revealed an excellent analgesic effect of the M-TAPA. At 7 and 24 h after the blockade, a complete sensory block of the T3-12 dermatomes from the posterior axillary line to the midline was demonstrated. At 36 h after the blockade, although the patient started to feel slight pain associated with movement, an almost complete sensory block of the T4-12 was still remained. Subsequently, the sensory block was observed to become incomplete (4 out of 10 as reported by the patient) and the affected area was cramped (T6-12) at 48 h. Finally, the effect of the sensory block disappeared at 56 h after the blockade.

This case has demonstrated an effective, broad, and long-lasting analgesic effect of the M-TAPA. Furthermore, by performing the repetitive pinprick tests, we demonstrated

the time course of the M-TAPA, which has not yet been investigated previously. The anesthetized area of the present patient was wider than previous reports [2, 4]. The larger volume of the local anesthetics used in the current case may be a reason behind such an observation. Although further research to reveal the spread of the local anesthetics and its mechanism is required, the M-TAPA can be a suitable option for LSG. Written informed consent was obtained from the patient.

91 **Declaration of interest**

92 None.

93

94 **Funding**

95 This research did not receive any specific grant from funding agencies in the public,

96 commercial, or not-for-profit sectors.

97

98 **Conflicts of interest**

99 None.

100

101 **Acknowledgements**

102

103

104

105

106

107

108

109 **References**

- 110 [1] Ruiz-Tovar J, Albrecht E, Macfarlane A, Coluzzi F. The TAP block in obese
 111 patients: pros & cons. *Minerva Anesthesiol.* 2019.
 112 <https://doi.org/10.23736/S0375-9393.19.13545-6>
- 113 [2] Tulgar S, Selvi O, Thomas DT, Deveci U, Özer Z. Modified
 114 thoracoabdominal nerves block through perichondrial approach (M-TAPA)
 115 provides effective analgesia in abdominal surgery and is a choice for opioid
 116 sparing anesthesia. *Journal of clinical anesthesia.* 2019;55:109.
 117 <https://doi.org/10.1016/j.jclinane.2019.01.003>
- 118 [3] Tulgar S, Senturk O, Selvi O, Balaban O, Ahiskalioğlu A, Thomas DT, et al.
 119 Perichondral approach for blockage of thoracoabdominal nerves: Anatomical
 120 basis and clinical experience in three cases. *Journal of clinical anesthesia.*
 121 2019;54:8-10. <https://doi.org/10.1016/j.jclinane.2018.10.015>
- 122 [4] Altıparmak B, Korkmaz Toker M, Uysal A, Turan M, Gümüş Demirbilek S.
 123 The successful usage of modified thoracoabdominal nerves block through
 124 perichondrial approach (M-TAPA) for analgesia of laparoscopic ventral hernia
 125 repair. *Journal of clinical anesthesia.* 2019;57:1-2.
 126 <https://doi.org/10.1016/j.jclinane.2019.02.016>

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145 **Figure titles and legends**

146 Ultrasound image and postoperative abdominal wall. A, B. Ultrasound image of the
147 perichondral area before (A) and after (B) local anesthetic solution was injected. C.
148 Image of the postoperative abdominal wall. (CC: costal cartilage, EOM: external
149 oblique muscle, IOM: internal oblique muscle, TAM: transversus abdominis muscle,
150 LA: local anesthetic)