



Title	Modification of semisolid medium for stab culture to isolate <i>Salmonella enterica</i> from rectal swabs
Author(s)	Fujihara, Masatoshi; Sakazume, Noriko; Tagaino, Yuuka
Citation	Japanese Journal of Veterinary Research, 70(1), 29-32
Issue Date	2022-02
DOI	10.14943/jjvr.70.1.29
Doc URL	http://hdl.handle.net/2115/84414
Type	bulletin (article)
File Information	JJVR70-1_29-32_MasatoshiFujihara.pdf



[Instructions for use](#)

Modification of semisolid medium for stab culture to isolate *Salmonella enterica* from rectal swabs

Masatoshi Fujihara^{1,*}, Noriko Sakazume¹⁾ and Yuuka Tagaino¹⁾

¹⁾ Hokkaido Kushiro Livestock Hygiene Service Center, 127-1 Otanoshike, Kushiro, Hokkaido 084-0917, Japan.

Received for publication, September 29, 2021; accepted, December 6, 2021

Abstract

Semisolid agar has shown good performance for selective isolation of *Salmonella enterica*. To control the occurrence salmonellosis within a herd, identification of asymptomatic carriers is necessary; however, rectal swab samples are not suitable for inoculation onto semisolid agar. In this study, we lowered the agar concentration of modified semisolid Rappaport agar for stab culture. *S. enterica* exhibited growth across the entirety of the medium. Other bacteria in bovine fecal samples were not observed to migrate, and the sensitivity for *S. enterica* isolation was greater than that of commercially available Hajna tetrathionate broth. As inoculation onto isolation agar plates is not necessary if no migration is observed, this stab culture method would drastically simplify the process of *S. enterica* isolation.

Key Words: modified semisolid Rappaport agar, *Salmonella enterica*, stab culture

Salmonella enterica, a rod-shaped gram-negative bacterium belonging to the family *Enterobacteriaceae*, is the causative agent of salmonellosis. It affects most animal species as well as humans and is a major public health concern. Its clinical presentation can range from a healthy chronic carrier state to patients with acute or chronic enteritis to patients with septicemia. The most common type of presentation is the carrier state, in which infected animals carry the pathogen for a variable period without showing any clinical symptoms⁷⁾. Carriers of salmonellosis, contaminated feedstuffs, and contaminated environments are major challenges.

To limit the spread within a herd, infected animals should be identified and either culled or treated. Treated animals must be rechecked several times to confirm that they are not carriers. Asymptomatic animals identified as carriers through repeated positive fecal cultures over a course of several months should be removed from the herd.

With respect to *S. enterica* isolation, several reports have shown that semi-solid agar has the highest sensitivity and specificity^{1,2,3,6,9,10)}. We previously reported an easy-to-prepare modified semi-solid Rappaport agar (MSRA), the composition of which is similar to that of

* Corresponding author:

Masatoshi Fujihara, Hokkaido Kushiro Livestock Hygiene Service Center, 127-1 Otanoshike, Kushiro, Hokkaido 084-0917, Japan

E-mail: fujihara.masatoshi@pref.hokkaido.lg.jp

TEL: +81-154-57-8775

FAX: +81-154-57-6125

doi: 10.14943/jjvr.70.1.29

Table 1. Composition of Rappaport broth (Eiken, Tokyo, Japan), modified semisolid Rappaport agar (MSRA) and MSRA for stab culture (MSRAsc) (per liter).

Rappaport broth	MSRA		MSRAsc		
	50% Rappaport broth		50% Rappaport broth		
Peptone	5.00 g	Peptone	2.50 g	Peptone	2.50 g
NaCl	8.00 g	NaCl	4.00 g	NaCl	4.00 g
KH ₂ PO ₄	1.60 g	KH ₂ PO ₄	0.80 g	KH ₂ PO ₄	0.80 g
MgCl ₂	20.30 g	MgCl ₂	10.15 g	MgCl ₂	10.15 g
Malachite green	0.12 g	Malachite green	0.06 g	Malachite green	0.06 g
		Casamino acid	5.00 g	Casamino acid	5.00 g
		Agar	3.00 g	Agar	1.20 g

Table 2. Migration of *Enterobacter* and *Salmonella* strains on modified semisolid Rappaport agar (MSRA) and MSRA for stab culture (MSRAsc) after 24 hr of incubation.

Organism	Strain	Migration zone diameter on MSRA (mm)		Migration in MSRAsc	
		at 40°C	at 42°C	at 40°C	at 42°C
<i>E. cloacae</i>	K13ec1	No migration	No migration	-	-
<i>E. cloacae</i>	K13ec2	61	No migration	+	-
<i>S. Dublin</i>	K13-R1-3	No growth	No growth	-	-
<i>S. Livingstone</i>	K13-H29-21	>90	68	+	+
<i>S. Nima</i>	K13-R3-8	>90	>90	+	+
<i>S. Typhimurium</i>	K13-H30-27	41	30	+	+
<i>S. O4:i:-</i>	K13-R1-9	55	35	+	+
<i>S. Umblio</i>	K13-R1-29	>90	>90	+	+

commercially available modified semisolid Rappaport Vassiliadis agar; MSRA exhibits advantages such as isolation sensitivity for fecal samples, earlier colony detection for serotyping and support for the growth of stressed *S. enterica*²⁾. However, as *S. enterica* must be in contact with the semi-solid agar in plates to grow, ensuring an adequate amount of inoculum is essential for reliable isolation, and thus, rectal swabs are considered unfavorable for semi-solid agar plates. In this study, we modified MSRA for stab culture (MSRAsc) and lowered the agar concentration of MSRA, to allow *S. enterica* to swim in the medium.

First, we determined bacterial growth and motility on MSRA and in MSRAsc. The composition of MSRA and MSRAsc is shown in Table 1. Six *S. enterica* strains were used in this study. All bacterial strains were isolated from cattle in Hokkaido prefecture from 2019 to 2021

Table 3. Isolation of *S. enterica* from artificially contaminated swabs.

Organism	Strain	Inoculated CFU ^a	Number of isolations positive for <i>S. enterica</i> out of three samples			
			HTTB ^b →ES2 ^c	MSRAsc ^d →ES2	HTTB →MLCB	MSRAsc →MLCB
<i>S. Livingstone</i>	K13-H29-21	4	0	2	1	2
		44	3	3	3	3
		440	3	3	3	3
<i>S. Nima</i>	K13-R3-8	11	2	3	2	3
		108	2	3	3	3
		1,080	2	3	3	3
<i>S. Typhimurium</i>	K13-H30-27	4	0	2	0	2
		40	2	3	3	3
		400	2	3	3	3
<i>S. O4:i:-</i>	K13-R1-9	2	0	2	2	2
		20	2	3	3	3
		200	3	3	3	3
<i>S. Umblio</i>	K13-R1-29	8	3	3	3	3
		76	3	3	3	3
		760	3	3	3	3

^aCFU, colony forming unit

^bHTTB, Hajna tetrathionate broth

^cES2, ES salmonellae agar II

^dMSRAsc, modified semisolid Rappaport agar for stab culture

and were identified using biochemical tests and *S. enterica* serotyping. Moreover, since *Enterobacter cloacae* is known to migrate on MSRA at 37°C in our previous study²⁾, two strains were isolated from healthy cattle feces by MSRA at 37°C incubation and investigated as the control. Other bacteria such as *Escherichia* and *Proteus* in cattle feces did not migrate on MSRA at 37°C. Approximately 2,000 CFU/10 µl of each bacterial suspension was directly inoculated on MSRA or instilled on cotton swabs, stabbed into MSRAsc and then incubated at 40°C or 42°C for 24 hr.

The results are presented in Table 2. In agreement with our previous study²⁾, *S. Dublin* did not grow on MSRA and MSRAsc. In MSRAsc, motile bacteria grew in the whole medium, especially around the aerobic region. This can be explained by oxygen taxis⁴⁾ and aerobic respiration⁵⁾. *E. cloacae* K13ec1 and Hi-ec²⁾ strains did not migrate at 40°C, while K13ec2 migrated at 40°C (but not at 42°C). In contrast, all the *S. enterica* strains migrated at 42°C. However, as *S.*




Fig. Migration of *S. enterica* from artificially contaminated swabs to MSRAsc.

S. enterica migrated from the inside of plastic straw and proliferated, exhibiting growth especially around the aerobic region, as observed in the two tubes on the left. The two tubes on the right were negative for motile bacteria. The diameters of test tubes and plastic straws are 16 and 8 mm, respectively.

Paratyphi B Ne-PB had migrated at 40°C, but not at 42°C, in our previous study (data not shown), the incubation temperature should be verified by culturing target *S. enterica* strains. Comparing MSRA with MSRAsc, if the tested bacterial strain spread over 30 mm in diameter on MSRA, migration was visible in MSRAsc.

Next, the efficacy of each selective enrichment medium for the detection of *S. enterica* strains in fecal samples was investigated. Each *S. enterica* strain was instilled on cotton swabs. After stabbing into bovine fecal samples, swabs were inoculated into 5 ml of Hajna tetrathionate broth (HTTB) (Eiken, Tokyo, Japan) or MSRAsc. In MSRAsc inoculation, swabs were passed through plastic straws to distinguish between bacterial migration and solely proliferation. After 24 hr of incubation at 37°C (HTTB) or 42°C (MSRAsc), each culture was plated onto ES salmonellae agar II (ES2) (Eiken, Tokyo, Japan) and MLCB agar (Nissui, Tokyo, Japan) for *S. enterica* isolation. As shown in Table 3, MSRAsc was superior to HTTB in sensitivity to isolate *S. enterica*. Among the isolation agars, MLCB showed a higher sensitivity of isolation than that of ES2 after

selective enrichment via culturing on HTTB. Agar media containing chromogenic substrates for salmonella-specific enzymes are known to be less sensitive but more specific, when compared to conventional media⁸⁾. On the other hand, after selective enrichment via culturing on MSRAsc, there was no difference in sensitivity between the two isolation agars because the migrated zone is pure *S. enterica* culture condition. Furthermore, it was easier to determine the culture as “motile *S. enterica* negative”, as the bacteria in the fecal samples had not migrated (Fig.).

In this study, we prepared MSRAsc for the selective enrichment of *S. enterica* from swab samples. The composition of MSRAsc is the same as that of MSRA, except for a lowered agar concentration. *S. enterica* exhibited growth across the entirety of the medium, with localization especially around the aerobic region. Other bacteria from bovine fecal samples did not migrate, and given that no migration was observed, inoculation onto isolation agar plates was not necessary. Therefore, the use of MSRAsc would drastically simplify the process of *S. enterica* isolation in clinical laboratories.

References

- 1) Baggesen DL, Nielsen LR, Sørensen G, Bødker R, Ersbøll AK. Growth inhibitory factors in bovine faeces impairs detection of *Salmonella* Dublin by conventional culture procedure. *J Appl Microbiol.* 103, 650–656, 2007.
- 2) Fujihara M, Tabuchi H, Uegaki K. Growth kinetics of *Salmonella enterica* in Hajna tetrathionate broth, Rappaport broth and modified semisolid Rappaport agar. *J Vet Med Sci* 78, 435–438, 2016.
- 3) Goossens H, Wauters G, de Boeck M, Janssens M, Butzler JP. Semisolid selective-motility enrichment medium for isolation of salmonellae from fecal specimens. *J Clin Microbiol.* 19, 940–941, 1984.

- 4) Grishanin RN, Bivikov SI. Mechanism of oxygen taxis in bacteria. *Biosci Rep.* 17, 77-83, 1997.
- 5) Guerrero T, Calderón D, Zepata S, Tueba G. *Salmonella* grows massively and aerobically in chicken faecal matter. *Microb Biotechnol.* 13, 1678-1684, 2020.
- 6) Hine EAS, Steffen EK, Wagner JE. New semisolid agar for the detection of motile salmonellae. *J Clin Microbiol* 26, 875-878, 1988.
- 7) Morisse JP, Cotte JP. Evaluation of some risks factors in bovine salmonellosis. *Vet Res.* 25, 185-191, 1994.
- 8) Nye KJ, Fallon D, Frodsham D, Gee B, Graham C, Howe S, Warren RE. An evaluation of the performance of XLD, DCA, MLCB, and ABC agars as direct plating media for the isolation of *Salmonella enterica* from faeces. *J Clin Pathol* 33, 802-804, 1995.
- 9) Rodríguez FI, Procura F, Bueno DJ. Comparison of 7 culture methods for *Salmonella* serovar Enteritidis and *Salmonella* serovar Typhimurium isolation in poultry feces. *Poult Sci* 97, 3826-3836, 2018.
- 10) Soria MC, Soria MA, Bueno DJ. Comparison of 2 culture methods and PCR assays for *Salmonella* detection in poultry feces. *Poult Sci.* 91, 616-626, 2012.