



Title	Preface of Lithuania-Japan seminar 2021
Citation	Eurasian Journal of Forest Research, 22, 1-4
Issue Date	2022
Doc URL	<a href="http://hdl.handle.net/2115/84974">http://hdl.handle.net/2115/84974</a>
Type	bulletin (editorial)
File Information	0) EJFR Lt-Jp preface_contents.pdf



[Instructions for use](#)

## Preface of Lithuania-Japan seminar 2021

### Introduction

The research collaboration started between Aleksandras Stulginskis University (ASU) (now the Agriculture Academy of Vytautas Magnus University: VMU) and Hokkaido University (HU) in 2012 with a sharing of knowledge and research collaboration about the cell and molecule biology of a wild berry (Hoshino, 2022, Silva, 2022) (Fig.1 & Fig. 2). Next, we discussed the impact of environmental changes on forests via IUFRO activity (International Union of Forest Research Organization) which led to an exchange of scientists between two universities.

To further increase our collaboration, we decided to organize an international seminar sponsored by the Lithuanian State Science and Studies Foundation (LSSSF) and the Japan Society for the Promotion of Science (JSPS). We planned to hold the seminar in the 2020 fiscal year; however, we were forced to postpone the seminar until 2021 due to the Covid19 and its varieties. Finally, we held the seminar remotely in August, 2021. Because discussion time was very limited, we decided to continue the discussions by synthesizing our knowledge in research papers based on the abstract (Watanabe et al. 2021). We have chosen to call these papers “Extended Abstracts”. The academic research bulletin of the Field Science Center for Northern Biosphere (FSC) of Hokkaido University offered to publish them as a special issue “the Lithuania-Japan Seminar 2021”. For this special issue we collected further references and synthesized topics; some papers include authors from each university.

### Social background

Under rapidly changing environments, we need to create sustainable agroforestry systems using green methods. We can achieve this goal by using environmental microbiology in horticulture, crop production, forest science, and environmental science. Since 1995, biodiversity conservation has been developing, and establishing the Access to Genetic Resources and Benefit Sharing (ABS) rules brings us stricter regulations to protect genes and species in the member countries (COP 2010, Aichi Prefecture). We should collaborate and exchange more students if we would like to use natural genetic resources more smoothly based on the ABS rules.

Lithuanians experienced a severe nuclear accident with the meltdown at Chernobyl, Ukraine, when it was a province of the former Soviet Union in 1986. Unfortunately, we also had a similar situation with the Fukushima Nuclear Power Plant accident. With utilizing the plant nutrition approach for the past decade, however, we avoided partly radioactive cesium (Cs) toxicity by applying massive amounts of potassium (K) to the soil. In the keynote lecture, Professor Takuro Shinano of the Plant Nutrition Laboratory shared the techniques to implement this successful process that improves the agriculture environment while eliminating Cs toxicity.

### A brief sketch of our collaboration

In 2012, cell and molecule biologists from our universities started to exchange research materials and their experiences in cell biology (polyploidy breeding) of a wild berry and the introduction of ectomycorrhizae fungi to low pH environments. An ASU member organized a Forest Health Workshop for the IUFRO and COST-EU meeting at the Kaunas campus of ASU (now VMU) in 2012. Three ecophysiologicalists from Hokkaido University (HU) joined to share their research on forest biological activities under changing environments. The rector of ASU visited our university in May 2014 to conclude a Memorandum of Understanding (MOU) between our schools in 2015

Both universities proposed to organize an international seminar between the two universities and related organizations in 2019. Based on the discussions between the two universities (VMU and HU), we selected the topics “biodiversity conservation and natural resources management based on a biotechnological approach to plant-microbe interaction under a changing environment.”

The goal of our seminar was 1) to continue the research collaboration and promote student exchanges

based on what we learned in this seminar, 2) to use genetic resources based on the ABS in the COP 2010. We hope to collaborate further to improve the quality and quantity of haskap and blueberry plants. Also, we will strive to identify characteristics of various forest tree species that will allow them to grow in special soil conditions (low pH and wet). 3) Based on the impressive record of this seminar, we will improve our knowledge of Sustainable Development Goals (SDGs), primarily related to sustainable agriculture and forest resource management using environmental microbiology under changing environments. Our seminar will contribute to developing research and education in both universities.

### **Social impact and future perspectives**

Lithuania, located in central Europe, is well known to the Japanese for its agriculture specialties such as honey and linen and for the humane work of Mr. Chiune Sugiura during WWII. Lithuanian agricultural sciences are unique in developing biological production and management under a changing environment, such as elevated ozone, nitrogen (N) deposition, and excessive phosphorous deposits in water basins. Global climate change is creating new challenges for agriculture and forestry.

The School of Agriculture of Hokkaido University and the Agriculture Academy of VMU have similar histories in science. Hokkaido University originated from the Sapporo Agricultural College, founded in 1876, making it the oldest bachelor-level college in Japan. At the same time, the Agriculture Academy of VMU started in 1924, making it the oldest agriculture school in Lithuania (Lithuania Agriculture University was re-established in 1996 after the country gained its independence from the former Soviet Union). Based on our close collaboration from 2012, we hope to increase our activities in the future.

### **Brief summary of the extended abstract**

At first, we can learn how to make wise use of the genetic resources of wild plants, such as haskap (honeysuckle blueberry) (Hoshino et al. and Žilinskaitė et al.). In addition, we summarized tree breeding and forest genetics in Japan (Nakada and Kita et al.). Moreover, we will be able to improve the methods of sustainable bio-resource management based on environmental microbiology including alien species (Qu et al. on larch; Kitaoka et al. on black locust). We hope to investigate ways to stabilize seacoast forests struggling under salt stress and increased nitrogen deposition (Marozas et al., Augustaitis et al.). Also, we expect to obtain new biotechnology from Lithuania's significant efforts at afforestation using two or three species of pine (genus *Pinus* spp.) and black locust on dunes along the seacoast (Fujita et al., Matsunami et al.). We propose the wise use of plant-microbe interactions to establish tree and shrub plantations, focusing mainly on ectomycorrhizae and wood-decaying fungi. Ectomycorrhizae are expected to be greening materials for conifer plantations, and wood-decaying fungi are candidates for regulating extraordinary growth in vegetation, especially in black locust.

Based on our research, we will synthesize our previous knowledge and techniques in resource management under increasing CO<sub>2</sub> (Takagi et al.), acid soil adaptation in shrub (Watanabe), N deposition (Fujita et al.), and ground-level ozone in plant and insect interaction (Masui et al.). We will offer the results of our research to individuals involved in the forestry industry (Sugai et al.), biological conservation (Brazaitis and Marozas), and landscape management in both countries.

We thank LSSSF and JSPS for supporting the fund of our seminar and for FSC of Hokkaido University, Hokkaido Forestry Research Institute of HRO and Forestry & Forest Products Research Institute for offering the chance to showing current activities.

We hope this special issue will contribute to the agriculture science of our respective countries for many years to come. The contents of the special issue are shown after this page.

*Toshihiro Watanabe, Vitas Marozas, Takayoshi Koike*

Toshihiro Watanabe, Vitas Marozas, Takayoshi Koike

## Reference

Watanabe T, Kitaoka S, Koike T, Marozas V (2021) Biodiversity Conservation and Natural Resources Management based on a Biotechnological approach to Plant-Microbe interaction under a Changing Environment. Abstract, Hakuyo Print, Sapporo, Japan.

## Contents

### Key note presentation

Shinano T, Maruyama H, Watanabe T, Fujimoto H, Suzuki M: The role of potassium on the remediation for the radiocesium contaminated soil 5-7

### Cell, Genetics, and Breeding

Žilinskaitė S, Naugžemys D, Patamsytė J, Hoshino Y, Žvingila D: Investigation of genetic diversity and relationships of blue-berried honeysuckle (*Lonicera caerulea* L.) cultivars, genetic lines and populations at the Vilnius University Botanical Garden 8-11

Hoshino Y.: An overview of the current research program for haskap (*Lonicera caerulea*), a useful genetic resource in Hokkaido, Japan 12-14

Nakada R.: Breeding and genetic resource conservation of forest trees in Japan 15-18

Kita K, Moriguchi Y, Kon H.: Breeding a hybrid larch in Hokkaido, northern Japan 19-22

### Restoration ecology and alien species

Qu LY, Wang XN, Mao QZ, Agathokleous E, Choi DS, Tamai Y, Watanabe T, Koike T.: Responses of ectomycorrhizal diversity of larch and its hybrid seedlings and saplings to elevated CO<sub>2</sub>, O<sub>3</sub>, and high nitrogen loading 23-27

Kitaoka S, Fujita S, Watanabe Y, Choi DS, Watanabe T, Shinano T, Satoh F, Koike T.: Growth and nitrogen use characteristics of black locust, an invasive alien species, grown under different light and CO<sub>2</sub> conditions. 28-32

Matsunami S, Kitaoka S, Koike T, Hirata T, Marozas V, Satoh F: Regulating the growth black locust seedlings by inhibiting the sprouting roots; An examination of effectiveness of root-growth regulation plates 33-38

Fujita S, Noguchi K, Tange T.: Root growth of *Pinus thunbergii* seedlings related to the restoration of Tohoku region coastal forests after the disastrous tsunami 39-44

### Environmental monitoring

Augustaitis A, Augustaitiene I, Sidabriene D, Koike T, Marozas V.: Integrated effect of environmental changes on forest ecosystems in Lithuania: Strategies for adaptation to and mitigation of the main threats of global climate change 45-48

Takagi K, Aguilos M, Liang NS, Takahashi Y, Saigusa N, Koike T, Sasa K.: Long-term monitoring on the dynamics of ecosystem CO<sub>2</sub> balance recovering from a clear-cut harvesting in a cool-temperate forest 49-51

Marozas V, Preikšal Z, Koike T, Watanabe T.: Impact of nitrogen pollution disturbances on forest vegetation and fungi near a fertilizer factory. 52-58

Watanabe T.: Basic understanding of aluminum accumulator plants 59-62

Masui N, Tani A, Matsuura H, Agathokleous E, Watanabe T, Koike T.: Elevated ozone disrupts the plant-insect communication; Changes of attractiveness of Japanese white birch leaves to *Agelastica coerulea* via Biogenic Volatile Organic Compounds (BVOCs) 63-68

### Practical Forestry and horticulture

Brazaitis G, Marozas V.: Ecological management forests in Lithuania 69-72

Kitaoka S, Shinano T, Suzuki T, Shoji S, Hasegawa Y.: Application of a new film for horticultural use to convert UV-light to photosynthetic active radiation 73-77

Sugai T, Yokoyama S, Tamai Y, Mori H, Marchi E, Watanabe T, Satoh F, Koike T.: Effects of soil compaction on the seedlings growth and ectomycorrhizal fungal community in hybrid larch 78-80

**Symbolic plant: Haskap (Honey berry: *Lonicera caerulea*)****Flower of Haskap in May (at Sapporo)**

Two flowers are attached to one ovary. (by Prof. Y. Hoshino)

**Fruits of Haskap in July (at Sapporo)**

Color of fruits becomes dark purple when the fruits are matured.  
Many fruits are arranged along with a branch. (by Prof. Y. Hoshino)