

# HOKKAIDO UNIVERSITY

Title	Changes in Cell Wall Structure During Rhizoid Formation of Silvetia babingtonii (Fucales, Phaeophyceae) Zygotes
Author(s)	Yonamine, Rina; Ichihara, Kensuke; Tsuyuzaki, Shiro; Hervé, Cécile; Motomura, Taizo; Nagasato, Chikako
Citation	Journal of phycology, 57(4), 1356-1367 https://doi.org/10.1111/jpy.13178
Issue Date	2021-05-01
Doc URL	http://hdl.handle.net/2115/85093
Rights	This is the peer reviewed version of the following article: Yonamine, R., Ichihara, K., Tsuyuzaki, S., Hervé, C., Motomura, T. and Nagasato, C. (2021), Changes in Cell Wall Structure During Rhizoid Formation of Silvetia babingtonii (Fucales, Phaeophyceae) Zygotes. J. Phycol., which has been published in final form at https://doi.org/10.1111/jpy.13178. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
Туре	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	text.pdf



1	CHANGES IN CELL WALL STRUCTURE DURING RHIZOID FORMATION OF
2	SILVETIA BABINGTONII (FUCALES, PHAEOPHYCEAE) ZYGOTES <sup>1</sup> .
3	
4	Rina Yonamine
5	Graduate School of Environmental Science, Hokkaido University, Sapporo 060-0810,
6	Japan
7	
8	Kensuke Ichihara
9	Muroran Marine Station, Field Science Center for Northern Biosphere, Hokkaido
10	University, Muroran 051-0013, Japan
11	
12	Shiro Tsuyuzaki
13	Graduate School of Environmental Earth Science, Hokkaido University, Sapporo 060-
14	0810, Japan
15	
16	Cécile Hervé
17	Sorbonne Universités, CNRS, Integrative Biology of Marine Models (LBI2M), Station
18	Biologique de Roscoff, 29680 Roscoff, France
19	
20	Taizo Motomura
21	Muroran Marine Station, Field Science Center for Northern Biosphere, Hokkaido
22	University, Muroran 051-0013, Japan
23	

24 Chikako Nagasato<sup>2</sup>

25	Muroran Marine Station, Field Science Center for Northern Biosphere, Hokkaido
26	University, Muroran 051-0013, Japan
27	
28	<sup>2</sup> Author for correspondence: e-mail nagasato@fsc.hokudai.ac.jp, phone: +81-143-22-
29	2846, fax: +81-143-22-4135
30	
31	
32	Running Title: Rhizoid formation of Silvetia zygotes
33	

34 ABSTRACT

35 We examined the ultrastructure of the cell wall and immunolocalization of 36 alginates using specific antibodies against M-rich alginates and MG blocks during 37 rhizoid formation in fucoid zygotes, Silvetia babingtonii. The thallus region of 24-hour 38 old zygotes had a cell wall made of three layers with different fiber distribution. In the 39 12-hour old zygotes, three layers in the thallus were observed before rhizoid formation, 40 namely the inner, middle and outer layers. During rhizoid elongation, only the inner 41 layer was apparent close to the rhizoid tip area. Immunoelectron microscopy detected 42 M-rich blocks of alginate on the inner half of the cell wall, irrespective of the number of 43 layers in the thallus and rhizoid regions. The MG blocks were seen to cover a slightly 44 wider area than M-rich alginate blocks. It was suggested that parts of M in mannuronan 45 would be rapidly converted to G, and MG-blocks are generated. Transcriptome analysis 46 was performed using 3 -, 10 -, and 24 h-old zygotes after fertilization to examine the 47 relationship between gene expression and alginate synthesis over time. The expression 48 of two mannuronan C5-epimerase homologs that convert mannuronic acid into 49 guluronic acid in alginates were upregulated or downregulated over the course of the 50 examination.

51

52 Keywords: cell wall, cryofixation, immunoelectron microscopy, RNA-seq, Silvetia

53 *babingtonii*, transmission electron microscopy

54

55 Abbreviations: AF, after fertilization; BAM, brown algae monoclonal antibody; MC5E,

56 mannuronan C5-epimerase; RFCW, Rhizoid-flank cell wall; G, Guluronic acid; M,

57 Mannuronic acid; RTCW, Rhizoid-tip cell wall; TCW, Thallus cell wall

## 59 INTRODUCTION

60 Many seaweeds grow in the intertidal region where environmental factors, 61 including temperature, wave shock, intensity and quality of light, salinity, and drying, 62 are changeable. The cell wall protects cells from physical shocks and unfavorable 63 factors, such as herbivores. Therefore, seaweeds develop characteristic polysaccharides 64 in the cell wall to enhance stress tolerance (Kloareg and Quatrano 1988, Popper et al. 65 2011). In brown algae, the main components of the cell wall are alginates, sulfated 66 fucans, and cellulose. The mature sporophyte of laminarialean species contains three 67 times more alginate than cellulose and sulfated fucan, whereas that of fucalean species 68 contains more than twice the alginate compared to cellulose and sulfated fucan (Mabeau 69 and Kloareg 1987, Skriptsova et al. 2012). Alginate consists of 1,4 linked mannuronic 70 acid (M) and guluronic acid (G) residues. M-rich blocks produce flexible gels, while G-71 rich blocks cause rigidity. Accordingly, the M/G ratio relates to the properties of the cell 72 wall, such as gel strength and stretching. Young sporophytes of *Costaria costata* show a 73 high M/G ratio, and the ratio gradually decreases with thallus growth (Wu et al. 2014). 74 The conversion of M residue into G residue is carried out by different mannuronan C5-75 epimerases (MC5E) expressed in different developmental stages and environmental 76 conditions (Nyvall et al. 2003, Tonon et al. 2008). The alginate biosynthesis pathway in 77 brown algae is not fully resolved, especially the first production of a polymer of M 78 residues (mannuronan) (Michel et al. 2010). It is predicted that, shortly after synthesis, 79 the alginate should consist of M-rich blocks; therefore, the M-rich region might hold the 80 clues to the understanding of alginate synthesis.

81	There are some ultrastructural observations of the cell walls of the brown algae.
82	The results showed that three or more cell wall layers were identified in the apical cell
83	of Sphacelaria rigidula (Karyophyllis et al. 2000), the meristematic epidermal cells of
84	Dictyota dichotoma (Evans and Horrigan 1972), zygotes of fucoid algae (Novotny and
85	Forman 1975, Callow et al. 1978, Bisgrove and Kropf 2001), and the lateral cell wall in
86	gametophytes and sporophytes of <i>Ectocarpus siliculosus</i> (Terauchi et al. 2016). The cell
87	wall layers were distinguished by the distribution of the cell wall fibers and the matrix.
88	Terauchi et al. (2016) clarified the quantification of the frequency of the electron-
89	opaque fibrils (alginate fibrils) and their junction based on three-dimensional analysis
90	using rapid freezing fixation and electron microscopy and tomography. The results
91	showed that the alginate fibers were distributed more densely and in a more complex
92	arrangement in the inner layer than in the middle layer.
93	Tip-growing apical cells of Sphacelaria showed a characteristic cell wall
94	structure. In Sphacelaria rigidula, the apical dome of the tip cell had a two-layered cell
95	wall. A very thin amorphous layer (L1) covered the cell, and the fibrous layer (L2)
96	existed just beneath the L1 layer. Other parts of the tip cell and the thallus cells were
97	covered with four layers of cell walls, with L3 and L4 added to L1 and L2
98	(Karyophyllis et al. 2000). In Sphacelaria furcigera, bud (branch) formation was
99	observed (Burns et al. 1982a, b). The axial cell wall is composed of four layers (CW1-
100	CW4), but the initial bud region is covered by two newly synthesized layers (CWO and
101	CWI). Burns et al. (1982a) suggested that alginates are abundant in CW2, CW4, and
102	CWI, while sulfated fucans were prominent in CW1, CW3, and CWO. Cell wall
103	structure on the rhizoid tip in fucoid zygotes has been observed (Novotny and Forman
104	1975, Callow et al. 1978, Bisgrove and Kropf 2001). Deposition of the highly sulfated

fucan occurs in the tip growth region (Quatrano and Crayton 1973, Quatrano and Steven
106 1976, Callow et al. 1978, Brawley and Quatrano 1979, Torode et al. 2016). However,
107 the changes occurring in the cell wall structure during rhizoid development remain
108 unclear.

109 Different appearances in the multilayered cell wall could reflect differences in 110 cell wall components, which would produce different physicochemical properties. To 111 examine the distribution of polysaccharides on the cell wall, immunoelectron 112 microscopy using anti-alginate antibody (Chi et al. 1999, Nagasato et al. 2010, Terauchi 113 et al. 2012, Terauchi et al. 2016), an anti-fucoidan antibody (Nagasato et al. 2010), and 114 a binding assay using a conjugation of cellulase or alginate lyase-colloidal gold were 115 applied (Nagasato et al. 2010, Terauchi et al. 2016). At the light microscopic level, 116 metachromatic analysis using toluidine blue and periodic acid-Schiff (PAS) was 117 performed to detect alginate and sulfated fucan (Novotny and Forman 1975, Burns et al. 118 1982a). Recently, monoclonal antibodies that recognize the different structures of 119 alginate and sulfated fucan were produced, and immunofluorescence data on the 120 localization of these epitopes in some brown algal species were collected (Torode et al. 121 2016, Linardić and Braybrook 2017, Rabillé et al. 2019, Linardić et al. 2020). 122 Antibodies recognizing different epitopes showed spatial and temporal differences in 123 localization. By analyzing the localization of these antibodies through immunoelectron 124 microscopy, it was expected that we could obtain more useful information relating to 125 the multilayered cell wall of the brown algae. 126 In brown algae, there is little existing information on how the cell wall changes 127 to generate tip-growth from the thallus cell. To address this question, we observed cell

128 wall structure during rhizoid growth in the fucoid alga *Silvetia babingtonii* zygotes,

129	using the rapid freezing and freeze substitution technique for transmission electron
130	microscopy (TEM). This method prevents loss of soluble polysaccharides, namely
131	alginate and fucan. In this study, we focused on localization of the alginate by
132	immunoelectron microscopy using antibodies that recognize different epitopes.
133	Moreover, to examine the synthesis regulation of alginate on rhizoid growth. We carried
134	out transcriptomic analysis using mRNA obtained from zygotes before and after
135	germination.
136	
137	MATERIALS AND METHODS
138	Materials
139	Mature Silvetia babingtonii (Harvey) E.A. Serrão, T.O. Cho, S.M. Boo &
140	Brawley was collected from September to November 2006-2010 at Charatsunai, and
141	2018 at Botofurinai, Muroran, Hokkaido, Japan. The collected mature thalli were
142	brought to the laboratory and wiped with gauze after washing with sterile seawater. The
143	zygote preparation procedure was the same as in the previous paper (Abe 1970,
144	Nagasato et al. 2010). They were placed in continuous light (30-40 $\mu$ mol photons $\cdot$ m <sup>-2</sup> ·s <sup>-</sup>
145	<sup>1</sup> ) at 18 °C overnight. Then, the thalli were transferred to a cool chamber at 4 °C for 2 h
146	after darkening with aluminum foil. Adding fresh seawater induced the release of eggs
147	and sperm into the medium. The age of the zygotes was counted from 1 h after pouring
148	seawater. Zygotes were cultured in PES medium (Provasoli 1968) containing 40 $\mu g \cdot m L^2$
149	<sup>1</sup> chloramphenicol at 18 °C in continuous light.
150	

*Electron microscopy and Immunoelectron microscopy* 

152	Zygotes were cultured on gel support films (ATTO Co., Tokyo, Japan). Before
153	fixation, the films were cut into small triangle shapes less than 1cm. Rapid freezing and
154	freeze substitution procedures were performed according to previous reports (Nagasato
155	and Motomura 2002, Nagasato et al. 2010). Finally, the samples were embedded in a
156	Spurr low-viscosity resin (Polysciences, Warrington, USA). Ultrathin sections were cut
157	using a diamond knife on an ULTRACUT ultramicrotome (Reichert-Jung, Vienna,
158	Austria). Sections were picked up on a formvar-coated one-slot copper grid, stained
159	with EM stainer (Nisshin EM, Tokyo, Japan), an alternative stain for uranyl acetate
160	based on lanthanide salts, and then lead citrate (Reynolds 1963). Observations were
161	performed using a JEM-1011 electron microscope (JEOL, Tokyo, Japan).
162	Localization of the alginate was examined using antibodies against various
163	alginate blocks (Torode et al. 2016). Nickel grids with sections were incubated with
164	blocking solution (2.5% skimmed milk, 5% normal goat serum, and 0.05% $NaN_3$ in
165	PBS; 137 mM NaCl, 2.7 mM KCl, 4.9 mM Na <sub>2</sub> HPO <sub>4</sub> , 1.5 mM KH <sub>2</sub> PO <sub>4</sub> , pH 7.4) for 30
166	min at 37 °C and then transferred to anti-alginate antibodies, BAM6 (M-rich epitopes)
167	and BAM7 (MG epitopes) diluted 1:5 with PBS for 1 h at 37 °C. The grids were then
168	washed with PBS three times and incubated for 1 h at 37 °C with a goat anti-rat IgG
169	(whole molecule) conjugated with 10 nm colloidal gold particles (Sigma, St. Louis,
170	Missouri, USA) diluted 1:20 with blocking solution. After washing with PBS and
171	distilled water, sections on the grids were treated with EM stainer in this case.
172	
173	Quantification of colloidal gold particles
174	

To quantify the distribution of alginate in the thallus cell wall (TCW) andrhizoid-tip cell wall (RTCW), the distance between each colloidal gold particle and

176 plasma membrane was measured using ImageJ 64 (NIH, Bethesda, MD, USA). 177 Measurements were made within three replicate rectangle regions of cell wall. Each 178 rectangle stretched from the plasma membrane to the outer edge of the cell wall and had 179  $0.75 \,\mu\text{m}$  wide sides over the plasma membrane and at the surface of the cell wall (Fig. 180 S1). Three different zygotes were measured (n=9). The thickness of the cell wall 181 differed in each sample; thus, the relative length between the plasma membrane and the 182 edge of the cell wall was set as 0-1. The distance of gold particles was assumed to 183 follow a beta distribution because the distance was continuous and ranged from 0 to 1. 184 Because the number of particles differed between the antibodies, the relative density of 185 distance was used to investigate the distributions. The kernel function and bandwidth 186 obtained the function to express the density. Assessment of the function was confirmed 187 by goodness of fit tests (GFT). Beta regression was used to investigate the effects of 188 antibodies (BAM6 and BAM7), and the cell wall regions (TCW and RTCW) used as 189 explanatory variables, on the distance, used as a response variable. Since interactions 190 between these three explanatory variables were unlikely, the analyses were conducted 191 on the respective explanatory variables. All statistical analyses were performed using R 192 software (version 4.0.2) (R Core Team 2020) with the libraries fitdistrplus (Delignette-193 Muller and Dutang 2015) and betareg (Cribari-Neto and Zeileis 2009).

194

## 195 *RNA-seq and differential expression analysis*

Total RNA was extracted from zygotes at 3, 10, and 24 h after fertilization
using cetyltrimethylammonium bromide (CTAB) buffer (Pearson et al. 2006). The
extract was purified with a chloroform-isoamyl alcohol solution (24:1 v/v). After
precipitation with 3 M LiCl, contaminating DNA was removed with an RNase-free

200 DNase Set (Qiagen, Hilden, Germany). Total RNA samples with OD260/280 >1.8,

OD260/230>2.0, were used for library construction.

201

202 The cDNA library construction, sequencing with Illumina Novaseq6000 (150 203 bp paired-ends), and generating clean reads by filtering raw reads were performed by 204 Filgen Inc. (Nagoya, Japan). The clean reads were assembled using Trinity software 205 (Haas et al. 2013). The assembled contigs were clustered and filtered out using CD-206 HIT-EST on CD-HIT suite server (Li and Godzik 2006) with 90% similarity, and the 207 longest open reading frame (ORF) sequences were identified by TransDecoder (v5.5.0) 208 (Haas et al. 2013). Sequences with fewer than 100 amino acids were discarded. The 209 longest ORF sequences were annotated using the annotation pipeline on Maser (Management and Analysis System for Enormous Reads) platform (https://cell-210 211 innovation.nig.ac.jp) (Kinjo et al. 2018). Clean reads were aligned onto the longest ORF 212 sequences and quantified using Salmon (Patro et al. 2017). Data normalization and 213 identification of differentially expressed genes (DEGs) without replicates were 214 performed using an iterative *deseq* pipeline in the TCC package (Sun et al. 2013). False 215 discovery rates (FDR)  $\leq 0.05$  were considered as DEGs. The Z-scores based on the 216 transcripts per million (TPM) were calculated using R software (3.6.2; http://www.R-217 project.org/), and the heatmap of genes for the biosynthetic pathway of alginates was 218 drawn using the gplots (3.0.1.2) package. The Bioconductor package topGO version 219 2.38.1 (Alexa et al. 2006) were used for Gene Ontology (GO) enrichment analysis in R 220 software. Over-representative GO terms in up-regulated or down-regulated genes in 221 three comparisons (3 h AF vs 10 h AF, 3 h AF vs 24 h AF and 10 h AF vs 24 h AF) 222 were identified by Fisher's exact test in combination with the 'weight' algorithm, using 223 a minimum p-value cut-off of 0.01.

*Real-Time qPCR analysis* 

226	To analyze the transcriptional expression of DEG, RT-qPCR was performed.
227	Synthesize of cDNA as template was performed by PrimeScript <sup>TM</sup> RT reagent Kit with
228	gDNA Eraser (TaKaRa, Shiga, Japan) using total RNA 50 ng in 20 $\mu$ l reaction mixture
229	according to the manufacturer's protocol. Specific primers for each contig were
230	designed using Primer-BLAST (Table S1). The reaction mixture contained 5 $\mu$ L SYBR
231	Premix Ex Taq II (TaKaRa), 0.4 $\mu$ l specific primers (10 $\mu$ M), and 1 $\mu$ l cDNA template
232	in a total of 10 $\mu$ l. qPCR was performed using Eco Real-Time PCR system (Illumina,
233	San Diego, CA, USA). After 95 °C for 30 s, the cycling parameters were 95 °C for 5 s,
234	and 60 $^{\circ}$ C for 1 min and were performed for 40 cycles. The melting curve analysis was
235	as follows: 95 °C for 15 s, 55 °C for 15 s, and 95 °C for 15 s. Transcript levels
236	corresponding to the comp32408_c2_seq24.p1 and comp33109_c0_seq1.p1 genes
237	(putative mannuronan C5-epimerase) were compared with the transcript levels of the
238	actin gene (actin; comp21090_c0_seq1.p1). The RT-qPCR was performed in triplicate
239	using three RNA samples from each zygote at the ages of 3, 10, and 24 h. The
240	differences in mean relative expression level between the developmental hours (3, 10,
241	and 24 h) were examined by Tukey's multiple comparison test.
242	
243	Identifying mannuronan C5-epimerase from S. babingtonii assembled contigs and

*molecular phylogenetic analysis* 

A reciprocal blast search against 24,486 proteins of the brown alga *Ectocarpus siliculosus* (http://bioinformatics.psb.ugent.be/orcae/overview/EctsiV2) was also
 conducted to identify genes for enzymes in the biosynthetic pathway of alginates using

249 To construct the MC5E phylogenetic tree, 36 sequences from *E. siliculosus* 250 (https://bioinformatics.psb.ugent.be/orcae/overview/EctsiV2), one sequence from 251 Saccharina japonica (Inoue et al. 2016), and seven sequences from assembled S. 252 babingtonii contigs were gathered. Three bacterial AlgG sequences from Azotobacter 253 vinelandii (P70805), Pseudomonas aeruginosa (Q51371), and P. fluorescens (P59828) 254 were added to the alignment file as an outgroup. Phylogenetic analysis of amino acid 255 sequences of MC5E was carried out using web service NGphylogeny.fr 256 (https://ngphylogeny.fr/) (Lemoine et al. 2019). The sequences were aligned using 257 MAFFT with the iterative refinement method and the scoring matrix Blosum62 (Katoh 258 et al. 2002). Ambiguously aligned non-informative regions were removed with BMGE 259 (Criscuolo and Gribaldo 2010). Twenty sequences were removed due to insufficient 260 sequence length or having identical sequences to the others from the phylogenetic 261 analysis. The model test for maximum likelihood (ML) method was carried out by SMS 262 (Lefort et al. 2017), and phylogenetic analysis was performed using PhyML (Guindon 263 et al. 2010). The best-fit model was Blosum62 + G + I + F based on Akaike information 264 criteria (AIC). Bootstrap values (Felsenstein 1985) were obtained from the analyses of 265 100 pseudoreplicates with Booster (Lemoine et al. 2018). Domain searches against the

BLAST+ version 2.7.1 (ftp://ftp.ncbi.nlm.nih.gov/blast/executables/blast+/LATEST).

- 266 InterPro server (https://www.ebi.ac.uk/interpro/) were also carried out.
- 267

248

#### 268 RESULTS

Immediately after fertilization, the *Silvetia* zygotes showed a spherical shape
(Fig. 1a). Rhizoid formation was initiated at 12 h after fertilization (AF) (Fig. 1b). The
first and second cytokinesis occurred at 20 h and 24 h AF, respectively (Fig. 1c, d). We

observed the ultrastructure and occurrence of alginate in cell walls of the zygotes at 12 h

and 24 h AF. Three cell areas were observed in terms of cell wall structures (Fig. 1e): (i)

the thallus cell wall (TCW), (ii) the rhizoid-flank cell wall (RFCW), and (iii) the

275 rhizoid-tip cell wall (RTCW).

276 At 12 h AF, the TCW had an electron-opaque structure with parallel-oriented 277 fibrils (Fig. 2a). In the RFCW, three layers with different contrasts, dark-light-dark, 278 were visible (Fig. 2b). The RTCW was less stained and had irregularly oriented fibrils 279 (Fig. 2c). The cell wall structure in the TCW at 24 h AF (Fig. 3a) was dependent on the 280 cutting angle of sections relative to the plasma membrane. Obliquely cut sections in 281 which the lipid bilayer of plasma membrane is unclear revealed three distinct layers 282 with different opacities and fiber orientations (Fig. 3b). The cell wall was roughly 283 divided into the inner layer (IL) having a rough fibrous net with porous, middle layer (ML) showing the fine fibrous net, and outer (OL) layers with a rough fibrous net. The 284 285 width of the layers was calculated. Each layer was almost the same width representing 286 32, 37, and 31% of the whole cell wall in IL, ML, and OL, respectively. The percentage 287 showed the mean value from the result by triplicated counting on each of three zygotes 288 (Table S2). However, it was difficult to define each layer on a perpendicular section of 289 the cell wall to the plasma membrane, because the difference of orientation among the 290 three layers was hard to be clarified by narrow fiber spacing and high electron-opacity 291 (Fig. 3c), as in 12 hour-old zygotes (Fig. 2a). In the RFCW, the outer and middle layers 292 were hardly observed (Fig. 3d, e). First, the outer layer was not apparent at the thallus 293 pole-facing side of the rhizoid tip (the site indicated by a double arrowhead in Fig. 3a) 294 and the middle layer disappeared toward the tip proper. The remaining inner layer 295 became thicker closer to the rhizoid tip (Fig. 3f). While well-structured in other

regions, the distribution of fibrils was irregular or wavy closer to the tip. The mucilage
material covered the outside of the rhizoid. The amorphous mucilaginous layer was not
included in the numbering of the cell wall layer in this study.

299 To locate alginate epitopes in the different cell wall regions, immunoelectron 300 microscopy using anti-alginate antibodies was used at 24 h AF (Fig. 4). Localization of 301 M-rich blocks with BAM6 antibody was evident in the inner half of the TCW, RFCW, 302 and RTCW (Fig. 4a, b, c). Fewer M-rich blocks were recognized in the RFCW (Fig. 4b, 303 Table 1), compared to the other two regions. With BAM7-conjugated gold particles, 304 MG-alginates appeared more uniformly and abundantly distributed in the cell wall than 305 M-rich blocks (Figs. 4d, e, f). The mean value of the number of gold particles using 306 BAM7 was shown in Table 1. The mean value of gold particle number with BAM7 was 307 higher as RTCW>RFCW>TCW. However, Tukey's multiple comparison tests showed 308 no significant difference in the number of gold particles with BAM6 and BAM7 among 309 TCW, RFCW, and RTCW.

310 To make comparisons across different antibodies, the width of the cell wall 311 between the plasma membrane and the edge of the cell wall was divided into ten 312 segments. Distributional patterns of gold particles in the cell wall over the relative 313 distance (0-1) were estimated by the function fitted at 10 segments (GFT, P < 0.01 for 314 all cases) (Fig. 5). The density of particles binding to BAM6 peaked at a distance of 315 approximately 0.3 in TCW, and it was flattened through 0.7 while decreasing. And it 316 dropped after 0.7 toward 1.0. The high density segments were at 0.1 to 0.7, which was 317 consistent with the region of the half of the inner layer and whole a middle layer. The 318 high density distribution with BAM7 in TCW was with a range of 0 to 0.7, which was 319 flatter than it with BAM6 in TCW. These results showed that the gold particles binding

320 to BAM7 were distributed closer to the plasma membrane compared with BAM6 in 321 TCW (beta regression, Z=-6.193, P<0.001). In RTCW, the density of particles binding 322 to BAM6 peaked at a distance of approximately 0.2, and became flatter between 0.5 to 323 0.9 after a peak. Compared with BAM6 in TCW, the high density segments were 324 shifted closer to plasma membrane (beta regression, Z=-6.726, P<0.001). In the density 325 distributions of gold particle with BAM7 in RTCW was flat with a range of 0 to 1.0, 326 and a plateau was observed from 0.1 to 0.4. There was no significant difference in the 327 distribution of density of gold particle between BAM7 in RTCW and BAM7 in TCW, 328 or BAM6 in RTCW.

329 To understand the genes involved in early development of the zygotes, 330 transcriptomic analysis was performed at three different developmental stages (3 h AF: 331 time of increased wall thickness, 10 h AF: time of initiation of rhizoid germination, and 332 24 h AF: time when the second cytokinesis finished and the rhizoid elongated) under 333 the current culture condition (Fig. S2). The timeline might change depending on the 334 cultural condition, so the event of characteristic morphological change in the early 335 development of fucoid zygotes was treated in this study. The short read data are 336 summarized in Table S3. *De novo* assembly produced 97,825 contigs. Filtering with 337 CD-HIT-EST reduced the size of the assembly to 78,268 contigs, and the 27,622 338 longest ORF contigs were detected by TransDecoder. A total of 11,087 contigs (40%) 339 were annotated by BLASTX with the UniProt database (Table S4), and 61-63% of reads 340 were mapped to the 25,890 longest ORF contigs. Then, differentially expressed genes 341 (DEGs) were detected for 107 contigs in 3 h AF vs. 10 h AF, 47 contigs in 3 h AF vs. 342 24 h AF, 125 contigs in 10 h AF vs. 24 h AF of the 25,890 contigs (Fig. S3; Tables S5-343 S7). GO enrichment analysis indicates that sulfotransferase activity (GO:0008146) and

extracellular region (GO:0005576) were significant in the regulated genes at 24 h AF(Table S8).

346 Reciprocal best-hit blast against *Ectocarpus siliculosus* reference proteins 347 recovered 39.6% (9,699) proteins in the group of longest ORFs in S. babingtonii 348 (Table S9). The BLAST search indicated that some alginate synthetic pathway 349 enzymes, two mannose-6-phosphate isomerase (MPI), one phosphomannomutase 350 (PMM), one GDP-mannose 6-dehydrogenase (GMD), and seven MC5E, were conserved 351 in S. babingtonii. The expression patterns of the enzymes' genes are summarized in Fig. 352 S4. The expression levels of MPI (comp32248 c1 seq4.p1) and GMD 353 (comp32002 c0 seq6.p1) remained stationary throughout the experiment. Five MC5Es 354 (comp12459 c0 seq1.p1, comp21318 c0 seq3.p1, comp25876 c0 seq1.p1, 355 comp32408 c2 seq24.p1 and comp32951 c0 seq4.p1) were up-regulated as the 356 developmental stage proceeded, and one (comp33109 c0 seq1.p1) was slightly 357 suppressed in 24 h AF. 358 To clarify the molecular function, domain search and phylogenetic analysis of 359 MC5Es to clarify the molecular function were performed. The catalytic MC5E domain 360 and the signal peptide region were detected in four contigs (Fig. S5). The phylogenetic 361 tree of MC5Es shows five clades supported with high to moderate value (42-100%) and 362 seven MC5Es homologs of S. babingtonii were scattered within these five clades (Fig. 363 S5). Two *MC5E* homologs (comp32408 c2 seq24.p1 and comp33109 c0 seq1.p1) 364 were detected as DEGs in the different developmental stages of the zygotes. RT-qPCR 365 was carried out on two contigs (comp32408 c2 seq24.p1 and comp33109 c0 seq1.p1) 366 (Fig. 6). The expression level of comp32408 c2 seq24.p1stayed lower until 10 h AF

- and increased eight times at 24 h AF. Comp33109\_c0\_seq1.p1 expression leveldeclined to one-fifth at 10 h AF from 3 h AF.
- 369

### 370 DISCUSSION

371 Ultrastructural comparisons of cell walls of the rhizoid and thallus parts of 372 fucoid zygotes have been reported (Novotny and Forman 1975, Vreugdenhil et al. 1976, 373 Callow et al. 1978, Bisgrove and Kropf 2001). Previous studies showed that the cell 374 wall of the thallus region consists of two layers, while the rhizoid cell wall had one or 375 two layers (Novotny and Forman 1975, Vreugdenhil et al. 1976, Callow et al. 1978, 376 Bisgrove and Kropf 2001). We observed three layers in the TCW in Silvetia babingtonii 377 zygotes at 24 h AF. Occasionally, it was difficult to define the number of layers in the 378 TCW, even in the same zygote. The cutting angle of the section affected the 379 ultrastructural appearance of the cell wall structure (Terauchi et al. 2016). To avoid this, 380 we prepared consecutive serial sections in each zygote and observed the cell wall 381 structure. Bisgrove and Kropf (2001) mentioned a porous area adjacent to the plasma 382 membrane within the thicker inner layer using chemical fixation. Although they did not 383 count it as another layer, we considered that the porous area corresponds to the inner 384 layer observed in the present study using rapid-freezing and freeze substitution (Fig. 385 3b).

Regarding the rhizoid cell wall, we observed that the two outer and middle layers from the cell wall of the thallus were not apparent along the rhizoid flank (RFCW), and only the inner layer surrounded the rhizoid tip. The inner layer became gradually thicker, and the cell wall fibrils showed an irregular orientation close to the rhizoid tip. This observation is the first to explain how cell wall structure changes

391 during rhizoid elongation in fucoid zygotes at the ultrastructural level. We considered 392 that the cleavage of the two outer layers occurred during rhizoid elongation, not during 393 its initiation. In the RTCW of the zygote at 12 h AF, namely, the stage of initiation of 394 rhizoid elongation, the cell wall layer could not be clearly seen (Fig. 2c); however, the 395 three layers were observed in the RFCW (Fig. 2b). At the beginning of tip growth, the 396 three layers were present; however, the outer two layers might break due to the 397 extension of the cell during further tip outgrowth. On the other hand, the inner layer 398 could be extended by the addition of new cell wall materials and remodeling. 399 Ultrastructural observation in the pollen tube growth of Lycopersicum peruvianum and 400 *Lilium longiflorum* showed that a part of the cell wall is broken on the germination 401 (Cresti et al. 1977, Miki-Hirosige and Nakamura 1982). During germination, new cell 402 wall subdivided into pectic outer layer and callosic inner layer is synthesized between 403 the intine and plasma membrane. The apical dome of pollen tube is always covered with 404 the callosic inner layer. The outer pectic layer breaks in lily pollen germination with the 405 intine. Newly synthesized polysaccharide is important for the tip growth in this case. In 406 the zygotes of S. babingtonii, the appearance timing of three layers is not known, 407 however, there is the possibility that a newly synthesized inner cell wall layer before 408 rhizoid formation would be essential for its elongation as in the pollen tube growth of 409 lily. 410

While, in the growing root hair of *Arabidopsis thaliana*, breaks of cell wall has
not been observed. The growing root hair cell wall shows a single layer, and the mature
one is surrounded by two layers with different textures and orientation of
microfilaments (MFs) (Akkerman et al. 2012). A similar pattern has been observed in *Equisetum hyemale* (Emons and Wolters-Arts 1983). Expansin, a protein that promotes

415 cell wall loosening without degradation of cell wall polymers, is involved in root hair 416 initiation and elongation in many land plants (Cho and Cosgrove 2002, Cosgrove 2015). 417 The apical dome of the pollen tube consists of an outer layer with pectin and a 418 translucent inner layer with callose and cellulose in A. thaliana (Dardelle et al. 2010, 419 Chebli et al. 2012) and Nicotiana tabacum (Geitmann et al. 1995, Ferguson et al. 1998). 420 In the apical dome of the outer layer, highly esterified pectin deposits are created by 421 exocytosis, while localization of low-esterified pectin increases from the transition area 422 between the apex and axial region in A. thaliana (Chebli et al. 2012). De-esterified 423 pectin is produced by pectin methyl esterase from methyl esterified pectin, and the 424 degree and pattern of methyl-esterification of pectin influence the elasticity of the gel 425 (Willats et al. 2001). In the transition area, remodeling of the cell wall by accumulation 426 of callose and cellulose with de-esterification of pectin occurs, and this affects the 427 aperture of the pollen tube and promotes tip-growth in A. thaliana (Chebli et al. 2012). 428 We had expected that elongation of the rhizoid cell wall would be caused by 429 secretion of new cell wall material and remodeling of existing components in the 430 zygotes of S. babingtonii. We focused on the spatial distribution of alginate because it is 431 the main polysaccharide of the brown algal cell wall and affects its mechanical 432 properties. Therefore, immunoelectron microscopy using antibodies against the 433 different epitopes, BAM6 and BAM7, was conducted against the TCW and the RTCW. 434 There was no significant difference in the distribution of gold particles of BAM7 435 between the thallus and rhizoid regions. The high density of the gold particle with 436 BAM6 was observed close to the plasma membrane in the RTCW, which differs from 437 the situation in the TCW (Fig. 5). Alginate synthesis pathways in brown algae are not 438 fully understood. However, it is considered that alginate is initially synthesized as a

439 polymer of mannuronic acid (mannuronan) by alginate synthase, and a part of the 440 mannuronic acid (M) is converted into guluronate acid (G) reside by mannuronan C5-441 epimerases (MC5Es) (Michel et al. 2010). In other words, the M residue-rich area 442 would be considered to be the relatively newly synthesized alginate. However, this 443 study showed that the distribution of gold particles binding to BAM7 was wider than 444 BAM6 at the sites adjacent to the plasma membrane at TCW and RTCW. From this 445 result, it was expected that parts of M in mannuronan would be rapidly converted to G 446 after polymerization. In this study, the outer layer was labeled as neither BAM6 nor 447 BAM7. In *Ectocarpus siliculosus*, the alginate lyase conjugated with gold particles 448 binds to the entire cell wall but labeling with an anti-alginate antibody is limited to the 449 inner two layers of the three (Terauchi et al. 2016). Therefore, it was considered that the 450 outer layer of S. babingtonii zygotes contained alginate with different types of MG 451 blocks, which are recognized by BAM6 and BAM7. We also performed 452 immunoelectron microscopy using the anti-alginates BAM8 to BAM11 antibodies 453 (Torode et al. 2016), however they were unable to label the outer layer of the cell wall 454 in S. babingtonii zygotes. 455 Transcriptomic analysis was performed in 3-, 10-, and 24-hour-old zygotes of 456 S. babingtonii and focused on gene expression related to alginate modification. The 457 enrichment of term "sulfotransferase activity" (GO:0008146) and "extracellular region" 458 (GO:0005576) indicate some cell wall components with sulfated fucan might be

- 459 actively modified during rhizoidal growth. By constructing a *de novo* transcriptome
- 460 assembly, we found homologous genes required for alginate synthesis and modification
- 461 (Michel et al. 2010). Previous genomic or transcriptomic studies revealed that there
- 462 were 31 MC5E genes in *Ectocarpus* (Michel et al. 2010), 105 genes in *Saccharina*

463 japonica (Ye et al. 2015), and 38 genes in Sargassum thunbergii (Liu et al. 2014). In 464 this study, 19 MC5E genes were identified by BLAST against the UniProt database 465 (Table S3), and the number was smaller than that of other brown algae. Because our 466 RNA-seq data are only from the early developmental stages (3-24 h AF), some MC5E 467 genes in S. babingtonii might not be expressed. Seven S. babingtonii's MC5Es included 468 a common MC5E domain, but each MC5E included a variety of domain structures 469 similar to genes in *Ectocarpus* (Fischl et al. 2016). This protein domain diversity in the 470 MC5Es family contributes to the regulation of alginate composition and remodeling in 471 the cell wall. Within the data, two MC5E homologs (comp32408 c2 seq24.p1 and 472 comp33109 c0 seq1.p1) showed significantly different gene expression during 473 development of the zygotes. This is consistent with the different transcriptional patterns 474 of MC5E genes related to developmental stage, thallus region, and different phases of 475 an algal life cycle (Nyvall et al. 2003, Tonon et al. 2008, Fischl et al. 2016, Linardić et 476 al. 2020). Some reports show that the transcripts required for rhizoid formation were 477 synthesized prior to 5 h AF (Quatrano 1968, Kropf et al. 1989). From these findings, 478 comp33109 c0 seq1.p1 may support rhizoid germination, while it is also possible that 479 the gene may be related to the expansion of the cell wall and the formation of the 480 multilayer before rhizoid formation. The expression level of comp32408 c2 seq24.p1 481 increased in 24 h AF, suggesting that the gene would involve cell wall remodeling for 482 rhizoid elongation.

There are some reports showing cross-linking of polyphenol to alginate in the cell wall in brown algae (Salgado et al. 2009, Deniaud-Bouët et al. 2014). It is considered that phenol involves in the adhesion of the zygotes on the substratum and rigidity of the cell wall. Vanadium-dependent bromoperoxidase is a candidate to

487	catalyze alginate-phenol complex. RNA-seq data in this study showed the 4 in 12
488	candidate vBPO genes were detected as DEG (Table S4, S7). During early development
489	of F. vesiculosus zygotes, vBPO activity increases gradually after fertilization
490	(Lemesheva et al. 2020). These results matched this study data. It is expected that how
491	the alginate-phenol complex affects the property of the cell wall and involved in the
492	morphogenesis becomes clear.
493	
494	ACKNOWLEDGEMENTS

495 We thank Ms. Theone van der Merwe, Daekin University for her critical

496 reading of the manuscript and helpful suggestions. This work was financially supported

497 by a grant from the Sasakawa Scientific Research Grant.

#### **498 REFERENCE**

- Abe, M. 1970. Method of inducing egg liberation in *Fucus evanescens*. *Bot Mag Tokyo*.
  83:254-5.
- 501 Akkerman, M., Franssen-Verheijen, M., Immerzeel, P., Hollander, L., Schel, J. &
- 502 Emons, A. 2012. Texture of cellulose microfibrils of root hair cell walls of
  503 *Arabidopsis thaliana, Medicago truncatula*, and *Vicia sativa. J. Microscop.*504 247:60-7.
- Alexa, A., Rahnenführer, J. & Lengauer, T. 2006. Improved scoring of functional
  groups from gene expression data by decorrelating GO graph structure. *Bioinformatics*. 22:1600-7.
- 508 Bisgrove, S. R. & Kropf, D. L. 2001. Cell wall deposition during morphogenesis in
  509 fucoid algae. *Planta*. 212:648-58.
- 510 Brawley, S. H. & Quatrano, R. S. 1979. Sulfation of fucoidin in Fucus embryos: IV.
- 511 Autoradiographic investigations of fucoidin sulfation and secretion during
- 512 differentiation and the effect of cytochalasin treatment. *Dev. Biol.* 73:193-205.
- 513 Burns, A. R., Oliveira, L. & Bisalputra, T. 1982a. A histochemical study of bud
- 514 initiation in the brown alga *Sphacelaria furcigera*. *New Phytol*. 92:297-307.
- 515 Burns, A. R., Oliveira, L. & Bisalputra, T. 1982b. A morphological study of bud
  516 initiation in the brown alga *Sphacelaria furcigera*. *New Phytol*. 92:309-25.
- 517 Callow, M. E., Coughlan, S. J. & Evans, L. V. 1978. The role of Golgi bodies in
  518 polysaccharide sulphation in *Fucus* zygotes. *J. Cell Sci.* 32:337-56.
- 519 Chebli, Y., Kaneda, M., Zerzour, R. & Geitmann, A. 2012. The cell wall of the
- 520 Arabidopsis pollen tube-spatial distribution, recycling, and network formation
  521 of polysaccharides. *Plant Physiol.* 160:1940-55.

- 522 Chi, E. S., Henry, E. C., Kawai, H. & Okuda, K. 1999. Immunogold-labeling analysis
  523 of alginate distributions in the cell walls of chromophyte algae. *Phycol. Res.*524 47:53-60.
- 525 Cho, H. T. & Cosgrove, D. J. 2002. Regulation of root hair initiation and expansin gene
  526 expression in Arabidopsis. *Plant Cell*. 14:3237-53.
- 527 Cosgrove, D. J. 2015. Plant expansins: diversity and interactions with plant cell walls.
  528 *Curr. Opin. Plant Biol.* 25:162-72.
- 529 Cresti, M., Pacini, E., Ciampolini, F. & Sarfatti, G. 1977. Germination and early tube
  530 development in vitro of *Lycopersicum peruvianum* pollen: ultrastructural
  531 features. *Planta*. 136:239-47.
- 532 Cribari-Neto, F. & Zeileis, A. 2009. Beta regression in R. J. Stat. Softw. 34:1-24.
- 533 Criscuolo, A. & Gribaldo, S. 2010. BMGE (Block Mapping and Gathering with
- 534 Entropy): A new software for selection of phylogenetic informative regions
  535 from multiple sequence alignments. *BMC Evol. Biol.* 10.
- 536 Dardelle, F., Lehner, A., Ramdani, Y., Bardor, M., Lerouge, P., Driouich, A. & Mollet,
- 537 J. C. 2010. Biochemical and immunocytological characterizations of
- 538arabidopsis pollen tube cell wall. Plant Physiol. 153:1563-76.
- 539 Delignette-Muller, M. L. & Dutang, C. 2015. fitdistrplus: An R package for fitting
  540 distributions. *J. Stat. Softw.* 64:1-34.
- 541 Deniaud-Bouët, E., Kervarec, N., Michel, G., Tonon, T., Kloareg, B. & Hervé, C. 2014.
- 542 Chemical and enzymatic fractionation of cell walls from Fucales: insights into
- 543 the structure of the extracellular matrix of brown algae. *Ann. Bot.* 114:1203-16.

Emons, A. M. C. & Wolters-Arts, A. M. C. 1983. Cortical microtubules and microfibril
deposition in the cell wall of root hairs of Equisetum hyemale. Protoplasma.
117:68-81.
Evans, L. V. & Holligan, M. S. 1972. Correlated light and electron microscope studies
on brown algae I. Localization of alginic acid and sulphated polysaccharides in
Dictyota. New Phytol. 71:1161-72.
Felsenstein, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap.
Evolution. 39:783-91.
Ferguson, C., Teeri, T. T., Siika-aho, M., Read, S. M. & Bacic, A. 1998. Location of
cellulose and callose in pollen tubes and grains of Nicotiana tabacum. Planta.
206:452-60.
Fischl, R., Bertelsen, K., Gaillard, F., Coelho, S., Michel, G., Klinger, M., Boyen, C.,
Czjzek, M. & Hervé, C. 2016. The cell-wall active mannuronan C5-epimerases
in the model brown alga Ectocarpus: from gene context to recombinant
protein. Glycobiology. 26:973-83.
Geitmann, A., Li, Y. Q. & Cresti, M. 1995. Ultrastructural immunolocalization of
periodic pectin depositions in the cell wall of Nicotiana tabacum pollen tubes.
Protoplasma. 187:168-71.
Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W. & Gascuel, O.
2010. New algorithms and methods to estimate maximum-likelihood
phylogenies: Assessing the performance of PhyML 3.0. Syst. Biol. 59:307-21.
Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J.,
Couger, M. B., Eccles, D., Li, B., Lieber, M., MacManes, M. D., Ott, M.,
Orvis, J., Pochet, N., Strozzi, F. Weeks, N., Westerman, R., William, T.,

568	Dewey, C. N., Henschel, R., LeDuc, R. D., Friedman N. & Regev, A. 2013. De
569	novo transcript sequence reconstruction from RNA-seq using the Trinity
570	platform for reference generation and analysis. Nat. Protoc. 8:1494-512.
571	Inoue, A., Satoh, A., Morishita, M., Tokunaga, Y., Miyakawa, T., Tanokura, M. &
572	Ojima, T. 2016. Functional heterologous expression and characterization of
573	mannuronan C5-epimerase from the brown alga Saccharina japonica. Algal
574	<i>Res.</i> 16:282-91.
575	Karyophyllis, D., Katsaros, C. & Galatis, B. 2000. F-actin involvement in apical cell
576	morphogenesis of Sphacelaria rigidula (Phaeophyceae): mutual alignment
577	between cortical actin filaments and cellulose microfibrils. Eur. J. Phycol.
578	35:195-203.
579	Katoh, K., Misawa, K., Kuma, K. I. & Miyata, T. 2002. MAFFT: a novel method for
580	rapid multiple sequence alignment based on fast Fourier transform. Nucleic
581	Acids Res. 30:3059-66.
582	Kinjo, S., Monma, N., Misu, S., Kitamura, N., Imoto, J., Yoshitake, K., Gojobori, T. &
583	Ikeo, K. 2018. Maser: One-stop platform for NGS big data from analysis to
584	visualization. Database. 2018:1-12.
585	Kloareg, B. & Quatrano, R. S. 1988. Structure of cell walls of marine algae and
586	ecophysiological functions of the matrix polysaccharides. Oceanogr. Mar. Bio.
587	Annu. Rev. 26:259-315.
588	Kropf, D. L., Hopkins, R. & Quatrano, R. S. 1989. Protein synthesis and morphogenesis
589	are not tightly linked during embryogenesis in Fucus. Dev. Biol. 134:452-61.
590	Lefort, V., Longueville, J. E. & Gascuel, O. 2017. SMS: Smart Model Selection in
591	PhyML. Mol. Biol. Evol. 34:2422-4.

592	Lemesheva, V., Birkemeyer, C., Garbary, D. & Tarakhovskaya, E. 2020. Vanadium-
593	dependent haloperoxidase activity and phlorotannin incorporation into the cell
594	wall during early embryogenesis of Fucus vesiculosus (Phaeophyceae). Eur. J.
595	<i>Phycol.</i> 55:275-84.
596	Lemoine, F., Correia, D., Lefort, V., Doppelt-Azeroual, O., Mareuil, F., Cohen-
597	Boulakia, S. & Gascuel, O. 2019. NGPhylogeny.fr: new generation
598	phylogenetic services for non-specialists. Nucleic Acids Res. 47:W260-5.
599	Lemoine, F., Entfellner, J. B. D., Wilkinson, E., Correia, D., Felipe, M. D., De Oliveira,
600	T. & Gascuel, O. 2018. Renewing Felsenstein's phylogenetic bootstrap in the
601	era of big data. Nature. 556:452-6.
602	Li, W. & Godzik, A. 2006. Cd-hit: A fast program for clustering and comparing large
603	sets of protein or nucleotide sequences. Bioinformatics. 22:1658-9.
604	Linardić, M. & Braybrook, S. A. 2017. Towards an understanding of spiral patterning in
605	the Sargassum muticum shoot apex. Sci. Rep. 7:13887.
606	Linardić, M., Cokus, S. J., Pellegrini, M. & Braybrook, S. A. 2020. Growth of the
607	Fucus embryo: insights into wall-mediated cell expansion through mechanics
608	and transcriptomics. bioRxiv. https://doi.org/10.1101/2020.01.29.925107
609	Liu, F., Sun, X., Wang, W., Liang, Z. & Wang, F. 2014. De novo transcriptome
610	analysis-gained insights into physiological and metabolic characteristics of
611	Sargassum thunbergii (Fucales, Phaeophyceae). J. Appl. Phycol. 26:1519-26.
612	Mabeau, S. & Kloareg, B. 1987. Isolation and analysis of the cell walls of brown algae:
613	Fucus spiralis, F. ceranoides, F. vesiculosus, F. serratus, Bifurcaria bifurcata
614	and Laminaria digitata. J. Exp. Bot. 38:1573-80.

615	Michel, G., Tonon, T., Scornet, D., Cock, J. M. & Kloareg, B. 2010. The cell wall
616	polysaccharide metabolism of the brown alga Ectocarpus siliculosus. Insights
617	into the evolution of extracellular matrix polysaccharides in Eukaryotes. New
618	<i>Phytol.</i> 188:82-97.
619	Miki-Hirosige, H. & Nakamura, S. 1982. Process of metabolism during pollen tube wall
620	formation. Microscopy. 31:51-62.
621	Nagasato, C., Inoue, A., Mizuno, M., Kanazawa, K., Ojima, T., Okuda, K. &
622	Motomura, T. 2010. Membrane fusion process and assembly of cell wall
623	during cytokinesis in the brown alga, Silvetia babingtonii (Fucales,
624	Phaeophyceae). Planta. 232:287-98.
625	Nagasato, C. & Motomura, T. 2002. Ultrastructural study on mitosis and cytokinesis in
626	Scytosiphon lomentaria zygotes (Scytosiphonales, Phaeophyceae) by freeze-
627	substitution. Protoplasma. 219:140-9.
628	Novotny, A. M. & Forman, M. 1975. The composition and development of cell walls of
629	Fucus embryos. Planta. 122:67-78.
630	Nyvall, P., Corre, E., Boisset, C., Barbeyron, T., Rousvoal, S., Scornet, D., Kloareg, B.
631	& Boyen, C. 2003. Characterization of mannuronan C-5-epimerase genes from
632	the brown alga Laminaria digitata. Plant Physiol. 133:726-35.
633	Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. 2017. Salmon
634	provides fast and bias-aware quantification of transcript expression. Nat.
635	Methods. 14:417-9.
636	Pearson, G., Lago-Leston, A., Valente, M. & Serrão, E. 2006. Simple and rapid RNA
637	extraction from freeze-dried tissue of brown algae and seagrasses. Eur. J.
638	<i>Phycol.</i> 41:97-104.

639	Popper, Z. A., Michel, G., Hervé, C., Domozych, D. S., Willats, W. G. T., Tuohy, M.
640	G., Kloareg, B. & Stengel, D. B. 2011. Evolution and diversity of plant cell
641	walls: from algae to flowering plants. Annu. Rev. Plant Biol. 62:567-90.
642	Provasoli, L. 1968. Media and prospects for the cultivation of marine algae, In:
643	Watanabe A, Hattori A (eds) Cultures and collections of algae. Proc US Japan
644	Conf Hakone, Jap. Soc. Plant Physiol. 63-75.
645	Quatrano, R. S. 1968. Rhizoid formation in Fucus zygotes: dependence on protein and
646	ribonucleic acid syntheses. Science. 162:468-70.
647	Quatrano, R. S. & Crayton, M. A. 1973. Sulfation of fucoidan in Fucus embryos: I.
648	Possible role in localization. Dev. Biol. 30:29-41.
649	Quatrano, R. S. & Stevens, P. T. 1976. Cell wall assembly in Fucus zygotes: I.
650	Characterization of the polysaccharide components. Plant Physiol. 58:224-31.
651	R Core Team. 2020. R: A language and environment for statistical computing. Version
652	4. 0. 2. R Foundation for Statistical Computing, Vienna, Austria
653	Rabillé, H., Torode, T. A., Tesson, B., Bail, A. L., Billoud, B., Rolland, E., Panse, S. L.,
654	Jam, M. & Charrier, B. 2019. Alginates along the filament of the brown alga
655	Ectocarpus help cells cope with stress. Sci. Rep. 9:12956.
656	Reynolds, E. S. 1963. The use of lead citrate at high pH as an electron-opaque stain in
657	electron microscopy. J. Cell Biol. 17:208-12.
658	Salgado, L. T., Cinelli, L. P., Viana, N. B., Tomazetto de Carvalho, R., de Souza
659	Mourao, P. A., Teixeira, V. L., Farina, M. & Filho, A. G. M. A. 2009. A
660	vanadium bromoperoxidase catalyzes the formation of high-molecular-weight
661	complexes between brown algal phenolic substances and alginates. J. Phycol.
662	45:193-202.

663	Skriptsova, A. V., Shevchenko, N. M., Tarbeeva, D. V. & Zvyagintseva, T. N. 2012.		
664	Comparative study of polysaccharides from reproductive and sterile tissues of		
665	five brown seaweeds. Mar. biotechnol. 14:304-11.		
666	Sun, J., Nishiyama, T., Shimizu, K. & Kadota, K. 2013. TCC: An R package for		
667	comparing tag count data with robust normalization strategies. BMC		
668	Bioinformatics. 14:219.		
669	Terauchi, M., Nagasato, C., Inoue, A., Ito, T. & Motomura, T. 2016. Distribution of		
670	alginate and cellulose and regulatory role of calcium in the cell wall of the		
671	brown alga Ectocarpus siliculosus. Planta. 244:361-77.		
672	Terauchi, M., Nagasato, C., Kajimura, N., Mineyuki, Y., Okuda, K., Katsaros, C. &		
673	Motomura, T. 2012. Ultrastructural study of plasmodesmata in the brown alga		
674	Dictyota dichotoma (Dictyotales, Phaeophyceae). Planta. 236:1013-26.		
675	Tonon, T., Rousvoal, S., Roeder, V. & Boyen, C. 2008. Expression Profiling of the		
676	Mannuronan C5-Epimerase multigenic family in the brown alga Laminaria		
677	Digitata (phaeophyceae) under biotic stress conditions. J. Phycol. 44:1250-6.		
678	Torode, T. A., Siméon, A., Marcus, S. E., Jam, M., Le Moigne, M. A., Duffieux, D.,		
679	Knox, J. P. & Hervé, C. 2016. Dynamics of cell wall assembly during early		
680	embryogenesis in the brown alga Fucus. J. Exp. Bot. 67:6089-100.		
681	Vreugdenhil, D., Dijkstra, M. L. & Libbenga, K. R. 1976. The ultrastructure of the cell		
682	wall of normal and apolar embryos of Fucus vesiculosus. Protoplasma.		
683	88:305-13.		
684	Willats, W. G. T., Orfila, C., Limberg, G., Buchholt, H. C., van Alebeek, G. J.,		
685	Voragen, A. G. J., Marcus, S. E., Christensen, T. M., Mikkelsen, J. D., Murray,		
686	B. S. & Knox, J. P. 2001. Modulation of the degree and pattern of methyl-		

687	esterification of pectic homogalacturonan in plant cell walls: implications for
688	pectin methyl esterase action, matrix properties, and cell adhesion. J. Biol.
689	Chem. 276:19404-13.
690	Wu, X., Wang, G. & Fu, X. 2014. Variations in the chemical composition of Costaria
691	costata during harvest. J. Appl. Phycol. 26:2389-96.
692	Ye, N., Zhang, X., Miao, M., Fan, X., Zheng, Y., Xu, D., Wang, J., Zhou, L., Wang, D.,
693	Gao, Y., Wang, Y., Shi, W., Ji, P., Li, D., Guan, Z., Shao, C., Zhuang, Z., Gao,
694	Z., Qi, J. & Zhao, F. 2015. Saccharina genomes provide novel insight into kelp
695	biology. Nat. commun. 6:1-11.
696	
697	
698	

700	Figure 1	Legends
	0	0

701 Fig. 1. Silvetia zygote development. (a-d) 30 min AF, 12 h AF, 20 h AF, 24 h AF, 702 respectively. Arrows and arrowheads indicate the first and second cytokinetic planes, 703 respectively. (e) Illustration indicating three cell wall regions observed by TEM; the 704 thallus cell wall (diagonal dot line area), the rhizoid-flank cell wall (striped area), and 705 the rhizoid-tip cell wall (plane area). Scale bars=50 µm (a-d). 706 707 Fig. 2. Ultrastructure of (a) TCW, (b) RFCW, and (c) RTCW in zygotes at 12 h AF. 708 Arrows show the outer edge of cell wall. Arrowheads show plasma membrane. IL; the 709 inner layer, ML; the middle layer, OL; the outer layer. Scale bars=500 nm. 710 711 Fig. 3. Zygotes at 24 h AF. (a) View of the Silvetia rhizoid. A double arrowhead indicates the point from which the two outer layers start to be reduced. (b) Oblique 712 713 section of TCW. The cell wall consists of three layers. (c) Perpendicular section of 714 TCW. (d) RFCW. The outer layer is disappearing. (e) RFCW. The middle layer fades 715 away. (f) RTCW has only one layer, which is made of rough fibrils. Thick mucilage 716 covers the rhizoid. Arrows indicate the outer edge of cell wall. Arrowheads show 717 plasma membrane. IL; the inner layer, ML; the middle layer, OL; the outer layer. Scale 718 bars=10 µm (a), 500 nm (b-f). 719 720 Fig. 4. Immunogold localization of alginate within cell wall in the zygotes at 24 h AF.

721 (a), (b), (c) show BAM6 localization. (d), (e), (f) indicate BAM7 localization. (a) (d)

TCW, (b) (e) RFCW, (c) (f) RTCW. Arrows show the outer edge of cell wall. Insets

- show the magnified images of dashed line rectangles. Arrowheads show plasma
- 724 membrane. Scale bars=500 nm (a-f), 100 nm (insets).
- 725
- Fig. 5. The distribution of the relative density of relative distance of gold particles
- 727 binding to BAM6 and BAM7 in TCW and RTCW at 24 h-old zygotes. The relative
- distance between plasma membrane and the cell-wall edge ranges from 0 to 1. The sum
- 729 of density is adjusted to 10. Grey lines show the function obtained by kernel function
- and bandwidth.
- 731
- 732 Fig. 6. Quantitative analysis of putative *Mannuronan C5-epimerase* genes transcript
- rain level in 3, 10, and 24 hour-old zygotes. (a) comp32408\_c2\_seq24.p1, (b)
- comp33109\_c0\_seq1.p1. The relative gene expression was set at one at 3 h AF. The
- ror bars indicate SDs of three separate experiments. Tukey's multiple comparison tests
- 736 were conducted to investigate significant differences between the treatment hours.
- 737 Asterisks (\*) indicate significant differences at P < 0.01.
- 738

Fig. S1. A diagram showing the measurement area of gold particles bound to antibodies

- on the cell wall. Measurements are made within three replicate rectangle regions of
- 741 TCW, RFCW, and RTCW of the cell wall. Each rectangle stretches from the plasma
- membrane to the outer edge of the cell wall and has 0.75 µm wide sides over the plasma
- 743 membrane and at the surface of the cell wall (Fig. S1). Three different zygotes are
- 744 measured (n=9).
- 745
- 746

749 second cytokinetic planes, respectively. 750 751 Fig. S3. MA plots (log expression ratio vs. mean average expression) comparing gene 752 expression during zygote germination. Differential expression genes were shown as 753 magenta points (TCC, FDR < 0.05). (a) 3 h AF vs. 10 h AF, (b) 3 h AF vs. 24 h AF, (c) 754 10 h AF vs. 24 h AF. 755 756 Fig. S4. Schematic representation of the alginic acid biosynthetic pathways with 757 expression levels of putative enzymatic genes during development of the zygotes. The 758 gene expression level was shown as TPM and Z-score. MPI: mannose-6-phosphate 759 isomerase, PMM: phosphomannomutase, GMD: GDP-mannose 6-dehydrogenase, 760 MC5E: mannuronan C5-epimerase. Asterisks indicate differential expression genes. 761 762 Fig. S5. Phylogenetic and putative protein structural analysis of mannuronan C5-763 epimerases in S. babingtonii. Numbers indicate the bootstrap values in the maximum 764 likelihood analysis. Asterisks indicate differential expression genes. Each shape on the 765 schematic protein sequences indicates the catalytic MC5E domain (purple box), a 766 transmembrane domain (cyan oval), a signal peptide region (orange box), and the WSC 767 domains (green box). 768 769

Fig. S2. Three different developmental stages of the zygotes for transcriptomic analysis.

(a) 3 h AF, (b) 10 h AF, (c) 24 h AF. Arrows and arrowheads indicate the first and

747

748