



Title	Secular trends of urinary phthalate metabolites in 7-year old children and association with building characteristics : Hokkaido study on environment and children's health
Author(s)	Ketema, Rahel Mesfin; Bamai, Yu Ait; Ikeda-Araki, Atsuko; Saito, Takeshi; Kishi, Reiko
Citation	International journal of hygiene and environmental health, 234, 113724 https://doi.org/10.1016/j.ijheh.2021.113724
Issue Date	2021-05
Doc URL	http://hdl.handle.net/2115/85207
Rights	© 2021. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	ijheh 234 113724.pdf



[Instructions for use](#)

1 **Secular trends of urinary phthalate metabolites in 7-year old children and association**
2 **with building characteristics: Hokkaido study on environment and children's health**

3

4 Rahel Mesfin Ketema, Yu Ait Bamai, Atsuko Araki, Takeshi Saito, Reiko Kishi

5

6 ¹ Center for Environmental and Health Sciences, Hokkaido University, Sapporo, Japan

7 ² Graduate School of Health Sciences, Hokkaido University, Sapporo, Japan

8

9

10

11 * Corresponding author: Reiko Kishi, MD, PhD, MPH

12 Hokkaido University Center for Environmental and Health Sciences

13 Kita 12, Nishi 7, Kita ku, Sapporo 060-0812, Japan

14 Phone: +81-11-706-4746

15 Email: rkishi@med.hokudai.ac.jp

16 **Abstract**

17 The widespread commercial production and use of phthalates as plasticizers in
18 consumer products have led to significant human exposure. Some phthalates are known to
19 disrupt the endocrine system and result in adverse health outcomes. As such, they have been
20 regulated in materials used for children’s items and food packages. In this study, we
21 examined the secular trend of urinary phthalate metabolites in children and the association
22 between metabolites and building characteristics. In total, 400 first-morning spot urine
23 samples of 7 years old children collected from 2012 to 2017 from an ongoing birth cohort
24 study were examined. Parents provided information on demographics and building
25 questionnaires. We analyzed 10 urinary phthalate metabolites from five phthalate diesters
26 using ultra-performance liquid chromatography tandem mass spectrometry (UPLC-MS/MS):
27 MiBP, MnBP, MBzP, MEHP, MEOHP, MEHHP, MECPP, MiNP, OH-MiNP, and cx-MiNP.
28 A multivariable regression model with creatinine-corrected metabolite levels was applied to
29 assess secular trends during 2012–2017. The association between metabolite levels and
30 building characteristics was investigated using a mutual-adjusted linear regression.

31 The metabolites MnBP, MEHP, MEOHP, MEHHP, MECPP, and OH-MiNP were
32 detected in all samples. The highest median concentration was for MECPP 37.4 ng/mL,
33 followed by MnBP and MEHHP at concentrations of 36.8 and 25.8 ng/mL, respectively.
34 Overall, DBP, BBzP, and DINP metabolite concentrations in this study were comparable to
35 or lower than those in previous studies from Japan and other countries in a similar study
36 period. Higher concentrations of DEHP metabolites were observed in this study than in

37 children from the USA and Germany, as per previous reports. Despite updated phthalate
38 regulations and reports of production volume change in Japan, all the measured metabolites
39 showed a stable trend between 2012 and 2017. Higher phthalate metabolite levels were
40 observed among children from households with low annual income, those who lived in old
41 buildings, and those with window opening habits of ≥ 1 h than ≤ 1 h. In contrast, children in
42 houses that vacuumed 4 or more days/week showed a lower level of MnBP than those in
43 houses that vacuumed ≤ 3 days/week.

44 This study demonstrates that the internal exposure level of phthalates in Japanese
45 children was stable from 2012 to 2017. Our findings suggest that phthalate exposure in
46 children is consistent. Thus, improvements in the indoor environment, such as frequent
47 vacuuming, may reduce exposure. Biomonitoring of phthalates is critical for elucidating their
48 possible health effects and developing mitigation strategies.

49

50

51

52 **Keywords**

53 Urinary phthalate metabolites, Secular trend, Children, Human biomonitoring, Building
54 characteristics

55 **1. Introduction¹**

56 Phthalates or phthalic acid esters are a group of synthetic chemicals with the chemical
57 structure of dialkyl or alkyl aryl esters of 1,2-benzene dicarboxylic acid (Cao et al., 2010).
58 Phthalates are widely used as plasticizers, solvents, and additives in products such as
59 polyvinyl chloride (PVC) materials, children's toys, food packaging, pharmaceuticals, and
60 personal care products (Shinohara et al., 2020; Wang et al., 2019; Hauser and Calafat, 2005).
61 The increased use of phthalates in several products results in its ubiquitous presence in the
62 environment and exposure to the general population through inhalation, ingestion, or dermal
63 contact (Anderson et al., 2018; Hauser and Calafat, 2005). Phthalate exposure has been
64 reported to have endocrine-disrupting effects in humans (Hauser and Calafat, 2005; WHO,
65 2012) and experimental studies (Lyche et al., 2009).

66 Owing to the reproductive toxicity of phthalates, the United States and European
67 government regulations were enacted to ban or restrict the use of phthalates, such as di(2-

¹ **Abbreviations:** ultra-performance liquid chromatography tandem mass spectrometry (UPLC-MS/MS), di(2-ethylhexyl) phthalate (DEHP), dibutyl phthalate (DBP), butyl benzyl phthalate (BBzP), Ministry of Health, Labor and Welfare (MHLW), di-iso-decyl phthalate (DiDP), di-isononyl-phthalate (DINP), di-octyl phthalate (DnOP), mono-n-butyl phthalate (MnBP), mono-isobutyl phthalate (MiBP), mono-benzyl phthalate (MBzP), mono (2-ethylhexyl) phthalate (MEHP), mono (2-ethyl-5-hydroxyhexyl) phthalate (MEHHP), mono (2-ethyl-5-oxohexyl) phthalate (MEOHP), mono (2-ethyl-5-carboxypentyl) phthalate (MECPP), mono-isononyl phthalate (MiNP), mono-hydroxy-isononyl phthalate (OH-MiNP), mono-carboxy-isononyl phthalate (cx-MiNP), UPLC charged surface hybrid (CSH), multiple-reaction monitoring (MRM), German external quality assessment scheme (G-EQUAS), limits of detection (LOD), limits of quantification (LOQ), standard deviation (SD), creatinine excretion (CE)

68 ethylhexyl) phthalate (DEHP), dibutyl phthalate (DBP), and butyl benzyl phthalate (BBzP),
69 in the production of items associated with children and cosmetics (DIRECTIVE 2005/84/EC,
70 2005; Public Law 110-314 H.R. 4040, 2008). Such government regulations aim to change
71 the production and use patterns of phthalates. Trend analysis studies in the general population
72 have been used to monitor changes in exposure, such as in Denmark, Germany, Italy, and the
73 US (Frederiksen et al., 2020; Koch et al., 2017; Tranfo et al., 2018; Zota et al., 2016). For
74 instance, a study from the US using the National Health Nutrition and Examination Survey
75 (NHANES) from 2001 to 2010 reported a decrease in DnBP, BBzP, and DEHP metabolite
76 concentrations and increased levels of DINP in the general population of children and adults
77 (CDC, 2019; Zota et al., 2014). Similar studies in Europe have also reported a decline in
78 urinary metabolites of DBP, BBzP, and DEHP in adults (Frederiksen et al., 2020; Tranfo et
79 al., 2018). Additionally, a biomonitoring study conducted in Germany reported a decline in
80 DBP, DEHP, and DINP metabolites in samples collected between 1988 and 2005, mainly
81 from students aged 20–29 years (Koch et al., 2017). Government regulations in the US and
82 European countries have been considered effective, as decreased phthalate exposure has been
83 attributed to them (Tranfo et al., 2018; Zota et al., 2014). In 2010, the Ministry of Health,
84 Labor and Welfare (MHLW) in Japan updated restrictions on the use of DBP, BBzP, DEHP,
85 di-iso-decyl phthalate (DiDP), di-isononyl-phthalate (DINP), and di-octyl phthalate (DnOP)
86 in children’s toys and food packaging materials (MHLW Notice No.370, 2010).

87 Our previous studies have shown higher urinary phthalate metabolite concentrations in
88 children than in adolescents or adults (Ait Bamai et al., 2015). Moreover, urinary phthalate
89 metabolite levels were positively correlated with phthalate concentrations in house dust

90 (AitBamai et al., 2016). Children are more vulnerable to the adverse effects of phthalate
91 exposure on asthma and allergies than adults (Ait Bamai et al., 2014). Previous studies have
92 indicated the importance of building characteristics and indoor environments on phthalate
93 exposure (Ait Bamai et al., 2014; Hsu et al., 2017). Moreover, to the best of our knowledge,
94 no prior study has been conducted on the biomonitoring of trends in consecutive years of
95 phthalate exposure in the Japanese population. Therefore, we aimed to investigate the secular
96 trend of phthalate exposure in Japanese children between 2012 and 2017 and examined the
97 association between internal phthalate exposure levels and the building characteristics of
98 their homes. Additionally, we estimated the daily intake of phthalates based on metabolite
99 levels in the urine.

100 **2. Materials and methods**

101 **2.1 Study population and data collection**

102 This study is part of the ongoing birth cohort study, Hokkaido Study of Environment
103 and Children's Health, Hokkaido Cohort. A detailed description of subject recruitment has
104 been previously described (Kishi et al., 2017, 2011). A total of 20,926 participants were
105 enrolled from February 2003 to March 2012. After accounting for the exclusion criteria of
106 spontaneous abortion, stillbirth, loss to follow-up, and withdrawal, 10,655 singleton children
107 aged 7 years until August 2017 were included. Since 2012, a follow-up questionnaire was
108 sent to these 10,655 children in their birth months, and parents of the 6,218 children returned
109 the questionnaire before September 2017 (response rate 58.4%). Among them, 2,451 children
110 provided urine samples and submitted questionnaires on demographic and building

111 characteristics. The participants were chosen based on a case-cohort study of wheeze, eczema,
112 and rhinoconjunctivitis. The sample size (n) was 100 for each symptom. A sub-cohort of 243
113 participants (11.1% of the original cohort) was randomly selected. Consequently, 83 cases
114 and 160 controls were included in this study. To make up the required 100 cases for each
115 symptom, participants with symptoms were randomly added from the original cohort with
116 urine and questionnaire data. After selecting 100 participants for each symptom, 60 children
117 had more than two symptoms. Finally, a total of 400 participants, 240 with symptoms and
118 160 without symptoms, were selected. Fourteen samples with insufficient urine volume were
119 excluded, leaving only 386 participants for this study (Table 1). Between 2012 and 2017, the
120 number of urine samples collected was randomly distributed as 62, 65, 54, 74, 86, and 45,
121 respectively. The selection details of the 400 participants were described in a previous report
122 (Ait Bamai et al., 2019).

123 The building questionnaire determined building age, annual household income,
124 number of residents, housing type (single-family house/multi-family house), structure
125 (wood/concrete), newly built or renovated within 1 year (yes/no), ventilation in living and/or
126 child room(s) (yes/no), condensation (yes/no), mold odor (yes/no), visible mold (yes/no),
127 water leakage (yes/no), humidity (yes/no), insecticide (yes/no), flooring (PVC/non-PVC),
128 wall material (PVC/non-PVC), vacuum-cleaning/week, duration of the window opening, and
129 whether the house was on the main road (yes/no).

130 Parents of children were asked to collect the first morning void urine samples of their
131 children in a polypropylene cup, and these were sent to Hokkaido University, Center for

132 Environmental and Health Sciences, using a cool delivery service. When the shipped urine
133 samples arrived at our center, the creatinine content was measured using an enzyme-linked
134 immunosorbent assay at SRL, Inc. (Tokyo, Japan). On the same day, samples were
135 transferred to glass test tubes with ground glass stoppers cleaned with acetone, sealed with
136 fluoroc tape, wrapped with aluminum foil, and kept at $-20\text{ }^{\circ}\text{C}$ until the day of analysis.

137 The research protocol regarding human sampling was reviewed and approved by the
138 Institutional Review Board of the Hokkaido University Center for Environmental and Health
139 Sciences before the study was conducted. The parents of all participants provided written
140 informed consent to confirm their participation in this study.

141 **2.2 Urinary chemical analysis**

142 Ten phthalate metabolites were measured in the first morning void urine samples of
143 children. The phthalate metabolites assessed included the DBP metabolites [mono-n-butyl
144 phthalate (MnBP), mono-isobutyl phthalate (MiBP)], BBzP metabolite [mono-benzyl
145 phthalate (MBzP)], DEHP metabolites [mono (2-ethylhexyl) phthalate (MEHP), mono (2-
146 ethyl-5-hydroxyhexyl) phthalate (MEHHP), mono (2-ethyl-5-oxohexyl) phthalate (MEOHP),
147 and mono (2-ethyl-5-carboxypentyl) phthalate (MECPP)], and the DINP metabolites [mono-
148 isononyl phthalate (MiNP), mono-hydroxy-isononyl phthalate (OH-MiNP), and mono-
149 carboxy-isononyl phthalate (cx-MiNP)]. For sample preparation, 500 μL urine sample was
150 spiked with 20 μL of a mixture of labeled internal standards and then buffered with 500 μL
151 100 mM ammonium acetate (pH 6.5), and deconjugated by 50 μL β -glucuronidase with
152 incubation at $37\text{ }^{\circ}\text{C}$ for 90 min. After incubation, 1 mL 0.5% ammonia water was added to

153 each sample. After this, sample extraction was performed using solid-phase extraction (SPE)
154 that was conditioned with 1 mL 0.05% nitric acid in 90% methanol, 1 mL methanol, and then
155 with 1 mL 0.5% ammonia water to activate the cartridge. Samples were loaded onto the
156 conditioned SPE cartridge and sequentially washed with 0.5 mL ultra-pure water, 0.5 mL
157 methanol, 0.5 mL ultra-pure water, and 0.5 mL 40% methanol containing 0.2% formic acid.
158 The samples were eluted using a mixture of 90% methanol containing 0.2% formic acid. The
159 eluted mixture (250 μ L) was transferred to a vial and diluted with 750 μ L ultrapure water. To
160 quantify phthalate metabolites, 40 μ L of the sample from the vial was injected into a UPLC-
161 MS/MS (ACQUITY UPLC H-class) equipped with a Xevo TQ-S micro mass spectrometer
162 (Waters Corporation, Milford, MA, USA). Detailed information including chemicals and
163 reagents, urine sample collection, sample preparation, instrumental analysis, and
164 chromatographic conditions can be found in the supporting information (Supplemental
165 Figure 1 and Supplemental Tables 1–3).

166 **2.3 Quality assurance**

167 For each batch, two procedural blanks were analyzed to control for background
168 contamination. For all target analytes, 12 calibration points ranging from 0 to 20 ng/mL were
169 used to construct the calibration curves. A satisfactory correlation coefficient of calibration
170 curves ≥ 0.998 was obtained for all measured metabolites. In each batch of 20 samples,
171 replicated analysis of the calibration standard at a concentration of 5 ng/mL and the reference
172 value of 63 G-EQUAS samples with known concentrations were conducted to determine both
173 inter-and intra-day precision and were within acceptable limits with a coefficient of variation

174 < 10%. The limits of detection (LOD) and limits of quantification (LOQ) of individual
175 phthalate metabolites were determined based on 7 repeated analyses of spiked ultra-pure
176 water with 0.16 ng/mL for MEOHP, OH-MiNP, and cx-MiNP; 0.32 ng/mL for MEHP, MiNP,
177 MEHHP, and MECPP; 0.8 ng/mL for MBzP; and 1.6 ng/mL for MiBP and MnBP. The
178 standard deviation (SD) of these repeated analyses was calculated using the following
179 formula: $LOD = 2 \times t(n-1, 0.05) \times SD$ and $LOQ = 10 \times SD$. Here, t is the student's t -value
180 for the 95th percentile of $n-1$ degree of freedom, where n is the number of repeated samples.
181 The metabolites LOD and LOQ ranged 0.05–0.95 ng/mL and 0.13–2.5 ng/mL, respectively.
182 The recovery percentages of native and labeled internal standards spiked in pooled urine
183 samples ranged from 81% to 120%. The detailed quality assurance (QA)/quality control (QC)
184 results are shown in Supplemental Table 4.

185

186 **2.4 Daily intake estimation**

187 Based on the measured concentrations of urinary phthalate metabolites, the daily
188 phthalate intake was estimated for each subject using the following equation (Wittassek et
189 al., 2007a).

$$190 \quad EDI = \frac{UE_{sum} \times CE_{smoothed}}{F_{UE} \times BW} \times MW_p$$

191 where EDI ($\mu\text{g}/\text{kg bw}/\text{day}$) is the estimated daily intake of phthalate; UE_{sum} is the sum of
192 the creatine-adjusted molar concentration of phthalate metabolites ($\mu\text{mol}/\text{g Cr}$); $CE_{smoothed}$
193 (g/day) is the smoothed creatinine excretion (CE) rate; BW (kg) is the bodyweight; MW_p

194 (g/mol) is the molecular weight of the respective parent phthalate; and F_{ue} is the urinary
195 excretion factor of the parent phthalate, DnBP, DiBP, BBzP, DEHP, and DINP, which were
196 set to 0.69, 0.69, 0.73, 0.62, and 0.39, respectively (Anderson et al., 2001; Koch et al., 2012;
197 Wittassek et al., 2011). Gender-based values for urinary CE were determined using the
198 following equations, where ht (cm) is the participant's height (Mage et al., 2008).

$$199 \quad CE = ht \times \{6.265 + 0.0564 \times (ht - 168)\} \times 10^3 \text{ (male)}$$

$$200 \quad CE = 2.045 \times ht \times \exp \{0.01552 \times (ht - 90)\} \times 10^3 \text{ (female)}$$

201

202 **2.5 Statistical analysis**

203 To ensure that the study population ($n = 386$) was representative of the original cohort,
204 we conducted sensitivity analyses with a sub-cohort that included 243 participants. The
205 results showed similarity in the distribution of demographic characteristics, building
206 characteristics, levels of phthalate metabolites, and their trends. Thus, we can anticipate
207 minimal probable bias for our study population ($n = 386$) (Supplemental Table 5 and 6; and
208 Supplemental Figure 2). For concentrations below the LOD, the LOD \times detection frequency
209 was assigned for statistical analysis (James et al., 2002). Additionally, the molar
210 concentrations of DBP metabolites (MiBP and MnBP), DEHP metabolites (MEHP, MECPP,
211 MEOHP, and MEHHP), and DINP metabolites (MiNP, OH-MiNP, and cx-MiNP) were
212 combined to estimate the parent compound exposure. The distribution of phthalate
213 metabolites is presented as minimum, percentiles (25, 50, 75, and 95), and maximum values.
214 A regression model with creatinine-corrected metabolite concentrations was applied to

215 assess the secular trend from 2012 to 2017. Additionally, Dunnett’s test, considering 2012
216 as a reference, was conducted to compare pairwise metabolite level mean differences by
217 year. The *p*-values were adjusted using the Bonferroni correction. We first conducted a
218 univariate analysis to analyze the distribution of urinary phthalate metabolite levels
219 according to different building characteristics. We then investigated associations between
220 phthalate metabolite concentrations and the building characteristics that showed a
221 significant difference in univariate analysis using mutual-adjusted linear regression. We
222 calculated the percent difference from the regression coefficient as $(e^{(\beta)} - 1) \times 100\%$ with
223 95% CIs estimated as $(e^{(\beta \pm \text{critical value} \times \text{SE})} - 1)$, where β and SE are the estimated regression
224 coefficient and standard error, respectively. Yes indicators regarding dampness;
225 condensation, mold odor, visible mold, water leakage, and humidity were assigned a value
226 of 1 to compute the dampness index (1–5). Statistical analysis was performed using JMP
227 Clinical 6.0, SAS.

228 **3. Results and discussion**

229 In this study, we presented the levels and secular trends of phthalate metabolites in
230 Japanese children from Hokkaido between 2012 and 2017. Moreover, we investigated the
231 association between building characteristics and levels of urinary phthalate metabolites and
232 estimated daily phthalate exposure in children based on urinary metabolite levels.

233 **3.1 Study population**

234 Due to insufficient sample volume or sample preparation error, 14 samples were
235 excluded, and 386 samples were included in this study. All children in this study were seven

236 years old; the gender participation was nearly balanced with males (52.6%) and girls (47.4%).
237 The data represent the phthalate exposure trend of six consecutive years as children's urine
238 samples were collected each year from 2012 to 2017. Nearly 60% of the participants lived in
239 a mechanically ventilated house. Compared to our previous study, more participants in this
240 study lived in houses with PVC flooring (16.9% vs. 7%) and slightly older buildings (median:
241 13 years vs. 10.5 years) (Kishi et al., 2018). The participants' demographic and building
242 characteristics with urine collection years are presented in Table 1.

243 **3.2 Concentration and secular trend of urinary phthalate metabolites**

244 The distribution of urinary phthalate metabolite concentrations along with creatinine
245 (Cr)-corrected levels in children is summarized in Table 2. Phthalate metabolites MnBP,
246 MEHP, MEOHP, MEHHP, MECPP, and OH-MiNP were detected in all samples. The
247 highest concentration was found among the DEHP metabolites MECPP and MEHHP,
248 followed by MnBP. The creatinine-corrected concentrations showed a similar trend. All
249 creatinine-corrected urinary phthalate metabolites in this study showed a significant positive
250 Spearman's correlation. The highest correlation was found between DEHP and DINP
251 metabolites (Supplemental Table 7). Although many studies have reported seasonal
252 variations in phthalate exposure (Bi et al., 2018; Li et al., 2019), in this study, no association
253 was observed between the sample collection seasons and phthalate metabolite levels
254 (Supplemental Table 8). The secular trend of the evaluated urinary phthalate metabolite
255 levels is shown in Figure 1. Regression analysis showed that all measured metabolites were
256 stable throughout the study period. To the best of our knowledge, this is the first human

257 biomonitoring study to investigate internal phthalate exposure trends in the Japanese
258 population. In 2010, the regulation of phthalates in children's toys and food packaging
259 materials was revised in Japan. Consequently, changes in the production and exposure to
260 phthalates have been reported in Japan, for example, from 2012 to 2017, the production of
261 DEHP decreased by 13.3% (135,000 to 117,000 t). In contrast, the production of DINP
262 increased by 43.2% (67,000 to 96,000 t) (IHS Markit, 2018; VEC, 2018). However, urinary
263 phthalate metabolites showed a stable trend. This suggests that the reported changes in
264 chemical production do not reflect children's exposure. A plausible explanation for the lack
265 of a trend in the current study could be the limited scope of phthalate regulation. The
266 regulation only concerns toys meant for children under 6 years of age and food containers
267 containing fats and oils but excludes materials such as PVC flooring, wall and ceiling
268 coverings (common in modern Japanese houses and apartments), and personal care products,
269 which have been reported as potential phthalate exposure sources (Bornehag et al., 2005;
270 Carlstedt et al., 2013; Husoy et al., 2020). Higher levels of DEHP in indoor dust in Japanese
271 households than in other EU countries and the USA have been reported (Ait Bamai et al.,
272 2014). Thus, the stable trend in exposure levels observed in this study can be attributed to
273 phthalates emitted from non-regulated materials (Ait Bamai et al., 2014; Carlstedt et al.,
274 2013; Husoy et al., 2020).

275 Comparing the results of our current and previous studies (Ait Bamai et al., 2015) in
276 Japanese children, we observed a relatively higher median concentration for MiBP, MBzP,
277 MEHP, MEOHP, and lower MnBP in the previous study than in the current study (Figure 2,
278 Supplemental Table 9). However, this comparison should be interpreted with caution because

279 the method of analysis was different for the two studies. The previous study used
280 derivatization and was measured by GC-MS, while the current study used LC-MS/MS. The
281 concentrations of urinary phthalate metabolites in children among comparable age groups
282 show variations similar to those in the findings from other countries over a similar period
283 (Figure 2, Supplemental Table 9) (Ait Bamai et al., 2015; Becker et al., 2009; CDC, 2019;
284 Hartmann et al., 2015; Liao et al., 2018; Schwedler et al., 2020; Song et al., 2013; Wang et
285 al., 2015; Weng et al., 2017). For instance, a similar level of MiBP was observed in children
286 from the US (CDC, 2019), while a 2- to 3-fold higher median concentration was observed in
287 Germany (Schwedler et al., 2020) and China (Liao et al., 2018; Wang et al., 2015). A
288 noticeably higher level of DEHP metabolites was observed in our study participants than in
289 Germany (Schwedler et al., 2020) and the USA (CDC, 2019) (Figure 2). This indicates that
290 Japanese children still have high exposure to DEHP despite production regulations and
291 efforts to replace DEHP (Rowdhwal and Chen, 2018; VEC, 2018). We previously reported
292 that DEHP and BBzP concentrations in house dust are positively correlated with urinary
293 metabolite concentrations (Bamai, 2016). Additionally, a review revealed that house dust
294 ingestion is a significant exposure pathway for phthalates such as DEHP in Japan (Takagi
295 and Yoshinaga, 2009). Thus, the high levels of urinary DEHP metabolites in this study might
296 be due to non-regulated consumer products, such as PVC building materials, which can
297 release phthalates.

298 Considering the DINP in this study, consistent trends in its metabolites were observed.
299 This might be due to the wide use of DINP in PVC wallpapers, wire, and cable insulation
300 jacketing in Japan, which are less likely to be changed/installed frequently (IHS Markit,

2018). Moreover, most of the children in this study lived in houses with a mean age of 13 years (Table 1), which was built before the regulation and increased production of DINP, which could explain the stable trend observed in this study. Biomonitoring studies conducted in the early 2010s reported increased levels of DINP in Germany, Italy, and the USA (Tranfo et al., 2018; Wittassek et al., 2007; Zota et al., 2016). DINP has been subjected to regulation due to its various health risks and is substituted with alternatives such as DEHTP and DINCH, resulting in a decline in DINP exposure in recent years (CDC 2019, Frederiksen et al., 2020, Schwedler et al., 2020). Since the current study did not measure urinary DINP metabolites before the 2010 revised phthalate regulations, exposure levels of DINP in Japanese children before the regulation are uncertain. This warrants follow-up studies with a large population size to elucidate exposure changes over time.

3.3 Urinary phthalate metabolite levels and building characteristics

Despite the government regulations regarding phthalates in Japan, our trend analysis showed a stable level of phthalate metabolites. Additionally, the high detection frequency of urinary phthalate metabolites indicates that children are still widely exposed, probably from non-regulated products such as building materials. Thus, we evaluated potential phthalate exposure based on household characteristics and daily habits (Table 3). Our mutually adjusted regression model revealed a significant positive association between lower household income and OH-MiNP ($\beta = 0.138$) and \sum DINP ($\beta = 0.127$). This result is in agreement with those of previous studies that reported that lower socioeconomic status (SES) families tend to show a higher level of urinary phthalate metabolites, such as MBzP and

322 DEHP metabolites (Koo et al., 2002; Navaranjan et al., 2020). This may have several
323 precipitating factors, such as the influence of SES on dwelling characteristics or fast-food
324 consumption habits resulting from parental education levels, which have previously been
325 reported as variables causing increased phthalate exposure (Ait Bamai et al., 2014; Zota et
326 al., 2016). In the present study, MnBP levels showed a significant positive association with
327 building age. Furthermore, our stratified analysis revealed an increased beta value of MnBP
328 ($\beta > 0$) for older buildings, indicating an increased level of MnBP as the building age
329 increased. Building age has been reported as a common predictor of indoor phthalate levels
330 in the dust (Bornehag et al., 2005). There is evidence that the DnBP parent compound of
331 MnBP was used as a plasticizer in PVC materials in the 1980s (Kavlock et al., 2002). Thus,
332 the use of DnBP in interior materials for older buildings could explain our finding of a
333 significant relationship between MnBP and building age. Metabolite MnBP was lower in
334 children who lived in houses that were vacuumed 4–7 times/week than in those that were
335 vacuumed ≤ 3 days/week. This result can be explained by the frequent vacuuming association
336 with decreasing dust accumulation, resulting in lower phthalate levels (Wilson and VanSnick,
337 2017), as phthalates emitted from vinyl building materials can be absorbed into house dust
338 (Liang and Xu, 2014).

339 Children in houses with ≥ 1 h/day window opening habits showed significantly
340 elevated levels of MiBP, MnBP, and DEHP metabolites compared to children in houses with
341 < 1 h/day window opening habits. This result is unpredicted, as opening windows are
342 expected to facilitate air exchange and decrease phthalate levels in the indoor environment
343 (Śmielowska et al., 2017). A possible reason for this contradictory result could be the

344 confounding effect of building age on the association between window opening and
345 metabolite levels. We observed higher metabolite levels in children living in older buildings
346 (Table 3). Hence, this finding raises the question: “Did the families with fewer open windows
347 live in newer houses with air-conditioning installed?”. Unfortunately, we did not have data
348 on the use/installation of air-conditioning, but in this study area Hokkaido, it was uncommon
349 for households to have air-conditioning due to the climate being cool. Moreover, we did not
350 find a significant difference ($p > 0.786$) in building age between less window opening houses
351 (median 15.9 year) and more window opening houses (median 16.8 year). Therefore,
352 building age was not found to be a confounding factor in the relationship between window
353 opening and metabolites. Another study that examined the impact of open and closed
354 windows on indoor air composition reported that emission of semi-volatile compounds such
355 as phthalates was enhanced when windows were open rather than closed (Fortenberry et al.,
356 2019). Additionally, Xu et al. (2010) revealed that the emission rate of DEHP from vinyl
357 flooring increased at a high ventilation rate because of the higher air velocity near the surfaces
358 and consequently results in an increase in the mass-transfer coefficients that promote the
359 emission of DEHP. This suggests that window opening enhances ventilation and increases
360 emissions to the indoor environment. Subsequently, the internal exposure level of the DEHP
361 increased.

362 **3.4 Estimated daily intake (EDI) of phthalates**

363 Hereafter, we estimated the daily intake of phthalates in children, as shown in Figure 3. Daily
364 intake of DEHP had the highest median EDI value of 3.7 $\mu\text{g}/\text{kg}/\text{day}$. The EDIs of DnBP were

365 slightly higher in boys than girls, with a mean value of 1.8 and 1.4 $\mu\text{g}/\text{kg}/\text{day}$, respectively.
366 Based on the European Food Safety Authority (EFSA), tolerable daily intake (TDI) reference
367 values for individual phthalates DnBP and DEHP, one child in each phthalate exceeded the
368 reference values of 10 and 50 $\mu\text{g}/\text{kg}/\text{day}$, respectively (EFSA, 2005a,2005c). Considering the
369 updated EFSA risk assessment of combined exposure to DBP, BBzP, DEHP, and DINP at a
370 group-TDI level of 50 $\mu\text{g}/\text{kg}/\text{day}$, two children with one child at a marginal level were
371 observed (EFSA 2019). Considering the US reference dose (RfD) of 20 $\mu\text{g}/\text{kg}/\text{day}$ for DEHP
372 (US. EPA., 1991), two children exceeded the RfD value and another 2 were on the reference
373 borderline, representing 1.03 % of the participants (Figure 3). Comparing this study's median
374 EDI values with those of other studies in children, DiBP, DnBP, BBzP, DEHP, and DINP
375 were lower or comparable (Ait Bamai et al., 2015; Kasper-Sonnenberg et al., 2014; Yoshida
376 et al., 2020). In contrast, the EDI value of DINP in this study was higher than that of
377 Taiwanese children with a median of 0.5 and 0.2 $\mu\text{g}/\text{kg}/\text{day}$, respectively (Chang et al., 2017).
378 However, caution should be taken when interpreting phthalate EDI comparisons, since
379 variations in participant characteristics or study methods may alter EDI among different
380 studies and countries.

381 In the future, the use of PVC gloves and rubbing alcohol during the COVID-19 pandemic is
382 likely to increase exposure risk among the general population, which further highlights the
383 need for phthalate biomonitoring.

384

385 **4. Strengths and limitations**

386 Selecting participants of the same age (7 years) in this study allowed for a better comparison
387 by eliminating age as a confounding factor. Additionally, the accuracy of our phthalate
388 metabolite measurement method was validated by the external quality assessment scheme G-
389 EQUAS, which strengthens the reliability of our results (Supplemental Table 10). The
390 building characteristics data in this study also strengthened our investigation by facilitating
391 the identification of possible phthalate exposure sources. The primary limitation of this study
392 is the small sample size. However, this data still provides valuable evidence on changes in
393 phthalate exposure during 2012–2017 in Japanese school-aged children. A secondary
394 limitation is the participants' selection bias of including children with allergies, which could
395 limit the generalizability of this study. However, since the distribution of children with
396 allergies in each year was similar, approximately 16%. Thus, we anticipated that the inclusion
397 of children with allergies would have a minimal effect on any probable bias from our trend
398 analysis. Another limitation is that the urine samples were collected only once, which may
399 not represent variance in urinary phthalate metabolite excretion based on individual activity,
400 diet, personal care product use, cleanliness, and seasonal dust accumulation with an open
401 window. Thus, to reduce the variability of metabolite levels during the day, we used the first
402 morning void urine samples.

403

404 **5. Conclusions**

405 This study is the first to document the consecutive stable trend of the internal exposure level
406 of phthalates in Japanese children between 2012 and 2017, indicating that consistent
407 phthalate exposure exists even after the regulation update of phthalates. Furthermore, we
408 identified correlations between a high exposure level of MnBP among children in old
409 buildings, DINP metabolites among those with lower household income, and MiBP, MnBP,
410 and DEHP metabolites among those with long window opening habits. Frequent vacuum
411 cleaning was associated with lower MnBP levels in children. Finally, other personal care and
412 protective equipment that are known to increase phthalate exposure risk should be evaluated
413 in future studies.

414

415

416

417

418

419

420

421 **Acknowledgments**

422 We are highly indebted to the study participants for taking part in our study.

423 **Funding**

424 This study was financially supported by the Ministry of Health, Labour and Welfare, Japan
425 (19LA1007), Environment Research and Technology Development Fund
426 (JPMEERF20175053), JSPS KAKENHI Grant (16H02645, 15H04780), and Rahel Mesfin
427 Ketema acknowledge that this work was supported by JSPS KEKENHI Grant Number
428 JP20J10766.

429

430 **Declaration of consent**

431 The authors have no conflicts of interest to report.

432

433

434

435

436

437

438 References

- 439 Ait Bamai, Y., Araki, A., Kawai, T., Tsuboi, T., Saito, I., Yoshioka, E., Cong, S., Kishi, R.,
440 2016. Exposure to phthalates in house dust and associated allergies in children aged
441 6–12years. *Environ. Int* 96, 16–23. <https://doi.org/10.1016/j.envint.2016.08.025>
- 442 Ait Bamai, Y., Araki, A., Kawai, T., Tsuboi, T., Saito, I., Yoshioka, E., Kanazawa, A., Tajima,
443 S., Shi, C., Tamakoshi, A., Kishi, R., 2014a. Associations of phthalate concentrations
444 in floor dust and multi-surface dust with the interior materials in Japanese dwellings.
445 *Sci Total Environ* 468–469, 147–157.
446 <https://doi.org/10.1016/j.scitotenv.2013.07.107>
- 447 Ait Bamai, Y., Araki, A., Kawai, T., Tsuboi, T., Yoshioka, E., Kanazawa, A., Cong, S., Kishi,
448 R., 2015. Comparisons of urinary phthalate metabolites and daily phthalate intakes
449 among Japanese families. *Int J Hyg Environ Health* 218, 461–470.
450 <https://doi.org/10.1016/j.ijheh.2015.03.013>
- 451 Ait Bamai, Y., Bastiaensen, M., Araki, A., Goudarzi, H., Konno, S., Ito, S., Miyashita, C.,
452 Yao, Y., Covaci, A., Kishi, R., 2019. Multiple exposures to organophosphate flame
453 retardants alter urinary oxidative stress biomarkers among children: The Hokkaido
454 Study. *Environ Int* 131, 105003. <https://doi.org/10.1016/j.envint.2019.105003>
- 455 Ait Bamai, Y., Shibata, E., Saito, I., Araki, A., Kanazawa, A., Morimoto, K., Nakayama, K.,
456 Tanaka, M., Takigawa, T., Yoshimura, T., Chikara, H., Saijo, Y., Kishi, R., 2014b.
457 Exposure to house dust phthalates in relation to asthma and allergies in both children
458 and adults. *The Science of the total environment* 485-486C, 153–163.
459 <https://doi.org/10.1016/j.scitotenv.2014.03.059>
- 460 Andersen C, Kraus AM., Eriksson AC., Jakobsson J., Löndahl J., Nielsen J., Lindh CH,
461 Pagels J., Gudmundsson A., Wierzbicka A., 2018. Inhalation and Dermal Uptake of
462 Particle and Gas-Phase Phthalates-A Human Exposure Study. *Environ Sci Technol*.
463 2018 Nov 6;52(21):12792-12800. doi: 10.1021/acs.est.8b03761. Epub 2018 Oct 12.
464 PMID: 30264993.
- 465 Anderson, W.A., Castle, L., Scotter, M.J., Massey, R.C., Springall, C., 2001. A biomarker
466 approach to measuring human dietary exposure to certain phthalate diesters. *Food*
467 *Addit Contam* 18, 1068–1074. <https://doi.org/10.1080/02652030110050113>.
- 468 Becker, K., Göen, T., Seiwert, M., Conrad, A., Pick-Fuss, H., Müller, J., Wittassek, M.,
469 Schulz, C., Kolossa-Gehring, M., 2009. GerES IV: phthalate metabolites and
470 bisphenol A in urine of German children. *Int J Hyg Environ Health* 212, 685–692.
471 <https://doi.org/10.1016/j.ijheh.2009.08.002>.

- 472 Bi C, Maestre JP, Li H, Zhang G, Givehchi R, Mahdavi A, Kinney KA, Siegel J, Horner SD,
473 Xu Y. Phthalates and organophosphates in settled dust and HVAC filter dust of U.S.
474 low-income homes: Association with season, building characteristics, and childhood
475 asthma. *Environ Int.* 2018 Dec;121(Pt 1):916-930. doi: 10.1016/j.envint.2018.09.013.
476 Epub 2018 Oct 20. PMID: 30347374.
- 477 Bornehag, C.-G., Lundgren, B., Weschler, C.J., Sigsgaard, T., Hagerhed-Engman, L.,
478 Sundell, J., 2005. Phthalates in indoor dust and their association with building
479 characteristics. *Environ Health Perspect* 113, 1399–1404.
480 <https://doi.org/10.1289/ehp.7809>
- 481 Carlstedt, F., Jönsson, B.A.G., Bornehag, C.-G., 2013. PVC flooring is related to human
482 uptake of phthalates in infants. *Indoor Air* 23, 32–39. [https://doi.org/10.1111/j.1600-](https://doi.org/10.1111/j.1600-0668.2012.00788.x)
483 [0668.2012.00788.x](https://doi.org/10.1111/j.1600-0668.2012.00788.x)
- 484 Cao X.L., 2010. Phthalate esters in foods: sources, occurrence, and analytical methods
485 *Compr Rev Food Sci Food Saf*, 9 (1), pp. 21-43
- 486 CDC, 2019. Fourth National Report on Human Exposure to Environmental Chemicals.
- 487 Chang, J.-W., Lee, C.-C., Pan, W.-H., Chou, W.-C., Huang, H.-B., Chiang, H.-C., Huang,
488 P.-C., 2017. Estimated Daily Intake and Cumulative Risk Assessment of Phthalates
489 in the General Taiwanese after the 2011 DEHP Food Scandal. *Sci Rep* 7, 45009.
490 <https://doi.org/10.1038/srep45009>
- 491 DIRECTIVE 2005/84/EC, 2005.
- 492 Duty, S.M., Ackerman, R.M., Calafat, A.M., Hauser, R., 2005. Personal care product use
493 predicts urinary concentrations of some phthalate monoesters. *Environ Health*
494 *Perspect* 113, 1530–1535. <https://doi.org/10.1289/ehp.8083>
- 495 EFSA (European Food Safety Authority), 2005a. Opinion of the Scientific Panel on Food
496 Additives, Flavourings, Processing Aids and Material in Contact with Food (AFC)
497 on a request from the Commission related to DiButylphthalate (DBP) for use in food
498 contact materials. *EFSA Journal* 2005;3(9):242, 17 pp.
499 <https://doi.org/10.2903/j.efsa.2005.242>.
- 500 EFSA (European Food Safety Authority), 2005b. Opinion of the Scientific Panel on Food
501 Additives, Flavourings, Processing Aids and Materials in Contact with Food (AFC)
502 on a request from the Commission related to Butylbenzylphthalate (BBP) for use in
503 food contact materials. *EFSA Journal* 2005;3(9):241, 14 pp.
504 <https://doi.org/10.2903/j.efsa.2005.241>.
- 505 EFSA (European Food Safety Authority), 2005c. Opinion of the Scientific Panel on Food
506 Additives, Flavourings, Processing Aids and Materials in Contact with Food (AFC)

507 on a request from the Commission related to Bis(2-ethylhexyl) phthalate (DEHP) for
508 use in food contact materials. EFSA Journal 2005;3(9):243, 20 pp. <https://doi.org/10.2903/j.efsa.2005.243>

510 EFSA (European Food Safety Authority), 2005d. Opinion of the Scientific Panel on Food
511 Additives, Flavourings, Processing Aids and Materials in Contact with Food (AFC)
512 on a request from the Commission related to Diisononylphthalate (DINP) for use in
513 food contact materials. EFSA Journal 2005;3(9):244, 18 pp. <https://doi.org/10.2903/j.efsa.2005.244>

515 EFSA: Updated risk assessment of five phthalates., 2019. Accessed on 2020/11/06;
516 [https://www.foodpackagingforum.org/news/efsa-updated-risk-assessment-of-five-](https://www.foodpackagingforum.org/news/efsa-updated-risk-assessment-of-five-phthalates)
517 [phthalates.](https://www.foodpackagingforum.org/news/efsa-updated-risk-assessment-of-five-phthalates)

518 Frederiksen, H., Nielsen, O., Koch, H.M., Skakkebaek, N.E., Juul, A., Jorgensen, N.,
519 Andersson, A.-M., 2020. Changes in urinary excretion of phthalates, phthalate
520 substitutes, bisphenols and other polychlorinated and phenolic substances in young
521 Danish men; 2009-2017. *Int J Hyg Environ Health* 223, 93–105.
522 <https://doi.org/10.1016/j.ijheh.2019.10.002>

523 Fortenberry C, Walker M, Dang A, Loka A, Date G, Cysneiros de Carvalho K, Morrison G,
524 Williams B. Analysis of indoor particles and gases and their evolution with natural
525 ventilation. <https://pubmed.ncbi.nlm.nih.gov/31264732/>. *Indoor Air*. 2019
526 Sep;29(5):761-779. doi: 10.1111/ina.12584. Epub 2019 Aug 1. PMID: 31264732.

527 Hartmann, C., Uhl, M., Weiss, S., Koch, H.M., Scharf, S., König, J., 2015. Human
528 biomonitoring of phthalate exposure in Austrian children and adults and cumulative
529 risk assessment. *Int J Hyg Environ Health* 218, 489–499.
530 <https://doi.org/10.1016/j.ijheh.2015.04.002>

531 Hauser, R., Calafat, A.M., 2005. Phthalates and human health. *Occup Environ Med* 62, 806–
532 818. <https://doi.org/10.1136/oem.2004.017590>

533 Hsu, N.-Y., Liu, Y.-C., Lee, C.-W., Lee, C.-C., Su, H.-J., 2017. Higher moisture content is
534 associated with greater emissions of DEHP from PVC wallpaper. *Environmental*
535 *Research* 152, 1–6. <https://doi.org/10.1016/j.envres.2016.09.027>

536 Husøy T, Martínez MA, Sharma RP, Kumar V, Andreassen M, Sakhi AK, Thomsen C,
537 Dirven H. Comparison of aggregated exposure to di(2-ethylhexyl) phthalate from diet
538 and personal care products with urinary concentrations of metabolites using a PBPK
539 model - Results from the Norwegian biomonitoring study in EuroMix. *Food Chem*
540 *Toxicol.* 2020 Sep;143:111510. doi: 10.1016/j.fct.2020.111510. Epub 2020 Jun 30.
541 PMID: 32615240.

542 IHS Markit, 2018. Chemical Economics Handbook.

- 543 James, R.A., Hertz-Picciotto, I., Willman, E., Keller, J.A., Charles, M.J., 2002. Determinants
544 of serum polychlorinated biphenyls and organochlorine pesticides measured in
545 women from the child health and development study cohort, 1963-1967. *Environ*
546 *Health Perspect* 110, 617–624. <https://doi.org/10.1289/ehp.02110617>
- 547 Kasper-Sonnenberg, M., Koch, H.M., Wittsiepe, J., Brüning, T., Wilhelm, M., 2014.
548 Phthalate metabolites and bisphenol A in urines from German school-aged children:
549 results of the Duisburg birth cohort and Bochum cohort studies. *Int J Hyg Environ*
550 *Health* 217, 830–838. <https://doi.org/10.1016/j.ijheh.2014.06.001>
- 551 Kavlock, R., Boekelheide, K., Chapin, R., Cunningham, M., Faustman, E., Foster, P., Golub,
552 M., Henderson, R., Hinberg, I., Little, R., Seed, J., Shea, K., Tabacova, S., Tyl, R.,
553 Williams, P., Zacharewski, T., 2002. NTP Center for the Evaluation of Risks to
554 Human Reproduction: phthalates expert panel report on the reproductive and
555 developmental toxicity of di-n-butyl phthalate. *Reprod Toxicol* 16, 489–527.
556 [https://doi.org/10.1016/s0890-6238\(02\)00033-3](https://doi.org/10.1016/s0890-6238(02)00033-3)
- 557 Kishi, R., Araki, A., Minatoya, M., Hanaoka, T., Miyashita, C., Itoh, S., Kobayashi, S., Ait
558 Bamai, Y., Yamazaki, K., Miura, R., Tamura, N., Ito, K., Goudarzi, H., 2017. The
559 Hokkaido Birth Cohort Study on Environment and Children’s Health: cohort profile-
560 updated 2017. *Environ Health Prev Med* 22, 46. [https://doi.org/10.1186/s12199-017-](https://doi.org/10.1186/s12199-017-0654-3)
561 [0654-3](https://doi.org/10.1186/s12199-017-0654-3)
- 562 Kishi, R., Ketema, R.M., Ait Bamai, Y., Araki, A., Kawai, T., Tsuboi, T., Saito, I., Yoshioka,
563 E., Saito, T., 2018. Indoor environmental pollutants and their association with sick
564 house syndrome among adults and children in elementary school. *Building and*
565 *Environment* 136, 293–301. <https://doi.org/10.1016/j.buildenv.2018.03.056>
- 566 Kishi, R., Sasaki, S., Yoshioka, E., Yuasa, M., Sata, F., Saijo, Y., Kurahashi, N., Tamaki, J.,
567 Endo, T., Sengoku, K., Nonomura, K., Minakami, H., 2011. Cohort profile: the
568 Hokkaido study on environment and children’s health in Japan. *Int J Epidemiol* 40,
569 611–618. <https://doi.org/10.1093/ije/dyq071>
- 570 Koch, H.M., Christensen, K.L.Y., Harth, V., Lorber, M., Brüning, T., 2012. Di-n-butyl
571 phthalate (DnBP) and diisobutyl phthalate (DiBP) metabolism in a human volunteer
572 after single oral doses. *Arch Toxicol* 86, 1829–1839. [https://doi.org/10.1007/s00204-](https://doi.org/10.1007/s00204-012-0908-1)
573 [012-0908-1](https://doi.org/10.1007/s00204-012-0908-1)
- 574 Holger M. Koch, Maria Rüther, André Schütze, André Conrad, Claudia Pälme, Petra Apel,
575 Thomas Brüning, Marike Kolossa-Gehring. Phthalate metabolites in 24-h urine
576 samples of the German Environmental Specimen Bank (ESB) from 1988 to 2015 and
577 a comparison with US NHANES data from 1999 to 2012. *Int J Hyg Environ Health*
578 220, 130-141. [https://doi: 10.1016/j.ijheh.2016.11.003](https://doi:10.1016/j.ijheh.2016.11.003). 2017 Mar.

- 579 Koo, J.-W., Parham, F., Kohn, M.C., Masten, S.A., Brock, J.W., Needham, L.L., Portier, C.J.,
580 2002. The association between biomarker-based exposure estimates for phthalates
581 and demographic factors in a human reference population. *Environ Health Perspect*
582 110, 405–410. <https://doi.org/10.1289/ehp.02110405>
- 583 Latini, G., 2005. Monitoring phthalate exposure in humans. *Clinica Chimica Acta* 361, 20–
584 29. <https://doi.org/10.1016/j.cccn.2005.05.003>.
- 585 Li J, Qian X, Zhao H, Zhou Y, Xu S, Li Y, Xiang L, Shi J, Xia W, Cai Z. Determinants of
586 exposure levels, metabolism, and health risks of phthalates among pregnant women
587 in Wuhan, China. *Ecotoxicol Environ Saf*. 2019 Nov 30;184:109657. doi:
588 10.1016/j.ecoenv.2019.109657. Epub 2019 Sep 14. PMID: 31526923.
- 589 Liao, C., Liu, W., Zhang, J., Shi, W., Wang, X., Cai, J., Zou, Z., Lu, R., Sun, C., Wang, H.,
590 Huang, C., Zhao, Z., 2018. Associations of urinary phthalate metabolites with
591 residential characteristics, lifestyles, and dietary habits among young children in
592 Shanghai, China. *Sci Total Environ* 616–617, 1288–1297.
593 <https://doi.org/10.1016/j.scitotenv.2017.10.189>
- 594 Lyche, J.L., Gutleb, A.C., Bergman, A., Eriksen, G.S., Murk, A.J., Ropstad, E., Saunders,
595 M., Skaare, J.U., 2009. Reproductive and developmental toxicity of phthalates. *J*
596 *Toxicol Environ Health B Crit Rev* 12, 225–249.
597 <https://doi.org/10.1080/10937400903094091>
- 598 Mage, D., Allen, R., Kodali, A., 2008. Creatinine corrections for estimating children’s and
599 adult’s pesticide intake doses in equilibrium with urinary pesticide and creatinine
600 concentrations. *Journal of exposure science & environmental epidemiology* 18, 360–
601 8. <https://doi.org/10.1038/sj.jes.7500614>
- 602 Ministry of Health and Welfare (MHLW) Notice No.370, 2010.
- 603 Navaranjan, G., Takaro, T.K., Wheeler, A.J., Diamond, M.L., Shu, H., Azad, M.B., Becker,
604 A.B., Dai, R., Harris, S.A., Lefebvre, D.L., Lu, Z., Mandhane, P.J., McLean, K.,
605 Moraes, T.J., Scott, J.A., Turvey, S.E., Sears, M.R., Subbarao, P., Brook, J.R., 2020.
606 Early life exposure to phthalates in the Canadian Healthy Infant Longitudinal
607 Development (CHILD) study: a multi-city birth cohort. *J Expo Sci Environ*
608 *Epidemiol* 30, 70–85. <https://doi.org/10.1038/s41370-019-0182-x>
- 609 Public Law 110-314 H.R. 4040, 2008.
- 610 Rowdhwal, S.S.S., Chen, J., 2018. Toxic Effects of Di-2-ethylhexyl Phthalate: An Overview.
611 *Biomed Res Int* 2018, 1750368–1750368. <https://doi.org/10.1155/2018/1750368>
- 612 Schwedler, G., Rucic, E., Lange, R., Conrad, A., Koch, H.M., Pälmeke, C., Brüning, T.,
613 Schulz, C., Schmied-Tobies, M.I.H., Daniels, A., Kolossa-Gehring, M., 2020.

614 Phthalate metabolites in urine of children and adolescents in Germany. Human
615 biomonitoring results of the German Environmental Survey GerES V, 2014–2017.
616 International Journal of Hygiene and Environmental Health 225, 113444.
617 <https://doi.org/10.1016/j.ijheh.2019.113444>

618 Shinohara N., Uchino K., 2020. Diethylhexyl phthalate (DEHP) emission to indoor air and
619 transfer to house dust from a PVC sheet. *Sci Total Environ.* Apr 1;711:134573. doi:
620 10.1016/j.scitotenv.2019.134573. Epub 2019 Nov 20. PMID: 32000312.

621 Śmiełowska, M., Marć, M., Zabiegała, B., 2017. Indoor air quality in public utility
622 environments-a review. *Environ Sci Pollut Res Int* 24, 11166–11176.
623 <https://doi.org/10.1007/s11356-017-8567-7>

624 Song, N.R., On, J., Lee, J., Park, J.-D., Kwon, H.-J., Yoon, H.J., Pyo, H., 2013.
625 Biomonitoring of urinary di(2-ethylhexyl) phthalate metabolites of mother and child
626 pairs in South Korea. *Environment International* 54, 65–73.
627 <https://doi.org/10.1016/j.envint.2013.01.007>

628 Takagi, M., Yoshinaga, J., 2009. Risk assessment of chemical exposure via house dust
629 ingestion in Japanese children. *Indoor Environment* 12, 103–114.
630 <https://doi.org/10.7879/siej.12.103>

631 Tranfo, G., Caporossi, L., Pigini, D., Capanna, S., Papaleo, B., Paci, E., 2018. Temporal
632 Trends of Urinary Phthalate Concentrations in Two Populations: Effects of REACH
633 Authorization after Five Years. *Int J Environ Res Public Health* 15.
634 <https://doi.org/10.3390/ijerph15091950>

635 US. EPA (US Environmental Protection Agency), Di (2-ethylhexyl) phthalate (DEHP)
636 CASRN 117-81-7, 1991.

637 VEC, 2018. 可塑劑出荷量, Vinyle Environmental Council.

638 Wang Y., Zhu H., Kannan K., 2019. A Review of Biomonitoring of Phthalate Exposures.
639 *Toxics* 7, 1–28. doi: 10.3390/toxics7020021.

640 Wang, B., Wang, H., Zhou, W., Chen, Y., Zhou, Y., Jiang, Q., 2015. Urinary excretion of
641 phthalate metabolites in school children of China: implication for cumulative risk
642 assessment of phthalate exposure. *Environ Sci Technol* 49, 1120–1129.
643 <https://doi.org/10.1021/es504455a>

644 Weng, T.-I., Chen, M.-H., Lien, G.-W., Chen, P.-S., Lin, J.C.-C., Fang, C.-C., Chen, P.-C.,
645 2017. Effects of Gender on the Association of Urinary Phthalate Metabolites with
646 Thyroid Hormones in Children: A Prospective Cohort Study in Taiwan. *Int J Environ*
647 *Res Public Health* 14, 123. <https://doi.org/10.3390/ijerph14020123>

- 648 WHO, 2012. State of the science of endocrine disrupting chemicals - 2012.
- 649 Wilson, H., VanSnick, S., 2017. The effectiveness of dust mitigation and cleaning strategies
650 at The National Archives, UK. *Journal of Cultural Heritage* 24, 100–107.
651 <https://doi.org/10.1016/j.culher.2016.09.004>
- 652 Wittassek, M., Heger, W., Koch, H.M., Becker, K., Angerer, J., Kolossa-Gehring, M., 2007.
653 Daily intake of di(2-ethylhexyl)phthalate (DEHP) by German children -- A
654 comparison of two estimation models based on urinary DEHP metabolite levels. *Int*
655 *J Hyg Environ Health* 210, 35–42. <https://doi.org/10.1016/j.ijheh.2006.11.009>
- 656 Wittassek, M., Koch, H.M., Angerer, J., Bruning, T., 2011. Assessing exposure to phthalates
657 - the human biomonitoring approach. *Mol Nutr Food Res* 55, 7–31.
658 <https://doi.org/10.1002/mnfr.201000121>
- 659 Xu Y, Cohen Hubal EA, Little JC. Predicting residential exposure to phthalate plasticizer
660 emitted from vinyl flooring: sensitivity, uncertainty, and implications for
661 biomonitoring. *Environ Health Perspect.* 2010 Feb;118(2):253-8. doi:
662 10.1289/ehp.0900559.
- 663 Yoshida, T., Mimura, M., Sakon, N., 2020. Intakes of phthalates by Japanese children and
664 the contribution of indoor air quality in their residences. *Environ Sci Pollut Res Int*
665 27, 19577–19591. <https://doi.org/10.1007/s11356-020-08397-w>
- 666 Zota, A.R., Calafat, A.M., Woodruff, T.J., 2014. Temporal trends in phthalate exposures:
667 findings from the National Health and Nutrition Examination Survey, 2001-2010.
668 *Environ Health Perspect* 122, 235–241. <https://doi.org/10.1289/ehp.1306681>
- 669 Zota, A.R., Phillips, C.A., Mitro, S.D., 2016. Recent Fast Food Consumption and Bisphenol
670 A and Phthalates Exposures among the U.S. Population in NHANES, 2003-2010.
671 *Environ Health Perspect* 124, 1521–1528. <https://doi.org/10.1289/ehp.1510803>
- 672

Table 1: Study population demographic and building characteristics

		N (%) or median (range)
Gender	Boys	203 (52.6)
	Girls	183 (47.4)
Height	cm	119.8 (102.0-150.0)
Weight	Kg	22.0 (14.8-42.3)
Urine sample collection year	2012	62 (16.1)
	2013	65 (16.8)
	2014	54 (14.0)
	2015	74 (19.1)
	2016	86 (22.2)
	2017	45 (11.6)
Annual household income (JPY)	< 3 Million	48 (12.4)
	≥ 3 Million	321 (83.2)
Number of residents	≤4	251 (65.0)
	≥5	135 (35.0)
Home type	Detached	269 (70.0)
	Apartment	116 (30.0)
House structure	Wooden	269 (69.5)
	Concrete	114 (29.5)
Renovation within the past 1 year	Yes	22 (5.7)
	No	364 (94.3)
Mechanical ventilation system in living and/or child room	Yes	229 (59.3)
	No	157 (40.7)
Use of insecticide	Yes	125 (32.4)
	No	261 (67.6)
PVC flooring	Yes	65 (16.9)
	No	321 (83.1)
PVC wall material	Yes	310 (80.3)
	No	76 (19.7)
Vacuum cleaning/week	≤3 times	199 (54.8)
	4-7 times	164 (45.2)
Duration of window opening/day	<1 hour	247 (64.0)
	≥1 hour	139 (36.0)
Main road	< 50 meters	75 (19.5)
	No or ≥ 50 meters	310 (80.5)
Building age (years)	Continuous	13 (<1- 50)
Dampness index (0-5)	Continuous	2 (1 - 5)

Table 2: Distribution of urinary phthalate metabolite concentrations in 7 years old children.

Phthalate metabolite (ng/mL)	LOD		Percentile					
	(ng/mL)	% > LOD	Min.	P25	P50	P75	P95	Max.
MiBP	0.95	99.7	<LOD	7.1	12.1	27.4	86	463.1
MnBP	0.78	100	2.5	20.7	35.1	58.8	117.1	1259.3
MBzP	0.10	98.9	<LOD	0.7	1.5	3.5	22.8	498.1
MEHP	0.15	100	0.4	2.4	4.1	7	16.1	31.6
MEOHP	0.05	100	1.3	12.3	20.5	33.2	65.1	158.7
MEHHP	0.15	100	1.8	16.4	26.7	43.8	84.8	219
MECPP	0.12	100	2.4	23.3	38.4	67.1	134.8	323.3
MiNP	0.09	96.9	<LOD	0.4	0.6	1.2	2.9	7.8
OH-MiNP	0.05	100	0.3	2.2	4.1	7.5	17	60
cx-MiNP	0.11	99.7	<LOD	1.3	2.4	4.6	10.8	35.3
Σ DEHP ^a (μ mol/L)	n.a	n.a	0.01	0.18	0.29	0.49	0.97	2.38
Σ DINP ^b (μ mol/L)	n.a	n.a	0.00	0.01	0.02	0.04	0.09	0.29
Creatinine corrected (μ g/g Cr)								
MiBP			1.6	8.4	13.3	25.0	96.6	341.4
MnBP			4.7	26.3	39.1	59.2	108.6	1516.4
MBzP			<LOD	0.7	1.7	3.9	37.1	533.9
MEHP			0.7	2.9	4.5	7.4	16.2	61.8
MEOHP			2.4	14.9	22.4	32.5	69.7	228.7
MEHHP			4.2	19.3	28.7	43.7	94.8	415.7
MECPP			4.9	27.0	42.8	68.5	136.1	554.4
MiNP			<LOD	0.4	0.7	1.3	3.2	15.0
OH-MiNP			0.7	2.8	4.5	7.4	18.3	113.7
cx-MiNP			<LOD	1.6	2.7	4.9	12.1	45.7
Σ DEHP ^a (μ mol/L)			0.04	0.22	0.34	0.51	1.01	4.16
Σ DINP ^b (μ mol/L)			0.00	0.01	0.02	0.03	0.08	0.49
Urinary creatinine (g/l)			0.1	0.6	0.9	1.2	1.7	2.2

^a Σ DEHP: sum of molar concentrations metabolites [MEHP + MEOHP + MEHHP + MECPP]

^b Σ DINP: sum of molar concentrations metabolites [MiNP + OH-MiNP + cx-MiNP]

Abbreviations; LOD: Limit of detection; Max: maximum; Min: minimum; P: percentiles; n.a: not applicable; MiBP: mono-isobutyl phthalate, MnBP: mono-n-buty phthalate, MBzP: mono-benzyl phthalate, MEHP: mono (2-ethylhexyl) phthalate, MEOHP: mono (2-ethyl-5-oxohexyl) phthalate, MEHHP: mono (2-ethyl-5-hydroxyhexyl) phthalate, MECPP: mono (2-ethyl-5-carboxypentyl) phthalate, MiNP: mono-isononyl phthalate, OH-MiNP: mono-hydroxy-isononyl phthalate, cx-MiNP: mono(carboxy-isononyl phthalate).

Table 3: Percent difference (95% CI) in phthalate metabolites concentrations with demographic and building characteristics of children house (N=386)

Variables	Categories	MiBP	MnBP	MBzP	∑DEHP	∑DINP
Annual household income, (in JPY)	≥ 3 Million	Ref	Ref	Ref	Ref	Ref
	<3 Million	8.2 (-7.0,25.9)	3.6 (-7.0,15.3)	-3.6 (-22.0,20.0)	10.3 (-0.6,22.4)	13.9 (1.1,28.3) *
Building age (years)	Continuous	-0.6 (-1.6,0.4)	1.0 (0.3,1.7) **	0.8 (-0.7,2.3)	0.1 (-0.6,0.8)	-0.5 (-1.3,0.2)
Vacuum cleaning/week	≤3 times	Ref	Ref	Ref	Ref	Ref
	4-7 times	-8.5 (-17.2,1.0)	-7.2 (-13.5, -0.4) *	-7.3 (-19.7,7.1)	-5.0 (-11.3,1.7)	4.7 (-3.2,13.3)
Duration of window opening	<1 hour	Ref	Ref	Ref	Ref	Ref
	≥1 hour	11.6 (0.6,23.8) *	9.7 (1.9,18.1) *	-1.4 (-15.2,14.6)	12.3 (4.6,20.6) **	-1.0 (-8.8,7.4)
Ventilation in living or child room	Yes	Ref	Ref	Ref	Ref	Ref
	No	3.0 (-7.6, 14.7)	-1.0 (-8.3,6.9)	-6.5 (-22.7,13.2)	0.4 (-6.7,8.0)	-4.0 (-11.8,4.5)
Dampness index (0-5)	Continuous	1.6 (-6.8,10.9)	5.3 (-1.0,12.0)	-4.7 (-16.0,8.1)	0.1 (-5.7,6.3)	0.8 (-5.9,8.0)

Ref: reference, *P<0.05, ** P<0.01, Phthalate metabolites in urine were natural log transformed and corrected for creatinine level before analysis. General regression analyses conducted with phthalate metabolites concentration as dependent variable and the all building characteristics were mutually adjusted. DEHP metabolites MEHP, MEOHP, MEHHP and MECPP showed similar estimates thus in this table ∑ DEHP is presented. DINP metabolites MiNP, OH-MiNP, and cx-MiNP showed similar direction estimate thus in this table ∑ DINP is presented.




Figure 1: Natural log transformed creatinine corrected concentration level of urinary phthalate metabolites. Bars represent interquartile ranges and median. Points on dotted line indicates samples with concentration limit of detection (<LOD).




Figure 2: Secular trends and comparison of phthalate metabolites concentration (median ng/mL) in children from different countries. Data sources: Japan from present study and (Ait Bamai et al., 2015); German from (Becker et al., 2009; Schwedler et al., 2020) ; USA from (CDC, 2019) ; China from references (Liao et al., 2018; Wang et al., 2015); Taiwan from reference Weng et al.,2017; Australia from (Hartmann et al., 2015) reference; South Korea from (Song et al., 2013). Points on zero represent concentrations with less than limit of detection (<LOD).




Figure 3: Estimated daily intake ($\mu\text{g}/\text{kg}/\text{day}$) of (I) di-n-butyl phthalate (DnBP) based on mono-n butyl phthalate, (II) di-isobutyl phthalate (DiBP) based on mono-iso butyl phthalate * TDI assumed by analogy to DnBP, (III) Butyl benzyl phthalate (BBzP) based on mono-benzyl phthalate MBzP, (IV) di(2-ethylhexyl) phthalate (DEHP) sum of molar concentrations metabolites mono(2-ethylhexyl) phthalate [(MEHP) + mono(2-ethyl-5-oxohexyl) phthalate (MEOHP)+ mono(2-ethyl-5-hydroxyhexyl) phthalate (MEHHP) + mono (2-ethyl-5-carboxypentyl) phthalate (MECPP)], (V) di-iso-nonyl-phthalate (DINP) sum of molar concentrations metabolites [mono-isononyl phthalate (MiNP) + mono (hydroxy-isononyl) phthalate +(OH-MiNP) mono (carboxy-isononyl) phthalate (cx-MiNP)]. (VI) grouped DBP, BBzP, DEHP and DINP. The horizontal dot-dashed lines represent EFSA (2005 and 2019) tolerable daily intake (TDI) and the U.S. EPA reference dose (RfD) and plots above these lines represent values that exceeded EFSA or RfD reference.