



Title	Continuous feeding of a combined high-fat and high-sucrose diet, rather than an individual high-fat or high-sucrose diet, rapidly enhances the glucagon-like peptide-1 secretory response to meal ingestion in diet-induced obese rats
Author(s)	Pinyo, Jukkrapong; Hira, Tohru; Hara, Hiroshi
Citation	Nutrition, 62, 122-130 <a href="https://doi.org/10.1016/j.nut.2019.01.004">https://doi.org/10.1016/j.nut.2019.01.004</a>
Issue Date	2019-06
Doc URL	<a href="http://hdl.handle.net/2115/85392">http://hdl.handle.net/2115/85392</a>
Rights	© 2019 This manuscript version is made available under the CC-BY-NC-ND 4.0 license <a href="http://creativecommons.org/licenses/by-nc-nd/4.0/">http://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Rights(URL)	<a href="http://creativecommons.org/licenses/by-nc-nd/4.0/">http://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Type	article (author version)
File Information	Nutrition 62_122.pdf



[Instructions for use](#)

1 **Continuous feeding with a combined high fat/high sucrose diet, rather than an individual high fat**  
2 **or high sucrose diet, rapidly enhances the GLP-1 secretory response to meal ingestion in diet-**  
3 **induced obese rats**

4  
5 **Abstract**

6       Glucagon-like peptide-1 (GLP-1) is secreted by enteroendocrine L-cells in response to nutrient  
7 ingestion. To date, GLP-1 secretion in diet-induced obesity is not well characterized. We aimed to  
8 examine GLP-1 secretion in response to meal ingestion during the progression of diet-induced obesity  
9 and to determine whether a combined high fat/high sucrose diet, an individual high fat or high sucrose  
10 diet affects adaptive changes in the postprandial GLP-1 response. Rats were fed a control diet, high fat  
11 diet (30% weight, HiFat), a high sucrose diet (40% weight, HiSuc), or a high fat/high sucrose diet (HFS,  
12 30% fat and 40% sucrose) for 5 weeks. Meal tolerance tests (MTTs) were conducted to determine  
13 postprandial glucose, insulin, and GLP-1 responses to standard (control) diet ingestion every 2 weeks.  
14 After 5 weeks, body weight gain of the HiFat ( $232.3 \pm 7.8$  g,  $P = 0.021$ ) and HFS groups ( $228.0 \pm 7.8$ ,  
15  $P = 0.039$ ), but not of the HiSuc group ( $220.3 \pm 7.9$ ,  $P = 0.244$ ), were significantly higher than the  
16 control group ( $200.7 \pm 5.4$  g). In MTT after 2 weeks, GLP-1 concentration was significantly elevated  
17 only in the HFS group ( $17.2 \pm 2.6$  pM,  $P < 0.001$ ) in response to meal ingestion, but the HiFat group  
18 ( $16.6 \pm 3.7$  pM,  $P = 0.156$ ) had a similar response to the HFS group. After 4 weeks, GLP-1  
19 concentrations were similarly elevated at 15 min in the HFS ( $14.1 \pm 4.4$ ,  $P = 0.010$ ), HiFat ( $13.2 \pm 2.0$ ,  
20  $P < 0.001$ ), and HiSuc groups ( $13.0 \pm 3.3$ ,  $P = 0.016$ ), but the HFS ( $9.8 \pm 1.0$ ,  $P = 0.019$ ) and HiFat  
21 groups ( $8.3 \pm 1.5$ ,  $P = 0.010$ ) had significant elevation also at 30 min. These results demonstrate that  
22 continuous ingestion of excessive fat and sucrose rapidly enhances the GLP-1 secretory response to  
23 luminal nutrients, and the high fat diet may have potent effect compared to the high sucrose diet on

24 GLP-1 secretory responses. The increment of postprandial GLP-1 and insulin secretion may have a role  
25 in normalizing postprandial glycaemia and slowing the establishment of glucose intolerance.

26

27 **Keywords:** Glucagon-like peptide-1; Luminal nutrients; L-cell; Meal tolerance test; Obesity

28

## 29 **Introduction**

30 Obesity is rapidly increasing in the global population and has become a major public health  
31 problem [1]. Dietary habits are now considered key factors related to obesity development, especially  
32 excessive energy consumption from a high fat and/or a high sugar diet [2,3]. Obesity leads to metabolic  
33 syndromes, such as hyperinsulinemia, hypertension, hyperlipidaemia, and type II diabetes mellitus  
34 [4,5]. Glucagon-like peptide-1 (GLP-1), a product of the proglucagon (*gcg*) gene, is a gut-derived  
35 hormone, which is produced and secreted from enteroendocrine L cells in response to nutrient ingestion,  
36 including protein, glucose, fatty acids, and dietary fibre [5-9]. GLP-1 has been recognized as an  
37 incretin hormone that reduces the postprandial glycaemic response by stimulating insulin secretion [10].  
38 Existing data on GLP-1 responses in humans to nutrient ingestion (enhanced, unchanged, and  
39 decreased) in obesity, prediabetes, and diabetes patients remains unclear [11,12]. GLP-1 secretion in  
40 response to oral glucose was enhanced in type II diabetic patients [13]. In contrast, GLP-1 secretion  
41 was impaired in obese subjects and type II diabetic patients [14,15], whereas GLP-1 secretion in  
42 response to oral glucose was unchanged in type II diabetic patients [16].

43 In order to understand how GLP-1 responses are modified in the process of diet-induced obesity,  
44 it is relevant to adopt animal models continuously fed an obesogenic diet, rather than using genetically  
45 obese/diabetic models. Continuous feeding (8 weeks) of a high fat and high sucrose diet increased  
46 postprandial GLP-1 secretion in response to normal meal administration, as previously reported [17].

47 Although the impact of feeding animal models with a high fat and/or high sucrose diet has been  
48 previously examined, the alteration of the postprandial GLP-1 response during obesity development  
49 has not been clearly characterized [18-20].

50 The aims of this study were to clarify the GLP-1 response to meal ingestion during the  
51 progression of diet-induced obesity and to determine whether a combined high fat/high sucrose diet or  
52 an individual high fat or high sucrose diet contributed to adaptive changes in postprandial GLP-1  
53 secretion. In this study, the rats were given a standard (control) diet as a meal tolerance test (MTT),  
54 instead of the typical oral glucose loading (OGTT).

55

## 56 **Experimental methods**

### 57 *Animals and diet*

58 Five-week-old male Sprague-Dawley rats (weighing 160-200 g) were purchased from Japan  
59 SLC, Inc. (Shizuoka, Japan) and were fed with the American Institute of Nutrition (AIN)-93G (control)  
60 diet for 1 week as an acclimation period [21]. Each rat was housed in an individual cage in a  
61 temperature- and humidity-controlled environment with a 12 h light-dark cycle (8.00 a.m.-8.00 p.m.  
62 light period) and was allowed free access to diet and water (*ad libitum*). After one-week acclimation  
63 period, fasting glucose and GLP-1 concentrations were measured. Then, the rats were divided into four  
64 groups to have matched body weight, plasma glucose, and GLP-1 concentrations. Rats in each group  
65 were given the control, high fat/high sucrose (HFS, 30% fat weight and 40% sucrose weight) [17], high  
66 fat (HiFat, 30% fat), or high sucrose (HiSuc, 40% sucrose) diet (Table 1) for 5 weeks. The initial body  
67 weight, fasting glucose, and GLP-1 concentrations were  $168.7 \pm 3.1$  g,  $73.2 \pm 2.4$  mg/dL,  $18.0 \pm 3.0$   
68 pM in the control group (n=10),  $169.3 \pm 2.5$  g,  $72.0 \pm 3.3$  mg/dL,  $17.7 \pm 3.1$  pM in HFS group (n=8),  
69  $169.1 \pm 4.3$  g,  $71.5 \pm 4.3$  mg/dL,  $17.8 \pm 1.8$  pM in HiFat group (n=10), and  $167.7 \pm 2.8$  g,  $71.5 \pm 2.0$

70 mg/dL,  $20.0 \pm 3.2$  pM in HiSuc group (n=8), respectively. The final compositions of 30% fat in the  
71 HiFat diet and 40% sucrose in the HiSuc diet were adjusted to be equal to those in the HFS diet. Body  
72 weight and food intake were measured every 2 days. All experimental animal procedures used in this  
73 study were approved by the institutional animal care committee, and the animals were maintained in  
74 accordance with the institutional animal guidelines for the care and use of laboratory animals.

#### 75 *Meal tolerance test (MTT)*

76 Meal tolerance test (MTT) was performed on rats after they were fed the test diets for 2 and 4  
77 weeks, in order to determine the postprandial glucose, insulin, and GLP-1 responses to standard meal  
78 ingestion. The rats were fasted for 16 h, and then basal (0 min) blood was collected from the tail vein.  
79 Then, the rats were given the control diet (10 g/kg body weight) for 30 min, and blood samples from  
80 the tail vein were collected at 15, 30, 60, 90 and 120 min after providing the diet. In this study, only the  
81 rats that ingested more than 90% of the given diet were used to assess the postprandial responses.  
82 Blood samples were collected in chilled tubes containing heparin (final concentration 50 IU/mL;  
83 Ajinomoto Company, Inc., Tokyo, Japan) and aprotinin (final concentration 500 Kallikrein inhibitor  
84 units (KIU)/mL; Wako Pure Chemical Industries, Ltd., Osaka, Japan). Plasma was separated by  
85 centrifugation at  $2300 \times g$  for 10 min at  $4^{\circ}\text{C}$  and then stored at  $-80^{\circ}\text{C}$  until analysis. The plasma  
86 glucose level was determined using the Glucose CII Test Kit (Wako Pure Chemical Industries, Osaka,  
87 Japan). Plasma insulin and GLP-1 concentrations were analysed using the Rat Insulin ELISA (AKRIN-  
88 010T; Shibayagi Company Limited, Gunma, Japan) and Multi Species GLP-1 Total ELISA  
89 (EZGLP1T-36K; Merck Millipore, Darmstadt, Germany), respectively. The GLP-1 Total ELISA  
90 detects both GLP-1 (7-36) and GLP-1 (9-36) and has no significant cross-reactivity with GLP-2, GIP,  
91 glucagon, oxyntomodulin. The minimum detection limit of the assay is 1.5 pM. The intra-assay  
92 precision is  $< 5\%$ , and the inter-assay precision is  $< 12\%$ , respectively. The homeostatic model

93 assessment of insulin resistance (HOMA-IR) was used to assess insulin resistance. It was calculated by  
94 the following equation [22], using glucose and insulin values at the fasting state and those values of the  
95 area under the curve during MTT:

$$96 \quad \text{HOMA-IR} = \text{insulin } (\mu\text{U/mL}) \times \text{glucose (mg/dL)} / 2430$$

97 Where: 1 mg insulin = 26 IU

### 98 *Blood and tissue collection*

99 After receiving the test diet for 5 weeks, the rats were fasted overnight (16 h). Blood samples  
100 from the portal vein were taken under sodium pentobarbital anaesthesia (50 mg/kg of body weight,  
101 somnopentyl injection; Kyoritsu Seiyaku Corporation, Tokyo, Japan) and collected in a chilled syringe  
102 containing heparin (final concentration 50 IU/mL), aprotinin (final concentration 500 KIU/mL), and  
103 DPP-IV inhibitors (final concentration 50  $\mu\text{mol/L}$ ; DPP4-010; Merck Millipore, Darmstadt, Germany).  
104 The rats were then sacrificed by exsanguination. Intestinal segments were carefully dissected and  
105 washed with a cold saline solution (0.9% NaCl); then, 2 cm segments of the jejunum, ileum, and colon  
106 were taken from the middle region to measure GLP-1 content. The caecal tissues were washed with a  
107 cold saline solution and divided equally into 2 parts; then, 2 cm from the middle region was collected  
108 for GLP-1 measurement. All intestinal segments were rapidly frozen in liquid N<sub>2</sub> and stored at -80°C  
109 until measurement of the GLP-1 content. The mesenteric, retroperitoneal, and epididymal adipose  
110 tissues were individually weighed and expressed as the visceral adipose tissue weight. Plasma was  
111 stored as described above in order to measure the glucose, insulin, total GLP-1, triacylglycerol, and  
112 total cholesterol levels. Plasma triacylglycerol and cholesterol concentrations were measured using the  
113 Triglyceride E Test and Cholesterol E Test kits (Wako Pure Chemical Industries, Osaka, Japan),  
114 respectively.

115 *Measurement of the glucagon-like peptide-1 (GLP-1) content in intestinal tissue*

116 Intestinal segments (2 cm length) were immersed in an ethanol-acid solution (absolute  
117 ethanol:water:12 M HCl = 74:25:1, 5 mL/g of intestinal tissue segments) [23] and then cut into small  
118 pieces. The tissue samples were homogenized at 25,000 rpm (Ultra Turrax homogenizer T18, IKA,  
119 Staufen, Germany) for 2 min and placed at 4°C for 24 h. The homogenates were then centrifuged at  
120 2000 x g for 20 min. The supernatant was collected for measuring the total GLP-1 and protein contents  
121 using the Multi Species GLP-1 Total ELISA (EZGLP1T-36K; Merck Millipore, Darmstadt, Germany)  
122 and Pierce BCA Protein Assay Kit (Thermo Fisher Scientific, Rockford, IL, USA), respectively. The  
123 supernatant was diluted 10-fold for protein measurement, 500-fold (jejunum, ileum, and colon  
124 segments), and 625-fold (colon segment) for GLP-1 measurement with normal saline solution.

125 *Statistical analysis*

126 The primary endpoint of this study was nutrient-induced GLP-1 responses in rats treated with  
127 various (obesogenic) diets. The secondary endpoint was to assess glycaemic and insulin responses,  
128 body weight and adipose tissue weight changes. The results were expressed as mean ± standard error of  
129 the mean (SEM). The sample size was calculated based on the experimental design (two-way repeated  
130 measure ANOVA) in MTTs for examining postprandial GLP-1 responses as the primary outcome  
131 measure, by using G\*Power software (version 3.1.9.2) as effect size  $f = 0.4$  and Power = 0.8.  
132 Significant effects of time (TI), treatment (TR) and the interactions of time and treatment (TI x TR)  
133 were assessed by two-way repeated measure ANOVA in the results of MTTs. One-way ANOVA and  
134 Tukey-Kramer's test were used to assess the significant difference among the treatments for parametric  
135 data (assessed by Goodness of Fit test). A Kruskal–Wallis test and Dunn's multiple range test were  
136 used for MTT data containing non-parametric data. Dunn's with control test was used to determine the  
137 significant differences from baseline (0 min) value within the same group. *P* values less than 0.05 was

138 considered to be statistically significant. Statistical analysis was performed using JMP Pro version 13  
139 software (SAS Institute, Inc., Cary, NC, USA).

140

## 141 **Results**

142 *Basal glucose, insulin, GLP-1 levels, and homeostasis model assessment of insulin resistance (HOMA-*  
143 *IR) after feeding with the test diet for 2 and 4 weeks.*

144 After receiving the test diet for 2 and 4 weeks, the rats were fasted overnight (16 h) before  
145 conducting the MTT. After 2 weeks of feeding, basal glucose ( $84.1 \pm 4.5 - 90.0 \pm 3.6$  mg/dL,  $P =$   
146  $0.396$ ), insulin ( $0.11 \pm 0.01 - 0.20 \pm 0.06$  nM,  $P = 0.329$ ), GLP-1 ( $12.1 \pm 1.5 - 20.4 \pm 2.6$  pM,  $P =$   
147  $0.070$ ), and HOMA-IR ( $0.54 \pm 0.07 - 1.11 \pm 0.34$ ,  $P = 0.287$ ) were not significantly different among all  
148 of the treatment groups (Fig 1A, B, C, and D). After 4 weeks, the basal insulin level and HOMA-IR  
149 tended to be higher in the HFS group ( $0.26 \pm 0.03$ ,  $P = 0.218$  and  $1.45 \pm 0.20$ ,  $P = 0.200$ ) compared to  
150 control group ( $0.17 \pm 0.02$  nM, and  $0.90 \pm 0.10$ ) (Fig 1F and H).

151 *Postprandial glycaemic, insulin, and GLP-1 responses during the MTT*

152 In this study, MTT was used instead of the oral glucose tolerance test (OGTT) in order to  
153 determine postprandial responses to ingestion of a standard diet (control diet). MTT can be used to  
154 mimic dietary exposure in normal life and voluntary ingestion is more relevant than enforced gavage  
155 feeding. Although HiSuc group had relatively lower concentrations, postprandial glycaemic responses  
156 were overall similar in all groups (Fig. 2A and E) after receiving the test diet for 2 weeks. The AUC of  
157 insulin increased largely in the HFS group ( $1.15 \pm 0.19$  nM,  $P = 0.015$ ), compared to that of the control  
158 group ( $0.46 \pm 0.04$  nM). The HiFat group ( $0.93 \pm 0.35$ ,  $P = 1.000$ ) also showed a similar trend, but less  
159 so than the HFS group (Fig. 2F). Insulin levels at 60 min were significantly higher in the HFS ( $0.62 \pm$   
160  $0.08$  nM) than those in the HiSuc group ( $0.23 \pm 0.03$  nM,  $P = 0.022$ ) and control group ( $0.24 \pm 0.03$  nM,



161  $P = 0.016$ ) (Fig. 2B). The basal levels of GLP-1 of control, HFS, HiFat, and HiSuc groups were  $20.4 \pm$   
162  $2.6$ ,  $12.1 \pm 1.5$ ,  $12.6 \pm 2.3$ , and  $18.2 \pm 3.8$  pM respectively. Due to varied basal GLP-1 levels in each  
163 treatment group (Fig. 1), changes in GLP-1 concentrations from the basal value ( $\Delta$ GLP-1) were  
164 presented to illustrate postprandial GLP-1 response in this study. The GLP-1 level significantly  
165 increased after 15 min in HFS ( $17.2 \pm 2.6$  pM,  $P < 0.001$ ), compared to basal level. HiFat groups also  
166 showed similar level ( $16.6 \pm 3.7$  pM,  $P = 0.156$ ) without significant difference, while the HiSuc group  
167 ( $6.6 \pm 4.8$  pM,  $P = 1.000$ ) was more similar to the control group ( $7.3 \pm 2.3$  pM,  $P = 0.169$ ) (Fig. 2C).  
168 The AUC of HOMA-IR was calculated using the glycaemic and insulin response (0-120 min) during  
169 the MTT to estimate the degree of insulin resistance in the postprandial state. The results revealed that  
170 the HFS group ( $22.4 \pm 1.0$ ,  $P = 0.008$ ) had the highest AUC of HOMA-IR, followed by the HiFat group  
171 ( $18.3 \pm 7.3$ ,  $P = 1.000$ ), whereas the HiSuc group ( $8.2 \pm 0.9$ ,  $P = 1.000$ ) showed a similar level to the  
172 control group ( $8.9 \pm 1.1$ ) (Fig. 2H).

173 After receiving the test diet for 4 weeks, AUC of postprandial glycaemia in the HFS ( $301.6 \pm$   
174  $6.1$  mg/dL), HiFat ( $293.0 \pm 5.9$  mg/dL), and HiSuc ( $291.6 \pm 7.4$  mg/dL) groups were almost similar to  
175 that in the control group ( $291.3 \pm 6.6$  mg/dL,  $P = 0.579$ ). The HFS group ( $1.53 \pm 0.18$  nM,  $P = 0.010$ )  
176 had a higher postprandial insulin AUC than the control group ( $0.77 \pm 0.06$  nM), while the HiFat group  
177 ( $1.17 \pm 0.14$  nM,  $P = 0.676$ ) showed an intermediate level between the HFS and control group (Fig.  
178 3B). The GLP-1 response and HOMA-IR AUC in the HiSuc group showed similar values to those of  
179 the HiFat group, in contrast to the result of the MTT after 2 weeks (Fig. 2). The basal levels of GLP-1  
180 of control, HFS, HiFat, and HiSuc groups were  $15.0 \pm 1.9$ ,  $13.7 \pm 3.7$ ,  $11.8 \pm 2.9$  and  $11.9 \pm 2.7$   
181 respectively. Postprandial GLP-1 levels at 15 min were significantly higher in all of the treatment  
182 groups (HFS, HiFat, and HiSuc,  $13.0 \pm 3.3$  -  $14.1 \pm 4.4$  pM,  $P < 0.05$ ), compared to the basal level (Fig.  
183 3C). Significant effects of treatment (TR) on postprandial GLP-1 secretion were also detected by two-

184 way repeated measure ANOVA in both MTTs ( $P = 0.031$  and  $P = 0.003$ , respectively) (Fig. 2C and  
185 3C). Although significant differences were not detected,  $\Delta$ AUC of GLP-1 in the HFS, HiFat, and  
186 HiSuc groups tended to increase when compared to the control group ( $P = 0.161$ ), as shown in Fig. 3G.

187 *Effect of feeding a chronic high fat and/or high sucrose diet on body weight gain, food intake, and fat*  
188 *accumulation*

189 After receiving the test diet for 5 weeks, body weight gain and total energy intake of the HiFat  
190 ( $232.3 \pm 7.8$  g,  $P = 0.021$  and  $2966.3 \pm 56.8$  kcal,  $P = 0.039$ ) and HFS groups ( $228.0 \pm 7.8$  g,  $P = 0.039$   
191 and  $2958.6 \pm 60.3$  kcal,  $P = 0.032$ ) were significantly higher than those of the control group ( $200.7$   
192  $\pm 5.4$  g and  $2736.6 \pm 61.7$  kcal), as shown in Table 2. Epididymal adipose tissue weight in the HFS  
193 group ( $9.7 \pm 0.5$  g,  $P = 0.040$ ) was significantly higher than the control ( $7.5 \pm 0.6$  g). The HiSuc  
194 feeding group showed similar values for these parameters as those seen in the control group.

195 *Peptide hormones, cholesterol, and triacylglycerol levels in the portal vein after feeding with the test*  
196 *diet for 5 weeks.*

197 On the final day of the experiment, blood samples were collected from the portal vein after  
198 overnight fasting. The results showed that the HFS group had the highest values for the glucose ( $107.3$   
199  $\pm 3.9$  mg/dL,  $P = 0.030$ ), insulin ( $1.38 \pm 0.20$  nM,  $P = 0.069$ ), total cholesterol ( $35.9 \pm 3.5$  mg/dL,  $P =$   
200  $0.023$ ), and TG levels ( $76.1 \pm 3.1$  mg/dL,  $P = 0.028$ ) (Fig. 4). The HiFat group overall showed similar  
201 results, but slightly lower values for these parameters, compared to the HFS group. The HiSuc group  
202 had results similar to the control group. The GLP-1 levels in the HFS ( $69.3 \pm 9.0$  pM,  $P = 0.325$ ) and  
203 HiFat groups ( $61.3 \pm 8.6$  pM,  $P = 0.812$ ) were slightly higher than the control group ( $51.8 \pm 6.2$  pM),  
204 but without significant differences (Fig. 4C).

205 *GLP-1 content in the intestinal tissues of rats after feeding with the test diet for 5 weeks.*

206 In the jejunum segment, the HFS group ( $4.8 \pm 0.4$  pmol/mg protein) had a significantly higher  
207 GLP-1 content than that of the control group ( $3.2 \pm 0.2$  pmol/mg protein,  $P = 0.004$ ), while in the other  
208 segments (ileum, cecum, and colon) significant differences were not observed (Fig. 5A, B, C, and D).

209

## 210 **Discussion**

211 Excess energy consumption, especially from a high fat and/or high sugar diet, is a major risk  
212 factor for the development of metabolic disorders, including obesity, insulin resistance, and type II  
213 diabetes in rodent [20,24-26]. Therefore, diets containing high fat and/or high sucrose are commonly  
214 used to establish obesity development in animal models [24,27-29]. In this study, a HFS diet (30% fat  
215 and 40% sucrose), HiFat diet (30% fat diet), or HiSuc diet (40% sucrose) were provided to rats for a  
216 total of 5 weeks in order to determine whether a combined high fat/high sucrose diet, an individual high  
217 fat or high sucrose diet contributed to adaptive changes in postprandial GLP-1 secretion and obesity  
218 development.

219 As expected, body weight gain and total energy intake of the HFS and HiFat groups were  
220 higher than the control group, while the HiSuc group was similar to the control group. The results were  
221 also consistent with a previous report, in which chronic feeding of HFS and HiFat caused additional  
222 body weight gain and obesity development [27-33]. Consumption of a high caloric diet directly relates  
223 to fat accumulation in various tissues and also obesity development [27,29,34]. Indeed, the epididymal  
224 adipose tissue weight of the HFS and HiFat groups increased largely when compared with the HiSuc  
225 group. These results suggest that the high fat diet, rather than high sucrose diet has a potent impact on  
226 obesity (adiposity) development in rodent. Nevertheless, a combination of a high fat and high sucrose  
227 diet had an intense impact on obesity development, compared to a high fat diet alone.

228           Nowadays, several methodologies, such as the OGTT and MTT, are available to assess  
229 postprandial glucose and insulin responses [35,36]. However, a previous study supported the fact that  
230 the MTT is more reflective of postprandial metabolic responses than the OGTT [37]. In addition, we  
231 considered that the voluntary ingestion was more appropriate than enforced gavage feeding to mimic  
232 the dietary exposure in normal life. Therefore, in the present study, MTT, using ingestion of the  
233 standard (control) diet at a dose of 10 g/kg body weight, was performed in order to evaluate  
234 postprandial glycaemia, insulin, and GLP-1. Unfortunately, some rats did not consume more than 90%  
235 weight of the provided diet (20.3 to 89.7% of provided diet) which could be a limitation of the current  
236 study. The data from these rats were not included since glycaemic and gut hormone responses primarily  
237 depend on amount of food ingested. Because it was difficult to interpret the data from all of rats with  
238 largely varied food consumption, we decided to omit the data from those rats. It is assumed that lower  
239 diet consumption resulted in lower GLP-1 secretion compared to rats consumed more than 90% of the  
240 diet since GLP-1 secretion depends on the caloric load administered [38-39.] Nevertheless, significant  
241 effects of treatment (diet) were detected by two-way repeated ANOVA in MTTs. Accordingly, we  
242 consider that oral administration of meal solution would be suitable to assess postprandial responses  
243 instead of voluntary feeding in MTT using animal models. Although it was difficult to control, the  
244 results obtained from the MTT with voluntary feeding should have significant meaning to understand  
245 postprandial glycaemic and gut hormone responses.

246           Continuous feeding of the HFS and HiFat diet for 2 weeks highly affected the postprandial  
247 insulin response and slightly enhanced postprandial GLP-1 response. Likewise, we found that the AUC  
248 of HOMA-IR was largely increased by feeding with the HFS diet for 2 weeks. This indicates that  
249 feeding with the HFS diet immediately triggered the development of insulin resistance. Interestingly,  
250 continuous consumption of the HiSuc diet for 4 weeks illustrated that the postprandial GLP-1, insulin,

251 and HOMA-IR parameters eventually matched those observed for the HiFat feeding group. It can be  
252 concluded that chronic feeding with the HiSuc diet gradually increased the postprandial insulin  
253 response and HOMA-IR index. These findings are supported by a previous study, which demonstrated  
254 that the HiFat diet caused severe metabolic dysfunction faster than the HiSuc diet [24].

255 After feeding with the test diet for 2 weeks, the HFS feeding groups showed significant  
256 elevation of the postprandial GLP-1 level, while the other groups did not. As all rats were given the  
257 identical control diet for the MTT, differences in the GLP-1 response should be attributed to the  
258 differences in the rat phenotype. In addition, basal GLP-1 levels did not differ significantly throughout  
259 the experimental period. One of limitations of the present study is that significant differences were not  
260 detected in absolute GLP-1 values between treatments (Fig. 2D and 3D). However, by observing  
261 incremental GLP-1 ( $\Delta$ GLP-1), it was found that postprandial GLP-1 responses differed between groups  
262 (assessed by two-way repeated measure ANOVA). These results suggest that the sensitivity of GLP-1-  
263 producing L-cells to luminal nutrients (protein, carbohydrates, or fatty acids) was enhanced by chronic  
264 feeding with the HFS diets. The HiFat and HiSuc feeding group also had a similar response after  
265 receiving the test diet for 4 weeks, suggesting that ingestion of excessive fat and sucrose rapidly  
266 enhances the sensitivity of GLP-1-producing cells to luminal nutrients rather than individual fat or  
267 sucrose alone. GLP-1 is secreted in response to macro-nutrient ingestion (proteins, carbohydrates, and  
268 triglycerides) [8,40,41], and we used a control diet containing all of the nutrients in the present study.  
269 Therefore, specific diet components that contributed to enhanced postprandial GLP-1 secretion in  
270 obese animals were not identified. It would be interesting to investigate which nutrient enhances the  
271 sensitivity of L cells under diet-induced obesity.

272 To date, it is still controversial whether GLP-1 is increased or diminished during obesity  
273 development. In this study, we found that the postprandial GLP-1 level was slightly increased in the

274 HFS and HiFat feeding groups, consistent with previous studies [17,31,32]. In contrast, other studies  
275 have demonstrated that chronic feeding with a high fat diet diminished the GLP-1 secretion response to  
276 oral glucose and impaired the function of GLP-1-producing L cells [19,42]. The different results  
277 observed by each research group might be due to differences in the experimental design, such as the  
278 treatment period, diet composition, rodent species, etc.

279 From the MTT experiments, postprandial glycaemic responses were apparently unchanged by  
280 chronic feeding with the high fat and/or high sucrose diet throughout the experimental period, whereas  
281 postprandial insulin and GLP-1 secretion increased progressively. Because significant differences were  
282 not detected in absolute GLP-1 levels among treatment groups, we then speculated that the incremental  
283 GLP-1 ( $\Delta$ GLP-1) levels played an important role in blood glucose regulation through enhanced insulin  
284 secretion. This may be partially explained by increased GLP-1 response as observed. In addition, a  
285 recent study suggest that treatment with GLP-1 receptor antagonist exendin (9-39) (Ex9) reduced body  
286 weight gain in rats fed a high fat diet without affecting food intake, compared to untreated rats  
287 suggesting that Ex9 increased energy expenditures [43]. Moreover, increasing of GLP-1 secretion in  
288 high fat fed rats lead to hyperinsulinemia, thus promoting energy storage (decreasing energy  
289 expenditure) and contributing to body weight gain [43]. We therefore speculate that HFS and HiFat  
290 groups had higher body weight gain, compared to control group might involve hyperinsulinemia and  
291 lowering energy expenditure. However, previous studies demonstrated that high fat diet-fed rats  
292 exhibited impaired- GLP-1 signaling [44] and reduced GLP-1 production [45], which possibly  
293 promoted hyperphagia and increased body weight gain. Those reports focused on intestinal GLP-1  
294 content and GLP-1 signaling instead of GLP-1 secretion. It is interesting to examine whether  
295 postprandial GLP-1 responses increased or decreased in other DIO models (obese-prone and -resistant  
296 rats). The GLP-1 content of the jejunum segment in the HFS group was higher than that of the control

297 group; whereas in other intestinal segments, significant differences were not detected, indicating that  
298 the jejunum was more prone to adapting to an obesogenic diet feeding than other intestinal segments. It  
299 can be assumed that the L cells in the jejunum might be the major source of enhanced postprandial  
300 GLP-1 secretion, based on the present result. Supporting our results, postprandial GLP-1 secretion is  
301 reported to be a direct action of the luminal contents on enteroendocrine L cells in the distal jejunum  
302 [46,47]. Although the majority of enteroendocrine L-cells are located in the distal intestine, our result  
303 suggests that L cells in the proximal intestine play a role in the rapid secretion of GLP-1 in response to  
304 meal ingestion. Moreover, L-cell numbers in jejunoileum (alimentary channel) are reportedly higher  
305 than the distal gut regions (colon) after Roux-en-Y Gastric Bypass (RYGB) [48], suggesting that L-cell  
306 numbers in upper jejunum part were adaptively increased after the surgery. In addition, the RYGB  
307 contributed to intestinal hypertrophy in rats, which might be due to nutrients-induced stimuli in the  
308 jejunum part [48]. It seems consistent with our findings that jejunum part is prone to adapt to stimulant  
309 rather than other intestinal segments.

310

## 311 **Conclusions**

312 Postprandial GLP-1 secretion was elevated in rats continuously fed the HFS diet within 2 weeks.  
313 Continuous feeding of the HiFat or HiSuc diet for 4 weeks, exhibited a similar effect on the  
314 postprandial GLP-1 response, suggesting that excessive ingestion of high fat/high sucrose diet rapidly  
315 caused adaptive changes in nutrient sensitivity in GLP-1-producing cells, rather than individual high fat  
316 or high sucrose diet alone. However, HiFat diet likely has relatively potent effect on the GLP-1  
317 response compared to HiSuc diet. This may have a role in the normalization of postprandial glycaemia  
318 and the slowing of the establishment of glucose intolerance.

319

320 **Funding:** This work was supported by JSPS KAKENHI Grant Number 16K07725

321

322 **Conflict of Interest:** The authors declare that they have no conflicts of interest.

323

## 324 **References**

- 325 1. Karimi G, Sabran MR, Jamaluddin R, Parvaneh K, Mohtarrudin N, Ahmad Z, et al. The anti-  
326 obesity effects of *Lactobacillus casei* strain Shirota versus Orlistat on high fat diet-induced obese  
327 rats. Food Nutr Res 2015;59:29273.
- 328 2. Shirai T, Shichi Y, Sato M, Tanioka Y, Furusho T, Ota T, et al. High dietary fat-induced obesity in  
329 Wistar rats and type 2 diabetes in nonobese Goto-Kakizaki rats differentially affect retinol binding  
330 protein 4 expression and vitamin A metabolism. Nutr Res 2016;36:262-70.
- 331 3. Bueno AA, Oyama LM, de Oliveira C, Pisani LP, Ribeiro EB, Silveira VL, et al. Effects of  
332 different fatty acids and dietary lipids on adiponectin gene expression in 3T3-L1 cells and  
333 C57BL/6J mice adipose tissue. Pflugers Arch 2008;455:701-9.
- 334 4. Kao YH, Chang HH, Lee MJ, Chen CL. Tea, obesity, and diabetes. Mol Nutr Food Res  
335 2006;50:188-210.
- 336 5. Sakar Y, Duca FA, Langelier B, Devime F, Blottiere H, Delorme C, et al. Impact of high-fat  
337 feeding on basic helix-loop-helix transcription factors controlling enteroendocrine cell  
338 differentiation. Int J Obes (Lond) 2014;38:1440-8.



- 339 6. De León DD, Li C, Delson MI, Matschinsky FM, Stanley CA, Stoffers DA. Exendin-(9-39)  
340 corrects fasting hypoglycemia in SUR-1<sup>-/-</sup> mice by lowering camp in pancreatic β-cells and  
341 inhibiting insulin secretion. *J Biol Chem* 2008;283:25786-93.
- 342 7. Carr RD, Larsen MO, Winzell MS, Jelic K, Lindgren O, Deacon CF, et al. Incretin and islet  
343 hormonal responses to fat and protein ingestion in healthy men. *Am J Physiol Endocrinol Metab*  
344 2008;295:E779-84.
- 345 8. Tolhurst G, Reimann F, Gribble FM. Nutritional regulation of glucagon-like peptide-1 secretion. *J*  
346 *Physiol* 2009;587:27-32.
- 347 9. Baggio LL, Drucker DJ. Biology of Incretins: GLP-1 and GIP. *Gastroenterology* 2007;132:2131-  
348 57.
- 349 10. Kim YO, Schuppan D. When GLP-1 hits the liver: a novel approach for insulin resistance and  
350 NASH. *Am J Physiol Gastrointest Liver Physiol* 2012;302:G759-61.
- 351 11. Smushkin G, Sathananthan A, Man CD, Zinsmeister AR, Camilleri M, Cobelli C, et al. Defects in  
352 GLP-1 response to an oral challenge do not play a significant role in the pathogenesis of  
353 prediabetes. *J Clin Endocrinol Metab* 2012;97:589-98.
- 354 12. Nauck MA, Vardarli I, Deacon CF, Holst JJ, Meier JJ. Secretion of glucagon-like peptide-1 (GLP-  
355 1) in type 2 diabetes: What is up, what is down? *Diabetologia* 2011;54:10-8.
- 356 13. Alssema M, Rijkelijhuizen JM, Holst JJ, Teerlink T, Scheffer PG, Eekhoff EM, et al. Preserved  
357 GLP-1 and exaggerated GIP secretion in type 2 diabetes and relationships with triglycerides and  
358 ALT. *Eur J Endocrinol* 2013;169:421-30.

- 359 14. Ranganath LR, Beety JM, Morgan LM, Wright JW, Howland R, Marks V. Attenuated GLP-1  
360 secretion in obesity: cause or consequence? *Gut* 1996;38:916-9.
- 361 15. Toft-Nielsen MB, Damholt MB, Madsbad S, Hilsted LM, Hughes TE, Michelsen BK, Holst JJ.  
362 Determinants of the impaired secretion of glucagon-like peptide-1 in type 2 diabetic patients. *J*  
363 *Clin Endocrinol Metab* 2001;86:3717-23.
- 364 16. Calanna S, Christensen M, Holst JJ, Laferrère B, Gluud LL, Vilsbøll T, et al. Secretion of  
365 glucagon-like peptide-1 in patients with type 2 diabetes mellitus: Systematic review and meta-  
366 analyses of clinical studies. *Diabetologia* 2013;56:965-72.
- 367 17. Nakajima S, Hira T, Hara H. Postprandial glucagon-like peptide-1 secretion is increased during the  
368 progression of glucose intolerance and obesity in high-fat/high-sucrose diet-fed rats. *Br J Nutr*  
369 2015;113:1477-88.
- 370 18. Wang F, Yoder SM, Yang Q, Kohan AB, Kindel TL, Wang J, et al. Chronic high-fat feeding  
371 increases GIP and GLP-1 secretion without altering body weight. *Am J Physiol Gastrointest Liver*  
372 *Physiol* 2015;309:G807-15.
- 373 19. Gniuli D, Calcagno A, Dalla Libera L, Calvani R, Leccesi L, Caristo ME, et al. High-fat feeding  
374 stimulates endocrine, glucose-dependent insulinotropic polypeptide (GIP)-expressing cell  
375 hyperplasia in the duodenum of Wistar rats. *Diabetologia* 2010;53:2233-40.
- 376 20. Zheng J, Xiao KL, Chen L, Wu C, Hu X, Zeng T, et al. Insulin sensitizers improve the GLP-1  
377 secretion and the amount of intestinal L cells on high-fat-diet-induced catch-up growth. *Nutrition*  
378 2017;39:82-91.

- 379 21. Reeves PG. Components of the AIN-93 diets as improvements in the AIN-76A diet. *J Nutr*  
380 1997;127(5 Suppl):838S-41S.
- 381 22. Cacho J, Sevillano J, de Castro J, Herrera E, Ramos MP. Validation of simple indexes to assess  
382 insulin sensitivity during pregnancy in Wistar and Sprague-Dawley rats. *Am J Physiol Endocrinol*  
383 *Metab* 2008;295(5):E1269-76.
- 384 23. Cani PD, Hoste S, Guiot Y, Delzenne NM. Dietary non-digestible carbohydrates promote L-cell  
385 differentiation in the proximal colon of rats. *Br J Nutr* 2007;98:32-7.
- 386 24. Long Z, Zhang X, Sun Q, Liu Y, Liao N, Wu H, et al. Evolution of metabolic disorder in rats fed  
387 high sucrose or high fat diet: Focus on redox state and mitochondrial function. *Gen Comp*  
388 *Endocrinol* 2015;242:92-100.
- 389 25. Avila-Nava A, Noriega LG, Tovar AR, Granados O, Perez-Cruz C, Pedraza-Chaverri J, et al. Food  
390 combination based on a pre-hispanic Mexican diet decreases metabolic and cognitive  
391 abnormalities and gut microbiota dysbiosis caused by a sucrose-enriched high-fat diet in rats. *Mol*  
392 *Nutr Food Res* 2017;6.
- 393 26. Pang J, Xi C, Huang X, Cui J, Gong H, Zhang T. Effects of excess energy intake on glucose and  
394 lipid metabolism in C57BL/6 mice. *PLoS One* 2016;11:e0146675.
- 395 27. Yang ZH, Miyahara H, Takeo J, Katayama M. Diet high in fat and sucrose induces rapid onset of  
396 obesity-related metabolic syndrome partly through rapid response of genes involved in lipogenesis,  
397 insulin signalling and inflammation in mice. *Diabetol Metab Syndr* 2012;4:32.
- 398 28. Sato-Mito N, Suzui M, Yoshino H, Kaburagi T, Sato K. Long term effects of high fat and sucrose  
399 diets on obesity and lymphocyte proliferation in mice. *J Nutr Health Aging* 2009;13:602-6.

- 400 29. Sumiyoshi M, Sakanaka M, Kimura Y. Chronic intake of high-fat and high-sucrose diets  
401 differentially affects glucose intolerance in mice. *J Nutr* 2006;136(3):582-7.
- 402 30. Qi Z, Xia J, Xue X, He Q, Ji L, Ding S. Long-term treatment with nicotinamide induces glucose  
403 intolerance and skeletal muscle lipotoxicity in normal chow-fed mice: compared to diet-induced  
404 obesity. *J Nutr Biochem* 2016;36:31-41.
- 405 31. Dusaulcy R, Handgraaf S, Skarupelova S, Visentin F, Vesin C, Heddad-Masson M, et al.  
406 Functional and molecular adaptations of enteroendocrine L-cells in male obese mice are associated  
407 with preservation of pancreatic alpha-cells function and prevention of hyperglycemia.  
408 *Endocrinology* 2016;157:3832-43.
- 409 32. Hira T, Suto R, Kishimoto Y, Kanahori S, Hara H. Resistant maltodextrin or  
410 fructooligosaccharides promotes GLP-1 production in male rats fed a high-fat and high-sucrose  
411 diet, and partially reduces energy intake and adiposity. *Eur J Nutr* 2018;57(3):965-979.
- 412 33. Hong SJ, Lee JH, Kim EJ, Yang HJ, Park JS, Hong SK. Anti-obesity and anti-diabetic effect of  
413 neoagarooligosaccharides on high-fat diet-induced obesity in mice. *Mar Drugs* 2017;15:90.
- 414 34. Hariri N, Thibault L. High-fat diet-induced obesity in animal models. *Nutr Res Rev* 2010;23:270-  
415 99.
- 416 35. Rijkkelijkhuizen JM, Girman CJ, Mari A, Alsema M, Rhodes T, Nijpels G, et al. Classical and  
417 model-based estimates of beta-cell function during a mixed meal vs. an OGTT in a population-  
418 based cohort. *Diabetes Res Clin Pract* 2009;83:280-8.
- 419 36. Staten M, Kelley DE. Using oral challenge testing to assess insulin action and secretion with  
420 mathematical modeling. *Diabetes* 2014;63:1188-90.

- 421 37. Brodovicz KG, Girman CJ, Simonis-Bik AM, Rijkeljkhuizen JM, Zelis M, Bunck MC, et al.  
422 Postprandial metabolic responses to mixed versus liquid meal tests in healthy men and men with  
423 type 2 diabetes. *Diabetes Res Clin Pract* 2011;94:449-55.
- 424 38. Steinert RE, Beglinger C, Langhans W. Intestinal GLP-1 and satiation: from man to rodents and  
425 back. *Int J Obes (Lond)*. 2016;40(2):198-205.
- 426 39. Ma J, Pilichiewicz AN, Feinle-Bisset C, Wishart JM, Jones KL, Horowitz M, Rayner CK. Effects  
427 of variations in duodenal glucose load on glycaemic, insulin, and incretin responses in type 2  
428 diabetes. *Diabet Med*. 2012;29(5):604-608.
- 429 40. Sakamoto E, Seino Y, Fukami A, Mizutani N, Tsunekawa S, Ishikawa K, et al. Ingestion of a  
430 moderate high-sucrose diet results in glucose intolerance with reduced liver glucokinase activity  
431 and impaired glucagon-like peptide-1 secretion. *J Diabetes Investig* 2012;3:432-40.
- 432 41. Shannon M, Green B, Willars G, Wilson J, Matthews N, Lamb J, et al. The endocrine disrupting  
433 potential of monosodium glutamate (MSG) on secretion of the glucagon-like peptide-1 (GLP-1)  
434 gut hormone and GLP-1 receptor interaction. *Toxicol Lett* 2017;265:97-105.
- 435 42. Richards P, Pais R, Habib AM, Brighton CA, Yeo GS, Reimann F, et al. High fat diet impairs the  
436 function of glucagon-like peptide-1 producing L-cells. *Peptides* 2015;77:21-7.
- 437 43. Krieger JP, Langhans W, Lee SJ. Novel role of GLP-1 receptor signaling in energy expenditure  
438 during chronic high fat diet feeding in rats. *Physiol Behav* 2018;192:194-199.
- 439 44. Duca FA, Katebzadeh S, Covasa M. Impaired GLP-1 signaling contributes to reduced sensitivity  
440 to duodenal nutrients in obesity-prone rats during high-fat feeding. *Obesity* 2015;23:2260-2268.
- 441 45. Duca FA, Swartz TD, Sakar Y, Covasa M. Decreased intestinal nutrient response in diet-induced  
442 obese rats: role of gut peptides and nutrient receptors. *Int J Obes (Lond)*. 2013;37:375-381.

443 46 Singh AK. Glucagon-like peptide 1 and dysglycemia: Conflict in incretin science. Indian J  
444 Endocrinol Metab 2015;19:182-187.

445 47. Wang X, Liu H, Chen J, Li Y, Qu S. Multiple factors related to the secretion of glucagon-like  
446 peptide-1. Int J Endocrinol 2015;2015:651757.

447 48. Hansen CF, Bueter M, Theis N, Lutz T, Paulsen S, Dalbøge LS, Vrang N, Jelsing J. Hypertrophy  
448 dependent doubling of L-cells in Roux-en-Y gastric bypass operated rats. PLoS One  
449 2013;8(6):e65696.

450

451

452

453

454

455

456

457

458

459

460

461

462 **Table 1.** Test diet composition

Ingredient	Control	HFS	HiFat	HiSuc
g/kg of diet				
Cornstarch	397.486	–	167.486	97.486
Casein <sup>1</sup>	200	200	200	200
Dextrinized cornstarch <sup>2</sup>	132	–	132	132
Sucrose	100	399.486	100	400
Soybean oil	70	70	70	70
Lard oil	–	230	230	–
Fibre (cellulose) <sup>3</sup>	50	50	50	50
Mineral mixture <sup>4</sup>	35	35	35	35
Vitamin mixture <sup>4</sup>	10	10	10	10
L-Cystine	3	3	3	3
Choline bitartrate	2.5	2.5	2.5	2.5
<i>tert</i> -Butylhydroquinone	0.014	0.014	0.014	0.014
Protein (energy %)	20.5	15.9	15.9	20.5
Carbohydrate	63.6	31.3	31.3	63.6
Fat (energy %)	15.9	52.8	52.8	15.9
Energy density (kcal/g)	3.96	5.11	5.11	3.96

463 <sup>1</sup> Acid Casein (Fonterra, Ltd., Auckland, New Zealand);

464 <sup>2</sup> TK-16 (Matsutani Chemical Industry Co., Ltd., Hyogo, Japan);

465 <sup>3</sup> Avicel PH102 (Asahi Kasei Chemicals Corporation, Tokyo, Japan);

466 <sup>4</sup> Mineral and vitamin mixtures were prepared according to the AIN-93G formulation.

467

468

469

470

471

472 **Table 2.** Initial body weight, body weight gain, visceral adipose tissue (mesenteric, epididymal, and  
 473 retroperitoneal) weight, and energy intake after feeding with the test diet for 5 weeks

	Control	HFS	HiFat	HiSuc
Initial weight (g)	168.7 ± 3.1 <sup>NS</sup>	169.3 ± 2.5	169.1 ± 4.3	167.7 ± 2.8
Body weight gain (g)	200.7 ± 5.4 <sup>b</sup>	228.0 ± 7.8 <sup>a</sup>	232.3 ± 7.8 <sup>a</sup>	220.3 ± 7.9 <sup>ab</sup>
Visceral fat (g)	25.0 ± 1.6 <sup>NS</sup>	29.7 ± 1.5	29.2 ± 2.0	23.6 ± 1.4
Mesenteric fat (g)	7.2 ± 0.5 <sup>NS</sup>	7.8 ± 0.4	7.8 ± 0.5	6.7 ± 0.4
Epididymal fat (g)	7.5 ± 0.6 <sup>bc</sup>	9.7 ± 0.5 <sup>a</sup>	9.3 ± 0.8 <sup>ab</sup>	6.9 ± 0.5 <sup>c</sup>
Retroperitoneal fat (g)	10.4 ± 0.6 <sup>NS</sup>	12.1 ± 0.7	12.1 ± 0.8	10.0 ± 0.6
Total energy intake (kcal)	2736.6 ± 61.7 <sup>b</sup>	2958.6 ± 60.3 <sup>a</sup>	2966.3 ± 56.8 <sup>a</sup>	2809.9 ± 40.0 <sup>ab</sup>

474 Visceral fat weight is the sum of the mesenteric, epididymal, and retroperitoneal fat weight. The values  
 475 were expressed as the mean ± SEM for n = 8-10 rats (control = 10, HFS = 10, HiFat = 8, HiSuc = 8).  
 476 The superscripts without the same letters differed significantly between treatments ( $P < 0.05$ , Tukey-  
 477 Kramer's test). NS indicates that there was no significant difference among the treatments.

478

479

480

481

482

483



484 **Figure legends**

485 **Fig. 1. Basal glucose, insulin, GLP-1 levels, and HOMA-IR after feeding with the test diet for 2**  
486 **and 4 weeks.**

487 Rats were given the control, HFS, HiFat, or HiSuc diet (*ad libitum*) for 2 and 4 weeks. On the  
488 day of the experiment, blood samples were collected from the tail vein after overnight fasting (16 h),  
489 before conducting the MTT. The values were expressed as the mean  $\pm$  SEM for n = 4-9 (A-D; control =  
490 9, HFS = 8, HiFat = 4, HiSuc = 7, E-H; control = 9, HFS = 6, HiFat = 5, HiSuc = 6) rats. NS indicates  
491 that there was no significant difference among the treatments (Tukey-Kramer's test,  $P < 0.05$ ).

492

493 **Fig. 2. Postprandial glucose, insulin, GLP-1 level, and HOMA-IR during the meal tolerance test**  
494 **(MTT), after feeding with the test diet for 2 weeks.**

495 Rats were given the control, HFS, HiFat, or HiSuc diet (*ad libitum*) for 2 weeks; then, the MTT  
496 was performed. After overnight fasting, basal blood (0 min) was taken, followed by feeding of the  
497 control diet (10 g/kg body weight) for 30 min, and blood samples were collected until 120 min. The  
498 values were expressed as the mean  $\pm$  SEM for n = 4-9 rats (control = 9, HFS = 8, HiFat = 4, HiSuc = 7).  
499 Two-way repeated measure ANOVA  $P$  values for time (TI), for treatment (TR), and for the interactions  
500 of time and treatment (TI x TR) are shown in each panel (A, B, C). The superscripts without the same  
501 letters differed significantly between treatments ( $P < 0.05$ , Dunn's multiple range test). Asterisks (\*)  
502 indicate significant differences from the basal value (0 min) in each group ( $P < 0.05$ , Dunn's with  
503 control). NS indicates that there was no significant difference among the treatments.

504

505 **Fig. 3. Postprandial glucose, insulin, GLP-1 level, and HOMA-IR during the meal tolerance test**  
506 **(MTT), after feeding with the test diet for 4 weeks.**

507 Rats were given the control, HFS, HiFat, or HiSuc diet (*ad libitum*) for 4 weeks; then, MTT was  
508 performed. After overnight fasting, basal blood (0 min) was taken, followed by feeding of the control  
509 diet (10 g/kg body weight) for 30 min, and blood samples were collected until 120 min. The values  
510 were expressed as the mean  $\pm$  SEM for n = 5-9 (control = 9, HFS = 6, HiFat = 5, HiSuc = 6) rats. Two-  
511 way repeated measure ANOVA *P* values for time (TI), for treatment (TR), and for the interactions of  
512 time and treatment (TI x TR) are shown in each panel (A, B, C). The superscripts without the same  
513 letters differed significantly between treatments ( $P < 0.05$ , Dunn's multiple range test). Asterisks (\*)  
514 indicate significant differences from the basal value (0 min) in each group ( $P < 0.05$ , Dunn's with  
515 control). NS indicates that there was no significant difference among the treatments.

516

517 **Fig. 4. Portal glucose, insulin, GLP-1, cholesterol, and triglyceride (TG) levels after feeding with**  
518 **the test diet for 5 weeks.**


519 Rats were given the control, HFS, HiFat, or HiSuc diet (*ad libitum*) for 5 weeks. After overnight  
520 fasting, blood samples were collected from the portal vein under sodium pentobarbital anaesthesia (50  
521 mg/kg of body weight). The values were expressed as the mean  $\pm$  SEM for n = 8-10 (control = 10, HFS  
522 = 10, HiFat = 8, HiSuc = 8) rats. The superscripts without the same letters differed significantly  
523 between treatments ( $P < 0.05$ , Tukey-Kramer's test). NS indicates that there was no significant  
524 difference among the treatments.

525

526 **Fig. 5. GLP-1 content in intestinal tissues after feeding with the test diet for 5 weeks.**

527           After feeding with the test diets for 5 weeks, each intestinal segment: (A) jejunum, (B) ileum,  
528 (C) cecum, and (D) colon was collected. The values were expressed as the mean  $\pm$  SEM for n = 8-10  
529 (control = 10, HFS = 10, HiFat = 8, HiSuc = 8) rats. The superscripts without the same letters differed  
530 significantly between treatments ( $P < 0.05$ , Tukey–Kramer’s test). NS indicates that there was no  
531 significant difference among the treatments.

2 weeks after feeding with the test diet



4 weeks after feeding with the test diet




Fig. 1.






Fig. 2.



**Fig. 3.**



**Fig. 4.**




Fig. 5.




**Continuous feeding obesogenic diets  
(HFS, HiFat, or HiSuc diet, vs control diet)**

↓ after 2 and 4 weeks


**MTTs** to assess postprandial GLP-1 response

Voluntary ingestion  
Control diet (30 min)



Rats fed **HFS** diet had enhanced GLP-1 response to luminal nutrients after 2 weeks

Combination of **fat and sucrose** rapidly enhanced GLP-1 secretory response during obesity development



**MTT**; Meal tolerance test      **HFS**; High fat/high sucrose diet  
**HiFat**; High fat diet          **HiSuc**; High sucrose diet